

Formation of Zn–Ca phyllomanganate nanoparticles in grass roots

B. Lanson, Matthew A. Marcus, S. Fakra, Frédéric Panfili, Nicolas Geoffroy,

Alain Manceau

► To cite this version:

B. Lanson, Matthew A. Marcus, S. Fakra, Frédéric Panfili, Nicolas Geoffroy, et al.. Formation of Zn–Ca phyllomanganate nanoparticles in grass roots. Geochimica et Cosmochimica Acta, 2008, 72 (10), pp.2478 à 2490. 10.1016/j.gca.2008.02.022 . insu-00334521

HAL Id: insu-00334521 https://insu.hal.science/insu-00334521

Submitted on 13 Jan 2009 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1	Formation of Zn-Ca Phyllomanganate Nanoparticles in Grass Roots
2	
3	Bruno Lanson ^a , Matthew A. Marcus ^b , Sirine Fakra ^b , Frédéric Panfili ^a ,
4	Nicolas Geoffroy ^a , Alain Manceau ^a
5	
6	^a Mineralogy & Environments Group, Maison des GéoSciences, Université Joseph Fourier –
7	CNRS, F-38041 Grenoble Cedex 9, France.
8	^b Advanced Light Source, Lawrence Berkeley National Laboratory, Berkeley, California
9	94720, USA
10	
11	* Corresponding author: <u>Bruno.Lanson@obs.ujf-grenoble.fr</u>
12	
13	Running title: Zn-Ca Phyllomanganate Nanoparticles in Grass Roots

Abstract. It is now well established that a number of terrestrial and aquatic 14 15 microorganisms have the capacity to oxidize and precipitate Mn as phyllomanganate. 16 However, this biomineralization has never been shown to occur in plant tissues, nor has the structure of a natural Mn(IV) biooxide been characterized in detail. We show that the 17 18 graminaceous plant Festuca rubra (red fescue) produces a Zn-rich phyllomanganate with 19 constant Zn:Mn and Ca:Mn ratios (0.46 and 0.38, respectively) when grown on a 20 contaminated sediment. This new phase is so far the Zn-richest manganate known to form in 21 nature (chalcophanite has a Zn:Mn ratio of 0.33) and has no synthetic equivalent. Visual 22 examination of root fragments under a microscope shows black precipitates about ten to 23 several tens of microns in size, and their imaging with backscattered and secondary electrons 24 demonstrates that they are located in the root epidermis. In situ measurements by Mn and Zn 25 K-edge extended X-ray absorption fine structure (EXAFS) spectroscopy and X-ray diffraction 26 (XRD) with a micro-focused beam can be quantitatively described by a single-phase model 27 consisting of Mn(IV) octahedral layers with 22% vacant sites capped with tetrahedral and 28 octahedral Zn in proportions of 3:1. The layer charge deficit is also partly balanced by 29 interlayer Mn and Ca. Individual crystallites have a domain radius of 33 Å in the *ab* plane and 30 contain only 1.2 layers (~8.6 Å) on average. Since this biogenic Mn oxide consists mostly of 31 isolated layers, basal 001 reflections are essentially absent despite its lamellar structure. 32 Individual Mn layers are probably held together in the Mn-Zn precipitates by stabilizing 33 organic molecules. Zinc biomineralization by plants likely is a defense mechanism against toxicity induced by excess concentrations of this metal in the rhizosphere. 34

1. INTRODUCTION

36

37 Graminaceous plants, like other so-called metal-tolerant plants, mostly sequester 38 metals in roots to protect reproductive and photosynthetic tissues (Li et al., 2000; Simon, 39 2005). The ability to store metals in underground tissues is used in phytoremediation to 40 reinstall a vegetation cover on heavily contaminated areas and limit the propagation of metals 41 into the food chain (Smith and Bradshaw, 1992; Cunningham et al., 1995; Vangronsveld et 42 al., 1995; Ma et al., 2003; Mench et al., 2003; Krämer, 2005). Panfili et al. (2005) showed that 43 the grass species Festuca rubra (red fescue) and Agrostis tenuis (colonial bentgrass) 44 accelerate the weathering of zinc sulfide when grown on contaminated dredged sediment, thus 45 increasing Zn bioavailability in the rhizosphere. After two years of plant growth, micrometer-46 sized Mn-Zn black precipitates were observed at the surface of *Festuca rubra* roots, but not 47 characterized (Panfili, 2004). Zinc precipitation may be a bioactive tolerance mechanism in 48 response to metal toxicity, or a passive mineralization at the soil-root interface (Cotter-49 Howells et al., 1999). Clarifying this issue and determining the mineralogy and structure of 50 this natural precipitate is important to enhance the effectiveness of using graminaceous plants 51 in phytoremediation. These questions are addressed here with electron microscopy and 52 synchrotron-based microanalytical tools, including X-ray fluorescence (µ-XRF), extended X-53 ray absorption fine structure (µ-EXAFS) spectroscopy and X-ray diffraction (µ-XRD) 54 (Manceau et al., 2002b). Micro-XRD was employed to determine the nanocrystalline structure 55 of the Mn-Zn precipitates and the nature of defects (layer stacking faults, cationic vacancies 56 and occupancies, Mn, Zn, and Ca site configuration, stoichiometry) through modeling of their 57 scattering properties (Villalobos et al., 2006; Drits et al., 2007). We show that the root 58 precipitates are present in the root epidermis (the outermost layer of root cells) and consist of 59 a poorly crystallized phyllomanganate with a constant Zn:Mn ratio higher than reported so far

for any natural and synthetic manganate. A structure model is proposed for this newbiomineral.

- 62
- 63

2. MATERIALS AND METHODS

64

65 2.1 Materials

The composition in major elements of the dredged sediment was 68.3 % SiO₂, 6.9 % 66 CaO, 4.8 % Al₂O₃, 2.4 % Fe₂O₃, 0.7 % P₂O₅, and 7.2% organic carbon, and the composition 67 in a trace metals was 4700 mg.kg⁻¹ Zn, 700 mg.kg⁻¹ Pb, and \sim 270 mg.kg⁻¹ Mn. Seeds of F. 68 rubra were sown in plastic pots filled with 40 kg of either the untreated sediment, the 69 70 sediment amended with 3 wt. % hydroxylapatite, or the sediment amended with 5 wt. % 71 Thomas basic slag. The pots were placed in a greenhouse without artificial lighting and daily 72 irrigated with tap water in an amount similar to the mean rainfall in northern France. After 73 two years of culture, the pots were dismantled to collect samples. The texture and color of the 74 sediment in areas colonized by the roots (upper 30 cm of the pots) were similar to a brown 75 silty soil, whereas the initial sediment was black and compact. Roots of F. rubra from the 76 treated and untreated pots were washed meticulously with distilled water to remove soil 77 particles from the surface and then freeze-dried.

The speciation of zinc in the initial sediment and in the rhizosphere of *F. rubra* after the two years of vegetation was described previously (Panfili et al., 2005). Briefly, in the untreated and unvegetated sediment, Zn was distributed as ~50% (mol ratio of total Zn) sphalerite, ~40% Zn-ferrihydrite, and ~10 to 20% (Zn-Al)-hydrotalcite plus Zn-phyllosilicate. In the presence of plants, ZnS was almost completely dissolved, and the released Zn bound to phosphate (~40–60%) and to Zn phyllosilicate plus (Zn,Al)-hydrotalcite (~20–40%). The coaddition of mineral amendment did not affect the Zn speciation in the vegetated sediment.

86 **2.2 Methods**

87 2.2.1 Electron microscopy

Regions of the roots rich in black precipitates were carbon-coated and examined in secondary and backscattered electron modes by high-resolution scanning electron microscopy (JEOL JSM-6320F with a field emission gun) and analyzed with energy dispersive X-ray spectroscopy (EDS, Tracor analyzer).

92

93 2.2.2 X-ray fluorescence, diffraction, and absorption spectroscopy

94 The Zn:Mn and Ca:Mn ratios were measured with an Eagle III µ-XRF spectrometer 95 (Röntgenanalytik Messtechnik GmbH) equipped with a Rh anode and a 40 µm poly-capillary. 96 The spectrometer was operated under vacuum at 20 kV and 400 µA, and fluorescence was 97 measured for 300 s per point. Micro XRF, XRD and EXAFS data were collected on beamline 10.3.2 at the Advanced Light Source (ALS, Berkeley - Marcus et al., 2004a). Short root 98 99 fragments were attached to the tips of glass capillaries and cooled down to 110-150 K 100 (Oxford CryoSystems Cryo-Stream) to minimize radiation damage (Manceau et al., 2002b). 101 X-ray fluorescence maps were taken at 10 keV incident energy, with a beam size ranging 102 from 5×5 µm to 16×7 µm (H×V). Fluorescence counts were collected for K, Ca, Mn, Fe and 103 Zn with a seven-element Ge solid-state detector and a counting time of 100 ms per pixel. For 104 μ-EXAFS measurements, the vertical beam size ranged from 5 to 7 μm. A maximum of two 105 spectra per precipitate were taken at either the Mn or the Zn K-edge to prevent the reduction 106 of tetravalent to divalent Mn and the increase of structural disorder under the beam (Manceau 107 et al., 2002b). Diffraction data were collected with a CCD camera (Bruker SMART6000, SMART software) at 17 keV ($\lambda = 0.729$ Å) and exposure times of 120-240 seconds. At this 108 109 energy, the incident flux and absorption cross-sections are low enough to make radiation 110 damage during an exposure negligible even at room temperature. A background pattern was

recorded next to each precipitate to subtract the scattering contribution from the root so as to obtain the precipitate pattern. Diffraction patterns collected on different precipitates were all statistically identical, and thus summed up to optimize data quality. Calibration of the energy and camera distance were obtained using an Al₂O₃ standard and Fit2D software (Hammersley, 1998). This software was also used to calculate the one-dimensional XRD traces from the radial integration of the two-dimensional patterns.

117

118 2.2.3 Data processing

119 The EXAFS data were analyzed according to standard procedure (Teo, 1986; Marcus 120 et al., 2004b). The µ-XRD patterns were simulated following the trial-and-error approach 121 developed by Drits and Tchoubar (1990), and applied previously to natural and synthetic 122 phyllomanganates (Chukhrov et al., 1985; Manceau et al., 1997; Drits et al., 1998; Lanson et 123 al., 2000, 2002a, b; Gaillot et al., 2003, 2005, 2007; Villalobos et al., 2006). Details on the 124 program and fitting procedure can be found in the articles by Drits et al. (1998) and Plancon (2002). The scattering background was considered to be linear in the 0.35-0.80 Å⁻¹ 1/d125 126 interval (2.86-1.25 Å). The fit quality was evaluated over this interval using the conventional 127 R_{WP} and R_{Exp} values (Howard and Preston, 1989).

128

129

3. RESULTS AND INTERPRETATION

130

131 **3.1 Optical and electron microscopy**

Under the optical microscope, the Mn-Zn precipitates appear as black stains about ten to several tens of micrometers in size on the root surface (Fig. 1a, EA-1). They are also observed in backscattered electron microscopy (Fig. 1b) due to the presence of high-Z elements (Mn, Zn, Ca, and minor Pb), but always are hardly noticeable in secondary electron imaging mode (Fig. 1c). This suggests that the precipitates are engulfed in the root epidermis
(Cotter-Howells et al., 1999) and do not coat the root surface as iron and manganese plaques
do (Otte et al., 1989; St-Cyr and Campbell, 1996). No differences were observed among
precipitates from plants grown in the untreated and mineral amended sediments. This result,
together with the compartmentation of the precipitates inside the roots, suggests a biological
origin. This interpretation is supported also by the absence of Zn-rich phyllomanganates in the
surrounding soil matrix (Panfili et al., 2005).

143

144 **3.2 Micro-XRF**

145 Elemental mapping of F. rubra roots shows that Zn is associated with Mn in 146 localized spots, and uniformly distributed without manganese in the vascular cylinder as 147 expected for this nutritive element (Fig. 2 - Rout and Das, 2003). All roots have Zn in their 148 central stele, but not all are speckled with Mn-Zn precipitates. Some root fragments are partly 149 covered by Zn-free Fe-rich plaques (Fig. EA-2). These plaques are made of ferric 150 oxyhydroxides, as indicated by their optical rusty color (Fig. 1a top right). In Zn-Mn-Ca 151 tricolor representation all Mn-Zn precipitates generally have the same color (Fig. 2), even 152 among different roots (Fig. EA-2), meaning that the relative proportions of Zn, Mn and Ca are 153 about the same. The correlation coefficient between Zn and Mn counts for the precipitates is 154 0.8, with P-value < 0.0001 for the Anova F-test (Fig. 2). The Zn:Mn atomic ratio was 155 calculated from the relative absorption jumps measured at the Mn and Zn K-edges on four 156 particles. For each particle, a pre-absorption edge background was removed first, and then a 157 linear fit to the post-edge region was extrapolated back to the edge to measure edge jumps. 158 The ratio of the Zn to Mn edge jumps is 0.310(7), which translates into a Zn:Mn ratio of 159 0.46(1) when taking into account the atomic absorptions of the two elements. A consistent 160 0.44 value was obtained independently with the Eagle III spectrometer. This analysis also

161 confirmed that root precipitates have a constant Ca:Mn ratio. An atomic ratio of 0.41 was162 calculated after correction of the Ca-fluorescence from the root.

163

164 **3.3 Micro-EXAFS spectroscopy**

165 The Mn and Zn μ -EXAFS spectra for all precipitates were indistinguishable from 166 each other, and thus averaged to improve the signal-to-noise ratio. If the precipitates consisted 167 of an assemblage of distinct Zn and Mn species, the proportions of these species would most 168 likely vary among the analyzed grains, and this variability would be detected by μ -EXAFS 169 (Panfili et al., 2005). However, this was not the case, suggesting that all root precipitates 170 consist of a single species, hereafter referred to as "Mn-Zn precipitate", in agreement with μ -171 XRF data.

172

173 3.3.1 Mn-EXAFS in Mn-Zn precipitate

Fig. 3 compares the EXAFS spectrum of Mn-Zn precipitate with those of reference 174 175 compounds. The references used are hollandite (Hol) and todorokite (Todo), two 176 tectomanganates with 2×2 (Hol) and 3×3 (Todo) tunnel structures, TcBi, a triclinic birnessite with 31% Mn^{3+} in the layers, $[Na^+_{0.31}(H_2O)_{0.40}(Mn^{4+}_{0.69}Mn^{3+}_{0.31})O_2 - (Silvester et al., 1997;$ 177 178 Lanson et al., 2002a)], Lit, a synthetic lithiophorite $[(Al_{0.67}Li_{0.32})(Mn^{4+}_{0.68}Mn^{3+}_{0.32})O_2(OH)_2 -$ 179 (Manceau et al., 2005)], HBi, a hexagonal birnessite prepared by equilibrating TcBi at pH 4 $[H_{0,33}^{+}Mn_{0,043}^{2+}Mn_{0,123}^{3+}(OH)_{0,013}^{-}(Mn_{0,122}^{4+}Mn_{0,111}^{3+}Vac_{0,167})O_2 - (Silvester et al., 1997;$ 180 181 Lanson et al., 2000)], chalcophanite (Chalco), a Zn-rich phyllomanganate with one in seven 182 octahedral sites vacant and capped on each side of the surface layer by interlayer octahedral Zn atoms [ZnMn₃O₇•3H₂O - (Wadsley, 1955)], and dBi, a turbostratic birnessite with no 183 interlayer Mn and no layer $Mn^{3+} [Na^+_{0.24}(H_2O)_{0.72}(Mn^{4+}_{0.94}Vac_{0.06})O_2 - (Villalobos et al., Na^+_{0.24}(H_2O)_{0.72}(Mn^{4+}_{0.94}Vac_{0.06})O_2 - (Villalobos et al., Na^+_{0.94}Vac_{0.06})O_2 - (Villalobos et al., Na^+_{0.94}Vac_{$ 184

185 2006)]. TcBi and Lit have a similar content of layer Mn³⁺, but different Mn⁴⁺-Mn³⁺ cation
186 ordering (Drits et al., 1997; Manceau et al., 2005).

187 Mn-Zn precipitate is not a tectomanganate because i) the second EXAFS oscillation of tectomanganates is split or has a shoulder at 6.5 Å^{-1} , depending on the tunnel size 188 189 (Manceau and Combes, 1988; Mckeown and Post, 2001; Manceau et al., 2007b), and ii) their [7.3-9.5 Å⁻¹] indicator region (Marcus et al., 2004b) does not match the data (Figs. 3a-b). In 190 the indicator region, Mn-Zn precipitate has a single maximum at 8.0-8.1 Å⁻¹ like dBi, HBi, 191 and Chalco. The shape and position of this maximum is diagnostic of Mn⁴⁺-rich manganate 192 193 layers with hexagonal symmetry (Gaillot et al., 2007). For example, this maximum is shifted to 7.9 Å⁻¹ (i.e., consistent with longer distances) in Lit (Manceau et al., 2004) due to the large 194 amount of Mn³⁺ in the hexagonal layer. TcBi has a distinctive double peak with a minimum at 195 7.9-8.0 Å⁻¹ as a result of the split of the Mn-Mn distances induced by the linear ordering of 196 Mn³⁺ and Mn⁴⁺ cations in the layer (Drits et al., 1997; Lanson et al., 2002a; Gaillot et al., 197 198 2003, 2007; Manceau et al., 2005). This cation ordering lowers the layer symmetry from 199 hexagonal to orthogonal. Thus, the Mn-Zn precipitate is a phyllomanganate having hexagonal layer symmetry and little layer Mn³⁺. The subtle differences between Mn-Zn precipitate and 200 201 some phyllomanganate references are more obvious when the data are Fourier transformed 202 (FT, Fig. 4). The phase of the Mn1_E peak matches those of dBi, Chalco, and HBi, and is 203 shifted to a shorter distance relative to Lit. In phyllomanganates, the Mn1_E peak represents 204 Mn atoms in the first-neighbor layer octahedra (Manceau and Combes, 1988). Therefore, its 205 phase depends on the nearest Mn-Mn distance (Mn-Mn1 pair), and is sensitive to the amount of Mn³⁺ in the layer similarly to the phase of the O peak. However, the sensitivity of the phase 206 of this peak to the presence of Mn^{3+} is not high enough to distinguish HBi (11% layer Mn^{3+} 207 208 per octahedral site) from dBi and Chalco (0% – Fig. EA-3). The $Mn3_E$ peak is twice as 209 sensitive as the Mn1_E peak because it arises from the third-neighbor octahedra at twice the

210 Mn-Mn1 distance (see for example Fig. 11 in Manceau et al., 2005). The sensitivity is now 211 high enough to differentiate HBi from dBi and Chalco, as shown in Fig. 4. Thus, using the 212 phase of the Mn1_E and Mn3_E peaks as chemical probes to the layer composition, we conclude that Mn-Zn precipitate has less Mn³⁺ in the octahedral layers than HBi (11%), if any. 213 214 Confirming evidence is found in the comparison of the frequency of EXAFS spectra. Fig. 3 215 shows that the overall EXAFS frequency, and hence interatomic distances, increase from 216 dBi/Chalco/Mn-Zn precipitate, to HBi, to Lit/TcBi, as a result of the increasing amount of layer Mn³⁺ (Gaillot et al., 2007). 217

In contrast to the Mn1 peak, the phase of the O peak (i.e., average Mn-O distance in the layer and interlayer) matches those of the two Mn³⁺-free references (dBi, Chalco) and is slightly shifted to lower distance relative to both HBi and Lit. This shift is barely visible in Fig. 4, but clearly apparent when the [1-3 Å] $R+\Delta R$ interval is expanded (Fig. EA-3). Thus, Mn-Zn precipitate has no detectable Mn³⁺ in the layer similarly to dBi and Chalco, nor in the interlayer, in contrast to HBi.

224 The FT of Mn-Zn precipitate also differs from that of HBi by the absence of the Mn1_{TC} peak from Mn_{laver}-Me pairs at ~3.5 Å ($R + \Delta R \sim 3.1$ Å), where Me is an interlayer 225 226 metal cation, such as Mn, Zn, or Pb, in triple-corner sharing position above or below vacant 227 layer sites (TC linkage – Manceau et al., 2002a). When there are interlayer Mn atoms, this peak is intense because each interlayer Mn is surrounded by as many as six Mn_{layer} neighbors. 228 229 In contrast, Mn_{laver} atoms near an octahedral vacancy have fewer interlayer Mn neighbors, 230 their exact number depending on the density of layer vacancies and composition of the 231 interlayer. Here, the lack of Mn1_{Tc} peak in Mn-Zn precipitate suggests that it has no 232 detectable Me cations on either side of the layer vacancies. The $Mn5_E$ peak is also absent in the FT of Mn-Zn precipitate, as in dBi, but not HBi nor Chalco. This peak arises from the 5th 233 Mn shell at about $2.90 \times 3 = 8.7$ Å, and is enhanced by forward scattering from two 234

intervening Mn_{layer} (Manceau et al., 2005). The absence of this peak in Mn-Zn precipitate indicates that the octahedral layers have a small lateral dimension. This peak is also absent in Lit, but in this case because the Mn-Mn5 shell is split as a result of the $Mn^{4+}-Mn^{3+}$ ordering in the layer (Manceau et al., 2005).

- 239
- 240

3.3.2 Zn-EXAFS in Mn-Zn precipitate

241 The best spectral match of Mn-Zn precipitate to our Zn species database was obtained with tetrahedrally coordinated Zn (^{IV}Zn) sorbed above/below octahedral vacancies of 242 a phyllomanganate (^{IV}TC site – Fig. 5a). This Zn complex is common in nature and its local 243 244 structure has been described previously (Manceau et al., 2000b, 2002a; Marcus et al., 2004b; Isaure et al., 2005; Toner et al., 2006). Although the ^{IV}Zn-sorbed phyllomanganate reference 245 246 provides a good approximation of Zn local structure in Mn-Zn precipitate, there are significant differences between the two spectra, particularly in the phase mismatches between 247 6.5 and 10 Å⁻¹, and the symmetry of the first and second oscillations. These differences have 248 249 been documented previously and result from the mixed occupancy of the TC site by 250 tetrahedral and octahedral Zn (Manceau et al., 2002a, 2007b). Adding the chalcophanite or VI/IVZn-sorbed dBi reference (Fig. 5b - Toner et al., 2005a, 2006) in a two-component linear 251 252 fit to the data compensated for the phase shift and asymmetry of the first oscillation, but not 253 entirely for that of the second (Fig. 5c). All attempts to suppress this residual by adding a 254 third component to the linear fit failed. Therefore, although there is some uncertainty on the exact configuration of the minor ^{VI}Zn site, the ^{IV}TC + ^{VI}TC model is the best description of 255 the data we can offer. Consideration of an edge-sharing complex (^{VI}TE) at layer edges did not 256 reproduce the phase as well as the ^{VI}TC surface species. The ^{VI}TC species may occur also at 257 258 the edge of the Mn-Zn precipitate layer, instead of the basal surface, in which case it has a 259 slightly different binding environment from that of dBi. Thus, the dBi and chalcophanite

standards may not be the best proxies for the ^{VI}TC species. The fraction of each Zn species derived from the two-component least-squares fit is $87 \pm 10\%$ ^{IV}TC and $27 \pm 10\%$ ^{VI}TC.

262

263 3.3.3 Zn-EXAFS in Zn-only precipitate

A few spots in the μ -XRF maps contained Zn and little else detectable. EXAFS spectroscopy identified zincite (ZnO) and sphalerite (ZnS), two mineral species originally present in the untreated sediment (Isaure et al., 2002). These grains are likely residual slag material stuck to the root surface that were not removed by washing. None of the Zn species formed in the rhizosphere of *F. rubra* after the two years of experiment were detected in or at the root surface.

270

271 *3.3.4 Zn-EXAFS in the root vascular cylinder*

272 Micro-EXAFS spectra were recorded in the vascular cylinder of four distinct roots, at 273 spots containing little Mn. All spectra were indistinguishable, indicating that Zn speciation is 274 uniform, and thus averaged. The resulting Root spectrum has the same frequency as the Mn-275 Zn precipitate spectrum, which suggests that Zn is also mostly tetrahedral in the roots (Fig. 276 5d). However, in contrast to Mn-Zn precipitate, the second and third oscillations of the Root 277 spectrum are not split, indicative of "light" backscatters from second-shell contributions. 278 Consistently, the best spectral match to our organic and inorganic database of the Root 279 spectrum was provided with Zn in a biofilm (Zncell7, Fig. 5e). This reference has $80 \pm 10\%$ 280 Zn complexed to phosphoryl groups and $20 \pm 10\%$ to carboxyl groups (Toner et al., 2005b). 281 Consistent with this other study, consideration of carboxyl (citrate, Fig. 5f) and phosphate 282 (phytate, Fig. 5g; Zn-sorbed *Penicillium chrysogenum*, Fig. 5h – Sarret et al., 1998a) ligands 283 alone, did not yield an optimal fit to the data. Zinc preferential binding to phosphate groups 284 has been reported also in the roots of Arabidopsis halleri and A. lyrata grown hydroponically,

on bacterial and fungi cells, and in biofilms (Sarret et al., 1998b, 2002; Fein et al., 2001).
These studies have shown that Zn has a higher affinity for phosphate than for carboxyl
groups, which is consistent with the predominance of the phosphate species in *F. rubra* roots.

200

3.4 Micro-XRD

290 3.4.1 Qualitative description of the data

291 According to µ-XRF and µ-EXAFS, Mn-Zn precipitate is a phyllomanganate with hexagonal layer symmetry, little to no layer Mn³⁺, less interlayer Mn³⁺ than HBi, and as much 292 293 as 0.46 interlayer Zn per total Mn, of which $87 \pm 10\%$ is tetrahedral and $27 \pm 10\%$ octahedral. 294 Despite its lamellar structure, the XRD pattern of Mn-Zn precipitate shows no patent basal 295 reflections (Fig. 6). The XRD trace is dominated by two reflections at ~2.45 and ~1.42 Å, the 296 first one being asymmetric towards higher 1/d values. This profile is characteristic of lamellar 297 compounds with turbostratic stacking, i.e., lacking well-defined displacement/rotation 298 between successive layers (Warren, 1941; Biscoe and Warren, 1942; Brindley, 1980). The 299 positions, profiles, and relative intensities of the two peaks match the [20,11] and [02,31] 300 scattering bands of turbostratic birnessite, choosing a C-centered cell for their crystallographic 301 assignment (Drits et al., 1997, 2007; Villalobos et al., 2006). Their d-spacings are in the ratio of 1.73, close to the $\sqrt{3}$ value for hexagonal symmetry. Their profiles are controlled by the 302 303 structure factor and can thus be used for structural determinations (Villalobos et al., 2006; 304 Drits et al., 2007), as shown below.

305

306 3.4.2 Simulation of the 0.35-0.80 Å⁻¹ 1/d interval

307 XRD simulations were performed with the C-centered unit-cell parameters b =308 2.850 Å, $a = b\sqrt{3} = 4.936$ Å, $\gamma = 90^{\circ}$ derived from the position of the [20,11] and [02,31] 309 bands, $d_{(001)} = 7.20$ Å, and a random layer stacking (Wr = 100%, Fig. EA-4). The atomic

coordinates for the ^{IV}TC and ^{VI}TC sites were considered to be the same as those in Zn-sorbed 310 311 birnessite (Table 1 – Lanson et al., 2002b). The Ca position was assumed to be close to that in 312 Ca-rich birnessite (Drits et al., 1998), that is in the mid-plane of the interlayer almost above a tetrahedral surface site (TE position - Fig. EA-5). Thus, parameters optimized in the 313 314 simulations are the amount of vacant layer sites, the amount, position, and coordination of interlayer Zn^{2+} and $Mn^{2+,3+}$, the xy coordinates of Ca^{2+} , the position of water molecules, and 315 316 the size of the coherent scattering domains (CSDs) in the ab plane. The CSDs were 317 considered to have a disk-like shape, whose average radius was constrained by fitting the 318 maximum at ~2.45 Å (Villalobos et al., 2006).

319 Since the phyllomanganate has 0.46 interlayer Zn per total Mn, at least 0.186 vacant 320 layer sites per octahedron are needed to accommodate all Zn $[(0.186 \times 2) / (1.00 - 0.186)]$ = 321 0.46]. In this case, every vacant site is capped by one Zn on either side. No satisfactory 322 agreement between theory and experiment could be obtained with this model, regardless of the amount and position of Ca^{2+} . In particular, the high 1/d tail of the [20,11] band at ~0.43-323 0.45 Å⁻¹ was poorly reproduced (Fig. 7a – R_{WP} = 3.57%; R_{Exp} = 1.57%). To reproduce this 324 325 feature it was necessary to increase the density of vacant sites to 0.22, and thus to introduce 326 interlayer Mn in TC and TE sites to keep the Zn:Mn ratio constant. Our best theoretical model to the data is shown in Fig. 7b ($R_{WP} = 3.49\%$; $R_{Exp} = 1.57\%$), and structural parameters are 327 328 listed in Tables The 1 and 2. structural formula is $[(Mn_{0.78}Vac_{0.22})O_2]^{TC}Mn_{0.010}^{TE}Mn_{0.046}^{VI}Zn_{0.088}^{IV}Zn_{0.300}Ca_{0.318}(H_2O)_{0.972}]$, and the 329 structure 330 model is schematized in Figs. 8 and EA-5. The refined model contains 0.38 Ca per Mn, in 331 agreement with μ -XRF data (0.41), at a position [(-0.410, 0, $\frac{1}{2}$) and symmetric positions] slightly shifted from the ideal TE position [(-0.333, 0, $\frac{1}{2}$) - Fig. EA-5]. Interlayer H₂O 332 333 molecules that are not bound to Zn and interlayer Mn are located in $(0.220, 0, \frac{1}{2})$ and 334 symmetric positions. This position is similar to ordered (Lanson et al., 2002a) and disordered

335 (Villalobos et al., 2006) Na-exchanged birnessite varieties. It allows for the formation of 336 strong H-bonds with O_{layer} atoms [d(O_{layer} -H₂ $O_{inter.}$) = 2.66 Å, Table 2]. Overall, only the Mn 337 TE position is new, all others have been described previously for other birnessite varieties.

338

339 3.4.3 Sensitivity of calculated XRD patterns to structural parameters

340 As the optimal fit to data was obtained using a trial-and-error approach, the 341 sensitivity of the XRD simulations to key structural parameters needs to be assessed. A key 342 parameter for birnessite's ability to sorb trace metals is the origin of the layer charge. In the 343 present model, the layer charge arises from layer vacancies only, not at all from substitution of Mn^{3+} for Mn^{4+} in the layer, because the hexagonal layer symmetry and small b unit-cell 344 dimension (2.85 Å) are incompatible with appreciable amount of Mn³⁺ in the layer (Gaillot et 345 al., 2005, 2007; Manceau et al., 2005). When birnessite contains ~25% layer Mn^{3+} per 346 octahedral site with the long Mn-O bonds oriented at random, b = 2.895 Å (Gaillot et al., 347 2007) and the [20,11] and [02,31] bands are shifted to lower 1/d values relative to data (R_{WP} = 348 7.15%; Fig. 7c). Mn-Zn precipitate has the same b value as HBi (2.850 Å) which contains 349 11% layer Mn³⁺ per octahedral site (Lanson et al., 2000). Based on EXAFS results, this 350 amount is regarded as an upper limit. Also, if the layer contained a high proportion of Mn³⁺ 351 cations ordered in rows, as in triclinic birnessite, a similar b value (2.84-2.85 Å) would be 352 353 measured but the [02,31] band would be a doublet due to the departure from hexagonal 354 symmetry (Drits et al., 2007; Gaillot et al., 2007). Calculations show that the band splitting 355 decreases with the layer dimension, but that it should still be observed for a CSD dimension of 33 Å, the optimal CSD value (Fig. $7d - R_{WP} = 6.79\%$). 356

The sensitivity of XRD profiles to the amount, coordination, and position of high-Z interlayer scatterers, and to the number of vacant layer sites is illustrated next. If interlayer Mn atoms are located only in ^{VI}TC sites (0.056 per octahedron) rather than distributed over

^{VI}TE and ^{VI}TC sites (0.046 and 0.010, respectively), the dip at ~0.52 Å⁻¹ is deeper and the 360 intensity of the broad hump at ~0.60 Å⁻¹ is increased (Fig. 7e – R_{WP} = 3.60%). When the 361 number of interlayer Ca^{2+} is decreased from 0.318 to 0.240, there is a deficit of intensity in 362 the 0.43-0.45 Å⁻¹ region (Fig. 7f – R_{WP} = 3.54%). Similarly, decreasing the number of 363 364 interlayer Zn from 0.388 to 0.244, and that of vacant layer sites from 0.22 to 0.15 broadens the high 1/d tail of the [20,11] band and shifts the hump at ~0.60 Å⁻¹ to lower 1/d values (Fig. 365 $7g - R_{WP} = 4.92\%$). Increasing the proportion of octahedrally coordinated interlayer Zn from 366 23% to 50% also broadens the high 1/d tail of the [20,11] band and increases the intensity of 367 the broad hump at ~0.60 Å⁻¹ (Fig. 7h – R_{WP} = 3.99%). Finally, the sensitivity of XRD profiles 368 to ^{IV}Zn, ^{VI}Zn, and Ca coordinates is shown in Figure EA-6. 369

370

371 3.4.4 Simulation of the 0.05-0.35 $Å^{-1}$ 1/d interval

372 Despite the lamellar structure of Mn-Zn precipitate, its XRD pattern lacks welldefined 00l reflections. To assess the possible origin of this oddity, the 001 and 002 373 374 reflections were calculated for crystallites with extremely small CSD dimension along the c^* 375 axis (Fig. 9). Calculations show that at least two layers are required to modulate the scattering 376 profile at low 1/d values, otherwise the X-ray intensity steadily decreases with increasing 1/d(Fig. 9). The best fit to data was obtained with a mixture of crystallites containing 1, 2, and 3 377 378 layers in the ratio of 20:3:1, respectively, leading to an average CSD dimension along the c^* 379 axis of 1.2 layers (~8.6 Å).

- 380
- 381

4. DISCUSSION

382

383 4.1 Structure model

385 4.1.1 Amount of vacant sites and Zn loading

386 Except for one sample obtained by metal sorption on poorly crystalline Mn oxides 387 (Nelson et al., 1999), the new Zn-rich phyllomanganate contains higher amounts of vacant 388 layer sites (0.22 per layer octahedron) and transition elements (0.44 total Mn and Zn per layer 389 octahedron) than any other natural and synthetic variety described so far (0.167 and 0.22, 390 respectively; Lanson et al., 2000, 2002b; Villalobos et al., 2006). The constant Zn:Mn ratio of 391 Mn-Zn precipitate suggests that Zn co-precipitated with Mn by a yet unknown mechanism to 392 form a chemically well-defined phase as natural solids formed by metal sorption on pre-393 existing mineral surfaces are chemically heterogeneous (see for example Manceau et al., 394 2007b).

395

396 4.1.2 Presence of heavy scatterers above and below vacant layers sites

According to XRD and Zn-EXAFS, Mn-Zn precipitate has 0.300 ^{IV}Zn and 0.088 397 ^{VI}Zn per octahedron in TC position (Table 1). The Zn-Mn pairs are detected at the Zn K-edge 398 399 (Fig. EA-7) but not at the Mn K-edge for two possible reasons. In a vacancy-free layer, each 400 Mn has six nearest-neighbor Mn atoms. In Mn-Zn precipitate, this number is 4.25 if two 401 vacancies cannot be adjacent which is always the case in layered structures (Manceau et al., 402 2000a). Thus, each Mn octahedron is surrounded statistically by 1.75 vacant sites. With 80% 403 of these vacant sites capped by Me cations on both sides, and 20% on one side only, Mn_{laver} atoms are coordinated on average to $1.75 \times (0.2 + (0.8 \times 2)) = 3.15$ Me in TC sites. This 404 number is about half the value for Zn, as each Zn is linked to six Mn_{layer}. Structural disorder 405 406 also can be invoked to explain the absence of Mn-Zn pairs in Mn-EXAFS. The TC position is occupied by at least three types of complex: 0.300 ^{IV}Zn, 0.088 ^{VI}Zn, and 0.010 ^{VI}Mn, and the 407 TE position by 0.046 Mn. Each Mn_{laver}-(TC,TE)Me pair has a different distance, and this broad 408 409 distribution results in an apparent loss of coordination.

411 **4.2 Intensity of basal reflections**

412 Birnessite and vernadite minerals were given different names because the basal reflections of birnessite at 7.2-7.0 Å (001) and 3.6-3.5 Å (002) were not observed originally in 413 414 the diffraction pattern of vernadite (Chukhrov and Gorshkov, 1980; Chukhrov et al., 1980). However, recent studies have shown that natural vernadite and its biogenic and chemical (δ-415 416 MnO₂) analogs most often display a 001 reflection when their XRD pattern is recorded on 417 modern diffractometers (Mandernack et al., 1995; Villalobos et al., 2003, 2005; Jurgensen et 418 al., 2004; Webb et al., 2005b; Bodeï et al., 2007; Manceau et al., 2007a, b), thus confirming 419 the view of Arrhenius et al. (1978) and Giovanoli (1980) that this mineral is a c-disordered 420 variety of birnessite. Villalobos et al. (2006) showed that basal reflections are present when 421 the diffracting crystallites have only 2-3 layers, on average. Here, this number is as low as 1.2 422 layers, meaning that Mn-Zn precipitate is essentially an assemblage of isolated layers. 423 Measurements of the Mn edge jumps on different Mn-Zn precipitates provide an inkling of 424 how the constitutive nanoparticles are joined at particle or so-called grain boundaries. The Mn 425 edge jump was typically between 0.2 and 0.3 for aggregates \sim 15-25 µm in diameter, which 426 indicates that the phyllomanganate represents only a small fraction (~20 volume %) of the 427 black precipitates, thus revealing a high micro-porosity. This porosity is possibly filled, at 428 least partly, by organics that may help disrupt the parallelism of the layers, but also to tie 429 them together.

430

431 **4.3 Biologically induced oxidation of manganese**

432 Many microorganisms have the capacity to oxidize and precipitate Mn as manganate 433 (Tebo et al., 2004, and references therein). Since biological oxidation of Mn is generally 434 faster than abiotic oxidation, most natural Mn oxides are considered to be biogenic.

Pseudomonas putida (Villalobos et al., 2003, 2006; Toner et al., 2005a), Leptothrix 435 436 discophora (Nelson et al., 1999, 2002; Nelson and Lion, 2003; Jurgensen et al., 2004; 437 Saratovsky et al., 2006), and Bacillus sp. strain SG-1 (Mandernack et al., 1995; Tebo et al., 438 1998; Webb et al., 2005a, b) are the three fairly-well characterized bacterial model systems 439 for the oxidation of manganese in the environment. Manganese oxidation and the subsequent 440 precipitation of Mn(III,IV) bioxides by microscopic fungi is also well documented (Krumbein 441 and Jens, 1981; Emerson et al., 1989; Schulze et al., 1995; Tani et al., 2003, 2004; Miyata et 442 al., 2004, 2006a, b; Thompson et al., 2005). Here, we showed that Mn can be biomineralized 443 also in higher living organisms, such as plants. Except for its atypical high Zn content and the 444 structural consequences thereof, this new manganese biomineral is no exception to the 445 intrinsic nanocrystalline nature of biogenic phyllomanganates.

446 Although the mechanism of Mn(II) to Mn(IV) oxidation is presently unknown, the 447 constant Zn:Mn ratio of the new Mn biooxide suggests the existence of a well-defined 448 bioactive process, likely in response to metal toxicity. The occurrence of Zn-Mn precipitate 449 only in the root epidermis and the absence in the roots of any Zn-rich species from the soil 450 matrix (Panfili et al., 2005) suggest that Mn oxidation did not occur in the rhizosphere, and 451 thus does not result from bacterial activity or abiotic reaction. Divalent manganese may have 452 been complexed and transported to the roots by phytosiderophores (Römheld, 1991), and then 453 oxidized by the plant itself or by endomycorrhizal fungi, as shown for wheat and soybean 454 (Schulze et al., 1995; Thompson et al., 2005, 2006).

Knowing how to stimulate the formation of this new phase in biological systems, or how to synthesize it abiotically, would be a significant progress towards Zn immobilization in contaminated environments and their (phyto-)remediation. Formation of this new phase could in particular facilitate the growth of plants in highly contaminated environments in lowering the concentration of bio-available Zn in the rhizosphere.

461

ACKNOWLEDGMENTS

462 The following persons are thanked for their contribution on various aspects of this 463 work: Fabienne Marseille and Bertrand Girondelot (CNRSSP, Douai) for the setup of pot 464 experiments, Clément Falk (LGIT, Grenoble) for optical microscopy on Mn-Zn precipitate, 465 Serge Nitsche, Damien Chaudanson, and Alain Baronnet (CRMCN, Marseilles) for SEM 466 analysis, Manuel Muñoz (LGCA, Grenoble) for XRF measurements on the Eagle 467 spectrometer, Didier Arnal, Nicole Paoletti, and Gérard Souche (R&S, Montpellier) for 468 growing Festuca rubra in hydroponic conditions, and Benoît Jaillard and Philippe Hinsinger 469 (R&S, Montpellier) for fruitful discussions about metal uptake by plants. This research was 470 supported by the ACI/FNS Ecco (EcoDyn) program. The ALS is supported by the Director, 471 Office of Energy Research, Office of Basic Energy Sciences, Materials Sciences Division of 472 the U.S. Department of Energy, under Contract No. DE-AC02-05CH11231.

473 REFERENCES 474 475 Arrhenius G., Cheung K., Crane S., Fisk M., Frazer J., Korkisch J., Melin T., Nakao S., Tsai 476 A., and Wolf G. (1978) Counterions in marine manganates. In La genèse des nodules de manganèse (ed. C. Lalou). Proc. Int. Coll. CNRS 289, 333-356. 477 478 Biscoe J. and Warren B.E. (1942) An X-ray study of carbon black. J. App. Phys. 13, 364-371. 479 Bodeï S., Manceau A., Geoffroy N., Baronnet A., and Buatier M. (2007) Formation of 480 todorokite from vernadite in Ni-rich hemipelagic sediments. Geochim. Cosmochim. 481 Acta 71, in press. 482 Brese N.E. and O' Keeffe M. (1991) Bond-valence parameters for solids. Acta Crystallogr. 483 **B47**, 192-197. 484 Brindley G.W. (1980) Order-disorder in clay mineral structures. In Crystal Structures of Clay 485 Minerals and their X-ray Identification (ed. G.W. Brindley and G. Brown), pp. 125-486 195. Mineralogical Society. 487 Chukhrov F.V. and Gorshkov A.I. (1980) Reply to R. Giovanoli's comment. Miner. Depos. 488 **15**, 255-257. 489 Chukhrov F.V., Gorshkov A.I., Rudnitskaya E.S., Berezovskaya V.V., and Sivtsov A.V. 490 (1980) Manganese minerals in clays: A review. Clays & Clay Miner. 28, 346-354. 491 Chukhrov F.V., Sakharov B.A., Gorshkov A.I., Drits V.A., and Dikov Y.P. (1985) Crystal 492 structure of birnessite from the Pacific ocean. Int. Geol. Rev. 27, 1082-1088. 493 Cotter-Howells J.D., Champness P.E., and Charnock J.M. (1999) Mineralogy of Pb-P grains 494 in the roots of Agrostis capillaris L. by ATEM and EXAFS. Miner. Mag. 63, 777-789. 495 Cunningham S.D., Berti W.R., and Huang J.W. (1995) Phytoremediation of contaminated 496 soils. Trends Biotech. 13, 393-397.

- 497 Drits V.A. and Tchoubar C. (1990) *X-ray diffraction by disordered lamellar structures:*498 *Theory and applications to microdivided silicates and carbons.* Springer-Verlag.
- Drits V.A., Silvester E.J., Gorshkov A.I., and Manceau A. (1997) The structure of monoclinic
 Na-rich birnessite and hexagonal birnessite. Part 1. Results from X-ray diffraction and
 selected area electron diffraction. *Amer. Mineral.* 82, 946-961.
- 502 Drits V.A., Lanson B., Gorshkov A.I., and Manceau A. (1998) Substructure and 503 superstructure of four-layer Ca-exchanged birnessite. *Amer. Mineral.* **83**, 97-118.
- 504 Drits V.A., Lanson B., and Gaillot A.C. (2007) Birnessite polytype systematics and 505 identification by powder X-ray diffraction. *Amer. Mineral.* **92**, 771-788.
- 506 Emerson D., Garen R.E., and Ghiorse W.C. (1989) Formation of *Metallogenium*-like 507 structures by a manganese-oxiding fungus. *Arch. Microbiol.* **151**, 223-231.
- Fein J.B., Martin A.M., and Wightman P.G. (2001) Metal adsorption onto bacterial surfaces:
 Development of a predictive approach. *Geochim. Cosmochim. Acta* 65, 4267-4273.
- Gaillot A.-C., Flot D., Drits V.A., Manceau A., Burghammer M., and Lanson B. (2003)
 Structure of synthetic K-rich birnessite obtained by high-temperature decomposition
 of KMnO₄. I. Two-layer polytype from 800°C experiment. *Chem. Mater.* 15, 46664678.
- Gaillot A.-C., Lanson B., and Drits V.A. (2005) Structure of birnessite obtained from
 decomposition of permanganate under soft hydrothermal conditions. 1. Chemical and
 structural evolution as a function of temperature. *Chem. Mater.* 17, 2959-2975.
- Gaillot A.-C., Drits V.A., Manceau A., and Lanson B. (2007) Structure of the synthetic K-rich
 phyllomanganate birnessite obtained by high-temperature decomposition of KMnO4 Substructures of K-rich birnessite from 1000°C experiment. *Micropor. Mesopor. Mater.* 98, 267-282.
- 521 Giovanoli R. (1980) Vernadite is random-stacked birnessite. *Mineral. Depos.* 15, 251-253.

- Hammersley A.P. (1998) FIT2D V9.129 Reference manual V3.1, Vol. 346. ESRF internal
 report ESRF98HA01T.
- Howard S.A. and Preston K.D. (1989) Profile fitting of powder diffraction patterns. In *Modern Powder Diffraction*, Vol. 20 (ed. D.L. Bish and J.E. Post), pp. 217-275.
 Mineralogical Society of America.
- Isaure M.P., Laboudigue A., Manceau A., Sarret G., Tiffreau C., Trocellier P., Lamble G.,
 Hazemann J.L., and Chateigner D. (2002) Quantitative Zn speciation in a
 contaminated dredged sediment by μ-PIXE, μ-SXRF, EXAFS spectroscopy and
 principal component analysis. *Geochim. Cosmochim. Acta* 66, 1549-1567.
- Isaure M.P., Manceau A., Geoffroy N., Laboudigue A., Tamura N., and Marcus M.A. (2005)
 Zinc mobility and speciation in soil covered by contaminated dredged sediment using
 micrometer-scale and bulk-averaging X-ray fluorescence, absorption and diffraction
 techniques. *Geochim. Cosmochim. Acta* 69, 1173-1198.
- Jurgensen A., Widmeyer J.R., Gordon R.A., Bendell Young L.I., Moore M.M., and Crozier
 E.D. (2004) The structure of the manganese oxide on the sheath of the bacterium *Leptothrix discophora*: An XAFS study. *Amer. Mineral.* 89, 1110-1118.
- 538 Krämer U. (2005) Phytoremediation: novel approaches to cleaning up polluted soils. *Curr*.
 539 *Opin. Biotechnol.* 16, 133-141.
- Krumbein W.E. and Jens K. (1981) Biogenic rock varnishes of the Negev Desert (Israel) an
 ecological study of iron and manganese transformation by cyanobacteria and fungi. *Oecologia* 50, 25-38.
- Lanson B., Drits V.A., Silvester E.J., and Manceau A. (2000) Structure of H-exchanged
 hexagonal birnessite and its mechanism of formation from Na-rich monoclinic
 buserite at low pH. *Amer. Mineral.* 85, 826-838.

546	Lanson B., Drits V.A., Feng Q., and Manceau A. (2002a) Structure of synthetic Na-birnessite:
547	Evidence for a triclinic one-layer unit cell. Amer. Mineral. 87, 1662-1671.
548	Lanson B., Drits V.A., Gaillot AC., Silvester E., Plancon A., and Manceau A. (2002b)
549	Structure of heavy-metal sorbed birnessite: Part 1. Results from X-ray diffraction.
550	Amer. Mineral. 87, 1631-1645.
551	Li Y.M., Chaney R.L., Siebielec G., and Kerschner B.A. (2000) Response of four turfgrass
552	cultivars to limestone and biosolids-compost amendment of a zinc and cadmium
553	contaminated soil at Palmerton, Pennsylvania. J. Environ. Qual. 29, 1440-1447.
554	Ma M., Lau P.S., Jia Y.T., Tsang W.K., Lam S.K.S., Tam N.F.Y., and Wong Y.S. (2003) The
555	isolation and characterization of Type 1 metallothionein (MT) cDNA from a heavy-
556	metal-tolerant plant, Festuca rubra cv. Merlin. Plant Sci. 164, 51-60.
557	Manceau A. and Combes JM. (1988) Structure of Mn and Fe oxides and hydroxides: A
558	topological approach by EXAFS. Phys. Chem. Miner. 15, 283-295.
559	Manceau A., Drits V.A., Silvester E.J., Bartoli C., and Lanson B. (1997) Structural
560	mechanism of Co ²⁺ oxidation by the phyllomanganate buserite. Amer. Mineral. 82,
561	1150-1175.
562	Manceau A., Lanson B., Drits V.A., Chateigner D., Gates W.P., Wu J., Huo D., and Stucki
563	J.W. (2000a) Oxidation-reduction mechanism of iron in dioctahedral smectites: I.
564	Crystal chemistry of oxidized reference nontronites. Amer. Mineral. 85, 133-152.
565	Manceau A., Lanson B., Schlegel M.L., Harge JC., Musso M., Eybert Berard L., Hazemann
566	J.L., Chateigner D., and Lamble G.M. (2000b) Quantitative Zn speciation in smelter-
567	contaminated soils by EXAFS spectroscopy. Amer. J. Sci. 300, 289-343.
568	Manceau A., Lanson B., and Drits V.A. (2002a) Structure of heavy metal sorbed birnessite.
569	Part III: Results from powder and polarized extended X-ray absorption fine structure
570	spectroscopy. Geochim. Cosmochim. Acta 66, 2639-2663.

571	Manceau A., Marcus M.A., and Tamura N. (2002b) Quantitative speciation of heavy metals
572	in soils and sediments by synchrotron X-ray techniques. In Applications of
573	Synchrotron Radiation in Low-Temperature Geochemistry and Environmental
574	Sciences, Vol. 49, pp. 341-428.
575	Manceau A., Marcus M.A., Tamura N., Proux O., Geoffroy N., and Lanson B. (2004) Natural
576	speciation of Zn at the micrometer scale in a clayey soil using X-ray fluorescence,

- absorption, and diffraction. *Geochim. Cosmochim. Acta* **68**, 2467-2483.
- Manceau A., Tommaseo C., Rihs S., Geoffroy N., Chateigner D., Schlegel M., Tisserand D.,
 Marcus M.A., Tamura N., and Chen Z.S. (2005) Natural speciation of Mn, Ni, and Zn
 at the micrometer scale in a clayey paddy soil using X-ray fluorescence, absorption,

and diffraction. *Geochim. Cosmochim. Acta* **69**, 4007-4034.

- Manceau A., Kersten M., Marcus M.A., Geoffroy N., and Granina L. (2007a) Ba and Ni
 speciation in a nodule of binary Mn oxide phase composition from Lake Baikal. *Geochim. Cosmochim. Acta* 71, 1967-1981.
- 585 Manceau A., Lanson M., and Geoffroy N. (2007b) Natural speciation of Ni, Zn, Ba, and As in 586 ferromanganese coatings on quartz using X-ray fluorescence, absorption, and 587 diffraction. *Geochim. Cosmochim. Acta* **71**, 95-128.
- Mandernack K.W., Post J., and Tebo B.M. (1995) Manganese mineral formation by bacterial
 spores of the marine *Bacillus*, strain SG-1: Evidence for the direct oxidation of Mn(II)
 to Mn(IV). *Geochim. Cosmochim. Acta* 59, 4393-4408.
- Marcus M.A., MacDowell A.A., Celestre R., Manceau A., Miller T., Padmore H.A., and
 Sublett R.E. (2004a) Beamline 10.3.2 at ALS: a hard X-ray microprobe for
 environmental and materials sciences. J. Synchrotron Radiat. 11, 239-247.
- Marcus M.A., Manceau A., and Kersten M. (2004b) Mn, Fe, Zn and As speciation in a fastgrowing ferromanganese marine nodule. *Geochim. Cosmochim. Acta* 68, 3125-3136.

- McKeown D.A. and Post J.E. (2001) Characterization of manganese oxide mineralogy in rock
 varnish and dendrites using X-ray absorption spectroscopy. *Amer. Mineral.* 86, 701713.
- Mench M., Bussiere S., Boisson J., Castaing E., Vangronsveld J., Ruttens A., De Koe T.,
 Bleeker P., Assuncao A., and Manceau A. (2003) Progress in remediation and
 revegetation of the barren Jales gold mine spoil after in situ treatments. *Plant and Soil*249, 187-202.
- Miyata N., Tani Y., Iwahori K., and Soma M. (2004) Enzymatic formation of manganese
 oxides by an *Acremonium*-like hyphomycete fungus, strain KR21-2. *FEMS Microbiol. Ecol.* 47, 101-109.
- Miyata N., Maruo K., Tani Y., Tsuno H., Seyama H., Soma M., and Iwahori K. (2006a)
 Production of biogenic manganese oxides by anamorphic ascomycete fungi isolated
 from streambed pebbles. *Geomicrobiol. J.* 23, 63-73.
- Miyata N., Tani Y., Maruo K., Tsuno H., Sakata M., and Iwahori K. (2006b) Manganese(IV)
 oxide production by *Acremonium sp.* strain KR21-2 and extracellular Mn(II) oxidase
 activity. *Appl. Environ. Microbiol.* 72, 6467-6473.
- Nelson Y.M., Lion L.W., Ghiorse W.C., and Shuler M.L. (1999) Production of biogenic Mn
 oxides by *Leprothrix discophora* SS-1 in a chemically defined growth medium and
 evaluation of their Pb adsorption characteristics. *Appl. Environ. Microbiol.* 65, 175180.
- Nelson Y.M., Lion L.W., Shuler M.L., and Ghiorse W.C. (2002) Effect of oxide formation
 mechanisms on lead adsorption by biogenic manganese (hydr)oxides, iron
 (hydr)oxides, and their mixtures. *Environ. Sci. Technol.* 36, 421-425.

- Nelson Y.M. and Lion L.W. (2003) Formation of biogenic manganese oxides and their
 influence on the scavenging of toxic trace elements. *Geochemical And Hydrological Reactivity Of Heavy Metals In Soils*, 169-186.
- Otte M.L., Rozema J., Koster L., Haarsma M.S., and Broekman R.A. (1989) Iron plaque on
 roots of *Aster tripolium* L.: Interaction with zinc uptake. *New Phytol.* 111, 309-317.
- Panfili F. (2004) Etude de l'évolution de la spéciation du zinc dans la phase solide d'un
 sédiment de curage contaminé induite par phytostabilisation. Ph.D. thesis, Univ. AixMarseille, France, .
- Panfili F.R., Manceau A., Sarret G., Spadini L., Kirpichtchikova T., Bert V., Laboudigue A.,
 Marcus M.A., Ahamdach N., and Libert M.F. (2005) The effect of phytostabilization
 on Zn speciation in a dredged contaminated sediment using scanning electron
 microscopy, X-ray fluorescence, EXAFS spectroscopy, and principal components
 analysis. *Geochim. Cosmochim. Acta* 69, 2265-2284.
- 632 Plançon A. (2002) CALCIPOW- a program for calculating the diffraction by disordered
 633 lamellar structures. *J. Appl. Cryst.* 35, 377.
- Römheld V. (1991) The role of phytosiderophores in acquisition of iron and other
 micronutrients in graminaceaous species: an ecological approach. *Plant & Soil* 130,
 127-134.
- Rout G.R. and Das P. (2003) Effect of metal toxicity on plant growth and metabolism: I. Zinc. *Agronomie* 23, 3-11.
- 639 Saratovsky I., Wightman P.G., Pasten P.A., Gaillard J.F., and Poeppelmeier K.R. (2006)
 640 Manganese oxides: Parallels between abiotic and biotic structures. *J. Am. Chem. Soc.*641 **128**, 11188-11198.

- Sarret G., Manceau A., Cuny D., VanHaluwyn C., Deruelle S., Hazemann J.L., Soldo Y.,
 EybertBerard L., and Menthonnex J.J. (1998a) Mechanisms of lichen resistance to
 metallic pollution. *Environ. Sci. Technol.* 32, 3325-3330.
- Sarret G., Manceau A., Spadini L., Roux J.C., Hazemann J.L., Soldo Y., EybertBerard L., and
 Menthonnex J.J. (1998b) Structural determination of Zn and Pb binding sites in *Penicillium chrysogenum* cell walls by EXAFS spectroscopy. *Environ. Sci. Technol.*32, 1648-1655.
- 649 Sarret G., Saumitou Laprade P., Bert V., Proux O., Hazemann J.L., Traverse A.S., Marcus
 650 M.A., and Manceau A. (2002) Forms of zinc accumulated in the hyperaccumulator
 651 *Arabidopsis halleri. Plant Physiol.* 130, 1815-1826.
- Schulze D.G., McCaybuis T., Sutton S.R., and Huber D.M. (1995) Manganese OxidationStates in *Gaeumannomyces*-Infested Wheat Rhizospheres Probed by Micro-Xanes
 Spectroscopy. *Phytopathology* 85, 990-994.
- Silvester E.J., Manceau A., and Drits V.A. (1997) The structure of monoclinic Na-rich
 birnessite and hexagonal birnessite. Part 2. Results from Chemical Studies and
 EXAFS Spectroscopy. *Amer. Mineral.* 82, 962-978.
- Simon L. (2005) Stabilization of metals in acidic mine spoil with amendments and red fescue
 (*Festuca rubra* L.) growth. *Environ. Geochem. Health* 27, 289-300.
- Smith R.A.H. and Bradshaw A.D. (1992) Stabilization of toxic mine wastes by the use of
 tolerant plant populations. *Trans. Inst. Min. Metall.* 81, A230-A237.
- St-Cyr L. and Campbell P. (1996) Metals (Fe, Mn, Zn) in root plaque of submerged aquatic
 plant collected in situ: relations with metal concentrations in adjacent sediments and in
 the root tissue. *Biogeochemistry* 33, 45-76.

- Tani Y., Miyata N., Iwahori K., Soma M., Tokuda S., Seyama H., and Theng B.K.G. (2003)
 Biogeochemistry of manganese oxide coatings on pebble surfaces in the Kikukawa
 River System, Shizuoka, Japan. *Appl. Geochem.* 18, 1541-1554.
- Tani Y., Miyata N., Ohashi M., Ohnuki T., Seyama H., Iwahori K., and Soma M. (2004)
 Interaction of inorganic arsenic with biogenic manganese oxide produced by a Mnoxidizing fungus, strain KR21-2. *Environ. Sci. Technol.* 38, 6618-6624.
- Tebo B.M., vanWaasbergen L.G., Francis C.A., He L.M., Edwards D.B., and Casciotti K.
 (1998) Manganese oxidation by spores of the marine *Bacillus sp.* strain SG-1 Application for the bioremediation of metal pollution. *New Developments In Marine Biotechnology*, 177-180.
- Tebo B.M., Bargar J.R., Clement B.G., Dick G.J., Murray K.J., Parker D., Verity R., and
 Webb S.M. (2004) Biogenic manganese oxides: Properties and mechanisms of
 formation. *Annu. Rev. Earth Planet. Sci.* 32, 287-328.
- 678 Teo B.K. (1986) EXAFS: basic principles and data analysis. Springer-Verlag.
- Thompson I.A., Huber D.M., Guest C.A., and Schulze D.G. (2005) Fungal manganese
 oxidation in a reduced soil. *Environ. Microbiol.* 7, 1480-1487.
- Thompson I.A., Huber D.M., and Schulze D.G. (2006) Evidence of a multicopper oxidase in
 Mn oxidation by *Gaeumannomyces graminis* var. *tritici. Phytopathology* 96, 130-136.
- Toner B., Fakra S., Villalobos M., Warwick T., and Sposito G. (2005a) Spatially resolved
 characterization of biogenic manganese oxide production within a bacterial biofilm. *Appl. Environ. Microbiol.* 71, 1300-1310.
- Toner B., Manceau A., Marcus M.A., Millet D.B., and Sposito G. (2005b) Zinc sorption by a
 bacterial biofilm. *Environ. Sci. Technol.* **39**, 8288-8294.

688	oner B., Manceau A., Webb S.M., and Sposito G. (2006) Zinc sorption to biogenic
689	hexagonal-birnessite particles within a hydrated bacterial biofilm. Geochim
690	<i>Cosmochim. Acta</i> 70 , 27-43.

- Vangronsveld J., Van Assche F., and Clijsters H. (1995) Reclamation of a bare industrial area
 contaminated by non-ferrous metals: *In situ* metal immobilization and revegetation.
 Environ. Pollut. 87, 51-59.
- Villalobos M., Toner B., Bargar J., and Sposito G. (2003) Characterization of the manganese
 oxide produced by *Pseudomonas putida* strain MnB1. *Geochim. Cosmochim. Acta* 67,
 2649-2662.
- 697 Villalobos M., Bargar J.R., and Sposito G. (2005) Mechanisms of Pb(II) sorption on a
 698 biogenic manganese oxide. *Environ. Sci. Technol.* 39, 569-576.
- Villalobos M., Lanson B., Manceau A., Toner B., and Sposito G. (2006) Structural model for
 the biogenic Mn oxide produced by *Pseudomonas putida*. *Amer. Mineral.* 91, 489502.
- Wadsley A.D. (1955) The crystal structure of chalcophanite, ZnMn₃O₇.3H₂O. *Acta Crystallogr.* 8, 165-172.
- Warren B.E. (1941) X-ray diffraction in random layer lattices. *Phys. Rev.* 59, 693-698.
- Webb S.M., Tebo B.M., and Bargar J.R. (2005a) Structural characterization of biogenic Mn
 oxides produced in seawater by the marine *Bacillus sp.* strain SG-1. *Amer. Mineral.*90, 1342-1357.
- Webb S.M., Tebo B.M., and Bargar J.R. (2005b) Structural influences of sodium and calcium
 ions on the biogenic manganese oxides produced by the marine *Bacillus sp.*, strain
 SG-1. *Geomicrobiol. J.* 22, 181-193.
- 711
- 712

FIGURE CAPTIONS

714

Fig. 1. Roots of *Festuca rubra* grown on a Zn-contaminated sediment. Optical
microphotograph (a), and scanning electron microscope image with (b)
backscattered electrons and (c) secondary electrons.

- 718Fig. 2. Tricolor (RGB) μ-XRF map of a root with Mn-Zn precipitates. Red codes for Ca,719green for Zn, and blue for Mn. Each pixel is colored in proportion to Ca-, Zn- and720Mn-Kα signals. Pixel size is $7 \times 7 \mu m^2$. The graph is a pixel-by-pixel scatterplot of721Zn counts vs. Mn counts, showing the constant Zn:Mn ratio.
- Fig. 3. Manganese K-edge EXAFS spectra of Mn-Zn precipitate (dotted lines) and the
 following references (solid lines): hollandite (Hol), todorokite (Todo), triclinic
 birnessite (TcBi), lithiophorite (Lit), low-pH hexagonal birnessite (HBi),
 chalcophanite (Chalco), and δ-MnO₂ (dBi, synthetic turbostratic birnessite).
- **Fig. 4.** Fourier transforms (FTs) of the *k*-weighted Mn-EXAFS spectra for the Mn-Zn precipitate (dotted lines) and synthetic ^{VI}Zn-sorbed δ -MnO₂ (dBi), chalcophanite (Chalco), hexagonal birnessite (HBi), and lithiophorite (Lit). E and TC subscripts denote Mn-Mn contributions from octahedral linked by edges and triple-corners, respectively.

Fig. 5. Zinc K-edge EXAFS spectra. (**a-c**) Overlay plots of the Mn-Zn precipitate spectrum (dotted line) with natural ^{IV}Zn-containing turbostratic birnessite (vernadite, sample 6KR) from Manceau et al. (2007b; **a**, solid line – Sum-sq = 0.12), ^{VI/IV}Zn-sorbed δ-MnO₂ (dBi) from Toner et al. (2006; **b**, solid line – Sum-sq = 0.83), and a leastsquares fit of the Mn-Zn precipitate spectrum to a linear combination of the model compounds in (**a**) and (**b**) (**c**, solid line – Sum-sq = 0.06). The arrows in (**a**) point out the phase misfit when the ^{VI}TC Zn species is omitted in the simulation. The dBi

sample contains 42 (±5) mol% ^{IV}TC Zn and 58 (±5) mol% ^{VI}TC Zn (Toner et al., 738 739 2006). Substituting chalcophanite for dBi in the two-component fit yielded the same 740 fraction of TC species and a similar Sum-sq residual (0.07). (d) Overlay plot of the 741 Mn-Zn precipitate spectrum (dotted line) and of the Root spectrum (solid line - Sum-742 sq = 0.10). (e-g) Overlay plots of the Root spectrum (dotted line) with organic model compounds (solid line): Zn sorbed on a biofilm at 5.6 10⁻⁴ mol of Zn per gram of 743 744 biosorbent from Toner et al. (2005b; e, Zncell7 – Sum-sq = 0.052), Zn citrate from 745 Sarret et al. (2002; \mathbf{f} , – Sum-sq = 0.089), Zn phytate from Sarret et al. (2002; \mathbf{g} , – Sum-sq = 0.089), Zn sorbed on *Penicillium chrysogenum* at 1.5 10^{-4} mol of Zn per 746 gram of biosorbent from Sarret et al. (1998b; \mathbf{h} , - Sum-sq = 0.13). The Sum-sq 747 values are calculated as the squares of the residuals, normalized to the sum of the 748 squares of the data values over the $[3.0 - 10.5 \text{ Å}^{-1}] k$ interval. 749

Fig. 6. XRD pattern of Mn-Zn precipitate after subtraction of the scattering from the root.

751 Fig. 7. Simulations of the [20,11] and [02,31] X-ray scattering bands (C-centered layer cell). 752 Black crosses are experimental data, and red lines are calculated profiles. Small but 753 significant misfits between experimental and calculated patterns are pointed out with 754 arrows. Diffracted intensities were calculated with a turbostratic layer stacking (no 755 interlayer correlation). (a) Initial model with 0.186 vacant layer sites and no 756 interlayer Mn. (b) Optimal model (Table 1; Figs. 8 and EA-5). (c) Model with a =5.014 Å and b = 2.895 Å (hexagonal layer symmetry). (d) Model with a = 5.100 Å 757 and b = 2.850 Å (orthogonal layer symmetry). (e) Model with all interlayer Mn 758 cations (0.056 per octahedron) located in ^{VI}TC sites, instead of 0.046 and 0.010 in 759 ^{VI}TE and ^{VI}TC sites, respectively, in the optimal fit. (f) Model with 0.240 Ca per 760 761 layer octahedron instead of 0.318 in the optimal model. (g) Model with 0.150 vacant sites and 0.244 interlayer Zn^{2+} (Zn:Mn = 0.27), compared to 0.220 and 0.388 762

- 763 (Zn:Mn = 0.46) in the optimal fit. (**h**) Model with 50% tetrahedral and 50% 764 octahedral Zn²⁺, instead of 77% and 23% in the optimal fit. Unless specified, all 765 parameters used in calculations are those of the optimal model.
- 766Fig. 8. Idealized structure for the Mn-Zn root precipitate with structural formula767 $[(Mn_{0.78}Vac_{0.22})O_2]Mn^{VITC}{}_{0.010}Mn^{VITE}{}_{0.046}Zn^{VITC}{}_{0.088}Zn^{IVTC}{}_{0.300}Ca_{0.318}(H_2O)_{0.972}].$ The768layer charge deficit due to vacant layer sites is balanced by interlayer Zn and Mn769above and/or below layer vacancies, and by Ca in the interlayer mid-plane (see Fig.770EA-5 for details).
- Fig. 9. Calculation of the 001 and 002 reflections for crystallites composed of one (dashed line), two (dotted-dashed line) and three (dot-dot-dashed line) parallel layers randomly stacked perpendicular to the *ab* plane, and optimal fit (red line) to the data (crosses) obtained with an assemblage of diffracting particles containing 1, 2, and 3 layers in the ratio of 2.0:0.3:0.1. Atomic coordinates of the optimal structure model, derived from the simulation of the [20,11] and [02,31] reflections, were used in all calculations (Figs. 7b and 8, Table 1).

Atom	x	у	ζ	Occ.	x	у	ζ	Occ.
Mn _{layer} (Mn1)	0	0	0	0.780	-	-	-	-
O _{layer} (O1)	0.333	0	1.00	1.00	-0.333	0	-1.00	1.00
^{TC} Mn _{inter} (Mn2)	0	0	2.20	0.005	0	0	-2.20	0.005
H ₂ O _{inter.} (O2)	-0.333	0	3.45	0.015	0.333	0	-3.45	0.015
^{TE} Mn _{inter} (Mn3)	-0.333	0	2.20	0.023	0.333	0	-2.20	0.023
H ₂ O _{inter.} (O3)	0	0	3.45	0.069	0	0	-3.45	0.069
^{VI} Zn _{inter} (Zn1)	0	0	2.20	0.044	0	0	-2.20	0.044
H ₂ O _{inter.} (O2)	-0.333	0	3.45	0.132	0.333	0	-3.45	0.132
^{IV} Zn _{inter} (Zn2)	0	0	1.77	0.150	0	0	-1.77	0.150
H ₂ O _{inter.} (O4)	0	0	3.70	0.150	0	0	-3.70	0.150
Ca _{inter} (Ca1)	-0.410	0	3.60	0.053	0.410	0	-3.60	0.053
Ca _{inter} (Ca1)	-0.295	0.115	3.60	0.053	0.295	0.115	-3.60	0.053
Ca _{inter} (Ca1)	-0.295	-0.115	3.60	0.053	0.295	-0.115	-3.60	0.053
H ₂ O _{inter.} (O5)	0.220	0	3.60	0.04	-0.220	0	-3.60	0.04
$H_2O_{inter.}$ (O5)	-0.110	0.330	3.60	0.04	-0.110	-0.330	-3.60	0.04
H ₂ O _{inter.} (O5)	-0.110	-0.330	3.60	0.04	0.110	-0.330	-3.60	0.04

Table 1. Structural parameters of Mn-Zn precipitate derived from the simulation of XRD data.

Note: b = 2.850 Å, $a = b\sqrt{3} = 4.936$ Å, $\gamma = 90^{\circ}$, and $d_{(001)} = 7.20$ Å. *x* and *y* coordinates are expressed as fractions of the *a* and *b* parameters, respectively. Coordinates along the *c**

axis, ζ , are expressed in Å to point out the thickness of layer and interlayer polyhedra. The average dimension of the coherent scattering domains (CSDs) along the *c** axis is 1.2 layers (8.6 Å). The average radius of the disk-like CSDs in the *ab* plane is 33 Å. This value was calculated by fitting the [20,11] reflection. Un-refined thermal B factors are 0.5 Å² for Mn_{layer}, 1.0 Å² for O_{layer}, interlayer Mn, and interlayer Zn, and 2.0 Å² for Ca²⁺ and H₂O.

Atomic pair	Distance (Å)	Atomic pair	Distance (Å)		
Mn _{layer} -O _{layer}	1.925				
${}^{TC,TE}Mn_{inter.}\text{-}O_{layer}$	2.04	$^{TC,TE}Mn_{inter.}$ -H ₂ O	2.07		
^{VI} Zn _{inter.} -O _{layer}	2.04	^{VI} Zn _{inter.} -H ₂ O	2.07		
^{IV} Zn _{inter.} -O _{layer}	1.82 ^a	^{IV} Zn _{inter.} -H ₂ O	1.93		
CainterOlayer	2.89	H_2O_{inter} - O_{layer}	2.66		
Mn _{layer} -VIZn _{inter}	3.60	Mn_{layer} - ^{IV} Zn_{inter}	3.35		
Mn _{layer} - ^{TC} Mn _{inter.}	3.60	Mn_{layer} -TE $Mn_{inter.}$	2.75 ^b		

Table 2. Selected interatomic distances for the optimal structure model.

^(a) The ^{IV}Zn_{inter.}-O_{layer} distance should be increased by ~0.1 Å to provide more realistic Zn-O bond valence (Table EA-1). This cannot be achieved by changing the *z* coordinate of ^{IV}Zn (Fig. EA-6b). Alternatively, moving apart the three oxygen atoms delimiting the underlying vacancy by 0.15 Å in the [110], [100], and [010] directions increases the ^{IV}Zn_{inter.}-O_{layer} distance from 1.82 Å to 1.95 Å, as discussed in Manceau et al. (2002a).

^(b) The Mn_{layer} -^{TE} $Mn_{inter.}$ distance can be increased by ~0.1 Å by shifting Mn_{inter} in the *ab* plane from its ideal TE position towards the nearest vacant site (see Fig. EA-5). This shift occurs along the $a \pm 120^{\circ}$ directions with equal probabilities resulting in an increased Debye-Waller B factor without significant alteration of the fit quality.

	O1 ^b	O1 ^c	$O1^d$	O1 ^e	02, 03	03	O4	Σ	Formal
									valence
	0.628 ×6→							3.8	4
Mini	$\times 3 \downarrow$	0.628 ×2↓	0.628 ×2↓	0.628 ×2↓					
Mn2,		0 469 ×3→			0 433 ×3→			27	3
Mn3		0.109 - 5 7			0.155 * 5 7			2.7	5
Zn1			0.403 ×3→			0.372 ×3→		2.3	2
Zn2				0.731 ×3→			0.543	2.7	2
H^{+}	0.11^{f}								
Σ	$1.9 - 2.0^{g}$	1.7	1.7	2.0					

Table EA-1. Empirical bond-valences for Mn-Zn precipitate^a.

^(a) Bond valences in valence unit (v.u.) were calculated using the Valence for Dos program (v. 2.0 -

http://www.ccp14.ac.uk/solution/bond_valence/index.html) and the parameters from Brese and O' Keeffe (1991).

^(b) O1 coordinated to 3 Mn⁴⁺ in Mn1 (Table 1).

 $^{(c)}$ O1 coordinated to 2 Mn^{4+} in Mn1 and 1 Mn^{3+} in Mn2 or Mn3.

 $^{(d)}$ O1 coordinated to 2 Mn^{4+} in Mn1 and 1 Zn^{2+} in Zn1.

 $^{(e)}$ O1 coordinated to 2 Mn^{4+} in Mn1 and 1 Zn^{2+} in Zn2.

^(f) O5-H-O1 H-bond.

 $^{(g)}$ Depending on whether this O1 receives additional valence from H⁺ through H-bond.





Lanson et al., Fig. 1







Lanson et al., Fig. 4







Lanson et al., Fig. 7









Fig. EA-1: Roots of *Festuca rubra* grown on a Zn-contaminated sediment. (**a**) Close-up photograph from Fig. 1a. (**b**) Lower magnification of Fig. 1c.



Fig. EA-2: Tricolor (RGB) μ -XRF map of a bundle of roots. Red codes for Fe, green for Zn, and blue for Mn. Each pixel is colored in proportion to Fe-, Zn- and Mn-K α signals. Pixel size is 18 × 15 μ m².



for the Mn-Zn root precipitate and synthetic VI Zn-sorbed δ -MnO₂ (dBi, synthetic turbostratic Fig. EA-3: Fourier transforms (FTs) of the manganese K-edge k-weighted EXAFS spectra birnessite), chalcophanite (Chalco), hexagonal birnessite (HBi), and lithiophorite (Lit). Expansion of the [1-3 Å] $R+\Delta R$ interval.



Fig. EA-4: Simulations of the [20,11] and [02,31] X-ray scattering bands (black crosses, C-centered layer cell). Intensities (red lines) were calculated for the optimal structure model (Table 1) with different occurrence probabilities of random stacking faults (W_r). Arrows indicate the positions of *hkl* reflections.





Fig. EA-5: Idealized structure of Mn-Zn precipitate. (a) Projection on the *ab* plane. The upper surface of the layer is shown as light shaded triangles, and the atomic notations are the same as in Table 1. (b) Projection along the *b* axis. Open and solid symbols indicate atoms at y = 0 and $y = \pm \frac{1}{2}$, respectively. Squares represent vacant layer octahedra. The Mn₃ and Zn₂ atoms can be shifted from their positions as indicated by the dashed arrows to obtain more realistic interatomic distances (see Table EA-1).



Fig. EA-6: Simulations of the [20,11] and [02,31] X-ray scattering bands (C-centered layer cell, black crosses). Small but significant misfits between experimental and calculated patterns are pointed out with arrows. Intensities (red lines) were calculated with a turbostratic layer stacking (no interlayer correlation). (a) Optimal model (Table 1; Figs. 8 and EA-5); $R_{wp} = 3.49\%$. (b) Model with ^{IV}Zn (Zn2) in (0, 0, 1.97 Å), instead of (0, 0, 1.77 Å) in the optimal fit, so as to increase the Zn-O bond length from 1.82 Å (Table 2) to 1.91 Å. Coordinated H₂O molecules (O4) were moved from (0, 0, 3.70 Å) to (0, 0, 3.90 Å); $R_{wp} = 3.92\%$. (c) Model with ^{VI}Zn (Zn1) in (0, 0, 2.30 Å), instead of (0, 0, 2.20 Å) in the optimal fit so as to decrease the sum valence of Zn1 from 2.3 (Table EA-1) to 2.0. Coordinated H₂O molecules (O2) were moved from (-0.333, 0, 3.65 Å); $R_{wp} = 3.55\%$. (d) Model with Ca in (-0.333, 0, 3.60 Å), instead of (-0.410, 0, 3.60 Å) and equivalent positions in the optimal fit; $R_{wp} = 3.51\%$. Unless specified, all parameters used in calculations are those of the optimal model.



Fig. EA-7: Fourier transform of the EXAFS spectra for the Mn-Zn precipitate at the Mn and Zn K-edges. The average Mn-O and Zn-O EXAFS distances (R values) are 1.90 Å and 2.00 Å, respectively.