



HAL
open science

Supporting evidence from the EPICA Dronning Maud Land ice core for atmospheric CO₂ changes during the past millennium

Urs Siegenthaler, Eric Monnin, Kenji Kawamura, Renato Spahni, Jakob Schwander, Bernard Stauffer, Thomas F. Stocker, Jean-Marc Barnola, Hubertus Fischer

► To cite this version:

Urs Siegenthaler, Eric Monnin, Kenji Kawamura, Renato Spahni, Jakob Schwander, et al.. Supporting evidence from the EPICA Dronning Maud Land ice core for atmospheric CO₂ changes during the past millennium. *Tellus B - Chemical and Physical Meteorology*, 2005, 57 (1), pp.51 à 57. 10.1111/j.1600-0889.2005.00131.x . insu-00374931

HAL Id: insu-00374931

<https://insu.hal.science/insu-00374931>


Submitted on 22 Jun 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Supporting evidence from the EPICA Dronning Maud Land ice core for atmospheric CO₂ changes during the past millennium

Urs Siegenthaler, Eric Monnin, Kenji Kawamura, Renato Spahni, Jakob Schwander, Bernhard Stauffer, Thomas F. Stocker, Jean-Marc Barnola & Hubertus Fischer

To cite this article: Urs Siegenthaler, Eric Monnin, Kenji Kawamura, Renato Spahni, Jakob Schwander, Bernhard Stauffer, Thomas F. Stocker, Jean-Marc Barnola & Hubertus Fischer (2005) Supporting evidence from the EPICA Dronning Maud Land ice core for atmospheric CO₂ changes during the past millennium, Tellus B: Chemical and Physical Meteorology, 57:1, 51-57, DOI: [10.3402/tellusb.v57i1.16774](https://doi.org/10.3402/tellusb.v57i1.16774)

To link to this article: <https://doi.org/10.3402/tellusb.v57i1.16774>



© 2005 The Author(s). Published by Taylor & Francis.



Published online: 18 Jan 2017.



[Submit your article to this journal](#)



Article views: 250



[View related articles](#)



Citing articles: 4 [View citing articles](#)

Supporting evidence from the EPICA Dronning Maud Land ice core for atmospheric CO₂ changes during the past millennium

By URS SIEGENTHALER^{1*}, ERIC MONNIN¹, KENJI KAWAMURA¹, RENATO SPAHNI¹, JAKOB SCHWANDER¹, BERNHARD STAUFFER¹, THOMAS F. STOCKER¹, JEAN-MARC BARNOLA² and HUBERTUS FISCHER³, ¹*Climate and Environmental Physics, Physics Institute, University of Bern, Bern, Switzerland;* ²*CNRS Laboratoire de Glaciologie et de Géophysique de l'Environnement (LGGE), Grenoble, France;* ³*Alfred Wegener Institut für Polar- und Meeresforschung, Bremerhaven, Germany*

(Manuscript received 17 January 2004; in final form 5 August 2004)

ABSTRACT

The most direct method of investigating past variations of the atmospheric CO₂ concentration before 1958, when continuous direct atmospheric CO₂ measurements started, is the analysis of air extracted from suitable ice cores. Here we present a new detailed CO₂ record from the Dronning Maud Land (DML) ice core, drilled in the framework of the European Project for Ice Coring in Antarctica (EPICA) and some new measurements on a previously drilled ice core from the South Pole. The DML CO₂ record shows an increase from about 278 to 282 parts per million by volume (ppmv) between AD 1000 and AD 1200 and a fairly continuous decrease to a mean value of about 277 ppmv around AD 1700. While the new South Pole measurements agree well with DML at the minimum at AD 1700 they are on average about 2 ppmv lower during the period AD 1000–1500. Published measurements from the coastal high-accumulation site Law Dome are considered as very reliable because of the reproducibility of the measurements, high temporal resolution and an accurate time scale. Other Antarctic ice cores could not, or only partly, reproduce the pre-industrial measurements from Law Dome. A comparison of the trends of DML and Law Dome shows a general agreement. However we should be able to rule out co-variations caused by the same artefact. Two possible effects are discussed, first production of CO₂ by chemical reactions and second diffusion of dissolved air through the ice matrix into the bubbles. While the first effect cannot be totally excluded, comparison of the Law Dome and DML record shows that dissolved air diffusing to bubbles cannot be responsible for the pre-industrial variation. Therefore, the new record is not a proof of the Law Dome results but the first very strong support from an ice core of the Antarctic plateau.

1. Introduction

CO₂ measurements of recent air extracted from ice cores (Neftel et al., 1985; Etheridge et al., 1996) overlap with direct atmospheric measurements (Keeling and Whorf, 2003) and are in good agreement, clearly confirming the reliability of the recent ice core CO₂ record. Records from Antarctic ice cores show a mean value of about 280 parts per million by volume (ppmv) for the last millennium before the anthropogenic increase started (Barnola, 1999). However, variations between 5 and 10 ppmv are observed in the pre-industrial epoch in Antarctic records from Siegenthaler et al. (1988), Barnola et al. (1995) and Etheridge et al. (1996). These variations, which are important for the understanding of the global carbon cycle, are significant con-

sidering the analytical uncertainties, but the time evolution of these records is quite different. A record from Central Greenland (Barnola et al., 1995) shows CO₂ concentrations of 283.0 ± 2.4 ppmv for the period AD 1600–1800 and 289.9 ± 2.9 ppmv between AD 1000 and 1600. These values are significantly higher than the ones from Law Dome (277.9 ± 2.0 ppmv and 281.6 ± 1.4 ppmv, respectively) and in view of *in situ* production of CO₂ in Greenland ice (Anklin et al., 1995; Smith et al., 1997) are not considered to represent the atmospheric CO₂ concentration. Ice cores from Antarctic high-plateau drill sites, generally exhibiting very low surface temperatures and low impurity concentrations, are better suited for reliable CO₂ measurements. However, the temperature at Law Dome is 10 °C higher than at Summit and the impurity concentrations of Ca²⁺ and H₂O₂, which are connected to *in situ* CO₂ production are about one-half of those at Summit (Table 1). Accordingly, significant atmospheric CO₂ changes as recorded in the Law Dome ice core during the

*Corresponding author.
e-mail: siegenthaler@climate.unibe.ch

Table 1. Details of the ice cores used in this study

Ice core	Location	Accumulation rate ($\text{kg m}^{-2} \text{ yr}^{-1}$)	Mean temperature ($^{\circ}\text{C}$)	Delta age (yr)	Age distribution width at half height (yr)	H_2O_2 (ppbw)	Ca^{2+} (ppbw)
DML	75°00'S, 00°04'E	64	-44.6	825	59 ± 5	10.1 ^a	1.3 ^a
South Pole	90°S	76.2	-51	950	95 ± 5	11.6/16.7 ^b	
DSS Law Dome	66°46'S, 112°48'E	600	-22	58		34/41 ^c	3.6 ^d
Summit (GRIP)	72°34'N, 37°37'W	200	-32	210		72.6 ^e	8.1 ^e

^aMean value of the continuous measurements on the B32 core covering the gas age period of AD 1300–1900. (Sommer, 2000).

^bMeasurements in a depth attributed to an age in the gas age scale used in this work of AD 1369 and AD 975 (Nefel et al., 1984).

^cMean value of samples with a mean age of AD 1648 and AD 1541 (Gillet et al., 2000).

^dMean value of continuous measurements covering the period of AD 1301–1997 (Mark Curran, personal communication).

^eMean value of continuous measurements covering the period of 9000–7800 BP (Führer et al., 1993).

last millennium (Etheridge et al., 1996) require an independent confirmation from high-altitude Antarctic sites. In addition the reason for the difference in Greenland and Antarctic ice core CO_2 records over the last millennium has not been ultimately resolved.

In this paper we present new CO_2 data derived from the EPICA (European Project for Ice coring in Antarctica) ice core currently being drilled in Dronning Maud Land (DML) as well as additional measurements on the South Pole ice core and compare those with the high-resolution Law Dome CO_2 record. In addition, we try to quantify potential artefacts which may be responsible for intersite differences in CO_2 concentrations. Those processes are chemical reactions within the ice as well as diffusion of high CO_2 trapped in microbubbles in the ice during snow formation into larger air bubbles, both processes having different magnitudes and rates due to different rates of snow accumulation, temperature and impurity content.

2. Methods

Generally, a dry extraction method is used where samples of 6–8 g are crushed by a needle cracker. The partial pressure of CO_2 is measured using infrared laser spectrometry (IRLS). The analytical uncertainty of standard gas introduced over bubble-free ice was 1.7 ppmv during the measurements on the DML core. More detailed information about the analytical method is given in Indermühle et al. (1999). We measured samples covering the time period from AD 950 to AD 1900 on 71 and 14 depth intervals of the DML and the South Pole ice core, respectively. At each depth interval of 60–100 mm, six samples were measured. Figure 1 shows the arithmetic means of six samples for each depth interval with a standard error of the mean in the range of 0.5–2.5 ppmv as error bars.

The DML and South Pole data are corrected for gravitational separation of gases in the firn column based on the barometric formula to get closer to the true atmospheric value. The close-off depths are calculated with the same model as in Barnola




Fig 1. Full circles: CO_2 data from DML. Open triangles: New measurements on the South Pole ice core. Both records were measured with dry extraction (mean of six samples; error bars, 1σ of the mean). Open circles: Law Dome data (Etheridge et al., 1996).

et al. (1991) with the close-off densities calculated according to Martinerie et al. (1994) (834 and 837 kg m^{-3} for DML and South Pole, respectively). For a concentration of 280 ppmv, the gravitational correction for DML and the South Pole is 1.9 and 2.6 ppmv, respectively.

No published timescale is available yet for the DML ice core. We construct a tentative timescale by adjusting the DML timescale to Dome C by comparing the records of electrical conductivity for both cores. The difference of gas age/ice age was assumed as 825 yr.

3. Results

The DML CO_2 record shows an increase from about 278 to 282 ppmv between AD 1000 and AD 1200 and a fairly continuous decrease to a mean value of about 277 ppmv around AD 1700

(Fig. 1). Variations of the atmospheric CO₂ concentrations on a centennial timescale were suggested many years ago based on measurements along the South Pole ice core (Siegenthaler et al., 1988). We remeasured samples from this core for 14 depth levels. The results are shown in Fig. 1. A new timescale was calculated, because an event recorded with the electrical conductivity method (ECM) attributed to a volcanic eruption in AD 1259 (Langway et al., 1988) indicates that the timescale used in Siegenthaler et al. (1988) is too young at this depth. This results in a lower average accumulation rate of $76.2 \text{ kg m}^{-2} \text{ yr}^{-1}$ compared with the previously used value of $84 \text{ kg m}^{-2} \text{ yr}^{-1}$. On average, the South Pole scale was about 200 yr too young over the last millennium. The new South Pole data follow the lower envelope of the data published in 1988 (not shown) and, similar to the DML results, also show a fairly continuous decrease between AD 1200 to a minimum concentration around AD 1700. While this minimum agrees well with DML, the South Pole data are on average about 2 ppmv lower than the DML record in the first half of the millennium. The most evident difference is at about AD 1200, where two South Pole measurements are 3 ppmv lower than the trend of the DML record. Therefore, the decrease from AD 1200–1700 seems to be less pronounced in the South Pole record.

The record from Law Dome (Etheridge et al., 1996) is considered as very reliable because of the reproducibility of the measurements of 1.2 ppmv and the high accumulation rate of $600 \text{ kg m}^{-2} \text{ yr}^{-1}$ at Law Dome leading to a higher temporal resolution and a smaller error in the ice age/gas age difference. For the DML and South Pole sites, the observed accumulation rates are about nine and eight times lower, respectively. Low accumulation rates cause an attenuation of fast variations in atmospheric CO₂ concentration which has to be taken into account. Processes for the attenuation are molecular diffusion in the open pores of the firn column and the gradual bubble close-off in the depth of the transition from firn to ice. The lower the accumulation rate, the wider is the age distribution of the enclosed air. The model by Spahni et al. (2003), which takes both molecular diffusion and gradual bubble close-off into account, has been used to determine the attenuation. An adjustment with the density profile and estimates of today's firn parameters in the model results in an air age distribution with a width at half height of 59 and 95 yr for the DML and South Pole cores, respectively. We use these age distributions for comparison of the new results with data from the high-accumulation site at Law Dome. To this end we smoothed the Law Dome data with the modelled age distribution of DML and the South Pole (Fig. 2).

The most evident difference between DML and Law Dome is in the period AD 1000–1500 where the Law Dome values are higher by about 0.5–2.5 ppmv. Although the decrease at about AD 1600 seems to be less pronounced because of lower values at about AD 1550 in the DML core, the remarkable minimum between AD 1600 and AD 1750, which was questioned by Stauffer et al. (2002), could apparently be confirmed.




Fig 2. Dashed thin curve: interpolated Law Dome data (Etheridge et al., 1996). Dashed thick and grey curves: signal of Law Dome as it would be smoothed in the ice cores from DML and the South Pole, respectively. The signal is calculated for both cores using the modelled age distribution after Spahni et al. (2003). Full black and dotted grey curves: spline with a cut-off period of 200 yr of the DML and South Pole data, respectively.

4. Discussion

The atmospheric CO₂ evolution recorded in the DML record confirms, within the uncertainties of the measurements and the air age resolution, the findings from the Law Dome record. It was speculated that the distinct minimum at about AD 1700 is related to the Little Ice Age. However, it would be very surprising if this climatic event, which is ubiquitous in the North Atlantic region but not well-defined temporally and of rather small temperature anomalies, would have such a distinct effect on CO₂. Another feature observed in the CO₂ data is difficult to explain. The model of Gerber et al. (2003), forced by the reconstructed variability in solar irradiance and radiative forcing by volcanoes, simulated a fairly continuous decrease of atmospheric CO₂ between AD 1200 and AD 1800, but the measured increase between AD 1000 and AD 1200 was not reproduced by the model.

Knowledge of the palaeoatmospheric carbon dioxide concentration is a key for a better understanding of the global carbon cycle. Therefore it has to be carefully checked if the good agreement of the records from DML and Law Dome could not be affected by the same artefact, despite different characteristics of the drilling sites.

Two possibilities for such artefacts are:

- CO₂ is produced by chemical reactions between impurities in the ice. The reactions need centuries to approach an equilibrium and do therefore not affect very young ice.
- CO₂ originally dissolved or enclosed as microbubbles in firn grains is not extracted in young ice by a dry extraction

method. In older ice the dissolved CO₂ will diffuse to the bubbles and will be extracted by dry extraction.

Both artefacts would not influence the good agreement of CO₂ records from ice cores with direct atmospheric measurements in the overlapping period.

CO₂ can be produced in the ice by acid-carbonate reactions or the oxidation of organic material (Tschumi and Stauffer, 2000). This production will certainly depend on the concentration of impurities (carbonates, acidic compounds, dissolved organic compounds, oxidation agents such as H₂O₂) and it will need a certain time to appear. It was observed that reaction compounds are still present after thousands of years in ice cores. One or several of the original reaction compounds can be concentrated in dust grains, in grain boundaries or in veins, or formed along the edges where three ice crystals meet. Tschumi and Stauffer (2000) suggest that the reaction rate is that low because it depends on the probability that molecules from each source compound will come close enough to react. The produced CO₂ will occur, like the dissolved CO₂ from microbubbles, somewhere within the ice matrix and has to diffuse to air bubbles to be extracted by a dry extraction method. Therefore, any dry extraction method would measure such surplus CO₂ with a delay which depends mainly on the temperature of the ice.

The concentrations of impurities in the coastal Law Dome ice (Gillet et al., 2000; Souney et al., 2002) are higher than in the ice of DML on the Antarctic Plateau (Sommer, 2000). Concentrations of Ca²⁺ (a proxy for carbonate concentration) and of the oxidation agent H₂O₂ are shown in Table 1.

However, if CO₂ produced by chemical reactions were responsible for the values above 276 ppmv before AD 1700 then it would be difficult to explain why the concentration would be low for all three records at the beginning of the millennium again.

Can we, based on these considerations, exclude any artefacts? Not necessarily—there is an additional observation which is puzzling. High-resolution records over short time intervals from the Dome C and DML record show a significantly higher scatter of CO₂ measurements for young ice (last millennium) than for older ice. This scatter is slightly higher than the analytical uncertainty, indicating the existence of some artefacts caused by processes in the ice sheet or during the extraction procedure. However, it is unlikely that the mean deviations exceed 1% of the measured concentration (Stauffer et al., 2003). In summary an explanation of variations or intersite differences in the three records based on *in situ* chemical production of CO₂ does not seem to be straightforward.

To quantify the effect of microbubbles we additionally performed extraction of air from ice cores with a sublimation method which allows us to also extract the CO₂ dissolved in the ice. Sublimation takes place below the triple point of water, which avoids dissolving CO₂ and a potential production of CO₂ by acid-carbonate reactions in the liquid phase (Stauffer and Berner, 1978). Furthermore, extraction by sublimation avoids fraction-

ation effects, because the extraction efficiency is close to 100% and independent of whether air is enclosed in bubbles, clathrates or dissolved in the ice.

The sublimation device, used at our laboratory for tests and not for routine measurements, is similar to the one of G ullik et al. (1998). Tests of the sublimation extraction were performed with bubble-free ice and the addition of standard gas. Tests during the time period of the DML measurements showed a reproducibility of 1.5 ppmv with a mean concentration which was 0.1 ppmv below the concentration of the standard gas. The eight measurements on the DML ice core during the time period AD 1000–1800 obtained with the sublimation method (not shown) are on average 1.0 ppmv higher than the results obtained with the dry extraction. Unfortunately, the sublimation results show a surprisingly large scatter (about 3 ppmv) and therefore a larger uncertainty than in the tests with bubble-free ice. However, the agreement of the two extraction methods allows the conclusion that a fractionation effect for the dry extraction of more than 5 ppmv is unlikely in the measured DML ice. But the results unfortunately do not allow us to confirm or disprove the hypothesis of dissolved CO₂ in young ice causing a fractionation effect of the order of a few ppmv.

Therefore, we try to quantify this effect using a simple diffusion model. The CO₂ content enclosed or dissolved in the firn is enriched compared with the air enclosed in bubbles after the firn–ice transition. Measurements with a wet extraction on evacuated firn samples from the North Central ice core (74°37'N, 39°36'W, *T* = –31.7°C) showed that especially the first 10 m of the firn core were enriched in CO₂ by more than 10 000 ppmv (Stauffer, 1981). While wet extraction of CO₂ is prone to acid-carbonate reactions, another hypothesis is that highly enriched CO₂ is enclosed in microbubbles, which would not be opened by a classical dry extraction method.

The question arises of how fast this possibly high level of CO₂ permeates through the ice matrix into the normal air bubbles and therefore influences the measured CO₂ concentration. In the Vostok ice core, it was observed that the volume fraction of microbubbles with respect to all bubbles is 0.3% (Lipenkov, 2000). For an estimate we assume that 0.5% of the total air content (100 ml air STP per kg ice) is enclosed in microbubbles. The composition of air in the normal bubbles is assumed to be atmospheric with a CO₂ concentration of 275 ppmv. The composition of the air in the microbubbles is certainly enriched in more soluble components. The partial pressure of CO₂ in air dissolved in water of 0 °C in equilibrium with an atmosphere of 275 ppmv is greater than 15 000 ppmv. A large part of the enriched air may escape through the open pores of the firn. Because of the growth of firn grains by sublimation–condensation, most of the microbubbles have contact with the firn grain boundaries. Using the parametrization of Gow (1969) for the firn grain growth, the fraction of microbubbles that have never had contact with grain boundaries in the firn zone, where the escape of enriched air through the open pores might be possible, is in the




Fig 3. Calculation of the CO₂ concentrations of air enclosed in normal bubbles with the boundary conditions explained in the text, $t = 0$ is at close-off. For every site, three calculations were made with different initial CO₂ concentrations in the microbubbles of 275, 500 and 1000 ppmv. The corresponding equilibrium concentrations (if all air from the microbubbles and normal bubbles is mixed up) are 275, 276.1 and 278.6 ppmv, respectively. Dashed curves: calculations for Law Dome DSS (mean temperature -22°C). Full curves: calculations for DML (-44.6°C). Dotted curves: calculations for the South Pole (-51°C).

range of 10% for the three cores. This result strongly depends on the assumptions about grain sizes at the surface of the firn zone.

Little is known about CO₂ concentrations of air included in microbubbles. We therefore started our model with three different CO₂ concentrations in microbubbles of 1000, 500 and 275 ppmv. The initial CO₂ mixing ratio of normal bubbles is 275 ppmv. We assume that the composition of O₂ and N₂ is that of the atmosphere.

The assumptions for the calculation of this diffusion or permeation are given in the Appendix. The results for the evolution of the CO₂ concentration in normal bubbles calculated for the sites of South Pole, DML and Law Dome are shown in Fig. 3. The concentration which would be measured, if the air of microbubbles and normal bubbles was mixed, are 278.6, 276.1 and 275 ppmv, respectively. However, these values are only reached after all gases from the microbubbles reach an equilibrium in their composition and pressure with normal bubbles by molecular diffusion. The results of the model show that the effect of diffusion through the ice matrix may influence the concentration in the normal bubbles by several ppmv. The partial pressure of CO₂ in the normal bubbles first decreases because of the larger ratio of N₂ and O₂ diffusing into the normal bubbles due to larger permeation coefficients compared with CO₂. Then, the partial pressure of CO₂ increases up to the equilibrium concentration when the concentrations in normal bubbles and microbubbles approach equilibrium, which needs more than 1000 yr. The time response of the minimum and the trend in general strongly depends on

the permeation coefficients and its temperature dependence, and also on the assumptions concerning bubble pressure. The results calculated for South Pole, DML and Law Dome show large differences, primarily due to the different temperatures of the sites. Therefore, we conclude that air dissolved in microbubbles cannot explain the distinct CO₂ minimum observed in all records (and also in the Greenland records) at AD 1700 which is of the order of about 5 ppmv and which is achieved on a timescale of just a few centuries and which, therefore, is clearly an atmospheric phenomenon.

5. Conclusions

Based on the good agreement of the general trend of the records from Law Dome and DML and also with a few values from the South Pole with very different ice characteristics, we conclude that the observed time evolution very likely describes the evolution of the atmospheric CO₂ concentration. In particular, the increase between AD 1000 and AD 1200 and the distinct CO₂ minimum between AD 1600 and AD 1750 is a robust feature which cannot be explained by diffusion of CO₂ from microbubbles or by production of CO₂ by chemical reactions. However, despite these considerations we cannot exclude any artefacts. No explanation for the observation of higher scatter of results in young compared with older ice has been found yet. Even if the general evolution of the atmospheric CO₂ concentration of the past millennium is not in question, there remain open questions concerning the concentration differences of up to 4 ppmv in the trends of the three records in the first half of the last millennium.

6. Acknowledgments

This work is a contribution to the “European Project for Ice Coring in Antarctica” (EPICA), a joint ESF (European Science Foundation)/EC scientific programme, funded by the European Commission and by national contributions from Belgium, Denmark, France, Germany, Italy, The Netherlands, Norway, Sweden, Switzerland and the United Kingdom. The measurements were supported by the Swiss NSF, University of Bern and the “Bundesamt für Energie”. This is EPICA publication no 107.

7. Appendix

CO₂ diffuses from microbubbles to normal bubbles through the ice matrix. A higher partial pressure surrounding a microbubble at close-off is caused by the higher CO₂ concentration of the air (except for the calculations with the same initial concentrations for microbubbles and normal bubbles), but also because the pressure in microbubbles is slightly higher than in normal bubbles. At Vostok, it was observed that the pressure of microbubbles and normal bubbles converges with increasing depth and becomes identical below a depth of 500 m (Lipenkov, 2000). The flux of CO₂ from microbubbles to normal bubbles is

Table 2. Parameters constant with time

Symbol	Definition	Value
n_n	Density of normal bubbles	480 cm ⁻³
n_a	Density of microbubbles	120 cm ⁻³
$D_{N_2}^{s0}$	Constant for diffusive permeation of N ₂	2.4 × 10 ⁻⁹ m ² s ⁻¹ MPa ⁻¹
$D_{O_2}^{s0}$	Constant for diffusive permeation of O ₂	2.4 × 10 ⁻⁸ m ² s ⁻¹ MPa ⁻¹
$Q_{N_2}^s$	Activation energy of diffusive permeation of N ₂	55 kJ mol ⁻¹
$Q_{O_2}^s$	Activation energy of diffusive permeation of O ₂	57 kJ mol ⁻¹
D_{g0}	Constant for diffusion of CO ₂	1.8 × 10 ⁻⁶ m ² s ⁻¹
E_{CO_2}	Activation energy for diffusion of CO ₂	0.43 eV
C_{CO_2}	Concentration of CO ₂ in ice matrix at 1 MPa	7.6 × 10 ⁻⁹ MPa ⁻¹

Table 3. Parameters varying with time (in the last column the numerical values for $t = 0$ are given).

Symbol	Definition	Value
\bar{r}_a	Mean radius of microbubbles	0.07 mm ³
\bar{r}_n	Mean radius of normal bubbles	0.34 mm ³
\bar{Z}_{CO_2}	Molar fraction of CO ₂ in normal bubble	275 ppmv
$\bar{Z}_{N_2}/\bar{Z}_{O_2}$	N ₂ /O ₂ ratio in normal bubble	3.71
$\bar{Z}_{CO_2}^a$	Mole fraction of CO ₂ in microbubbles	275, 500, 1000 ppmv
$\bar{Z}_{N_2}^a/\bar{Z}_{O_2}^a$	N ₂ /O ₂ ratio in microbubbles	3.71
$\langle P_a \rangle$	Mean pressure in microbubbles	$\langle P_a \rangle = 2.5 \times 10^5$ Pa + $\Delta P_a t^b$
$\langle P_n \rangle$	Mean pressure in normal bubbles	$\langle P_n \rangle = 10^5$ Pa + $\Delta P_n t^b$

^aThe mean radii are calculated assuming an air content of 99.5 and 0.5 ml air STP per kg ice for normal bubbles and microbubbles, respectively and using the bubble densities and bubble pressures n_n , n_a and $\langle P_n \rangle$, $\langle P_a \rangle$. The value in the table is calculated for the mean temperature of DML.

^b ΔP_a is estimated to be 628, 748 and 5886 Pa yr⁻¹ for DML, South Pole and Law Dome respectively using the accumulation rates of the appropriate sites. ΔP_n is 680, 810 and 6376 Pa yr⁻¹ for DML, South Pole and Law Dome respectively. ΔP_n is calculated so that at 200 m below close-off, both mean bubble pressures agree. At lower depths $\langle P_n \rangle = \langle P_a \rangle$ is valid.

calculated according to Ikeda-Fukazawa et al. (2001), who calculated a fractionation between O₂ and N₂ in ice from the Dome Fuji ice core due to a diffusion of O₂ and N₂ from microbubbles to normal bubbles. We also include the equations for CO₂ with a higher concentration for microbubbles at enclosure depth and, different from their assumptions, we simplify the calculation by selecting a linear decreasing pressure difference against depth between microbubbles and normal bubbles which becomes equal at a depth of 200 m below close-off. The flux of the gas compound m from microbubbles to normal bubbles is:

$$q_m^{(a)} = 4\pi\gamma\bar{r}_a D_m \frac{M_m}{M_{air}} \left(\bar{Z}_m - \frac{\langle P_a \rangle}{\langle P_n \rangle} \bar{Z}_m^a \right). \quad (A1)$$

The flux from normal bubbles to microbubbles is:

$$q_m^{(n)} = -4\pi(1-\gamma)\bar{r}_n D_m \frac{M_m}{M_{air}} \left(\bar{Z}_m - \frac{\langle P_a \rangle}{\langle P_n \rangle} \bar{Z}_m^a \right),$$

where M_m is the molar mass of the gas compound m and M_{air} is the molar mass of an average air molecule.

The averaging parameter γ , ranges from 0 to 1 and is given by:

$$\frac{\gamma}{1-\gamma} = \frac{\bar{r}_n}{\bar{r}_a} \frac{n_n}{1-n_n}. \quad (A2)$$

The mass balance of the gas compound m is given by:

$$\frac{d(\langle P_a \rangle \bar{Z}_m^a \langle V_a \rangle)}{dt} = p_{iw} q_m^{(a)}, \quad (A3)$$

$$\frac{d(\langle P_n \rangle \bar{Z}_m \langle V_n \rangle)}{dt} = p_{iw} q_m^{(n)},$$

where $\langle V_a \rangle$ and $\langle V_n \rangle$ are the mean volumes of microbubbles and normal bubbles. $p_{iw} = z_g R \rho_i T / M_w$, where z_g is the non-ideal compressibility coefficient, ρ_i is the density of ice, and M_w is the molar mass of water.

The diffusive permeation coefficient of molecule m ($m = N_2, O_2$) in ice in equilibrium with air bubbles in m² s⁻¹ is given after

Ikeda-Fukazawa et al. (2001):

$$D_m = D_m^{s0} \langle P_n \rangle \exp\left(-\frac{Q_m^s}{RT}\right). \quad (\text{A4})$$

A permeation coefficient for CO₂ can be calculated with the values given in Hondoh (1996):

$$D_{\text{CO}_2} = C_{\text{CO}_2} \langle P_n \rangle D_{g0} \exp\left(-\frac{E_{\text{CO}_2}}{kT}\right). \quad (\text{A5})$$

The other symbols have the meanings and numerical values given in Tables 2 and 3. For example, the initial permeation coefficients D_{O_2} , D_{N_2} and D_{CO_2} calculated with (A4) and (A5) using the parameters given in the tables and the mean temperature of DML are 2.3×10^{-22} , 6.5×10^{-23} and $4.5 \times 10^{-25} \text{ m}^2 \text{ s}^{-1}$, respectively.

References

- Anklin, M., Barnola, J.-M., Schwander, J., Stauffer, B. and Raynaud, D. 1995. Processes affecting the CO₂ concentrations measured in Greenland ice. *Tellus* **47B**, 461–470.
- Barnola, J.-M. 1999. Status of the atmospheric CO₂ reconstruction from ice core analyses. *Tellus* **51B**, 151–155.
- Barnola, J.-M., Anklin, M., Porcheron, J., Raynaud, D., Schwander, J. et al. 1995. CO₂ evolution during the last millennium as recorded by Antarctic and Greenland ice. *Tellus* **47B**, 264–272.
- Barnola, J.-M., Pimienta, P., Raynaud, D. and Korotkevich, Y. S. 1991. CO₂-climate relationship as deduced from the Vostok ice core: a re-examination based on new measurements and on a re-evaluation of the air dating. *Tellus* **43B**, 83–90.
- Etheridge, D. M., Steele, L. P., Langenfelds, R. L., Francey, R. J., Barnola, J.-M. and Morgan, V. I. 1996. Natural and anthropogenic changes in atmospheric CO₂ over the last 1000 years from air in Antarctic ice and firn. *J. Geophys. Res.* **101**, 4115–4128.
- Fuhrer, K., Neftel, A., Anklin, M. and Maggi, V. 1993. Continuous measurements of hydrogen peroxide, formaldehyde, calcium and ammonium concentrations along the new GRIP ice core from Summit, Central Greenland. *Atmos. Environ.* **12**, 1873–1880.
- Gerber, S., Joos, F., Brügger, P., Stocker, T. F., Mann, M. E. et al. 2003. Constraining temperature variations over the last millennium by comparing simulated and observed atmospheric CO₂. *Clim. Dyn.* **20**, 281–299.
- Gillet, R. W., van Ommen, T. D., Jackson, A. V. and Ayers, G. P. 2000. Formaldehyde and peroxide concentrations in Law Dome (Antarctica) firn and ice cores. *J. Glaciol.* **46**, 15–19.
- Gow, A. J. 1969. On the rates of growth of grains and crystals in South Polar firn. *J. Glaciol.* **8**, 241–252.
- Güllük, T., Slemr, F. and Stauffer, B. 1998. Simultaneous measurements of CO₂, CH₄, and N₂O in air extracted by sublimation from Antarctica ice cores: confirmation of the data obtained using other extraction techniques. *J. Geophys. Res.* **103**, 15971–15978.
- Hondoh, T. 1996. Clathrate hydrates in polar ice sheets. *Proc. 2nd Int. Conf. on Natural Gas Hydrates* 131–138.
- Ikeda-Fukazawa, T., Hondoh, T., Fukumura, T., Fukazawa, H. and Mae, S. 2001. Variation in N₂/O₂ ratio of occluded air in Dome Fuji Antarctic ice. *J. Geophys. Res.* **106**, 17799–17810.
- Indermühle, A., Stocker, T. F., Joos, F., Fischer, H., Smith, H. J. et al. 1999. Holocene carbon-cycle dynamics based on CO₂ trapped in ice at Taylor Dome, Antarctica. *Nature* **398**, 121–126.
- Keeling, C. D. and Whorf, T. P. 2003. Atmospheric CO₂ records from sites in the SIO air sampling network. In: *Trends: A Compendium of Data on Global Change*. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, TN.
- Langway, C. C., Clausen, H. B. and Hammer, C. U. 1988. An inter-hemispheric volcanic time-marker in ice cores from Greenland and Antarctica. *Ann. Glaciol.* **10**, 102–108.
- Lipenkov, V. Y. 2000. Air bubbles and air-hydrate crystals in the Vostok ice core. In: *ICSI Workshop on Physics of Ice-core Records* (ed. T. Hondoh). Hokkaido University Press, Sapporo, 327–358.
- Martinier, P., Lipenkov, V. Y., Raynaud, D., Chappellaz, J., Barkov, N. I. et al. 1994. Air content paleo record in the Vostok ice core (Antarctica): a mixed record of climatic and glaciological parameters. *J. Geophys. Res.* **99**, 10565–10576.
- Neftel, A., Jacob, P. and Klockow, D. 1984. Measurements of hydrogen peroxide in polar ice samples. *Nature* **311**, 43–45.
- Neftel, A., Moor, E., Oeschger, H. and Stauffer, B. 1985. Evidence from polar ice cores for the increase in atmospheric CO₂ in the past two centuries. *Nature* **315**, 45–47.
- Siegenthaler, U., Friedli, H., Loetscher, H., Moor, E., Neftel, A. et al. 1988. Stable-isotope ratios and concentration of CO₂ in air from polar ice cores. *Ann. Glaciol.* **10**, 151–156.
- Smith, H. J., Wahlen, M., Mastroianni, D., Taylor, K. C. and Mayewski, P. A. 1997. The CO₂ concentration of air trapped in Greenland Ice Sheet Project 2 ice formed during periods of rapid climate change. *J. Geophys. Res.* **102**, 26577–26582.
- Sommer, B. 2000. *Klima-Informationen von chemischen Spurenstoffkonzentrationen in polaren Eisbohrkernen*. PhD Thesis. Physikalisches Institut, Universität Bern, Bern.
- Souney, J. M., Mayewsky, P. A., Goodwin, I. D., Meeker, L. D., Morgan, V. et al. 2002. A 700-year record of atmospheric circulation developed from the Law Dome ice core, East Antarctica. *J. Geophys. Res.* **107**, doi:10.1029/2002JD002104.
- Spahni, R., Schwander, J., Flückiger, J., Stauffer, B., Chappellaz, J. et al. 2003. The attenuation of fast atmospheric CH₄ variations recorded in polar ice cores. *Geophys. Res. Lett.* **30**, doi:10.1029/2003GL017093.
- Stauffer, B. 1981. Mechanismen des Lufteinschlusses in natürlichem Eis. *Z. Gletscherkunde Glazialgeol.* **17**, 17–56.
- Stauffer, B. and Berner, W. 1978. CO₂ in natural ice. *J. Glaciol.* **21**, 291–300.
- Stauffer, B., Flückiger, J., Monnin, E., Nakazawa, T. and Aoki, S. 2003. Discussion of the reliability of CO₂, CH₄ and N₂O records from polar ice cores. In: *International Symposium on the Dome Fuji Ice Core and Related Topics* (eds. H. Shoji and O. Watanabe). National Institute of Polar Research, Tokyo, 139–152.
- Stauffer, B., Flückiger, J., Monnin, E., Schwander, J., Barnola, J.-M. et al. 2002. Atmospheric CO₂, CH₄ and N₂O records over the past 60,000 years based on the comparison of different polar ice cores. *Ann. Glaciol.* **35**, 202–208.
- Tschumi, J. and Stauffer, B. 2000. Reconstructing past atmospheric CO₂ concentration based on ice-core analyses: open questions due to in situ production of CO₂ in the ice. *J. Glaciol.* **46**, 45–53.