



HAL
open science

**Geochemistry and tectonic setting of Matakaoa
Volcanics (East Coast Allochthon, New Zealand);
supra-subduction zone affinity, regional correlations and
origin.**

Dominique Cluzel, P.M. Black, Christian Picard, K.N. Nicholson

► **To cite this version:**

Dominique Cluzel, P.M. Black, Christian Picard, K.N. Nicholson. Geochemistry and tectonic setting of Matakaoa Volcanics (East Coast Allochthon, New Zealand); supra-subduction zone affinity, regional correlations and origin.. *Tectonics*, 2010, 29 (TC2013), 21p. 10.1029/2009TC002454 . insu-00435464

HAL Id: insu-00435464

<https://insu.hal.science/insu-00435464>

Submitted on 24 Nov 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Geochemistry and tectonic setting of Matakaoa Volcanics (East Coast Allochthon, New Zealand); supra-subduction zone affinity, regional correlations and origin.

D. Cluzel¹⁻², P.M. Black³, C. Picard⁴ and K.N. Nicholson⁵

¹ Pôle Pluri-disciplinaire de la Matière et de l'Environnement – EA 3325, The University of New Caledonia, BP R4, 98851 Nouméa cedex, New Caledonia.
dominique.cluzel@univ-nc.nc

² ISTO – UMR 6113, Universities of Orleans and Tours, France. BP 6759, 45067 Orleans cedex 2, France

³ Geology Department, School of Geography, Geology & Environmental Science, The University of Auckland, Private Bag 92019, Auckland, New Zealand.

⁴ Université de Franche-Comté, UMR 6249 Chrono-environnement - Laboratoire des Géosciences, 16 route de Gray, 25030 Besançon - France

⁵ Dept. of Geology Ball State University, 2000 W. University Ave. Muncie, IN 47306 USA.

Abstract

In north-eastern New Zealand, the Late Cretaceous to Eocene submarine volcanic rocks of the East Coast Allochthon referred to as Matakaoa Volcanics formed a dominantly mafic melange with minor abyssal sediments during the subduction of "oceanic" crust. Accreted and melanged pillow lavas and dolerites display the geochemical and isotopic features of tholeiites with some diversity referred to as mid-ocean ridge basalts, island-arc tholeiites and back-arc basin basalts. The association of supra-subduction tholeiitic magmas, abyssal sediments and polymetallic volcanogenic sulphide deposits is typical of many back-arc basins; therefore, a correlation with the alkaline and transitional basalts of the Hikurangi-Manihiki-Ontong Java Plateau may be ruled out. These petrological features and the fossil ages as well, are closely similar to those of the Tangihua Complex (Northland Allochthon) and allow Matakaoa and Tangihua mafic volcanic rocks, now separated by the modern Havre Trough, to be correlated within one single Late Cretaceous to Eocene

basin. The consideration of possible younger ages (Oligocene) is also discussed. The occurrence of Late Cretaceous-Eocene "oceanic" crust with supra-subduction zone affinities to the east of the Norfolk Ridge poses the problem of the intervening southwest-dipping subduction, the missing Late Cretaceous-Eocene arc and their relationship with Southeast Gondwana marginal break-up.

Key words: New Zealand, Southwest Pacific, Late Cretaceous-Eocene, back-arc basins, subduction, peel-off mélange.

Introduction

The evolution of the Southwest Pacific apparently underwent a marked change from the active margin setting that prevailed during the Early Permian to Early Cretaceous into the widespread and synchronous marginal basins opening that isolated large slices of the south-eastern Gondwana margin during the Late Cretaceous and Paleocene (ca. 90-55 Ma). These continental fragments now form New Zealand, New Caledonia and their undersea extensions, the Lord Howe Rise and Norfolk Ridge (Fig. 1). The end of the long-lived active margin activity in the New Zealand part of southeast Gondwana at ca. 100 Ma (Laird and Bradshaw, 2004) is probably due to the collision of the Hikurangi Plateau with the active margin at ca. 105 Ma (Hoernle et al., 2004; Mortimer et al., 2006; Davy et al., 2008) although some uncertainty remains about the precise timing of active margin extinction (Mortimer, 2004). In contrast, the driving mechanism of the marginal break-off is still debated and many hypotheses have been considered: i) marginal rifting triggered by a mantle plume (Bryan et al., 1997); ii) subduction of a spreading ridge (Mortimer et al., 2006); or, iii) marginal basin opening controlled by eastward slab roll-back and arc migration in a continuously converging environment (Lister and Etheridge, 1989; Veevers et al., 1991, Cluzel et al., 1999; Veevers, 2000a,b; Cluzel et al., 2001; Betts et al., 2002; Schellart et al., 2006). Thus, the nature of the magmas that have been generated in the Southwest Pacific during the Late Cretaceous and Paleocene is crucial to constrain its geodynamic evolution. However, most of the geologic evidence is now below sea level and the bulk of relevant information is restricted to the mafic allochthons of northern New Zealand and New Caledonia.

A feature of the mid-Tertiary geology of the northern and eastern parts of the North Island is the occurrence of extensive tracts of sedimentary and volcanic material of Cretaceous to Oligocene age that were thrust and/or slid onto the in-place Mesozoic basement and its early Tertiary sedimentary cover in the earliest Miocene. The oldest but uppermost part of the allochthons consists of disrupted pillow-lavas, dolerites, and associated abyssal sediments of Late Cretaceous to early Eocene age. These thrust complexes are known as the East Coast and Northland Allochthons (Fig. 2) and the mafic rocks contained within them as the Matakaoa Volcanics and the Tangihua Complex respectively. While the Tangihua Complex rocks have been subjected to detailed geochemical and geochronological studies in recent years (Nicholson et al., 2000 a, b; Whattam et al., 2005, 2006), the Matakaoa rocks are poorly known and except for the recent paper of Brathwaite et al (2008) there has been no geochemical data for them. In this paper, we describe the geochemical and isotopic features of Matakaoa Volcanics, and compare them to the mafic rocks of the Tangihua Complex, in order to establish correlations and discuss the tectonic conditions that were involved in their generation and emplacement.

Nature and geological setting of the Matakaoa Volcanics.

The East Coast Allochthon forms a 120 km long SSW trending lobe that is exposed over a large part of north-eastern North Island (Raukumara Peninsula; Fig. 3). It is formed of sliced sediments (Late Cretaceous-Oligocene) that are not dissimilar to the autochthonous basement on one hand, and mafic volcanic rocks and pelagic/abyssal sediments (the Matakaoa Volcanics) of undoubtedly “oceanic” origin on the other. The allochthon is covered by the overlying Miocene-Pliocene sequence and is almost certainly more extensive at depth (Mazengarb and Speden, 2000). The Matakaoa Volcanics which form two massifs (Mangaroa and Pukemaru) separated by a Neogene graben (Fig. 3), are considered to be the oldest but uppermost and hindmost parts of the East Coast Allochthon that was emplaced in the earliest Miocene.

The bulk of Matakaoa Volcanics is formed of sliced massive and pillowed basalt associated with hyaloclastite; pelagic or abyssal sediments, and many small-scale mafic hypabyssal bodies. Volcanic breccia of basaltic to andesitic composition

with angular gabbro boulders and tachylite are not uncommon and suggest explosive eruption, or almost synchronous tectonics and/or submarine collapse. No evidence for tidal zone or sub-aerial evolution such as rounded pebbles or shallow water fossils may be reported. Basalts may be either fresh; or alternatively, affected by low temperature alteration (ocean floor metamorphism ?). The general attitude is upright, generally top-to-the-north but in detail, there is no continuity and slices are commonly separated by strike-slip faults. Near the boundary with Pukemaru Massif (Fig. 3), the overlying Late Miocene marl and limestone locally pass into monogenetic matrix-supported cobble conglomerates. The occurrence of cm to dm scale angular blocks of basalt included within a marine limestone matrix suggests the existence of basalt cliffs at the time of sediment deposition (rock fall deposits at the base of Miocene fault scarps?) rather than a tidal or sub-aerial environment. In addition, in the vicinity of the two massifs, cobbles of Matakaoa Volcanics are enclosed in Miocene and in many other late Tertiary sediments throughout Raukumara Peninsula (e.g. Black, 1980; Kenny, 1984).

While the Matakaoa Volcanics in the Mangaraoa Massif have good coastal exposures, the inland Pukemaru Massif is poorly exposed and thus much less well known; it seems to be mainly composed of massive basalt, scarce intrusive rocks and pillow lavas with no, or only a few, associated sediments.

Sediments are abundant in the Matakaoa Volcanics with lenses and slivers having an apparent thickness of a few tens of cm to 100s of metres (max about 800 m). There is a considerable variety of sedimentary lithologies including shale and mudstone (green, grey or purple), red and green chert, cream siliceous volcanogenic siltstone and pink micrite. All these lithologies are consistent with pelagic or abyssal environments. The pillow lavas rarely have interpillow material but when it does occur it is usually limestone rather than chert. Assessing the time relationship of volcanics and associated sediments is not always possible because of the high degree of tectonic disruption of most outcrops; however, in a few less deformed locations, clear sedimentary contacts may be observed between submarine volcanic rocks and deep water sediments. Pink pelagic limestone often contain glass shards and tiny basalt clasts; in addition, basalt intrusion within unconsolidated and water saturated sediments have been observed by previous authors (e.g. Gifford, 1970; Strong 1976, 1980) providing robust evidence for syn-depositional relationship. Some of the sediments are fossiliferous, and macrofossils (*Inoceramus*) have been recorded on the

west side of Cape Runaway, together with a radiolarian and foraminifera micro-fauna that indicate a Late Cretaceous age (Strong, 1976). On the western side of Lottin Point, radiolarian and foraminifera assemblages in sedimentary inliers also indicate a Late Cretaceous age for the IAT-like basalts (see below) that are closely associated with them. However sediments between these two localities contain Paleocene to mid-Eocene foraminifera and radiolarian faunas; thus, MORB-like basalts that rest below Paleocene sediments to the south-east of Cape Runaway have most probably the same age (Fig. 3). Foraminifera from sedimentary float in the Pukemaru Massif indicate that rocks in this massif also have Paleocene - Eocene ages (Strong, 1980).

The coastal succession of Mangaroa has been affected by km-scale gentle steeply plunging folds associated with semi-brittle wrench faulting (Spörli and Aita, 1994). Sedimentary inliers are often severely disrupted, boudinaged and display melange features. It is worth noting that melange only contains Late Cretaceous-Eocene pelagic or bathyal sediments and no terrigenous or younger sediments have been hitherto reported.

The Matakaoa Volcanics show a significant range of igneous material which includes basaltic hyaloclastites, tuffs, massive and pillowed lavas, and doleritic intrusives. Lava and dolerite compositions are mainly basaltic. The lavas are mainly non-vesicular dominantly augite-pyroxene basalts with little interpillow material and pillow breccia complexes. There are some substantial exposures in the middle of the massif, on the west side of Lottin Point and in the Matakaoa Point area, composed of resedimented hyaloclastite (tachylite), and basalt – dolerite – gabbro breccias which also contain tachylite blocks. The breccias are locally hydrothermally altered and contain abundant prehnite (and less commonly pumpellyite). Prehnite also appears in early veins (sometimes with quartz) crosscut by zeolite veins with laumontite, or analcime + natrolite + stilbite, the last veins in all cases containing calcite. Dykes and gabbros cut the pillow complex sequences and the breccias. Olivine is not common in the basalts where it usually occurs as chloritised relics; however some fresh olivine-rich rocks, called picrites by earlier workers (Gifford, 1970; Rutherford, 1980; Pirajno, 1980), occur together with olivine gabbros and “teschenites” (actually amphibole-bearing sub-alkaline rocks) in a fault bounded block near Lottin Point. Dolerite dykes are common cutting all the volcanic lithologies including the breccias and the picrite – gabbro – “teschenite” complexes (Pirajno, 1980).

Geochemistry

A collection of 39 samples of predominately volcanic and minor intrusive rocks from the Matakaoa Volcanics have been analysed for major oxides, trace and rare earth elements. This has enabled a characterization and recognition of magma sources not possible in previous studies. The locality and chemical affinity details are given for representative samples in Table 1 and their major, minor and trace element data in Table 2.

Sample preparation: External surfaces of fresh rocks were removed by splitting and the samples split into chips which were then ground to a powder in a tungsten carbide ring grinder. H_2O^+ , H_2O^- were determined by gravimetry, the remaining major oxides and trace elements were first determined by XRF (after Norrish & Chappel, 1977) at the University of Auckland, New Zealand. The precision (2σ) of the major oxide determinations is generally less than 2%, and less than 5% for the trace elements at ten times the detection limit of 1-2 ppm. HFS and rare earth element abundances were secondly determined by inductively coupled plasma mass spectroscopy (ICP-MS) at the Laboratoire de Géodynamique des Chaines Alpines, Université Joseph Fourier, Grenoble, France (UMR 5025) using the method of Barrat et al. (1996). The precision (2σ) for ICP-MS analyses varies from 2-10% for the elements analysed as determined using the standards BHVO and Br24 (Barrat et al., 1996). Nd and Sr isotopic ratio were determined using mass spectrometry in Toulouse University (France) after separations on columns in LGCA – Grenoble University.

In the following paragraphs, although the difference is nowhere meaningful, (X/Y)_{nC} represent elemental ratios normalised to the chondrite, and (X/Y)_{nM} elemental ratios normalised to the average MORB. The following abbreviations will be used throughout the text: MORB: Mid-Oceanic Ridge Basalt; N-MORB: Normal MORB; T-MORB: Transitional MORB; E-MORB: Enriched MORB; IAT: Island-Arc Tholeiite; BABB: Back-arc Basin Basalt; CAB: calc-alkali Basalt; OIB: Ocean Island Basalt; LILE: Large Ion Lithophile Element; HFSE: High Field Strength Element; LREE: Light Rare Earth Elements; HREE: Heavy Rare Earth Elements.

The presence of alteration is normally reflected in the abundances of the more mobile elements and LOI (loss of ignition), and only two samples present LOI > 4% (LOI = 4.43 and 7.45); so, 39 samples with total volatile content < 4wt% were chosen

for the study of Matakaoa volcanic rocks (Table 1). With regard to major elements composition, lavas and dolerites are mainly basaltic ($\text{SiO}_2 = 47\text{-}53$ wt %) with medium to low MgO contents (7.5-1.7 wt%), and some rare silica-rich rocks ($\text{SiO}_2 = 55.8\text{-}60.3$ wt%, $\text{MgO} = 4.5\text{-}3.3$ wt%) occur in the north of the Matakaoa Massif. However, an alkali-silica (not presented) or alkali-MgO plot (see annex I) shows a prominent scatter in the total alkali content that contrasts with the relatively narrow field in which Matakaoa Volcanics plot on classification diagrams based upon "immobile" trace elements, indicating that the effects of alteration on mobile elements have to be considered and that rock classification should not be established upon them. On the Zr/TiO_2 vs. Nb/Y classification diagram (Winchester and Floyd, 1977) which is used instead of the total alkali vs. silica diagram in the case of metamorphic or slightly altered rocks, the Matakaoa Volcanics plot within the field of andesitic basalts with only some scatter of the "alkalinity ratio" represented by Nb/Y (Fig 4). In addition, the Zr/TiO_2 ratio is not correlated with SiO_2 variation; therefore, high SiO_2 contents are likely to be a secondary feature. It is worth noting that the typology established on the basis of REE contents and Nb-Ta depletion (see below) is fully consistent with the variation of the Nb/Y ratio. Similarly, on the Th/Yb vs Ta/Yb diagram (Pearce, 1983) (Fig 5), the Matakaoa Volcanics have similar Th/Yb ratios consistent with the high degree of melting of a moderately depleted mantle source, but may be distinguished upon the variation of the Ta/Yb ratio (equivalent to Nb/Y) into uncontaminated (MORB-like), poorly contaminated (BABB-like) and contaminated (IAT-like) basalt types. Two samples which have $\text{Ta}/\text{Yb} > 0.1$ will be thereafter referred to as "transitional basalt" (TB in figure captions).

The Matakaoa basalts and dolerites are all strongly depleted in LREE ($(\text{La}/\text{Sm})_{\text{nC}} = 0.48$ to 0.79 ; $(\text{La}/\text{Yb})_{\text{nC}} = 0.47\text{-}1.0$) (Fig. 6a), a feature that suggests a depleted shallow mantle source. However, they show variable LILE contents and also a variable Nb-Ta depletion ($(\text{Nb}/\text{La})_{\text{nM}} = 0.37$ to 0.97 ; $(\text{Ta}/\text{La})_{\text{nM}} = 0.45$ to 1.19) (Fig. 6b). These features suggest the coexistence of lavas types similar to mid-oceanic ridge basalt (MORB-like), back-arc basin basalts (BABB-like) and island arc tholeiites (IAT-like) that can be distinguished as follows:

- MORB-like volcanics ($\text{SiO}_2 = 47.7\text{-}50.6$ wt%, $\text{MgO} = 4.98\text{-}7.55$ wt%) are depleted in LREE ($(\text{La}/\text{Sm})_{\text{nC}} = 0.55\text{-}0.79$, $(\text{La}/\text{Yb})_{\text{nC}} = 0.68\text{-}1.0$) with relatively high TiO_2 contents (1.75-2.73 wt%). They show moderate LILE enrichment, almost no Nb-Ta

depletion ($(\text{Nb/La})_{\text{NM}} = 0.73\text{-}0.97$ and $(\text{Ta/La})_{\text{NM}} = 0.81\text{-}1.19$), and are characterised by high Nb/Th ratios (12.8 -17).

- BABB-like volcanics ($\text{SiO}_2 = 48.5\text{-}50.2$ wt%, $\text{MgO} = 6.26\text{-}9.06$ wt%) are also strongly depleted in LREE ($(\text{La/Sm})_{\text{NC}} = 0.55\text{-}0.75$; $(\text{La/Yb})_{\text{NC}} = 0.62\text{-}0.89$). They show lower TiO_2 contents (1.35-1.79 wt%), intermediate Nb-Ta depletion ($(\text{Nb/La})_{\text{NM}} = 0.56\text{-}0.72$; $(\text{Ta/La})_{\text{NM}} = 0.70\text{-}0.93$) and Nb/Th ratio 9 to 14.
- IAT-like basalts have medium TiO_2 (1.05-1.73 wt%) and MgO (3.31-7.08 wt%) contents, and variable SiO_2 (48.2 to 60.3 wt%). They are all enriched in LILE and show LREE and Nb-Ta depletion ($(\text{La/Sm})_{\text{NC}} = 0.48\text{-}0.72$; $(\text{La/Yb})_{\text{NC}} = 0.47\text{-}0.84$; $(\text{Nb/La})_{\text{NM}} = 0.37\text{-}0.62$; $(\text{Ta/La})_{\text{NM}} = 0.44\text{-}0.74$). They typically display the lowest Nb/Th ratios (4.7 to 9.5).
- andesitic basalts transitional towards alkali basalt (two samples only) hereafter referred to as TB (transitional basalt) typically display higher Nb/Y ratios, high TiO_2 and much higher LREE contents compared with the rest of Matakaoa Volcanics.

The three main geochemical types are petrographically very similar and cannot be distinguished in the field; however, the IAT are usually porphyritic (augite and plagioclase phenocrysts) and the rare orthopyroxene recorded in the Matakaoa volcanics always occurs in the IAT. MORB-like basalts are dominated modally by clinopyroxene and are often non-porphyritic; and the BABB occasionally contain pink - brown Ti-rich pyroxene and a reddish brown hornblende. These chemically distinct rock types are found in many locations in the western part of Matakaoa massif; in contrast, IAT are the major volcanic type to the east of the massif where they occur as pillow lavas. Elsewhere in the massif, rocks of IAT affinity are dolerites and diorites which suggest that at least some of the IAT are younger than the MORB and BABB; however there is no sensible chemical difference between IAT lavas and sub-volcanic intrusions.

Magma sources and geodynamic setting

One of the most obvious features of the Matakaoa Volcanics is the continuous range of composition between IAT, MORB and BABB end-members which is very apparent on the Ta/Yb vs. Th/Yb diagram of Pearce (1983, Fig. 5), and on many other "discriminant" diagrams (not represented) this feature is also apparent on trace and

rare earth element spidergrams (Fig 6). A common feature of these rocks is the depletion in LREE (Fig 6a) which is consistent with low pressure melting of depleted mantle. In contrast, the trace element abundance patterns differ with a relative depletion of Nb and Ta in BABB and lower bulk REE content (Fig 6b). Distinguishing back-arc from mid-ocean ridge basalts is possible because the depletion of Nb and Ta in BABB reflects formation from a subduction-modified mantle that contains Nb and Ta receptors such as ilmenite or pargasitic amphibole in the refractory phase. However, in the case of Matakaoa Volcanics, MORB-like lavas also show a small Nb depletion. All the rock types are variably enriched in LILE which may result either from the input of fluids derived from slab dehydration, selective extraction during hydrous melting (Langmuir et al., 2006); or, alternatively, from elemental mobility during low grade metamorphism and/or hydrothermal alteration. The occurrence of IAT is also generally associated with juvenile intra-oceanic arcs and/or oceanic basins that form in connexion with metasomatised supra-subduction mantle (Keller et al., 2008; Metcalf and Shervais, 2008). It is worth noting that Nb-Ta depletion decreases when the bulk REE content increases. The bulk REE content of basalt may be related either to the degree of partial melting; or, alternatively to fractional crystallisation; as the three rock types show no evidence for prominent differentiation, it appears that Nb-Ta depletion is correlated with a higher melting degree, which may be due to higher water content of the source, as is the case of metasomatised mantle. A close association of MORB, BABB and IAT-like basalts is thus a product of low pressure melting of a variably metasomatised mantle source in a supra-subduction environment (Marsh et al., 1980; Wood et al., 1981; Perfit et al., 1987; Vallier et al., 1991; Jenner et al., 1991).

The Nb-Ta depletion which is apparent on REE and trace elements spiderdiagrams is uncorrelated with an increase of Pb and Sr contents which are generally associated with volcanic-arc magmas (Fig. 7); therefore, a prominent input of subduction or continental crust-derived material is unlikely as emphasised by Nd and Sr isotopic ratios (see below).

ϵ_{Nd} isotopic ratios of the Matakaoa volcanic rocks remain within a very narrow range of variation ($+7.06 < \epsilon_{\text{Nd}} < +8.12$) and plot within the depleted mantle (MORB) field (Fig. 8). In contrast, $(^{87}\text{Sr}/^{86}\text{Sr})_i$ ratios display a wider range (0.7021 to

0.7044). Moderately variable ($^{87}\text{Sr}/^{86}\text{Sr}$)_i and invariable $\epsilon_{\text{Nd}i}$ suggest post-magmatic water-rock interaction (ocean floor metamorphism or hydrothermal alteration) rather than contamination by subducted sediments, continental crust, or subduction fluids. $\epsilon_{\text{Nd}i}$ values fit with a moderately depleted mantle source for these volcanic rocks which is consistent with either a juvenile intra-oceanic arc, or a back-arc environment.

Distinguishing tectonic environments on the basis of geochemistry alone is not possible because similar magma forming processes may actually occur in different settings. In the present case, a similar association could appear in an incipient arc, a back-arc or in a fore-arc setting as well; however, a close association of magmas formed from variably metasomatised sources and abyssal sediments is a feature of basins formed in an extensional setting in a supra-subduction zone environment, e.g. a back-arc basin environment (Keller et al., 2008; Metcalf and Shervais, 2008). This interpretation is compatible with that of Brathwaite et al (2008) who additionally considered the occurrence of polymetallic volcanogenic massive sulphide (VMS-type) mineralisation closely associated with Matakaoa Volcanics.

Comparison of Matakaoa and Tangihua mafic rocks

The Matakaoa and Tangihua Volcanics display significant petrographic and geochemical similarities. The comparison below is based on the dataset already published by Mortimer et al. (1998), and Nicholson et al. (1999, 2000a, b). It is worth noting that both the Matakaoa and Tangihua volcanic rocks contain MORB, BABB and IAT-like rocks with similar REE patterns and Ta-Nb depletion (Fig 9 and Fig 10) which often coexist at the same locality. The Tangihua Complex rocks similarly display rather variable Sr isotopic ratio ($(^{87}\text{Sr}/^{86}\text{Sr})_i = 0.7027$ to 0.7036 ; Table 3 - Fig. 9) within the same range of variation, and positive and constant ϵ_{Nd} values ($+7.8 < \epsilon_{\text{Nd}i} < +8.3$). In contrast with Tangihua Volcanics, there is apparently in the Matakaoa Volcanics no occurrence of mafic rocks with high compatible elements contents (Mg, Cr, V), and referred to as "boninite-like" by Whattam et al. (2006).

Both mafic complexes are very similar with respect to the age of enclosed pelagic sediments (Campanian to mid-Eocene), timing of tectonic emplacement, lithology, mineralogy, and geochemistry. The very close similarity suggests that they have a common origin and were once parts of the same slice of Late Cretaceous to Early Eocene “ocean” floor, the bulk of which has been consumed along one of the convergent plate boundaries in the northern New Zealand region in the mid Tertiary. The geochemical characteristics of the Matakaoa Volcanics, and the features of associated sediments and polymetallic mineralisation indicate that they have formed in an extensional back-arc basin, which is similar to the back-arc setting proposed for the Tangihua Complex in the Northland Allochthon (Thompson et al., 1997; Nicholson, 2002a, b).

Comparison of Northland and East Coast allochthons

Stonely (1968) was the first to recognise that the Cretaceous to Tertiary beds of the East coast of the North Island were an allochthon emplaced in a south-westerly direction by gravity sliding onto a structurally simple Cretaceous autochthon and discordantly overlain by upper Tertiary sediments. In this seminal paper he recognised at least 20 slices of coherent sediments, each emplaced before the next (older) slice was slid on top of it and a trend for the oldest, highest thrust slices in the sequence to be the most deformed. While there has been some refinement of Stonely’s concept as the thrust sequences have been examined in detail and particularly traced to the north of the Raukumara Peninsula, the basic concepts are now accepted. The East Coast Allochthon is now considered to be composed of 5 groups of thrusts, in which the individual slices are closely related. Many sediments in the thrust sequence have lateral equivalents to the Southwest and represent deposition at greater depths than equivalents in the in-place sequence (Mazengarb and Speden, 2000) and there is some evidence for eastward younging of the deformed sediment / cover basin unconformity (Kenny, 1984). Rait (2000a) estimates that Cretaceous to Oligocene rocks in the East Coast Allochthon have been moved possibly 300 kilometres from their original site of deposition. The Matakaoa Volcanics constitute the sheet at the top and back of the thrust pile and are thus believed to be the oldest slice in the sequence (Mazengarb and Speden, 2000).

The Northland Allochthon extends at least 300 km along the centre and west of the Northland Peninsula (Fig. 2) reaching estimated thicknesses of up to 4000m in central Northland. The sedimentary units in the Northland Allochthon are typically highly deformed compared with the underlying in situ sequence, and melange and broken formation are widespread. At least six thrust slices have been recognised and the Tangihua Volcanics occur in the structurally highest sheet (Isaac et al., 1994). As is the case in the East Coast Allochthon, the higher level thrust slices in the Northland Allochthon are more deformed and show evidence of reimbrication by continued thrusting, particularly on the sub-ophiolite thrust (Rait, 2000b). From north to south in the Northland Allochthon the slices become more disrupted and in its southern part it is composed of small masses as well as dismembered blocks of Tangihua Volcanics and slices of sediment intercalated with Miocene sediments.

The timing and direction of emplacement, and the age of the sediments in the two allochthons are similar (see Hollis and Hanson, 1991). The thrust sequences in the East Coast Allochthon appear to be more coherent than those in the Northland Allochthon. The sediments in the two units are all marine but there are differences in the nature of the sedimentary material in the two allochthons. Allochthonous sediments are in general more pelagic than those of the same age in the autochthon; however, in the East Coast region sediments in the allochthon are sometimes so similar to those in the autochthon that they can be distinguished only on differences in structural style (Mazengarb and Speden, 2000). In Northland, the sediments in the allochthon have clear facies (environment of deposition) differences compared with those in the autochthon. In spite of this and the fact that the East Coast and Northland Allochthons are over 300 km apart, because of their strong similarities they have frequently been correlated and it is assumed that the same regional tectonic event has been responsible for their emplacement. Both allochthons have been emplaced from a north-easterly direction and are displaced about 300 km (Rait, 2000a). The geochemical features presented in this paper confirm and reinforce the assessments of previous authors.

Both the Tangihua Complex and Matakaoa Volcanics contain deformation that was sustained prior to emplacement. In the case of the Tangihua Volcanics there is additional evidence for delamination and disruption, although without substantial displacement, and that they were then remobilised at about 30 Ma (Nicholson &

Black, 2004). There is no evidence of a metamorphosed sole either in the Matakaoa Volcanics or in the Tangihuas; this indicates that no hot, e.g. young lithosphere, was directly involved in the obduction process (see below). We presume that the Matakaoa Volcanics like the Tangihuas have had a complex post-formational but pre-emplacement history that may be related to pre-obduction tectonic accretion.

Hence, the term "ophiolite" which has been sometimes used for the Matakaoa Volcanics (and Tangihua Complex as well) should be avoided because these rocks do not represent obducted parts of a coherent oceanic lithosphere, but rather a mafic melange probably formed by tectonically sliced parts of an upper "oceanic" crust and the associated bathyal sediments. The lack of terrigenous deposits is strong evidence for the melange to have occurred away from any continental source well before the final thrusting over Miocene rocks. All these lithologies may have been scraped off the down going plate, accreted in a fore-arc setting, and finally thrust "en bloc" onto the continental margin during the final closure of the basin.

Age problems

In recent papers (Whattam et al., 2005; 2006, 2008), the correlation of the Northland "Ophiolite" with the Late Cretaceous-Paleogene has been questioned, mainly on the basis of new Ar-Ar ages. Dating of mafic rocks by the Ar-Ar whole rock method provides evidence for consistent Oligocene apparent ages (25-30 Ma) for some parts of the mafic allochthon. In addition, Oligocene U-Pb zircon ages (ca. 30 Ma) have been obtained from one felsic dyke (Whattam et al., 2005) and a gabbro (Whattam et al., 2006). Thus, the bulk of the Northland Complex was reassigned to the Oligocene and obduction was thought to have followed oceanic accretion very closely (Whattam et al., 2008). However, this interpretation is problematical, especially because no large Oligocene basin is known to have existed to the North of New Zealand and also because obduction did not involve hot lithosphere (see above). In addition, Oligocene ages are inconsistent with the available stratigraphic data, as no fossils younger than Early Eocene have been hitherto found within the mafic terranes (cf. section above). It has been pointed out by Nicholson et al. (2007) that some of these new dates have been obtained from basalts that contain much older interpillow sediments already dated by radiolarian fossils. Large volumes of basalt are affected to various degrees by low to medium grade (ocean floor?) metamorphism

(Pirajno, 1980; Black, 1989; Nicholson & Black, 2004) resulting in some elemental mobility and radiogenic argon input, giving unreliable Ar-Ar apparent ages at ca. 300 Ma (Nicholson, 1999). Such older apparent ages are likely due to the massive absorption of hydrous fluids containing radiogenic Ar during water-rock interaction; in contrast, much younger ages (ca. 25-30 Ma) may be related to argon loss during a subsequent thermal event (e.g. intrusion of Oligocene dolerite and/or to the end of hydrothermal circulation) (Nicholson, 2007).

If we consider some of the Oligocene Ar-Ar ages reliable and unaffected by a possible post-eruption reset, the inconsistency between paleontological and radiochronological ages may be solved only if the Late Cretaceous-Paleogene basalt formed the basement through and upon which Oligocene basalts erupted. It is worth noting that the Oligocene gabbro contains reworked Early Cretaceous zircons (Whattam et al., 2006), a feature which is not consistent with the "oceanic" and uncontaminated character of Tangihua volcanics. Zircons would not travel from the subduction zone into supra-subduction melts unless the zircon-bearing slab rocks were melting. But this is not such a common occurrence and it is obviously not the case as adakite-like and other slab-melt rocks are not found in either the Tangihua or in the Matakaoa Volcanics. The Oligocene (30 Ma) "plagiogranite" and gabbro dykes dated by U-Pb zircon dating (Whattam et al., 2005; 2006) may thus be minor fore-arc magmas that have been emplaced through an older basement. This interpretation suggests the existence to the north of the North Island of an Oligocene volcanic arc partly buried below Miocene to Recent volcanic rocks and sediments. This Oligocene volcanic arc, remnants of which had been first identified in offshore dredges (Mortimer et al., 2007), has been recently explored in more detail (Herzer et al., 2009), and it could be coeval with and extend northward into the Three Kings Ridge and farther north into the older Loyalty volcanic-arc.

Regional correlation: Late Cretaceous back-arc basins of the Southwest Pacific

Based upon bulk geochemical similarities, it has been proposed earlier that Matakaoa and Tangihua basalts might be correlated with Hikurangi Plateau rocks (Hayward et al., 1989; Mortimer and Parkinson, 1996); however, this correlation does

not account for the critical differences between them as evidenced by modern geochemical data (Mortimer et al., 1998, this study). Hikurangi Plateau basement rocks formed between ca. 120-100 Ma and the seamounts that overlie it between ca. 99-86 Ma (Hoernle et al., 2005). The basement E-MORB display isotopic signatures ($(^{87}\text{Sr}/^{86}\text{Sr})_i = 0.70361-0.70374$; $\epsilon\text{Nd}_i = 5.7-6.2$) which are consistent with magma generation in a fertile asthenospheric source without any evidence for supra-subduction features (Mortimer and Parkinson, 1996). Thus, the Hikurangi basalts are much older (ca. 120-90 Ma) than the Matakaoa Volcanics (ca. 95-50 Ma), have been generated by contrasting mantle sources (Fig. 8) and owing to their present location on the Pacific Plate, have most probably erupted farther east. Thus the correlation is not supported by our new geochemical data and should be ruled out.

In other places of the Southwest Pacific, the occurrence of Late Cretaceous-Paleocene back-arc basins to the east or north-east of the Norfolk Ridge has been firmly established on geochemical (Eissen et al., 1998; Cluzel et al., 2001), paleontologic (Aitchison et al., 1995; Cluzel et al., 2001), paleomagnetic (Ali and Aitchison, 2000), and geophysical bases (Mauffret et al., 2001; Bernardel et al., 2002). The existence of such fragments is corroborated by the occurrence of very deep (ca. 4,500-5,000 m; i.e. old) oceanic crust forming parts of the South Norfolk Basin (Mauffret et al., 2001). This is evidence for Late Cretaceous "oceanic" basins having existed to the east of the Norfolk Ridge. These basins, that probably extended 500 km to the north and north-east of New Caledonia (Ali and Aitchison, 2000; Schellart et al., 2006), have almost completely disappeared as the result of north-eastward subduction and slab roll-back during the Eocene and Oligocene. This subduction produced the Loyalty, Three Kings (in parts) and possibly Northland Plateau volcanic-arcs.

In New Caledonia, Late Cretaceous to Paleocene mafic oceanic rocks form a 400 km long complex, referred to as the Poya Terrane (Cluzel et al., 1994), underneath the ultramafic ophiolite. The Poya Terrane is a peel-off melange, i.e. a matrix-free juxtaposition of fragments and slivers of the upper oceanic crust, characteristically lacking any terrigenous component, which was formed in the Late Eocene in an intra-oceanic fore-arc setting. It is composed of generally upright km-scale slices of pillowed and massive basalt associated with thin abyssal argillite/radiolarite and rare pelagic limestone. The accreted material originated in a

marginal basin located to the east of the Norfolk Ridge and referred to as the South Loyalty Basin (Cluzel et al., 2001). The Poya Terrane rocks are dominantly undepleted MORB (E-MORB), rare BABB and a few younger OIB volcanics that have been interpreted to represent a marginal basin in which magma was generated in conditions similar to that of the North Fiji Basin (see Eissen et al., 1996; Cluzel et al., 2001). In such back-arc basins, the supra-subduction signature is generally weak, and only magmas formed from a metasomatised upper mantle are likely to display BABB (i.e. supra-subduction) features.

Thus, there are significant differences between the Tangihua/Matakaoa back-arc basin that opened to the north of New Zealand and the South Loyalty Basin that formed to the east and northeast of New Caledonia (Nicholson et al., 2000b). The synchronous opening of two dissimilar back-arc basins to the east of the Norfolk Ridge during the Late Cretaceous suggests the existence of a right-lateral transform fault zone that separated the Tangihua/Matakaoa and South Loyalty basins. Following the kinematic model for the opening of the Southeast Gondwana marginal basins (Gaina et al., 1998; Sutherland, 1999; Hall, 2002; Schellart et al., 2006), we suggest that the transform fault extended north-eastward, was inherited from the break-up fracture set (see model below), and allowed a differential opening of the two marginal basins and the overall north-eastward migration of the volcanic arc (Fig. 11). The Northland volcanic-arc/back-arc system was probably limited to the east by another pre-existing transform fault while the northern boundary of the system and its connexions with the Solomon-Papua arc-basins system are still unclear.

South Gondwana break-off: arc- or non arc-related ?

It is generally considered that the end of the South Gondwana active margin activity in New Zealand that occurred at ca. 100-105 Ma (Mortimer, 2004) may be related to the subduction of a spreading ridge (Bradshaw, 1989); and/or the attempted subduction of the Hikurangi oceanic plateau on the Southeast Gondwana active margin (Mortimer and Parkinson, 1996; Mortimer et al., 2006; Davy et al., 2008). Thus, the opening of the Southwest Pacific marginal basins at ca. 90 Ma (Tasman Sea, New Caledonia Basin, etc.) has been interpreted in terms of a Large Igneous Province (Bryan et al., 1997), a feature generally triggered by a mantle plume.

Evidence for Early to mid-Cretaceous (120-110 Ma) marginal rifting is widespread in Southeast Australia (Bryan et al., 1997) and in the South Island of New Zealand (Tulloch and Kimbrough, 1989; Spell et al., 2000; Deckert et al., 2002). Logically, the rift activity predates the marginal basin opening at 90 Ma by some 30 Ma; however, extensive intraplate volcanism and development of Otway, Gippsland and other intra-continental basins was almost synchronous with the end of subduction over a wide area. Intrusion of large amounts of adakitic plutons in the 120-105 Ma interval indicates the melting of an eclogitized mafic source (Muir et al., 1995; Mortimer et al., 1999). Therefore, it may be postulated that the slab rollback that followed the end of subduction triggered a prominent eastward flow of the asthenospheric mantle that heated the base of the lithosphere and generated the adakitic magmas by partial melting of previously underplated mafic rocks. It was also responsible for the large-scale boudinage and extensional break-off of the southeast Gondwana margin (Fig. 12).

In Australia and in New Zealand as well, no post-Early Cretaceous active margin magmatism has hitherto been evidenced and this has been taken as an evidence for a purely extensional tectonic setting. However, the occurrence of 100 Ma old detrital zircons in the volcanoclastic greywackes of "basement" terranes of New Zealand (Cawood et al., 1999), and New Caledonia (Adams and Cluzel, in press; Cluzel, et al. submitted) indicates the persistence of subduction until the upper Early Cretaceous (Albian), at least in the northern part of the system, after a short period of quiescence that may be related to the "collision" of Hikurangi Plateau. In addition, younger Late Cretaceous (ca. 90-80 Ma) high-K calc-alkaline and IAT-like volcanic activity is reported in New Caledonia (Noumea and Diahot regions respectively) (Black, 1995; Picard, 1995), and also in New Zealand (Mount Camel Terrane, Northland and the Three Kings Islands) (Nicholson and Black, 2004). Therefore, we suggest that the eastward push caused by the stretching of the Southeast Gondwana margin reactivated the west-dipping subduction along the eastern end of the system, except in the south where it was blocked by the Hikurangi plateau. Once this subduction was reactivated, the slab roll-back of the old, dense Pacific plate generated two new back-arc basins, the South Loyalty basin to the north, and the Matakaoa/Tangihua basin to the south, after which they evolved independently (Fig. 11).

A model for the obduction of Northland and East Coast Allochthons.

For a long time it has been commonly accepted that the Northland Allochthon and its East Coast correlative were obducted onto the North Island continental margin as a result of south-dipping subduction (Ballance and Spörli, 1979; Parrot and Dugas, 1980; Brothers and Delaloye, 1982; Spörli, 1989; Malpas et al., 1992; Nicholson et al., 2000a; Herzer et al., 2000; Whattam et al., 2005); however, this view is now questioned and an increasing number of authors consider the possibility of north or north-east dipping subduction having occurred prior to obduction. (Cluzel et al., 1999; Bernardel et al., 2002; Sdrolias et al., 2003; Crawford et al., 2003; Bradshaw, 2004; Schellart et al., 2005; Whattam et al., 2006, 2008).

The model of south dipping subduction suggests delamination and accretion of slivers of oceanic upper crust onto the former passive margin (flaking). However, it is difficult to reconcile this model, that would have resulted in a relatively narrow, fan-shaped, mainly north-verging accretion complex, with the shortening that results from computing the throws of all parts of the allochthon (Rait, 2000a) and paleomagnetic data (Cassidy, 1993; Whattam et al., 2005), all of which indicate a southward relative motion of ca. 300 km. Advocating southward gravitational sliding apparently solves this inconsistency but requires a large pre-existing slope which is not substantiated on paleogeographic grounds.

Sea-floor spreading data for the Late Eocene – Early Miocene Pacific – Australia plate motion indicates that the Pacific - Australian plate boundary motion was slightly oblique, with the Pacific plate moving south-westward relative to the Australian plate at c. 25-30mm/year. It has been suggested that it was this movement that tipped ocean floor material onto the autochthonous sediments (Rait, 2000b). However, this model implies that the Northland basin was at that time part of the Pacific plate, but this was not the case because the Vitiaz-Fiji-Tonga subduction system was already active.

The obduction of the fore-arc region of a north- or northeast-dipping subduction zone within the Australian plate can account for all the features of the Northland and East Coast allochthons and allow integration in a coherent regional model. The existence of an Eocene to Oligocene north- or northeast-dipping subduction zone to the east of the Norfolk ridge is now firmly established by the

occurrence of: i) Late Eocene subduction-related accretion, high-pressure metamorphism and obduction in New Caledonia (Aitchison et al., 1995a; Cluzel et al., 1999; Cluzel et al., 2001); ii) Eocene andesite drilled on the Bougainville seamount (Andrews et al., 1975); and, iii) mid- to Late Eocene andesitic volcanoclastic turbidites that have been drilled in the North-Loyalty Basin, at the DSDP 286 site (Andrews et al., 1975). Convergence was still active in the Late Oligocene and generated the 27-24 Ma post-obduction granodiorites of New Caledonia (Cluzel et al., 2005; Paquette and Cluzel, 2007). Recently, the discovery of ophiolite remnants, Early Oligocene boninite (Bernardel et al., 2002) and 31 Ma old blueschist on the western edge of the Three Kings Ridge (Meffre et al., 2006) provide further evidence of a subduction/obduction system that extended from New Caledonia to the Norfolk basin. We suggest that this subduction zone extended southward and that mafic allochthons of Northland and East Coast represent the obducted fore-arc of the now buried Oligocene Northland Plateau arc, and are back-arc lithosphere (Late Cretaceous-Eocene) in a fore-arc position (Oligocene) (Fig 12). According to our model, some of the IAT pillow lavas (if any) and the Oligocene intrusive rocks that crosscut Late Cretaceous-Eocene basalts may be evidence of the Oligocene volcanic-arc/fore-arc activity. The pre-obduction melange features observed in the East Coast Allochthon thus represent remnants of a peel-off melange similar to the Poya terrane of New Caledonia that developed in an early stage of intra-oceanic convergence away from any terrigenous source. The sedimentary components of the allochthon are docked in an in-sequence order and mainly display southward verging tectonic features that are consistent with the progressive duplexing and underplating of sediments that originally accumulated on a northward deepening passive margin. The topography that resulted from the flexural bend due to overburden on one hand, and tectonic imbrication on the other, allowed the final gravitational sliding of the southernmost parts of the allochthon as observed in Northland.

Conclusion

The mafic allochthonous terranes of northern New Zealand (Matakaoa Complex and Tangihua Volcanics) have most probably been generated in a single Late Cretaceous-Early Eocene back-arc basin. Correlation of the Matakaoa and

Tangihua volcanic rocks with those of the giant Cretaceous Hikurangi-Manihiki-Ontong Java Plateau (Taylor, 2005), which is significantly older and erupted in a quite different geodynamic setting, can be ruled out. It is proposed that the Hikurangi Plateau, the southernmost fragment of the giant Early Cretaceous plateau, choked the subduction zone and was responsible for the fan-shaped opening of the easternmost Late Cretaceous marginal basins of the Southwest Pacific. A direct connection between the back-arc basin in which the Matakaoa and Tangihua Volcanics were generated, and the synchronous South Loyalty back-arc basin (New Caledonia) is not possible, as they display arguably different geochemical features and transform faults are likely to have operated during their opening. Late Paleocene/Early Eocene subduction reversal was responsible for westward and thereafter south-westward arc/trench migration that finally resulted in a diachronous obduction that propagated southwards. In northern New Zealand, the Late Cretaceous-Early Eocene mafic rocks are crosscut and possibly overlain by some younger, Early Oligocene volcanic rocks which represent the arc/fore-arc activity that closely preceded obduction.

Acknowledgements

The authors acknowledge the accurate review of the ms. and very constructive comments by N. Mortimer (GNS) and an anonymous reviewer. The University of New Caledonia, the University of Grenoble 1 (LGCA, UMR 5025), and the University of Auckland have supported the analytical work on rocks of New Caledonia and New Zealand. The French Ministry of Foreign Affairs has funded the fieldwork in New Zealand (PMB, DC and CP) within the framework of the New Zealand-France Collaborative Research Programme in Earth Sciences (Fonds Pacifique).

REFERENCES

- Adams C.J., Cluzel D., and Griffin W.L., 2009. Detrital zircon ages and geochemistry of sedimentary rocks in basement Mesozoic terranes and their cover rocks in New Caledonia, and provenances at the Eastern Gondwanaland margin. *Australian Journal of Earth Sciences*. In press.
- Aitchison J. C., Clarke G. L., Cluzel D. and Meffre S. 1995. Eocene arc-continent collision in New Caledonia and implications for regional southwest Pacific tectonic evolution. *Geology*, 23, 161-164.
- Aitchison, J.C., Meffre, S., Cluzel, D., 1995. Cretaceous/Tertiary Radiolarians from New Caledonia. *Geological Society of New Zealand, Miscellaneous Publication 81A*, 70.
- Ali, J.R., Aitchison, J.C., 2000. Significance of palaeomagnetic data from the oceanic Poya Terrane, New Caledonia, for SW Pacific tectonic models. *Earth Planet. Sci. Lett.*, 177, 153-161.
- Andrews J.E., Packham G.H. et al., 1975. Initial reports of the Deep Sea Drilling Project; Washington D.C., U.S. Government Printing Office, 30, 753 p.
- Ballance, P.F. and Spörli, K.B., 1979, Northland Allochthon, *Journal of the Royal Society of New Zealand* 9, 259-275.
- Barrat J.A., Keller F., Amossé J., Taylor R.N., Nesbitt R.W., Hirata J., 1996. Determination of rare earth elements in sixteen silicate samples by ICP-MS after Tm addition and ion exchange separation, *Geostand. Newslett.* 20 (1996) 133–139.
- Bernardel, G., Carson, L., Meffre, S., Symonds, P.; Mauffret, A. 2002. Geological and morphological framework of the Norfolk Ridge to the Three Kings Region. *Geoscience Australia Record* 2002/08.
- Ben Othman, D., White W. M., and Patchett J., 1989. The geochemistry of marine sediments, island arc magma genesis, and crust-mantle recycling, *Earth Planet. Sci Lett*, 94, 1-21.
- Betts, P.G., Giles, D., Lister, G.S., Frick, L.R., 2002. Evolution of the Australian lithosphere. *Australian Journal of Earth Sciences* 49, 661–695.
- Billen M. I. & Stock J., 2000. Morphology and Origin of the Osbourn Trough. *Journal of Geophysical Research*, 105, 13482-13489.
- Black, P.M., 1989. Hydrothermal metamorphism in Tangihua Volcanics, Northland, New Zealand: a review and synthesis, In: *Geology of Northland: accretion, allochthons and arcs at the edge of the New Zealand microcontinent*, Eds. Spörli, K.B. and Kear D., *Royal Society of New Zealand Bulletin* 26.
- Black, P.M., 1995. High-Si rhyolites and shoshonitic volcanics; a late Cretaceous bimodal association, Nouméa basin, New Caledonia, *Pacrim'95 Congress, Auckland 11/95, Proceedings Vol.*, 55-58.
- Black, R.D. 1980. Upper Cretaceous and Tertiary geology of the Mangatu State Forest, Raukumara Peninsula, New Zealand, *New Zealand Journal of Geology and Geophysics*, 23: 293-312.
- Bradshaw J.D., 1989. Cretaceous geotectonic patterns in the New Zealand region. *Tectonics*, 8, 803-820.
- Bradshaw, J.D., 2004. Northland Allochthon: an alternative hypothesis of origin. *New Zealand Journal of Geology and Geophysics* 47, 375-382.
- Brathwaite R.L., Sewell R.J., and Christie A. B., 2008. Nature and tectonic setting of massive sulfide mineralisation and associated sediments and volcanics in the Matakaoa Volcanics, Raukumara Peninsula, New Zealand. *New Zealand Journal of Geology and Geophysics*, 51, 349-366, doi: 0028–8306/08/5104–0349

- Brothers, R. N. and Delaloye, M., 1982. Obducted ophiolites of North Island, New Zealand: origin, age, emplacement and tectonic implications for Tertiary and Quaternary volcanism, *New Zealand Journal of Geology and Geophysics* 25: 257-274.
- Bryan S.E., Constantine A.E., Stephens C.J., Ewart A., Schon R.W., Parianos J., 1997. Early Cretaceous volcano-sedimentary successions along the eastern Australian continental margin: Implications for the break-up of eastern Gondwana, *Earth Planet. Sci. Let.* 153, 85–102.
- Burns R.E., Andrews J.E. and the Shipboard Scientific Party, 1973. Initial rept. of the Deep Sea Drilling Project, Washington (U.S. Government Printing Office), 21, 33-56. doi:10.2973/dsdp.proc.21.105.1973
- Cassidy J., 1993. Tectonic implications of paleomagnetic data from the Northland ophiolite, New Zealand. *Tectonophysics*, 223, 199-211.
- Cawood, P.A., Nemchin, A.A., Leverenz, A., Saeed, A. and Ballance, P.F., 1999. U/Pb dating of detrital zircons: implications for the provenance record of Gondwana margin terranes. *Geol. Soc. Am. Bull.*, 111, 1107-1119.
- Cluzel D., Aitchison J.C., Clarke G., Meffre S. and Picard C., 1994. Point de vue sur l'évolution tectonique et géodynamique de la Nouvelle Calédonie. *C.R. Acad. Sci. Paris*, 319, 683-690.
- Cluzel D., Aitchison J.C., Black P.M. and Picard C., 1999. Origin and fate of Southwest Pacific marginal basins; an appraisal from New Caledonia. *Geological Society of America*, '99 Penrose Conference Arthur's Pass, South Island, New Zealand, Abstr. Vol., 21-22.
- Cluzel, D., Aitchison, J.C. Picard, C., 2001. Tectonic accretion and underplating of mafic terranes in the Late Eocene intraoceanic fore-arc of New Caledonia (Southwest Pacific): geodynamic implications. *Tectonophysics* 340, 23-59.
- Cluzel, D., Bosch D., Paquette J.L., Lemennicier Y., Montjoie Ph., Ménot R.P., 2005. Late Oligocene post-obduction granitoids of New Caledonia: a case for reactivated subduction and slab break-off. *The Island Arc*, 14, 254-271.
- Cluzel D., Adams C.J., Meffre S., Campbell H., and Maurizot P., 2009. Discovery of Early Cretaceous rocks in New Caledonia (Southwest Pacific). New U-Pb zircon age constraints on the transition from subduction to marginal breakup in south-eastern Gondwanaland. Submitted to the *Journal of Geology*.
- Crawford, A.J., Meffre, S., Symonds, P.A., 2003. 120 to 0 Ma tectonic evolution of the southwest Pacific and analogous geological evolution of the 600 to 220 Ma Tasman Fold Belt System. *Geological Society of America Special Paper* 372, 383-403.
- Davey, F.J., Henrys, S., Lodolo, E. 1997. A seismic crustal section across the East coast convergent margin, New Zealand. *Tectonophysics* 269, 199-215.
- Davy B., Hoernle K. and Werner R., 2008. Hikurangi Plateau; crustal structure, rifted formation and Gondwana subduction history. *Geochem., Geophys., Geosyst.*, 9, Q07004, doi:10.1029/2007GC001855.
- Deckert H., Ring U., Mortimer N., 2002. Tectonic significance of Cretaceous bivergent extensional shear zones in the Torlesse accretionary wedge, central Otago Schist, New Zealand. *New Zealand Journal of Geology & Geophysics*, 45, 537–547
- Eissen, J.P., Nohara, M., Cotten, J., Hirose, K., 1994. The North Fiji Basin basalts and their magmatic sources. I: Incompatible elements constraints, *Mar. Geol.*, 116, 163-178.

- Eissen, J.P., Crawford, A.J., Cotten J., Meffre S., Bellon H., Delaune M., 1998. Geochemistry and tectonic significance of basalts in the Poya Terrane, New Caledonia. *Tectonophysics*, 284, 203-219.
- Gaina C., Müller D.R., Royer J.Y., Stock J., Hardebeck J., Symonds P., 1998. The tectonic history of the Tasman Sea: a puzzle with 13 pieces. *Jour. of Geophysical Research*, 10, 312,413-12,433.
- Gifford, W.R.G. 1970. The Igneous geology of the Mangaroa Range, Hicks Bay area. Unpublished MSc thesis, University of Auckland Library.
- Gill J.B., 1981. Orogenic andesites and plate tectonics. Springer-Verlag, Berlin.
- Hall R., 2002. Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations. *J. of Asian Earth Sci.*, 20, 353-431.
- Hayward B.W., Brooke F.J., and Isaac M.J., 1989. Cretaceous to middle Tertiary stratigraphy, paleogeography, and tectonic history of Northland, New Zealand. *Royal Society of New Zealand Bull.* 26, 47-64.
- Herzer, R.H., 1995. Seismic stratigraphy of a buried volcanic arc, Northland, New Zealand and implications for Neogene subduction. *Marine and Petroleum Geology*, 12, 511-531.
- Herzer, R.H. & Mascle, J., 1996. Anatomy of a continent – backarc transform – the Vening-Meinesz Fracture Zone Northwest of New Zealand. *Marine geophysical researches* 18, 401-427.
- Herzer R.H., Mascle J., Davy B., Ruellan E., Mortimer N., Laporte C. and Duxfield A., 2000. New constraints on the New Zealand – South Fiji Basin continent back-arc margin. *C. R. Acad. Sci. Paris, Earth and Planetary Sciences*, 330, 701-708.
- Herzer R.H., Davy B.W., Mortimer N., Quilty P.G., Chaproniere G.C.H., Jones C.M., Crawford A.J., Hollis C.J., 2009. Seismic stratigraphy and structure of the Northland Plateau and the development of the Vening Meinesz transform margin, SW Pacific Ocean. *Mar Geophys Res* 30, 21-60, DOI 10.1007/s11001-009-9065-1
- Hoernle, K., Hauff, F., Werner, R., and Mortimer, N., 2004. New Insights into the Origin and Evolution of the Hikurangi Oceanic Plateau (Southwest Pacific) from Multi-beam Mapping and Sampling. *EOS, Transactions AGU*, 85 (41), 401-408.
- Hoernle K., Werner R., Hauff F., Bogaard P.v.d., 2005. The Hikurangi Oceanic Plateau: A Fragment of the Largest Volcanic Event on Earth, In: *IFM-GEOMAR Yearbook 2002-2004*, 51-54.
- Hollis, C.J. & Hanson, J.A., 1991. Well-preserved late Palaeocene radiolaria from Tangihua Complex, Camp Bay, eastern Northland. *Tane* 33: 65-76.
- Huang Y., Hawkesworth C., Smith I., van Calsteren P., Black P.M., 2000. Geochemistry of late Cenozoic basaltic volcanism in Northland and Coromandel, New Zealand: implications for mantle enrichment processes. *Chemical Geology* 164, 219–238.
- Isaac, M.J., Herzer, R.H., Brook, F.J. and Hayward, B.W., 1994, Cretaceous and Cenozoic sedimentary basins of Northland, New Zealand. *Institute of Geological and Nuclear Sciences Monograph* 8.
- Jenner G.A., Dunning G.R., Malpas J., Brown M. and Brace T., 1991. Bay of Islands and Little Port complexes, revisited : age, geochemical and isotopic evidence confirm suprasubduction-zone origin. *Canadian Jour. of Earth Sci.*, 28, 1635-1652.
- Keller N. S., Arculus R. J., Hermann J., Richards S. (2008). "Submarine back-arc lava with arc signature: Fonualei Spreading Center, northeast Lau Basin, Tonga". *JGR* 113:B08S07 doi:10.1029/2007JB005451

- Kenny J.A., 1984. Petrography of Cretaceous and Tertiary sedimentary rocks of the Ihungia catchment, Raukumara Peninsula, North Island, New Zealand. *New Zealand journal of Geology and Geophysics*, 27, 291-298.
- Laird M.G. and Bradshaw J.D., 2004. The break-up of a long-term relationship: the Cretaceous separation of New Zealand from Gondwana. *Gondwana Research*, 7, 273-286.
- Langmuir, C. H., A. Bezos, et al. (2006). "Chemical Systematics and Hydrous Melting of the Mantle in Back-Arc Basins".in D. M. Christie, C. R. Fisher, S.-M. Lee and S. Givens: *Back-Arc Spreading Systems: Geological, Biological, Chemical, and Physical Interactions: Geophysical Monograph*. American Geophysical Union.
- Leterrier, J., Maury, R.C., Thonon, P. Girard, D., Marchal, M. 1982. Clinopyroxenes composition as a method of identification of magmatic affinities of paleo-volcanic series. *Earth and planetary science letters* 59, 567-600.
- Lister, G.S., Etheridge, M.A., 1989. Towards a general model; Detachment models for uplift and volcanism in the eastern highlands, and their application to the origin of passive margin mountains. In: Johnson, R.W., Knutson, J., Taylor, S.R. (Eds.), *Intraplate Volcanism in Eastern Australia and New Zealand*. Cambridge Univ. Press, Cambridge, pp. 297–313.
- Malpas, J., Spörli, K. B., Black, P. M. and Smith, I. E. M., 1992. Northland ophiolite, New Zealand, and implications for plate-tectonic evolution of the southwest Pacific: *Geology*, 20, 149-152.
- Marsh N.G., Saunders A.D., Tarney J., and Dick H.J.B., 1980. Geochemistry of basalts from the Shikoku and Daito basins, DSDP Leg 58. In: Stout L.N. (Ed.), *Initial Repts. Deep Sea Drilling Project 58*, US Government Printing Office, Washington DC, 805-842.
- Mauffret A., Symonds P., Benkhelil J, Bernardel G., Buchanan C., D'Acremont E., Gorini C., Lafoy Y., Nercessian A., Ryan J., Smith N., and Van de Beuque S., 2001. Collaborative Australia-France Multibeam Seafloor Mapping Survey – Norfolk Ridge to Three Kings Ridge region: FAUST 2, Preliminary results. Australian Geological Survey Organisation Record, 2001/27.
- Mazengarb, C. and Speden, I.G. (compilers), 2000. Geology of the Raukumara area. Geological and Nuclear Sciences 1:250 000 geological map 6. Lower Hutt, New Zealand. Institute of Geological and Nuclear Sciences Limited.
- Meffre S., Crawford A.J., Quilty P.G., 2006. Arc-continent collision forming a large island between New Caledonia and New Zealand in the Oligocene. AESC2006, Melbourne, Australia.
- Metcalf, R. V. and J. W. Shervais (2008). Subrasubduction-zone ophiolites: Is there really an ophiolite conundrum? *GSA Special Paper: Ophiolites, Arcs, and Batholiths: A Tribute to Cliff Hopson*. J. E. Wright and J. W. Shervais, Geological Society of America. 438: 191-222.
- Moore, P.R., 1985. Distribution, age and relationship of volcanic rocks ("East Coast volcanics") in the Gisborne – East coast Region. *New Zealand Geological Survey Record* 8, 16-23.
- Moore, P.R., Brathwaite, R.L. and Roser, B. 1987 Correlation of Early Cretaceous volcanic-sedimentary sequences at Red Island and Hinemahanga Rocks, southern Harkes Bay. *New Zealand Geological Survey Records* 18, 27-31.
- Mortimer, N. 2004 New Zealand's geological foundations. *Gondwana Research*, 7, 261-272

- Mortimer, N. and Parkinson, D. 1996. Hikurangi Plateau: a Cretaceous large igneous province in the southwest Pacific Ocean. *Journal of geophysical research* 101, No B1, 687-696.
- Mortimer N., Herzer R.H., Gans P.B., Parkinson D.L., and Seward D., 1998. Basement geology from Three Kings Ridge to West Norfolk Ridge, southwest Pacific Ocean; evidence from petrology, geochemistry and isotopic dating of dredge samples. *Marine Geology*, 148, 135-162.
- Mortimer N., Tulloch A.J., Spark R.N., Walker N.W., Ladley E., Allibone A. and Kimborough D.L., 1999. Overview of the Median Batholith, New Zealand: a new interpretation of the geology of the Median Tectonic Zone and adjacent rocks. *Journal of African Earth Sciences*, 29, 257-268.
- Mortimer N., Herzer R.H., Walker N.W., Calvert A.T., Seward D., Chapronniere, G.C.H., 2003. Cavalli seamount, Northland Plateau, SW Pacific ocean: a Miocene metamorphic core complex? *Journal of the Geological Society of London*, 160, 971-983.
- Mortimer N., Hoernle K., Hauff F., Palin J.M., Dunlap W.J., Werner R. and Faure K., 2006. New constraints on the age and evolution of the Wishbone Ridge, southwest Pacific Cretaceous microplates, and Zealandia–West Antarctica breakup *Geology* 34; 3; 185-188; DOI: 10.1130/G22168.1
- Mortimer N., Herzer R.H., Gans P.B., Laporte-Magoni C., Calvert A.T., Bosch D., 2007. Oligocene-Miocene tectonic evolution of the South Fiji Basin and Northland Plateau, SW Pacific Ocean: Evidence from petrology and dating of dredged rocks. *Marine Geology*, 237, 1-24.
- Muir R.J., Weaver S.D., Bradshaw J.D., Eby G.N. and Evans J.A., 1995. The Cretaceous Separation Point batholith, New Zealand: granitoid magmas generated by partial melting of mafic lithosphere. *J. Geol. Soc. London*, 152, 689-701.
- Nicholson, K.N. 1999. The Tangihua complex, New Zealand: implication for Cretaceous-Oligocene convergent margin process in the SW Pacific from comparison with the Poya terrane, New Caledonia. Unpublished PhD thesis, Universities of Auckland and Grenoble I libraries.
- Nicholson, K.N., Black, P.M. and Picard, C., 2000a. Geochemistry and tectonic significance of the Tangihua Ophiolite Complex, New Zealand. *Tectonophysics* 321, 1-15.
- Nicholson, K.N., Picard, C. and Black, P.M., 2000b. A comparative study of Late Cretaceous ophiolitic basalts from New Zealand and New Caledonia: implications for the tectonic evolution of the SW Pacific. *Tectonophysics*, 327, 157-171.
- Nicholson K.N. and Black P.M., 2004. Cretaceous to early Tertiary basaltic volcanism in the Far North of New Zealand: geochemical associations and their tectonic significance *New Zealand Journal of Geology & Geophysics*, 47, 437–446.
- Nicholson, K.N., Black, P.M., 2004. Cretaceous to early Tertiary basaltic volcanism in the Far North of New Zealand: geochemical associations and their tectonic significance, *New Zealand Journal of Geology & Geophysics*, 47, 437-446.
- Nicholson K.N., Black, P.M., Picard C., Cooper P., Hall C.M., Itaya T., 2007. Alteration, age, and emplacement of the Tangihua Complex Ophiolite, New Zealand. *New Zealand Journal of Geology and Geophysics*. 50, 151-164, 0028-8306/07/5002-0151
- Norrish K. and Chappel B.W., 1977. X-Ray fluorescence spectrography. In: Zussman J. (ed.) *Physical methods in determinative mineralogy*. Academic Press, New York, 201-272.

- Paquette J.L. and Cluzel D., 2007. Timing and crustal heritage of post-obduction volcanic-arc granitoids and a granulite xenolith from New Caledonia. Evidence from U-Pb zircon geochronology. Inference on Southwest Pacific Geodynamic Models. *Intern. J Earth Sci (Geol Rundsch)* 96:613-622. DOI, 10.1007/s00531-006-0127-1.
- Parrot J.F., and Dugas F., 1980. The disrupted ophiolite belt of the Southwest Pacific: evidence for an Eocene subduction zone. *Tectonophysics*, 66, 349-372.
- Pearce J.A., 1980. Geochemical evidence for the genesis and eruption settings of lavas from Tethyan ophiolites, *In: Panayiotou A. (ed.), Ophiolites*, Geological Survey Dept. Cyprus, 261-272.
- Perfit M.R., Langmuir C.H., Baekisapa M., Chappell B., Johnson R.W., Staudigel H. and Taylor S.R., 1987. Geochemistry and petrology of rocks from the Woodlark basin: addressing the questions of ridge subduction. In : Taylor S.R. and Exxon N.F. (Eds), *Marine geology, geophysics and geochemistry of the Woodlark – Solomon Islands*, Circum Pacific Council for Energy and Mineral Resources, Houston Texas, Earth Sci. Series, 7, 113-154.
- Picard, M., 1995. Ouverture de bassins marginaux dans le Pacifique sud-ouest. Paléovolcanisme et reconstitution des environnements géodynamiques. Extrémité nord de la Nouvelle-Calédonie, Unpubl. MSc thesis, Univ. Grenoble, 124 p.
- Pirajno, F. 1979. Geology, geochemistry, and mineralisation of a spilite-keratophyre association in Cretaceous flysch, East coast area, New Zealand. *New Zealand Journal of Geology and Geophysics*, 22, 307-328.
- Pirajno, F. 1980. Subsea floor alteration in rocks of the Matakaoa Volcanics around Lottin Point, East coast, New Zealand *New Zealand Journal of Geology and Geophysics*, 23, 313-334.
- Rait, G.J. 2000a. Amounts and rates of transport of the Northland and East Coast Allochthons (abstract). Geological Society of New Zealand annual conference, Wellington. *Geological Society of New Zealand miscellaneous publication* 108A: 129.
- Rait, G.J. 2000b. Thrust transport directions in the Northland Allochthon, New Zealand. *New Zealand Journal of Geology and Geophysics*, 43, 271-288.
- Rutherford, P.G. 1980. Geology of the Matakaoa Volcanic Group, Cape Runaway area. Unpublished MSc thesis, University of Auckland Library.
- Saunders A.D. and Tarney J., 1984. Geochemical characteristics of basaltic volcanism within back-arc basins. In: Kokelaar B.P. and Howells M.F. (Eds.), *Marginal Basin Geology*, Geol. Soc. Spec. Pub. 16, 59-76.
- Schellart W.P., Lister G.S. and Toy V.G., 2006. A Late Cretaceous and Cenozoic reconstruction of the Southwest Pacific region: tectonics controlled by subduction and slab rollback processes. *Earth Science Reviews*, 76, 191-233.
- Sdrolias, M., Muller, R.D., Gaina, C., 2003. Tectonic evolution of the Southwest Pacific using constraints from back-arc basins. *Geological Society of America Special Paper* 372, 343-359.
- Sdrolias, M., Mueller, R.D., Mauffret, A., Bernadel, G. 2004. Enigmatic formation of the Norfolk Basin, SW Pacific: a plume influence on back-arc extension. *Geochemistry, Geophysics, Geosystems* 5(6), Q06005, doi:10.1029/2003GC000643.
- Spell T.L., McDougal I., and Tulloch A.J., 2000. Thermochronological constraints on the break-up of the Pacific Gondwana margin: the Paparoa metamorphic core complex, South Island, New Zealand. *Tectonics*, 19, 433-451.
- Spörl, K. B., 1989. Tectonic framework of Northland, New Zealand, *In: , Spörl, K. B. and Kear, D. (eds), Geology of Northland: accretion, allochthons and arcs at the*

- edge of the New Zealand micro-continent, *Royal Society of New Zealand Bulletin* 26, 3-14.
- Spörli, K.B. and Aita, Y. 1994. Tectonic significance of Late Cretaceous radiolaria from the obducted Matakaoa Volcanics, East coast, North Island, New Zealand. *Geoscience reports of Shizuoka University* 20, 115-133.
- Stonely, R. 1968. A lower Tertiary decollement on the East Coast, North Island, New Zealand. *New Zealand Journal of Geology and Geophysics*, 11, 128-156.
- Strong, C.P. 1976. Cretaceous foraminifera from the Matakaoa Volcanics (note). *New Zealand Journal of Geology and Geophysics*, 19, 140-143.
- Strong, C.P. 1980. Early Paleogene foraminifera from Matakaoa Volcanics. *New Zealand Journal of Geology and Geophysics*, 23, 267-272.
- Sun, S.S., McDonough, W.I., 1989. Chemical and isotopic systematics of oceanic basalts: implications for mantle composition and processes. In : Saunders A.D. et Norry M.D. (eds), *Magmatism in the Ocean Basins*, Geological Society Special Publication, 42, 313-345.
- Sutherland R., 1999. Basement geology and tectonic development of the greater New Zealand region: an interpretation from regional magnetic data. *Tectonophysics*, 308, 341-362.
- Taylor B., 2005. The single largest oceanic plateau: Ontong Java–Manihiki–Hikurangi. *Earth and Planet. Sci. Let.* 241, 372-380 doi:10.1016/j.epsl.2005.11.049
- Thompson G.M., Malpas J., and Smith I.E.M., 1997. The geochemistry of tholeiitic and alkalic plutonic suites within the Northland ophiolite, northern New Zealand; magmatism in a back-arc basin. *Chemical Geology* 142, 213-223.
- Tulloch A.J. and Kimbrough D.L., 1989. The Paparoa metamorphic core complex, New Zealand: Cretaceous extension associated with fragmentation of the Pacific margin of Gondwana. *Tectonics*, 8, 1217-1234.
- Vallier T.L., Jenner G.A., Frey F.A., Gill J.B., Davis A.S., Volpe A.M., Hawkins J.W., Morris J.D., Cawood P.A., Morton J.L., Scholl D.W., Rautenschlein M., White W.M. and Williams M., 1991. Subalkaline andesites from Valu Fa Ridge, a back-arc spreading center in southern Lau Basin: Petrogenesis, comparative chemistry and tectonic implications. *Chemical Geology*, 91, 227-256.
- Veevers, J.J., 2000a. Change of tectono-stratigraphic regime in the Australian plate during the 99 Ma (mid-Cretaceous) and 43 Ma (mid-Eocene) swerves of the Pacific. *Geology* 28, 47–50.
- Veevers, J.J., 2000b. Chapter 16. Mid-Cretaceous (99 Ma) and mid-Eocene (43 Ma) events in the Indo-Australian and Antarctic plate and coeval swerves of the Pacific Plate. In: Veevers, J.J. (Ed.), *Billion-year Earth History of Australia and Neighbours in Gondwanaland*. GEMOC Press, Sydney, pp. 102–109.
- Veevers, J.J., 2004. Gondwanaland from 650–500Ma assembly through 320Ma merger in Pangea to 185–100Ma breakup: supercontinental tectonics via stratigraphy and radiometric dating. *Earth-Science Reviews* 68, 1–132.
- Whattam S.A., Malpas J., Ali, J.R., Lo C.H., Smith I.E., 2005. Formation and emplacement of the Northland ophiolite, northern New Zealand: SW Pacific tectonic implications. *Journal of the Geological Society of London*, 162, 225-241.
- Whattam S.A., Malpas J., Smith I.E.M., Ali J.R., 2006. Link between SSZ ophiolite formation, emplacement and arc inception, Northland, New Zealand: U–Pb SHRIMP constraints; Cenozoic SW Pacific tectonic implications. *Earth Planet. Sci. Let.* 250, 606–632.
- Whattam, S. A., J. Malpas, J. R. Ali, and I. E. M. Smith, 2008. New SW Pacific tectonic model: Cyclical intraoceanic magmatic arc construction and near-coeval

emplacement along the Australia-Pacific margin in the Cenozoic, *Geochem. Geophys. Geosyst.*, 9, Q03021, doi:10.1029/2007GC001710.

Figure captions

Figure 1: Sketch map of the Southwest Pacific (oceanic crust in grey, continental crust in white, volcanic-arc crust dotted). B: Bougainville Seamount.

Figure 2: Geologic sketch map of North Island (New Zealand) to show the location of Northland and East Coast allochthons (dashed) with their Late Cretaceous-Early Eocene oceanic crust components (in black) (adapted from Davey et al., 1997). VMFZ: Veining-Meinesz fracture zone, TVZ: Taupo Volcanic Zone.

Figure 3: Geologic map of the East Coast Allochthon in the northern Raukumara peninsula (simplified after Mazengarb & Speden, 2000), with sample location (grid numbers refer to the 1:50,000 map of New Zealand Y 14 sheet), and local names cited in the text. For map location, see Fig. 2. Insert shows the location of Matakaoa Volcanics within the East Coast Allochthon.

Figure 4: Zr/TiO₂ vs. Nb/Y classification diagram for altered or metamorphic rocks of Winchester and Floyd (1977) to show a limited range of variation of Zr/TiO₂ which represents differentiation; in contrast, a wide range of Nb/Y values is related to contrasting sources and reflects a variable Nb depletion.

Figure 5: Th/Yb vs. Ta/Yb discrimination diagram of Pearce (1983) to show the range of variation of Ta/Yb which allows the discrimination of the four rock types discussed in the text.

Figure 6: Chondrite-normalised REE diagram (a) and N-MORB-normalised expanded REE-trace elements spiderdiagram (b) of Matakaoa volcanic rocks (East Coast Allochthon) – Normalization values are from Pearce (1980) and Sun and McDonough (1989).

Figure 7: Plotted against the (Nb/La)_n ratio which represents the increasing subduction influence in Matakaoa Volcanics from MORB-like to IAT-like end-members, Sr and Pb contents do not show any correlation; a feature which suggest only a minor or negligible influence of subduction-derived or continental crust material.

Figure 8: Sr and Nd isotopic ratios of Tangihua and Matakaoa volcanic rocks (Table 3) after Ben Othman et al. (1989). A shift towards higher ⁸⁷Sr/⁸⁶Sr ratios for constant εNdi values (especially prominent for the "teshenite") is a common feature of volcanic rocks mildly altered by "hydrothermal" processes (e.g. sea water-rock interaction). A plot of three leached samples of Hikurangi Plateau basalts with distinctively lower εNdi (Mortimer and Parkinson, 1996) is shown for comparison.

Figure 9 : Chondrite-normalised REE diagram (a) and MORB-normalised expanded REE-trace elements spiderdiagram (b) of Tangihua volcanic rocks (Northland Allochthon) – same references as Fig 5. Data from Nicholson et al (2000a).

Figure 10 : Th/Yb vs. Ta/Yb discrimination diagram of Pearce (1983) to compare the range of variation of Mataoa Volcanics and Tangihuas Complex basalts that both plot within the same restricted field and the fields of Lau, Mariana and East Scotia back-arc basin basalts (data from Langmuir et al., 2006). New Zealand's allochthonous basalts appear to have been generated in conditions intermediate between the juvenile Lau Basin and the more mature Mariana and East Scotia basins.

Figure 11: A tentative reconstruction of the evolution of south-eastern Gondwana margin in the Late Early Cretaceous (ca. 105 Ma, 10a) (modified from Gaina, et al., 1998, Sutherland, 1999, and Hall, 2002), and Paleocene (ca. 70 Ma, 10b) (modified from Cluzel et al., 1999; this study), to show the marginal basins in which the mafic allochthons of New Caledonia and New Zealand have been generated. Plate boundaries are only indicative due to syn-rift stretching and subsequent Tertiary tectonics. NC: New Caledonia, LHR: Lord Howe rise, MBL: Mary Byrd Land, NLB: North Loyalty Basin, SFB: South Fiji Basin, Camp: Campbell Plateau, Chall: Challenger Plateau, Chath: Chatham Plateau, Hiku: Hikurangi Plateau.

Figure 12 : A model for Late Cretaceous to Miocene geodynamic evolution of the Southwest Pacific (modified from Cluzel et al., 1999, no scale). 12a: Early Cretaceous (c. 105 Ma) end of Phoenix/Pacific plate subduction, marginal accretion and Median Batholith activity; 12b: Early Late Cretaceous (c. 90 Ma) slab break-off and incipient boudinage of the lithosphere (note that section 11a is located at a more southerly latitude as 11b); 12c: Campanian-Paleocene (80-65 Ma) Gondwana marginal rifting, subduction reactivation, oceanward arc migration and marginal basin opening; 12d: Eocene-Oligocene (52-30 Ma) subduction flip initiated at New Caledonia's latitude and propagating southward, formation of an Oligocene island arc; 12e: Early Miocene (ca. 22-20 Ma) attempted subduction of the North Island margin and obduction of Northland and East Coast Allochthons. Boxes legend: 1 oceanic crust; 2 continental crust, 3 lithospheric mantle, 4 asthenosphere.

Annex I: A series of binary diagrams to show the variation of selected major (in wt%), trace elements (in ppm) and elemental ratios versus MgO content of Matakaoa Volcanics which is known to represent the differentiation process. Note the prominent variation of mobile elements (K_2O), the poor correlation of SiO_2 and MgO; and in contrast, the relatively good fit of the four rock types defined in this paper with the contents of selected "immobile" trace elements (Ti, REE), and some elemental ratios.

Table 1: location, geologic setting and chemical affinity of the analysed samples of Matakaoa Volcanics.

| Sample n° | rock name | setting | Affinity | Location | NZMS 260 1 : 50,000 map / grid reference |
|-----------|--------------------|---------------|----------|--------------|--|
| PMB32043 | basalt | flow | MORB | Potikirua Pt | Y14 / 598935 |
| PMB32046 | basalt | flow | IAT | Runaway E | Y14 / 530933 |
| MK-34 | doleritic basalt | flow | IAT | Runaway E | Y14 / 522935 |
| MK-38 | doleritic basalt | flow | IAT | Runaway E | Y14 / 525933 |
| MK-33 | dolerite | dyke | IAT | Runaway E | Y14 / 523934 |
| PMB32008 | dolerite | dyke | IAT | Runaway E | Y14 / 504950 |
| PMB32005 | basalt | flow | TB | Runaway E | Y14 / 508946 |
| PMB32111 | basalt | pillow | TB | Runaway E | Y14 / 508946 |
| PMB32048 | dolerite | dyke | BABB | Runaway E | Y14 / 505947 |
| MK-36 | doleritic basalt | flow | BABB | Runaway E | Y14 / 517937 |
| MK-37 | basalt | flow | BAB | Runaway E | Y14 / 520936 |
| MK-35 | basalt | pillow | MORB | Runaway E | Y14 / 514940 |
| MK-30 | basalt | pillow | MORB | Runaway W | Y14 / 504920 |
| MK-29 | basalt | pillow | MORB | Runaway W | Y14 / 505918 |
| MK-32 | basalt | pillow | MORB | Runaway W | Y14 / 506917 |
| PMB15789 | variolithic basalt | breccia | MORB | Midway Pt | Z14 / 709933 |
| PMB15831 | basalt | dyke | IAT | Midway Pt | Z14 / 710930 |
| MK-21 | teschenite | sill | MORB | Lottin Pt | Y14 / 666931 |
| MK-23 | teschenite | sill | BAB | Lottin Pt | Y14 / 665925 |
| MK-24 | basalt | pillow | IAT | Lottin Pt | Y14 / 649926 |
| MK-14 | dolerite | dyke | IAT | Lottin Pt | Y14 / 662925 |
| MK-16 | basalt | pillow | IAT | Lottin Pt | Y14 / 662925 |
| MK-27 | dolerite | dyke | IAT | Lottin Pt | Y14 / 633278 |
| MK-45 | gabbro | dyke | MORB | Lottin Pt | Y14 / 645930 |
| PMB15856 | basalt | glass | IAT | Lottin Pt E | Y14 / 692938 |
| MK- 2 | dolerite | breccia | IAT | Matakaoa pt | Z14 / 782909 |
| MK- 3 | andesite | flow | IAT | Matakaoa pt | Z14 / 764917 |
| MK- 4 | basalt | hyaloclastite | IAT | Matakaoa pt | Z14 / 767917 |
| MK- 8 | dolerite | volc. breccia | IAT | Matakaoa pt | Z14 / 773916 |
| MK-13 | basalt | hyaloclastite | IAT | Matakaoa pt | Z14 / 784915 |
| MK- 7 | andesitic basalt | clast | IAT | Matakaoa pt | Z14 / 773916 |
| MK-10 | andesitic basalt | flow | IAT | Matakaoa pt | Z14 / 779916 |
| MK-12 | basalt | breccia | IAT | Matakaoa pt | Z14 / 784915 |
| MK- 1 | basalt | flow | BABB | Pukeamaru E | Z14 / 782874 |
| MK-48 | basalt | clast | IAT | Pukeamaru E | Z14 / 796864 |
| MK-40 | basalt | flow | BABB | Pukeamaru N | Y14 / 650853 |
| MK-46 | basalt | volc. breccia | MORB | Pukeamaru | Z14 / 803869 |
| MK-41 | dolerite | flow | BABB | Pukeamaru S | Z14 / 760803 |
| MK-42 | basalt | pillow | MORB | Pukeamaru S | Z14 / 759802 |

Table 2: Major and trace element compositions of Matakaoa volcanic rocks (major elements in wt.%, trace elements and REE in ppm)

| Sample n° | PMB32043 | MK-46 | MK-42 | MK-35 | MK-30 | MK-29 | MK-32 | MK-21 | PMB15789 | MK-1 | MK-40 | PMB32048 | MK-36 |
|---|--------------|------------------|---------------|---------------|---------------|---------------|---------------|------------------|------------------|-------------|-------------|-----------|------------------|
| Name | basalt | basaltic breccia | pillow basalt | pillow basalt | pillow basalt | pillow basalt | pillow basalt | tesche-nite sill | variolit. basalt | basalt | basalt | dolerite | dolerite. basalt |
| Affinity | MORB | MORB | MORB | MORB | MORB | MORB | MORB | MORB | MORB | BAB | BAB | BAB | BAB |
| Location | Potikirua Pt | Pukeamaru | Pukeamaru S | Runaway E | Runaway W | Runaway W | Runaway W | Lotin Pt | Midway Pt | Pukeamaru E | Pukeamaru N | Runaway E | Runaway E |
| SiO ₂ | 49.52 | 47.76 | 50.46 | 48.92 | 47.73 | 49.22 | 50.62 | 50.67 | 48.55 | 48.97 | 48.89 | 48.43 | 50.20 |
| TiO ₂ | 2.14 | 2.25 | 1.75 | 1.75 | 2.04 | 2.52 | 2.73 | 1.94 | 2.39 | 1.38 | 1.79 | 1.45 | 1.35 |
| Al ₂ O ₃ | 14.01 | 14.63 | 14.26 | 14.59 | 13.87 | 13.57 | 13.78 | 15.36 | 13.57 | 14.89 | 13.99 | 12.78 | 15.46 |
| Fe ₂ O ₃ _T | 11.96 | 12.45 | 11.92 | 11.22 | 12.40 | 12.30 | 11.16 | 12.05 | 13.67 | 10.72 | 11.83 | 10.74 | 10.54 |
| MnO | 0.22 | 0.20 | 0.20 | 0.17 | 0.21 | 0.28 | 0.18 | 0.23 | 0.22 | 0.19 | 0.18 | 0.17 | 0.17 |
| MgO | 6.08 | 6.60 | 5.83 | 7.55 | 6.95 | 5.77 | 5.76 | 4.98 | 6.26 | 7.14 | 6.26 | 9.06 | 6.97 |
| CaO | 9.56 | 10.78 | 10.25 | 10.13 | 9.08 | 10.90 | 9.06 | 7.54 | 9.92 | 9.26 | 8.36 | 11.29 | 11.65 |
| Na ₂ O | 3.58 | 3.07 | 2.76 | 3.61 | 3.93 | 3.05 | 4.02 | 5.10 | 2.93 | 3.92 | 4.70 | 2.96 | 2.23 |
| K ₂ O | 0.31 | 0.30 | 0.19 | 0.15 | 0.36 | 0.19 | 0.76 | 0.42 | 0.31 | 0.62 | 0.11 | 0.17 | 0.13 |
| P ₂ O ₅ | 0.27 | 0.24 | 0.17 | 0.21 | 0.19 | 0.24 | 0.26 | 0.22 | 0.26 | 0.13 | 0.16 | 0.13 | 0.14 |
| Loi+ | 1.18 | 1.10 | 2.14 | 1.78 | 2.91 | 1.40 | 1.14 | 1.19 | 0.82 | 2.01 | 2.38 | 2.23 | 1.11 |
| H ₂ O- | 0.86 | 0.77 | 0.14 | 0.08 | 0.40 | 0.49 | 0.47 | 0.28 | 0.46 | 0.72 | 1.17 | 0.34 | 0.05 |
| Total | 99.69 | 100.14 | 100.06 | 100.18 | 100.05 | 99.93 | 99.93 | 99.97 | 99.34 | 99.95 | 99.83 | 99.76 | 100.00 |
| Co | 44.6 | 41.6 | nd | 41.2 | 42.7 | 64.8 | 42.0 | 31.8 | 61.2 | 36.6 | 39.0 | 53.1 | 40.1 |
| Cr | 230 | 218 | 102 | 318 | 268 | 253 | 158 | 55 | 199 | 318 | 172 | 702 | 290 |
| Ni | 61.7 | 65.4 | 42.9 | 108.4 | 75.6 | 77.3 | 67.5 | 33.2 | 84.5 | 72.2 | 51.5 | 134.7 | 81.1 |
| Ga | 19.4 | 20.1 | 20.9 | 18.8 | 21.0 | 20.6 | 21.3 | 21.5 | 20.5 | 15.9 | 17.3 | 14.7 | 17.3 |
| Sc | 38.2 | 41.0 | 38.0 | 37.2 | 41.5 | 46.1 | 48.9 | 41.8 | 44.7 | 43.6 | 43.1 | 56.4 | 35.1 |
| V | 362 | 404 | 364 | 309 | 413 | 441 | 430 | 352 | 397 | 316 | 365 | 322 | 291 |
| Ba | 29.1 | 13.7 | 30.8 | 17.0 | 54.4 | 21.9 | 61.4 | 353.3 | 20.1 | 33.4 | 571.1 | 23.9 | 16.5 |
| Rb | 3.70 | 1.98 | 2.59 | 1.80 | 1.29 | 3.05 | 15.25 | 5.98 | 3.05 | 3.88 | 1.58 | 2.15 | 2.21 |
| Sr | 155 | 170 | 127 | 120 | 110 | 119 | 127 | 173 | 125 | 253 | 166 | 181 | 128 |
| Y | 57.9 | 54.2 | 42.2 | 42.8 | 43.7 | 57.6 | 56.4 | 48.6 | 56.8 | 31.7 | 41.5 | 37.6 | 32.3 |
| Zr | 199 | 184 | 129 | 143 | 135 | 177 | 187 | 119 | 198 | 89 | 120 | 73 | 94 |
| Nb | 5.50 | 4.54 | 3.03 | 4.12 | 3.90 | 4.44 | 4.97 | 4.09 | 4.96 | 1.67 | 1.86 | 2.72 | 2.68 |
| Cs | 0.03 | 0.02 | nd | 0.16 | 0.08 | 0.05 | 0.19 | 0.12 | 0.05 | 0.15 | 0.82 | 0.23 | 0.07 |
| Hf | 4.83 | 4.43 | nd | 3.73 | 3.50 | 4.49 | 4.98 | 3.30 | 5.03 | 2.40 | 3.14 | 2.21 | 2.49 |
| Ta | 0.360 | 0.308 | nd | 0.281 | 0.249 | 0.303 | 0.343 | 0.259 | 0.400 | 0.151 | 0.132 | 0.160 | 0.177 |
| Pb | 1.43 | 0.47 | 2.37 | 0.39 | 0.53 | 0.87 | 0.55 | 0.61 | 0.89 | 0.89 | 0.24 | 0.64 | 0.39 |
| Th | 0.390 | 0.324 | 0.000 | 0.306 | 0.230 | 0.278 | 0.324 | 0.241 | 0.386 | 0.182 | 0.159 | 0.126 | 0.280 |
| U | 0.175 | 0.187 | 2.280 | 0.151 | 0.102 | 0.107 | 0.293 | 0.222 | 0.152 | 0.122 | 0.116 | 0.139 | 0.104 |
| La | 7.53 | 6.18 | 1.55 | 5.79 | 4.31 | 5.33 | 5.47 | 6.05 | 7.25 | 3.06 | 3.58 | 4.10 | 3.96 |
| Ce | 22.45 | 18.78 | 14.16 | 17.17 | 13.45 | 17.38 | 19.73 | 18.12 | 20.89 | 9.53 | 11.77 | 12.88 | 11.70 |
| Pr | 3.72 | 3.14 | nd | 2.84 | 2.35 | 2.99 | 3.35 | 3.05 | 3.58 | 1.69 | 2.10 | 2.04 | 1.97 |
| Nd | 18.49 | 16.67 | nd | 14.42 | 12.64 | 16.29 | 18.04 | 15.85 | 18.83 | 9.22 | 11.69 | 10.87 | 10.14 |
| Sm | 6.22 | 5.59 | nd | 4.65 | 4.45 | 5.67 | 6.27 | 5.24 | 5.81 | 3.17 | 4.13 | 3.96 | 3.35 |
| Eu | 1.95 | 1.80 | nd | 1.65 | 1.53 | 1.88 | 2.17 | 1.75 | 1.96 | 1.15 | 1.45 | 1.25 | 1.22 |
| Gd | 7.66 | 6.93 | nd | 5.84 | 5.90 | 7.45 | 8.19 | 6.83 | 8.13 | 4.12 | 5.81 | 5.00 | 4.28 |
| Tb | 1.366 | 1.290 | nd | 1.071 | 1.059 | 1.409 | 1.481 | 1.197 | 1.442 | 0.780 | 1.005 | 0.884 | 0.801 |
| Dy | 9.50 | 8.56 | nd | 6.81 | 7.08 | 9.17 | 9.63 | 7.56 | 8.74 | 5.09 | 6.61 | 5.65 | 5.18 |
| Ho | 1.97 | 1.88 | nd | 1.49 | 1.55 | 2.03 | 2.08 | 1.66 | 1.89 | 1.15 | 1.50 | 1.24 | 1.16 |
| Er | 5.85 | 5.44 | nd | 4.28 | 4.54 | 5.89 | 5.81 | 4.75 | 5.71 | 3.28 | 4.30 | 3.68 | 3.31 |
| Tm | 3.40 | nd | nd | nd | nd | nd | nd | nd | 3.55 | nd | nd | 2.65 | nd |
| Yb | 5.38 | 4.87 | nd | 3.91 | 4.25 | 5.28 | 5.38 | 4.37 | 5.37 | 3.00 | 3.88 | 3.29 | 3.08 |
| Lu | 0.830 | 0.748 | nd | 0.588 | 0.662 | 0.779 | 0.799 | 0.649 | 0.815 | 0.462 | 0.593 | 0.515 | 0.479 |

| Sample | MK-23 | MK-24 | MK-14 | MK-16 | MK-27 | MK-2 | MK-13 | MK-7 | MK-10 | MK-12 | PMB15831 | MK-48 | PMB32046 |
|--------------------|---------------------|------------------|--------------|------------------|--------------|---------------------|------------------|--------------------|-------------------------|---------------------|------------------|-------------------|-----------|
| Name | Tesch- nite sill | pillow basalt | dolerit e | pillow basalt | dolerit e | dolerite breccia | vitric nodule | andesitic clast | andesit ic basalt | basaltic breccia | basaltic dyke | basaltic clast | basalt |
| Affinity | BAB | IAT | IAT | IAT | IAT | IAT | IAT | IAT | IAT | IAT | IAT | IAT | IAT |
| Location | Lottin Pt | Lottin Pt | Lottin Pt | Lottin Pt | Lottin Pt | Matakaoa pt | Matakaoa pt | Matakaoa pt | Matakaoa pt | Matakaoa pt | Midway Pt | Pukeamaru E | Runaway E |
| SiO2 | 49.94 | 50.62 | 50.89 | 52.64 | 49.37 | 52.23 | 49.90 | 55.79 | 56.80 | 51.61 | 48.28 | 52.85 | 51.53 |
| TiO2 | 1.54 | 1.46 | 1.12 | 1.59 | 1.31 | 1.13 | 1.33 | 1.59 | 1.26 | 1.36 | 1.47 | 1.12 | 1.51 |
| Al2O3 | 14.67 | 15.02 | 15.41 | 14.40 | 18.58 | 15.53 | 15.56 | 14.66 | 14.81 | 15.44 | 14.98 | 14.84 | 14.84 |
| Fe2O3 _T | 10.16 | 10.62 | 9.64 | 11.50 | 9.27 | 9.08 | 9.91 | 10.43 | 8.31 | 9.99 | 11.31 | 10.99 | 11.64 |
| MnO | 0.19 | 0.15 | 0.17 | 0.17 | 0.15 | 0.15 | 0.16 | 0.15 | 0.12 | 0.17 | 0.18 | 0.13 | 0.19 |
| MgO | 6.37 | 5.46 | 7.08 | 4.75 | 4.81 | 6.42 | 6.07 | 4.06 | 4.52 | 6.23 | 6.49 | 6.47 | 5.85 |
| CaO | 11.30 | 7.51 | 9.64 | 8.21 | 10.23 | 9.63 | 11.29 | 8.79 | 9.47 | 10.90 | 10.57 | 7.28 | 9.44 |
| Na2O | 3.50 | 5.29 | 3.69 | 3.70 | 4.00 | 3.59 | 2.74 | 3.35 | 2.91 | 2.79 | 3.03 | 3.38 | 2.69 |
| K2O | 0.19 | 0.35 | 0.34 | 0.25 | 0.15 | 0.87 | 0.38 | 0.53 | 0.21 | 0.34 | 0.20 | 0.42 | 0.37 |
| P2O5 | 0.14 | 0.14 | 0.10 | 0.15 | 0.12 | 0.13 | 0.13 | 0.16 | 0.12 | 0.14 | 0.14 | 0.12 | 0.14 |
| Loi+ | 1.90 | 1.89 | 1.04 | 1.22 | 1.90 | 0.55 | 0.93 | 0.34 | 0.22 | 0.43 | 0.75 | 2.06 | 1.51 |
| H2O- | 0.29 | 1.49 | 0.80 | 1.38 | 0.26 | 0.78 | 1.70 | 0.41 | 1.42 | 0.77 | 2.16 | 0.97 | 0.26 |
| Total | 100.19 | 100.00 | 99.93 | 99.96 | 100.14 | 100.09 | 100.09 | 100.24 | 100.18 | 100.16 | 99.55 | 100.64 | 99.98 |
| Co | 36.7 | 30.6 | 37.8 | 33.3 | 35.2 | 37.2 | 42.4 | 31.1 | 35.4 | 46.9 | 51.1 | 29.3 | 39.0 |
| Cr | 201 | 125 | 270 | 93 | 133 | 266 | 334 | 142 | 190 | 287 | 129 | 228 | 154 |
| Ni | 65.3 | 31.6 | 87.0 | 30.8 | 37.0 | 73.7 | 99.6 | 25.0 | 47.7 | 63.0 | 64.3 | 68.5 | 35.2 |
| Ga | 18.2 | 19.0 | 16.6 | 19.7 | 19.1 | 19.3 | 17.2 | 20.3 | 17.4 | 17.4 | 19.1 | 17.6 | 18.1 |
| Sc | 44.4 | 30.5 | 37.4 | 36.6 | 29.5 | 30.1 | 38.1 | 31.7 | 38.4 | 39.1 | 39.2 | 31.8 | 37.9 |
| V | 323 | 356 | 300 | 377 | 267 | 275 | 313 | 370 | 307 | 320 | 343 | 297 | 356 |
| Ba | 26.9 | 23.9 | 15.5 | 12.4 | 161.9 | 45.0 | 21.2 | 27.4 | 20.2 | 23.6 | 9.6 | 32.1 | 77.6 |
| Rb | 2.33 | 7.62 | 3.93 | 2.19 | 2.45 | 7.86 | 4.84 | 14.77 | 1.43 | 5.96 | 2.16 | 8.10 | 5.45 |
| Sr | 162 | 102 | 205 | 139 | 169 | 149 | 152 | 154 | 131 | 155 | 140 | 125 | 131 |
| Y | 37.9 | 34.8 | 27.7 | 38.7 | 37.1 | 40.6 | 32.2 | 40.5 | 31.7 | 40.4 | 35.5 | 28.2 | 34.0 |
| Zr | 111 | 107 | 75 | 116 | 106 | 138 | 88 | 124 | 93 | 116 | 100 | 84 | 97 |
| Nb | 2.73 | 1.76 | 1.22 | 1.66 | 1.83 | 1.75 | 1.32 | 1.81 | 1.71 | 2.22 | 1.80 | 1.34 | 1.86 |
| Cs | 0.06 | 0.28 | 0.05 | 0.02 | 0.15 | 0.05 | 0.08 | 0.47 | 0.04 | 0.08 | 0.01 | 0.33 | 0.26 |
| Hf | 2.82 | 2.68 | 1.99 | 3.10 | 2.89 | 3.21 | 2.27 | 2.85 | 2.42 | 2.54 | 2.81 | 2.26 | 2.62 |
| Ta | 0.180 | 0.110 | 0.081 | 0.113 | 0.129 | 0.106 | 0.090 | 0.111 | 0.117 | 0.121 | 0.146 | 0.091 | 0.127 |
| Pb | 0.33 | 0.40 | 0.53 | 0.74 | 0.22 | 0.51 | 0.60 | 0.58 | 0.90 | 0.86 | 0.80 | 0.37 | 0.75 |
| Th | 0.195 | 0.255 | 0.185 | 0.296 | 0.193 | 0.295 | 0.210 | 0.308 | 0.251 | 0.242 | 0.227 | 0.247 | 0.215 |
| U | 0.081 | 0.106 | 0.197 | 0.184 | 0.091 | 0.236 | 0.701 | 0.105 | 0.100 | 0.180 | 0.097 | 0.109 | 0.091 |
| La | 4.29 | 3.85 | 2.82 | 4.02 | 3.48 | 4.51 | 3.23 | 4.38 | 3.45 | 4.06 | 3.75 | 2.89 | 3.23 |
| Ce | 12.93 | 11.53 | 8.57 | 12.61 | 10.96 | 13.98 | 10.09 | 13.61 | 10.58 | 12.25 | 11.31 | 8.82 | 9.77 |
| Pr | 2.17 | 1.97 | 1.48 | 2.15 | 1.94 | 2.35 | 1.76 | 2.26 | 1.78 | 2.07 | 1.96 | 1.50 | 1.72 |
| Nd | 11.34 | 10.29 | 8.00 | 11.52 | 10.61 | 12.05 | 9.52 | 12.08 | 9.51 | 10.88 | 10.54 | 8.13 | 9.32 |
| Sm | 3.84 | 3.55 | 2.71 | 3.96 | 3.68 | 4.14 | 3.21 | 3.99 | 3.37 | 3.68 | 3.56 | 2.79 | 3.26 |
| Eu | 1.42 | 1.27 | 1.01 | 1.38 | 1.32 | 1.38 | 1.20 | 1.39 | 1.11 | 1.26 | 1.28 | 1.02 | 1.16 |
| Gd | 4.92 | 4.53 | 3.64 | 5.01 | 4.86 | 5.04 | 4.22 | 5.11 | 4.21 | 4.60 | 4.95 | 3.76 | 4.51 |
| Tb | 0.926 | 0.834 | 0.659 | 0.924 | 0.897 | 0.919 | 0.772 | 0.952 | 0.802 | 0.897 | 0.898 | 0.683 | 0.798 |
| Dy | 6.00 | 5.30 | 4.26 | 5.98 | 5.94 | 5.94 | 5.00 | 5.87 | 5.00 | 5.46 | 5.69 | 4.49 | 5.22 |
| Ho | 1.29 | 1.16 | 0.94 | 1.32 | 1.34 | 1.32 | 1.10 | 1.30 | 1.15 | 1.21 | 1.20 | 1.01 | 1.14 |
| Er | 3.72 | 3.37 | 2.71 | 3.84 | 3.91 | 3.83 | 3.25 | 3.92 | 3.29 | 3.57 | 3.61 | 2.94 | 3.43 |
| Tm | nd | nd | nd | nd | nd | nd | nd | nd | nd | nd | 2.68 | nd | 3.29 |
| Yb | 3.47 | 3.25 | 2.54 | 3.67 | 3.53 | 3.62 | 2.95 | 3.65 | 3.02 | 3.37 | 3.47 | 2.74 | 3.25 |
| Lu | 0.514 | 0.489 | 0.381 | 0.556 | 0.553 | 0.545 | 0.442 | 0.561 | 0.463 | 0.512 | 0.535 | 0.404 | 0.482 |

| Sample | MK-34 | MK-38 | MK-33 | PMB32008 | PMB32005 | PMB32111 | MK-37 | PMB15856 | MK-45 | MK-3 | MK-4 | MK-8 | MK-41 |
|---------------------------------|------------------|------------------|-----------|-----------|-----------|---------------|-----------|--------------|-----------|-------------|-------------|-------------|------------------|
| Name | doleritic basalt | doleritic basalt | dolerite | dolerite | basalt | pillow basalt | basalt | Basalt glass | dolerite | andesite | basalt | dolerite | doleritic basalt |
| Affinity | IAT | IAT | IAT | IAT | TB | TB | BAB | IAT | MORB | IAT | IAT | IAT | BAB |
| Location | Runaway E | Runaway E | Runaway E | Runaway E | Runaway E | Runaway E | Runaway E | Lottin Pt | Lottin Pt | Matakaoa Pt | Matakaoa Pt | Matakaoa Pt | Pukeamaru S |
| SiO ₂ | 51.16 | 51.90 | 52.03 | 50.72 | 53.31 | 48.43 | 50.18 | 48.61 | 49.45 | 60.26 | 50.17 | 56.28 | 48.59 |
| TiO ₂ | 1.76 | 1.05 | 1.30 | 1.27 | 1.37 | 0.89 | 1.51 | 1.18 | 1.66 | 1.42 | 1.52 | 1.76 | 1.59 |
| Al ₂ O ₃ | 14.65 | 15.13 | 14.77 | 15.13 | 18.11 | 22.82 | 15.19 | 14.71 | 14.45 | 13.01 | 14.43 | 15.03 | 14.66 |
| Fe ₂ O _{3T} | 12.65 | 10.95 | 11.44 | 11.27 | 7.58 | 6.72 | 11.36 | 10.13 | 12.08 | 8.46 | 11.75 | 9.75 | 10.84 |
| MnO | 0.20 | 0.18 | 0.19 | 0.18 | 0.09 | 0.11 | 0.17 | 0.15 | 0.22 | 0.12 | 0.18 | 0.17 | 0.18 |
| MgO | 5.58 | 6.48 | 6.09 | 6.08 | 1.67 | 2.51 | 6.55 | 5.85 | 5.96 | 3.31 | 5.14 | 3.73 | 6.22 |
| CaO | 9.80 | 10.20 | 9.88 | 9.66 | 9.71 | 8.66 | 10.84 | 8.89 | 8.25 | 7.80 | 9.43 | 8.53 | 8.06 |
| Na ₂ O | 3.08 | 2.80 | 3.00 | 3.24 | 2.67 | 4.54 | 2.59 | 1.80 | 4.40 | 2.99 | 2.90 | 3.65 | 4.61 |
| K ₂ O | 0.09 | 0.17 | 0.18 | 0.17 | 0.74 | 1.51 | 0.16 | 0.19 | 0.17 | 0.18 | 0.43 | 0.41 | 0.60 |
| P ₂ O ₅ | 0.14 | 0.08 | 0.09 | 0.10 | 0.34 | 0.12 | 0.17 | 0.11 | 0.14 | 0.13 | 0.14 | 0.19 | 0.16 |
| Loi+ | 0.99 | 1.06 | 1.08 | 1.43 | 3.68 | 1.81 | 1.35 | 4.57 | 2.87 | 0.73 | 2.75 | 0.40 | 1.75 |
| H ₂ O- | 0.09 | 0.06 | 0.13 | 0.39 | 0.21 | 0.35 | 0.08 | 2.89 | 0.55 | 1.72 | 1.01 | 0.13 | 2.68 |
| Total | 100.19 | 100.06 | 100.18 | 99.66 | 99.49 | 98.47 | 100.14 | 99.08 | 100.22 | 100.16 | 99.85 | 100.02 | 99.94 |
| Co | 37.1 | 36.3 | 38.8 | 34.5 | 30.8 | 38.3 | 36.9 | 42.9 | nd | 37.1 | 34.6 | 25.3 | 36.2 |
| Cr | 151 | 146 | 137 | 179 | 134 | 253 | 241 | 215 | 129 | 125 | 166 | 75 | 223 |
| Ni | 39.5 | 42.1 | 40.8 | 43.1 | 21.5 | 56.8 | 69.5 | 63 | 44.5 | 26.8 | 39.2 | 16.9 | 55 |
| Ga | 20.6 | 16.8 | 17.6 | 17.9 | 20.6 | 17.0 | 18.2 | 16.00 | 19.2 | 17.3 | 18.7 | 21.3 | 18.13 |
| Sc | 38.9 | 41.3 | 43.0 | 44.2 | 18.8 | 22.1 | 35.7 | 42 | 41.4 | 29.9 | 34.73 | 31.9 | 36.0 |
| V | 380 | 313 | 346 | 333 | 182 | 86 | 324 | 330 | 362 | 349 | 352 | 435 | 333 |
| Ba | 21.7 | 32.0 | 31.9 | 51.4 | 133.1 | 60.9 | 28.1 | 32.2 | 183 | 24.7 | 32.3 | 30.3 | 72.1 |
| Rb | 0.86 | 2.49 | 2.69 | 2.18 | 13.47 | 27.88 | 2.33 | 2.72 | 2.38 | 1.62 | 3.50 | 17.0 | 4.72 |
| Sr | 123 | 104 | 118 | 156 | 422 | 331 | 141 | 114 | 176 | 141 | 134 | 149 | 126 |
| Y | 38.1 | 25.4 | 31.8 | 30.9 | 55.4 | 32.4 | 35.6 | 30.16 | 38.3 | 33.20 | 37.52 | 43.48 | 37.2 |
| Zr | 95 | 62 | 77 | 79 | 139 | 64 | 115 | 83.7 | 114 | 106 | 113 | 134 | 115 |
| Nb | 1.43 | 0.79 | 0.88 | 0.97 | 19.36 | 4.57 | 2.497 | 1.30 | 2.77 | 1.500 | 1.604 | 1.718 | 2.29 |
| Cs | 0.09 | 0.14 | 0.14 | 0.06 | 0.25 | 0.96 | 0.100 | 0.09 | nd | 0.032 | 0.052 | 0.510 | 0.29 |
| Hf | 2.71 | 1.76 | 2.26 | 2.21 | 2.38 | 1.53 | 2.689 | 2.35 | nd | 2.689 | 2.866 | 3.395 | 2.82 |
| Ta | 0.103 | 0.051 | 0.064 | 0.067 | 1.032 | 0.238 | 0.170 | 0.44 | nd | 0.091 | 0.108 | 0.122 | 0.15 |
| Pb | 0.55 | 0.08 | 0.31 | 1.10 | 3.36 | 0.60 | 0.302 | 25.8 | 2.17 | 0.535 | 0.757 | 0.446 | 0.34 |
| Th | 0.175 | 0.136 | 0.158 | 0.180 | 1.350 | 0.205 | 0.279 | 0.23 | nd | 0.233 | 0.280 | 0.364 | 0.21 |
| U | 0.072 | 0.061 | 0.060 | 0.104 | 0.633 | 0.325 | 0.83 | 0.16 | 1.15 | 1.96 | 2.05 | 1.53 | 0.09 |
| La | 3.17 | 1.90 | 2.15 | 2.52 | 34.33 | 10.06 | 4.389 | 2.98 | 0.74 | 3.833 | 3.967 | 4.961 | 3.90 |
| Ce | 10.33 | 6.13 | 7.32 | 8.49 | 71.99 | 9.88 | 12.72 | 9.30 | 10.93 | 11.73 | 12.28 | 14.93 | 12.2 |
| Pr | 1.86 | 1.11 | 1.36 | 1.48 | 9.16 | 2.70 | 2.095 | 1.57 | nd | 2.01 | 2.10 | 2.53 | 2.11 |
| Nd | 10.07 | 6.19 | 7.65 | 8.30 | 36.67 | 12.33 | 11.12 | 8.56 | nd | 10.43 | 11.19 | 13.43 | 10.86 |
| Sm | 3.70 | 2.34 | 2.82 | 3.02 | 8.12 | 3.19 | 3.70 | 3.00 | nd | 3.60 | 3.73 | 4.35 | 3.78 |
| Eu | 1.34 | 0.82 | 1.09 | 1.11 | 2.32 | 1.26 | 1.25 | 1.08 | nd | 1.25 | 1.32 | 1.54 | 1.31 |
| Gd | 4.76 | 3.23 | 3.93 | 4.09 | 8.55 | 4.14 | 4.83 | 4.03 | nd | 4.54 | 4.88 | 5.64 | 4.75 |
| Tb | 0.919 | 0.582 | 0.735 | 0.747 | 1.43 | 0.64 | 0.86 | 0.75 | nd | 0.82 | 0.89 | 1.05 | 0.88 |
| Dy | 6.17 | 3.97 | 5.03 | 4.81 | 8.68 | 4.07 | 5.71 | 4.71 | nd | 5.30 | 5.92 | 6.74 | 5.65 |
| Ho | 1.33 | 0.92 | 1.10 | 1.06 | 1.78 | 0.91 | 1.26 | 1.05 | nd | 1.16 | 1.29 | 1.49 | 1.24 |
| Er | 3.90 | 2.73 | 3.24 | 3.23 | 5.01 | 2.69 | 3.60 | 3.22 | nd | 3.43 | 3.85 | 4.30 | 3.62 |
| Tm | nd | nd | nd | 2.65 | 2.97 | 2.89 | nd | 2.62 | nd | nd | nd | nd | nd |
| Yb | 3.60 | 2.54 | 3.09 | 3.02 | 4.13 | 2.42 | 3.33 | 3.01 | nd | 3.21 | 3.60 | 4.04 | 3.43 |
| Lu | 0.546 | 0.388 | 0.475 | 0.456 | 0.617 | 0.364 | 0.501 | 0.45 | nd | 0.488 | 0.544 | 0.595 | 0.52 |

Table 3: Sr and Nd isotopic compositions of Matakaoa (East Coast) and Tangihua (Northland) mafic rocks. All the isotopic ratios are back-calculated at 80Ma which is the average fossil age of Matakaoa Volcanics; using a younger age does not significantly change the results.

| | | | | | | | | | | | | | |
|--|------------|------------|---------------|------------|------------|------------|---------------|------------|------------|------------|------------|---------------|---------------|
| Sample n° | MK- 4 | MK-12 | MK-16 | MK-38 | MK-37 | MK-21 | MK-35 | 49211 | 49164 | 49183 | 49186 | 47648 | 47662 |
| Rock type | Basalt | Basalt | Pillow basalt | Basalt | Basalt | Teschenite | Pillow basalt | Basalt | Glass | Basalt | Basalt | Pillow basalt | Pillow basalt |
| Geochemical type | IAT | IAT | IAT | IAT | BAB | MORB | MORB | IAT | IAT | BAB | BAB | BAB | BAB |
| Terrane | Matakaoa | Matakaoa | Matakaoa | Matakaoa | Matakaoa | Matakaoa | Matakaoa | Tangihua | Tangihua | Tangihua | Tangihua | Tangihua | Tangihua |
| Assumed age (Ma) | 80 | 80 | 80 | 80 | 80 | 80 | 80 | 80 | 80 | 80 | 80 | 80 | 80 |
| Sm (ppm) | 3.73 | 3.68 | 3.96 | 2.34 | 3.67 | 5.24 | 4.65 | 2.32 | 4.09 | 3.08 | 3.55 | 4.69 | 4.39 |
| Nd (ppm) | 11.2 | 10.9 | 11.5 | 6.18 | 11.1 | 15.8 | 14.4 | 6.19 | 11.5 | 9.13 | 11.0 | 14.8 | 13.9 |
| Rb (ppm) | 3.50 | 5.96 | 2.19 | 2.49 | 2.33 | 5.98 | 1.80 | 0.67 | 2.51 | 3.63 | 2.18 | 2.13 | 1.95 |
| Sr (ppm) | 134 | 155 | 138 | 104 | 141 | 172 | 120 | 120 | 86.4 | 143 | 206 | 130 | 105 |
| (¹⁴³ Nd/ ¹⁴⁴ Nd) _m | 0.51300 | 0.51306 | 0.51305 | 0.51304 | 0.51304 | 0.51306 | 0.51305 | 0.51306 | 0.51305 | 0.51306 | 0.51306 | 0.51306 | 0.51304 |
| (¹⁴³ Nd/ ¹⁴⁴ Nd) _i | 0.51290 | 0.51295 | 0.51294 | 0.51292 | 0.51293 | 0.51295 | 0.51294 | 0.51293 | 0.51293 | 0.51295 | 0.51295 | 0.51296 | 0.51294 |
| εNdi | 7.1 | 8.1 | 7.8 | 7.5 | 7.8 | 8.1 | 8.0 | 7.8 | 7.8 | 8.1 | 8.2 | 8.3 | 7.9 |
| (⁸⁷ Sr/ ⁸⁶ Sr) _m | 0.70298 | 0.70222 | 0.70304 | 0.70284 | 0.70284 | 0.70452 | 0.70280 | 0.70361 | 0.70299 | 0.70320 | 0.70349 | 0.70295 | 0.70268 |
| (⁸⁷ Sr/ ⁸⁶ Sr) _i | 0.70290 | 0.70210 | 0.70299 | 0.70277 | 0.70279 | 0.70441 | 0.70276 | 0.70359 | 0.70289 | 0.70311 | 0.70345 | 0.70290 | 0.70262 |




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig.5






Fig.6








Fig.7






Fig.9




Fig.10

