

The "Bois du Peu" thrust sheets (external French Jura mountains):re-examining the concept of "Fault-Fold"

Grégory Bièvre, Eric Mercier

► To cite this version:

Grégory Bièvre, Eric Mercier. The "Bois du Peu" thrust sheets (external French Jura mountains):re-examining the concept of "Fault-Fold". 2010. insu-00442191v2

HAL Id: insu-00442191 https://insu.hal.science/insu-00442191v2

Preprint submitted on 4 Oct 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1	The "Bois du Peu" thrust sheets (external French Jura mountains): re-examining
2	the concept of "fault-folding"
3	
4	Grégory Bièvre (1,2,*) and Éric Mercier (3)
5	
6	
7	1. Centre d'Études Techniques de l'Équipement de Lyon, Laboratoire Régional d'Autun, BP 141, 71404 Autun
8	cedex, France.
9	2. Laboratoire de Géophysique Interne et Tectonophysique, Université de Grenoble, BP 53, 38041 Grenoble
10	cedex 9, France
11	3. Laboratoire de Planétologie et Géodynamique, Université de Nantes, Géodynamique, CNRS, 2 rue de la
12	Houssinière, BP 92208, 44322 Nantes cedex 3, France.
13	
14	* Corresponding author (gregory.bievre@equipement.gouv.fr; Tel: +33-385-86-67-90)
15	
16	
17	Abstract
18	Significant reconnaissance field work along a road project which crosses the site of the "Bois du Peu" thrust
19	sheets (near Besançon, Eastern France), provides us the opportunity to re-examine the concept of "fault-folding"
20	(in french, "faille-pli") which was introduced by Glangeaud (1944) to account for the observed tectonics of the
21	Jura mountains and, more specifically, external Jura. From a theoretical point of view, we contend that this
22	concept is incompatible with general principles of balanced cross-sections and has to be rejected. We show that
23	in the "Bois de Peu" area, data fit with a deformation model which associates several modes of folding (fault-
24	propagation fold and fault-bend fold). The decollement level related to these folds is located into Keuper strata,
25	Oxfordian-Argovian levels being used locally as a secondary decollement level.
26	
27	Keywords: Jura mountains, "fault-fold", décollement tectonics, balanced cross- sections, forward modelling.

29 1 Introduction

The concept of "faille-pli" (in english "fault-fold"; not to be mistaken with the french term of "pli-faille" which means fault-related fold) was introduced by Glangeaud (1944) to account for the observed tectonics of the Jura mountains and, more specifically, external Jura. The latter consists of kilometer-wide strips (called bundles) which are densely folded and faulted. Relatively tabular sections (called plateaus) are found on both sides of these bundles, with the topographically higher ones lying to the East (Fig. 1).

Glangeaud (1944) proposed that the bundles in these carbonate strata resulted from a process he named "faultfold" and which is illustrated in figure 2. According to him, a pre-existing normal fault (inherited from a previous distension phase) would have elevated the eastern bloc (relative to the western bloc). Then, during the following compression phase, this eastern bloc covered the topographic surface below on the hanging wall block. Hence, the normal fault turns into a thrust fault and the distorted strata draw a fold structure which spills in the direction of the deformation.

Following a field-trip to the Jura that took place in 1951, this morpho-structural concept became very popular in the French geological community. Glangeaud (1951) illustrated his proposed concept on the cross-sections of the Besançon bundle in the "Bois du Peu" area, 2 km south of the city of Besançon. Later authors such as Caire (1963), Chauve and Perriaux (1974), Chauve (1975) then worked on these sections and further detailed them, thus helping to make the "Bois du Peu" area the reference location for the concept of "fault-folding".

46

The concept of "fault-folding" remained quite popular for a long time in French geological literature. Several reference books applied the concept to the Jura mountains without discussion (Aubouin 1973, Mattauer 1973, Foucault and Raoult 1980, Dercourt et al. 2006). The concept remained dominant for the Jura mountains until at least the end of the 1980s (Chauve 1987).

51 During the same period, however, the anglophone (in the sense of non-francophone) community completely 52 ignored the fault-fold concept. Since the end of the 1960s, their approach to the tectonics of the external zones 53 has been based on the concept of balanced cross-sections (Wilson and Stearns 1958, Bally et al 1966, 54 Dahlstrom 1969, to cite only a few pioneering works). Applying this concept, some authors (Laubscher 1961, 55 Mugnier and Vialon 1986, Endignoux and Mugnier 1990, Zoetemeijer and Sassi 1992, Martin and Mercier 1996, 56 Meyer 2000) proposed balanced cross-sections of the Jura bundles without taking into account the concept of 57 "fault-folding". Moreover, Madritsch et al. (2008, 2010) recently proposed that the Besançon zone is affected by 58 thin-skinned tectonics only.

59

De facto, even if this concept has never been strongly discussed in the literature, it appears inconsistent with the principle of balanced cross-sections and therefore must be abandoned. Since Glangeaud's work (Glangeaud 1951), the "Bois du Peu" area has never been reinterpreted. Geological study and civil engineering works conducted for the Besançon higway by-pass project provide new outcrops and subsurface drillings (corings and tunnels) in the "Bois du Peu" area. This new data provides a good opportunity to re-examine and re-interpret this reference location.

66

The aim of this paper is first to re-examine the concept of "fault-fold" and to show why it is inconsistent with the concept of balanced cross-sections. The implications of this on Glangeaud's concept will be re-examined. Secondly, new data concerning the "Bois du Peu" area will be presented. This data allows a new detailed geological map to be created. Finally, it will be proposed that a new fault-related fold model is consistent with field observations.

72

73 2 Geological context of the study area

74 The Jura is an arched moutain belt located NW of the Swiss molasse basin (Fig. 1a). The study zone is located 75 in the outer Jura chain and is mainly made of Jurassic carbonate formations (Fig. 3). These sedimentary strata 76 are arranged in relatively tabular plateaus separated by severely folded and faulted narrow elongated bundles 77 (Fig. 1b). This arrangement is the result of the "multi-phased" tectonic history of the area, where two main phases 78 can be distinguished. The first phase was extensional, with an E-W sense and has an Oligocene age. This 79 resulted in a general westward downstepped blocks geometry which corresponds to the present-day look of the 80 massif. The second phase was compressive, directed towards the NW and is Miocene. The deformation related 81 to the compressive phase was principally located at the boundaries between the plateaus, thus generating 82 bundles, characterized by folds and thrust faults (Glangeaud 1951, Caire 1963, Bergerat et al 1990, Guellec et al 83 1990, Lacombe and Angelier 1993, Martin and Mercier 1996, Homberg et al 1999). Furthermore, the major 84 oligocene-inherited meridian faults induced a leftward strike-slip motion which allowed the panels to slide. 85 According to palaemoagnetic recordings, this translation movement did not induce significant rotation of the 86 structures (Gehring et al 1991).

The Besançon bundle constitues one of the narrow strips. In the study area, this bundle is oriented SW-NE and is about 4 km wide. It is bounded by the Besançon/Thise plateau to the NW and by the topographically higher Montrond plateau to the SE (Fig. 1b). At the outcrop, this bundle is made of two parallel antinclines. To the NW, the Citadel anticline shows a symetric or slightly SE overturned geometry. To the SE, the Mercureaux anticline is strongly dyssimetric and overturned towards the NW (Fig. 1b). It is the Mercureaux anticline, along with its associated Montfaucon fault (Dreyfuss and Kuntz 1968), which have been interpreted as one of the best examples for the "Fault-fold" structure by Glangeaud (1951).

95

96 The local geological series consists of alternating soft (clay and marls) and hard levels (limestones, dolomites 97 and sandstones) and is shown in figure 3. The main decollement level is located within the Triassic gypsum-rich 98 strata. Secondary decollement levels may be found within soft Jurassic strata (Fig. 3): Pliensbachian-Aalenian, 99 Oxfordian sensu stricto-Argovian and middle Sequanian. Regional sub-stages denominations (Dreyfuss and 100 Kuntz 1968) have been kept to distinguish between the mechanically variable strata.

101

102 The Besançon highway by-pass (Maurin 2001, Bièvre 2007) crosses the Mercureaux anticline in the "Bois du 103 Peu" area. The geological complexity of the site lead to dense prospectings: several kilometers of boreholes, and 104 logging (gamma-ray, microseismics, digital camera) as well as mechanical in situ and laboratoray tests. 105 Furthermore, a reconnaissance gallery has been drilled that crosses the base of one of the bundles. The 106 integration of these new data allow to detail the orientation and dip of faults as well as the lithology of the bedrock 107 underlying the top-soil layer, especially along the Mercureaux anticline axis (Fig. 1b) made of soft clayey and 108 marly formations. The data allows the construction of a detailed geological map of the area as well as an original 109 synthetic cross-section of the bundle which was crossed by the reconnaissance gallery.

110

111 3 The concept of "fault-fold" in the light of balanced cross-sections theory

The theory of balanced cross-sections is based on the assumption that thanks to the law of conservation of matter, the amount of material remains constant during tectonic deformations. In faults and folds belts, it is often possible to work on cross-sections (see complete discussion in Marshak and Woodward (1988) for example) and, in this way, to transform the law of conservation of matter into a law of conservation of surfaces (i.e. during deformation, the surface of each bed remains constant). To verify this conservation, it is necessary to set boundaries on the system studied and, consequently, to discuss the conditions of these boundaries.

Figure 4 shows that surface conservation requires that, during the growth of a "fault-fold", the boundaries of the system undergo differential simple shear. This diagram shows that only the upper layers of the upper block suffer simple shear during horizontal shortening. More specifically, only the beds which are in elevation over the top of lower block after the first deformation (normal fault) suffer simple shear. This is problematic because, obviously, the boundary conditions are controled by rear area deformation conditions and not by internal parameters (normal fault offset) as the figure might falsely imply.

Furthermore, Figure 5 shows that the deformations are amortized into the structure and are not transfered forward from one block to another. Accordingly, this model fails to explain the succession of several "fault-fold" strutures on a unique cross-section as is the case in the Jura (Chauve 1987).

128

A high proportion of authors who recently worked on the Jura (see above), appear to have abandoned the concept of "fault-fold". None of these authors have justified this abandonment however, and the concept of faultfolding therefore goes unmentioned. We have shown that the concept of "fault-folding" is incompatible with the theory of balanced cross-sections, this incoherence being the reason for the abandonement. The "Bois de Peu "area, the reference location for the concept of "fault-folding", therefore has to be explained otherwise. The following sections show that it is possible to reinterpret this structure using the concepts of "fold and fault-belts" tectonics.

136

137 4 Results

138 4.1 Geological field data

Surveys conducted for the highway by-pass study provided a large amount of geological data (Maurin 2001, Bièvre 2007). Combined with detailed field observations, these data allow us to produce a new geological map based on a previously established one (Dreyfuss and Kuntz 1968). There is no fundamental change between the two maps, but the one that is proposed here is much more detailed (Fig. 6) due to new available data. In combination with these field observations, a reconnaisance gallery was drilled through the base of one of the "Bois du Peu" thrust sheets and an interpretative cross-section was built (Fig. 7; modified from CETU 1999).

145

The proposed cross-section for this work is located one km SW of the reference location. The geological map reveals that fault F2 dips towards the SE (as it has been revealed by corings and gamma-ray logging; Maurin 148 2001). Fault breccia was found in corings conducted along the road project to define the dip of F3 (Bièvre 2007). 149 Along with the gamma-ray logging in surrounding drillings, this reveals that the F3 fault (Montfaucon fault of 150 Dreyfuss and Kuntz 1968) slighlty dips towards the SE (Fig. 6). These two faults were previsously considered to 151 be subvertical (Fig. 8a; Glangeaud 1951, Caire 1963, Chauve and Perriaux 1974, Chauve 1975). Moreover, the 152 Mercureaux anticline axis is located a few hundred metres SE of F3 (Fig. 6; Bièvre, 2007). These two initial 153 observations are inconsistent with the "fault-fold" interpretation of the area ".

154

A vast outcrop composed of Triassic strata is present in contact with F3 along the road project, 200 m SE of the tunnel (location on Fig. 6). This Triassic outcrop is bordered by Pliensbachian strata (Belemnites as well as Ammonite Amaltheus margaritatus were found in corings). Associated with the presence of fault breccia, cartography and orientations of dips, these elements allow this Triassic outcrop to be considered as a tectonic flake forced against the Montfaucon fault (F3). The presence of such a flake constitutes an important argument to interpret F3 like a thrust fault seated within Triassic strata (Fig. 8b).

161

The hinge and inverted limb, situated NW of the Mercureaux anticline overthrust the very competent Jurassic beds. In first approximation, despite some irregularities (Fig. 7 and Fig. 8), this thrust is parallel to the autochton stratification. Previous authors had considered implicitly (compare Fig. 3 and Fig. 8a) or explicitly that the thrust surface was the Pontian topographic surface (surface of "Montrond"; Dreyfuss and Glangeaud 1950). This hypothesis is discussed further in section XXX.

167

Thrust F1 is not the only one to be locally parallel with the stratification. For example, thrust F2 is, from NW to SE, successively sub-parallel, oblique and again sub-parallel with the stratification(Fig. 8b). These particular relations refer to the ramp geometry. This observation, along with previous works on other bundles in the Jura (Endignoux and Mugnier 1990, Zoetemeijer and Sassi 1992, Martin and Mercier 1996, Meyer 2000), lead us to propose a kinematic scheme characterized by the development of folds-related ramps for the "Bois de Peu" area.

173

174 4.2 Kinematic modelling

175 Cross-section balancing has become a standard method for testing viability and admissibility of hypothetical 176 deep geometry. Many theoretical and applied works have focused on this method in thrust and fold belts. Several 177 approaches have been developed but, according to most of the authors, in the Jura, the "forward" method is the 178 most appropriate (Endignoux and Mugnier 1990, Zoetemeijer and Sassi 1992, Martin and Mercier 1996).

Martin and Mercier (1996) proposed a comprehensive discussion on the application of this method to a bundle of the Jura. To summarise, this method provides a viable and admissible kinematic pathway between an initial state (undeformed) and a final state (deformed). The need to respect the law of conservation of matter (1) between initial stage and final stage, and (2) between each kinematic step, strongly limits the number of possible solutions.

184

In practice, a trial and error process was used to build an image of the finite deformation which is consistent with field data. With this kind of problem solving process, there is a risk of neglecting alternative solutions. Hence, many tests were carried out to assess the influence of changes in calibration parameters.

188

In this study, we chose to work on an "average" cross-section. This section can be considered as representative of the whole area, and allows the elimination of local variations that can not be taken into account by modelling. The numerical solution is shown in figure 9.

192

Steps a, b and c: a "fault-propagation fold" (in the sense of Suppe 1985) gradually grows over a ramp deeplyseated in a decollement level located at the top of the Triassic strata. It has long been known (e.g. Glangeaud 1951, Caire 1963) that the Jura bundles are the result of the superposition of an Oligocene tectonic distension and a Miocene tectonic compression. Previous modelling works (Martin and Mercier 1996) showed that in the bundle, ramp initiation occurs systematically at the intersection between a decollement level and inherited normal faults. Surprisingly, there are no arguments in these works to link the initiation of the ramp with a normal fault.

199

Step d: the fault-propagation fold suffers a standard late evolution: transport on the flat (Mercier 1992; Mercier et al 1997). Usually, such evolution occurs when the ramp can no longer propagate upwards (when the ramp crosses very competent beds, for example), and seeps into an interbed level. It does not seem to be the case here, the very competent Jurassic series is already crossed and the Cretaceous series, thin and weakly competent, can not stop the propagation of the ramp. The simplest is to assume that the allochton slept upon the paleo-surface topography ("Montrond" surface; Dreyfuss and Glangeaud 1950).

206

207 Step e: The transport on the flat becomes increasingly difficult (friction increasing, blocking on local micro-

208 topography) and the mechanical conditions in autochton change because of the tectonic overload. A new thrust 209 plane is established.

The movement over this thrust creates a duplex which is transported under the fold. Modeling suggests that this duplex is deep-seated in a secondary detachment level in the Oxfordian-Argovian strata. This hypothesis is fully consistent with mechanical properties of these levels (Fig. 3) and field data (Fig. 6).

213

Step f: After a significant displacement, this new thrust is blocked in turn. Out-of-sequence thrusts occur from the existing ramps and through weakened areas of the structure (Mercier and Mansy 1995). The northern out-ofsequence fault corresponds to the "forelimb breakthrough" of fault-propagation folds (Mercier 1992). It isolates, between F1 and F3, a small thrust sheet with reverse polarity. We suggest that this thrust sheet is torn into several elements that are more or less carried forward in reponse to the movement of the allochton.

Finally, synchronically or not, growth of the Citadel anticline, located just NW of the section studied, affects the whole structure which is partly integrated into its SE limb. The final bending is not really taken into account by our modeling. The surface topography drawn in figure 9f is distorted from reality, but this adjustment imprecision does not affect the principle and the conclusions of our model. In fact, it only introduces an uncertainty on the geometric modeling of the out-of-sequence faults F4 and F4'.

224

The total shortening, of about 50 % (4 km), is significantly higher than what was calculated on the sections located further north accross the same bundle (Martin and Mercier 1996). This difference suggests small rotations of the plateaus during deformation.

228

229 5 Discussion and conclusions

Major geological reconnaissance for the Besançon by-pass took place in the reference location for the concept of "fault-folding" ("Bois du Peu" area; Glangeaud 1951). It provides firstly an opportunity to discuss this concept of "fault-folding", and to show its incompatibility with the theory of balanced cross-sections. Secondly, it allows us to propose a new structural evolution for this area (Fig. 8b and Fig. 9).

234

We show that available field data are consistent with a typical scenario of folds and thrust belt evolution, particularly characterized by the growth and evolution of fault-related folds deeply seated within Triassic strata. This is very similar to scenarios already proposed for other Jura bundles (e.g. Guellec et al 1990, Martin and Mercier 1996, Meyer 2000). In particular, we note the combination of various folding modes (fault-related folds with late evolution, duplex, etc.) in the same sector. This work, among many others, confirms the utility of the "forward" method in the study of the Jura tectonics. However, without syntectonic sedimentary markers, the sequence of deformation proposed remains, in the study case as elsewhere, poorly constrained. In the study area, the shortening is about 50% which is higher than what is known in other bundles. Finally, we note the importance of earlier morphological evolution in the development of some thrust faults wich slip onto a paleoerosion surface.

- 245
- 246
- 247
- 248

250

251 Aubouin J (1973) Précis de géologie, t.3. Tectonique, tectonophysique, morphologie. Dunod, Paris, France.

Bally AW, Gordy PL, Stewart GA (1966) Structure, seismic data and orogenic evolution of southern canadian
 rocky mountains. Bulletin of Canadian Petroleum Geology 14(3):337–381.

254 Bergerat F, Mugnier JL, Guellec S, Truffert C, M C, Damotte B, F R (1990) Deep structure of the Alps, Société

Géologique de France, chap Extensional tectonics and subsidence of the Bresse basin: an interpretation from
 ECORS data, pp 145–156.

257 Bièvre G (2007) élaboration d'un modèle géologique au coeur d'un anticlinal : contribution aux études

258 géotechniques d'un tracé routier et de ses ouvrages d'art (voie des mercureaux, besançon, est de la france).

259 Bulletin des Laboratoires des Ponts et Chaussées 267:1–16.

260 Caire A (1963) Livre à la mémoire de Pierre Fallot, Société Géologique de France, chap Problèmes de 261 tectonique et de morphologie Jurassiennes, pp 105–158.

262 CETU (1999) Galerie de reconnaissance du tunnel du bois du peu. document synthétique par 25 mètres. Tech.

rep., Center for Tunnel Studies, unpublished geological report.

264 Chauve P (1975) Jura. In: Guides Géologiques Régionaux, Masson, Paris. 216 pp.

265 Chauve P (1987) Aspect et évolutions géologiques du Bassin Parisien, Associations des Géologues du Bassin

Parisien, chap Le Jura, du Bassin Parisien à la Chaîne alpine, pp 133–141.

^{249 6} References

- 267 Chauve P, Perriaux J (1974) Géologie de la France, vol 2, Doin, chap Le Jura, pp 443–464.
- 268 Dahlstrom CDA (1969) Balanced cross-sections. Canadian Journal of Earth Sciences 6:743–757.
- Dercourt J, Paquet J, Thomas P, Langlois C (2006) Géologie, objets, méthodes et modèles, 12th edn. Dunod,
 Paris, France.
- 271 Dreyfuss M, Glangeaud L (1950) La vallée du doubs et l'évolution morphotectonique de la région bisontine.
- 272 Annales Scientifiques de l'Université de Besançon, Géologie V:1–16.
- 273 Dreyfuss M, Kuntz G (1968) Besançon. In: Carte géologique de la France au 1/50 000, BRGM, Orléans, France.
- Endignoux L, Mugnier JL (1990) The use of a forward kinematical model in the construction of balanced cross sections. Tectonics 9:1249–1262.
- 276 Foucault A, Raoult JF (1980) Dictionnaire de Géologie. Masson, Paris.
- 277 Gehring A, Keller P, Heller F (1991) Paleomagnetism and tectonics of the jura arcuate mountain belt in france 278 and switzerland. Tectonophysics 186:269–278.
- Glangeaud L (1944) Le rôle des failles dans la structure du jura externe (pincée, faille-plis et gouttières). Bulletin
 de la Société d'Histoire Naturelle du Doubs 51:17–33.
- Glangeaud L (1951) Réunion de la société dans le jura. Bulletin de la Société Géologique de France 6(1):763–
 872.
- Guellec S, Mugnier JL, Tardy M, F R (1990) Deep structure of the Alps, Société Géologique de France, chap
 Neogene evolution of the western Alpine foreland in the light of ECORS data and balanced cross-section, pp
- 285 165–184.
- Homberg C, Lacombe O, Angelier J, Bergerat F (1999) New constraints for indentation mechanisms in arcuate
 belts from the jura mountains, france. Geology 27:827–830.
- 288 Lacombe O, Angelier J (1993) Evolution tectonique du jura externe au cénozoïque et perturbations de 289 contraintes dans la zone transformante rhin-saône. Comptes Rendus de l'Académie des Sciences 290 317(2):1113–1120.
- Laubscher H (1961) Die fernschubhypothese der jurafaltung. Eclogae Geologicae Helvetiae 54:212–280.
- 292 Madritsch, H., Schmid, S. M. And Fabbri, O. (2008) Interactions between thin- and thick-skinned tectonics at the
- 293 northwestern front of the Jura fold-and-thrust belt (eastern France). Tectonics, 27, TC5005.
- 294 Marshak S, Woodward N (1988) Basic methods of Structural Geology, Prentice Hall, chap Introduction to cross-

- section balancing, pp 303–332.
- 296 Martin J, Mercier E (1996) Héritage distensif et structuration chevauchante dans une chaî ne de couverture:
- 297 apport de l'équilibrage par modélisation géométrique dans le jura nord-occidental. Bulletin de la Société

298 Géologique de France 167:101–110.

- 299 Mattauer M (1973) Les déformations des matériaux de l'écorce terrestre. Hermann, Paris, France.
- 300 Maurin P (2001) Viaduc des mercureaux, étude de la fondation d'un ouvrage d'art exceptionnel au droit d'une
- 301 faille majeure du faisceau bisontin. Bulletin des Laboratoires des Ponts et Chausées 233:77–86.
- 302 Mercier E (1992) Une évolution possible des chevauchements associés aux plis de propagation: le transport sur
- 303 le plat (modélisation et exemple). Bulletin de la Société Géologique de France 163(6):713–720.
- 304 Mercier E, Mansy JL (1995) Le blocage du transport sur le plat des plis de propagation: une cause possible des
- 305 chevauchements hors séquence. Geodynamica Acta 8(4):199–210.
- 306 Mercier E, Outtani F, Frizon de Lamotte D (1997) Late evolution of fault-propagation folds: principles and 307 example. Journal of Structural Geology 19:185–193.
- 308 Meyer M (2000) Structure du "diapir" de champfromier (jura, ain, france). Eclogae geologicae Helvetiae 93:221–
 309 229.
- 310 Mugnier JL, Vialon P (1986) Deformation and displacement of the jura cover on its basement. Journal of 311 Structural Geology 8:373–387.
- 312 Suppe J (1985) Principles of structural geology. Prentice Hall.
- Wilson CW, Stearns RG (1958) Structure of the cumberland plateau, tennessee. Geological Society of America
 Bulletin 69:1283–1296.
- Zoetemeijer R, Sassi W (1992) Thrust tectonics, Chapman and Hall, chap 2-D reconstruction of thrust evolution
 using the fault-bend method, pp 133–140.
- 317
- 318

320

321 List of figures

322

Figure 1: Location of the study area and strutural map of the Besançon area. a) Strutural map of the Jura mountains around Besançon showing the organization in plateaus and bundles (simplified from Madritsch et al. 2008). b) Detailed structural map of the study area (location on Fig. 1a) showing the main faults and folded structures (adapted from Dreyfuss and Kuntz 1968). CA: Citadelle anticline; MA: Mercureaux anticline; MF: Mercureaux fault; TF: Trochatey fault.

328

329 Figure 2: The kinematics of a "fault-fold" according to Glangeaud (1944).

330

Figure 3: Synthetic lithologic log of the Besançon area showing the alternation of clays/marls soft levels with hard
limestone layers. The main decollement level is located within the upper Keuper layers; Pliensbachian-Aalenian,
Oxfordian-Argovian and middle Sequanian layers may serve as secondary décollement levels. Oxf. s.s.:
Oxfordian sensu stricto. Arg.: Argovian. Raur.: Rauracian. Seq.: Sequanian.

335

Figure 4: An attempt to integrate the concept of "fault-fold" in a balanced cross-section. To balance the structure (same as step b on Fig. 2), layers have to be subject to a simple shear whose characteristics depend on the inherited fault net slip.

339

Figure 5: Non-balanced cross-section showing that a "fault-fold" is unable to transmit forward the deformation necessary to the growth of a second "fault-fold". Examination of this diagram shows that the upper part of the central flat can not undergo at the same 1) a moderate shear resulting from the deformation coming from the back and 2) a significant shear necessary to generate the forward (left) structure.

344

Figure 6: Geological map of the study area, location of the road works (bold dashed black line) and of the "Bois du Peu" tunnel (black rectangle). Coordinates are metric according to the French Lambert II system. BdP.T.: Bois de Peu tunnel and cross-section of figure 7. faults are named after Dreyfuss and Kuntz (1968). F1 to F5: faults. Map adapted from Dreyfuss and Kuntz (1968) and Bièvre (2007).

Figure 7: Geological cross-section of the "Bois du Peu' tunnel. Faults are labelled in the same manner as in
Fig. 6. Modified from CETU (1999).

352

Figure 8: Cross-sections of the "Bois du Peu" area. a: Interpretation in terms of a "fault-fold" (according to Glangeaud 1951, Chauve and Perriaux 1974). Faults are labelled in the same manner as in Fig. 6. The fault associated to the "fault-fold" structure corresponds to F1 and to the lower part of F3. The uppert part of F3 would be the result of a late reactivation Chauve (1975). b: Proposed interpretation according to new data and balanced cross-sections. Position of Figure 7 is indicated. See text for details.

358

Figure 9: Kinematic evolution of the Besançon bundle in the study area based on balanced cross-sections and using a forward modelling approach. Faults are labelled in the same manner as in Fig. 6. M.A.: Mercureaux anticline. BdP.T.S.: "Bois du Peu" thrust sheets. Step a to f: see text for details.

362





Figure 1





Figure 3









North



Figure 9