

Dynamic constraints on the crustal-scale rheology of the Zagros fold belt, Iran

Philippe Yamato, Boris J.P. Kaus, Frédéric Mouthereau, Sébastien Castelltort

► To cite this version:

Philippe Yamato, Boris J.P. Kaus, Frédéric Mouthereau, Sébastien Castelltort. Dynamic constraints on the crustal-scale rheology of the Zagros fold belt, Iran. Geology, 2011, 39 (9), pp.815-818. 10.1130/G32136.1 . insu-00623034

HAL Id: insu-00623034 https://insu.hal.science/insu-00623034

Submitted on 13 Sep 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Dynamic constraints on the crustal-scale rheology of the Zagros

2 fold belt, Iran

3 Philippe Yamato^{1,2}*, Boris J.P. Kaus^{2,3}, Frédéric Mouthereau⁴, and Sébastien Castelltort²

4 ¹Geosciences Rennes, UMR CNRS 6118, Université de Rennes 1, F-35042 Rennes Cedex,

5 France

6 ²Department of Earth Sciences, E.T.H. Zürich, Sonnegstrasse 5, 8092 Zürich, Switzerland

7 ³Department of Earth Sciences, University of Southern California, Los Angeles, USA

8 ⁴UPMC Univ. Paris 06, UMR CNRS 7193, ISTeP, 75252 Paris Cedex 5, France

9

10 *E-mail: philippe.yamato@univ-rennes1.fr

11 ABSTRACT

12 Thin-skinned fold-and-thrust belts are generally considered as the result of contractional 13 deformation of a sedimentary succession over a weak decollement layer. The resulting surface 14 expression frequently consists of anti- and synclines, spaced in a fairly regular manner. It is thus 15 tempting to use this spacing along with other geological constraints to obtain insights in the 16 dynamics and rheology of the crust on geological time scales. Here we use the Zagros Mountains 17 of Iran as a case study as it is one of the most spectacular, well-studied thin-skinned fold-and-18 thrust belts in the world. Both analytical and numerical models are employed to study what 19 controls fold-spacing and under which conditions folding dominates over thrusting. The models 20 show that if only a single basal décollement layer is present underneath a brittle sedimentary 21 cover, deformation is dominated by thrusting which is inconsistent with the data of Zagros Fold 22 Belt. If we instead take into account additional décollement layers that have been documented in

the field, a switch in deformation mode occurs and crustal-scale folding is obtained with the correct spacing and timescales. We show that fold spacing can be used to constrain the friction angle of the crust, which is ~5 degrees in Zagros Fold Belt. This implies that on geological timescales, the upper crust is significantly weaker than previously thought, possibly due to the effect of fluids.

28 INTRODUCTION

It is often assumed that fold belts can be explained by folding of a sedimentary layer above a basal detachment formed by a weak layer. As the spacing between folds in such belts is quite regular, we can consider them as a large-scale natural experiment of crustal deformation. Ideally, it should be possible to combine fold spacing with other geological data and theory to constrain parameters such as crustal rheology that are difficult or impossible to constrain from field observations alone.

The classical explanation of folds in fold belts is that they are due to a folding instability, which is well known for a homogeneous sedimentary sequence with either a power-law viscous or an elastic rheology (Schmalholz et al., 2002; Burg et al., 2004; Schmalholz, 2006). The dominant wavelength λ_{dom} , for a viscous power-law layer of viscosity η_{sed} and with exponent *n* overlying a linear viscous layer of viscosity η_{salt} , is given by (Schmalholz et al., 2002):

40
$$\lambda_{dom} = 3.1 \left(\frac{\eta_{sed}}{n\eta_{salt}}\right)^{\frac{1}{6}} \sqrt{\frac{H_{salt}}{H_{sed}}} H_{sed}$$
(1)

41 where H_{sed} and H_{salt} are the thicknesses of the sedimentary cover and of the salt, respectively. 42 The growth rate (q_{dom}) of this instability non-dimensionalized over the background strain rate 43 $\dot{\varepsilon}$ is given by (Schmalholz et al., 2002):

44
$$\frac{q_{dom}}{\dot{\varepsilon}} = 2.5n \left(\frac{\eta}{n\eta_{salt}}\right)^{\frac{1}{3}} \frac{H_{salt}}{H_{sed}} = 2.5n \left(\frac{\lambda_{dom}}{3.1H_{sed}}\right)^{2}$$
(2)

and a combination of numerical and analytical studies have shown that $q_{dom}/\dot{\epsilon}$ should be larger than ~20 for folding to form observable folds, rather than homogeneous thickening (e.g., Kaus et. al, 2008).

48 The Zagros Fold Belt of Iran constitutes a classical example of such a folded belt that is 49 geologically (e.g., Stocklin, 1968; Alavi, 2004; McQuarrie, 2004; Sherkati and Letouzey, 2004; 50 Mouthereau et al., 2007) and geophysically (e.g., Jahani et al., 2009; Hatzfeld and Molnar, 2010; 51 Nissen et al., 2010) well studied due to excellent exposure and extensive seismic and borehole 52 data from exploration. The main tectonic and stratigraphic units are summarized on Figure 1 and 53 show that a particular feature of the Zagros Fold Belt is a consistent spacing of folds with a 54 wavelength (λ_{dom}) of 14 ± 3 km. These folds are generally explained as detachment folding of 55 the post-Cambrian sedimentary sequence above a basal weak layer constituted by the Hormuz 56 salt. 57 The centroid depths of waveform-modeled earthquakes indicate that faulting is restricted 58 to two structural levels located in the competent sediment cover units at 5–6 km depth and within 59 the Precambrian basement at depth larger than 11 km down to depths of 30 km (e.g., Talebian 60 and Jackson, 2004; Nissen et al., 2010). Seismic reflection profiles (Jahani et al., 2009) and field

observations in the Fars region (Mouthereau et al., 2007) show a lack of major thrust faults
cutting the folded cover up to the surface. This confirms that detachment folding rather than
thrusting is the dominant deformation mode in the Zagros Fold Belt. In this aspect, the Zagros
Mountains differ from other fold-and-thrust belts such as the Jura Mountains, where large-offset

65 faults are continuous across the stratigraphic sequence, well imaged through seismic studies66 (Simpson, 2009).

67 Detachment folding theory should thus be perfectly applicable to the Zagros Fold Belt. 68 Equations (1) and (2) show that fold spacing depends strongly on the rheology of the overburden 69 and on the thickness of the basal salt layer. In the Zagros, a linear viscous overburden (n = 1) and 70 a viscosity contrast of 100 between salt and overburden, requires a salt layer thickness of \sim 7.8 71 km to fit the observed spacing of folds (Equation 1). Yet, seismic data indicates that the 72 thickness of the Hormuz salt is no more than 1 or 2 km (Jackson et al., 1990; Mouthereau et al., 73 2006; Jahani et al., 2009). If the sedimentary cover has a power law rheology instead, its power 74 law exponent should be $n \sim 23$ (Equation 2) to explain the data, which is considerably larger than 75 estimates from rock creep experiments (Ranalli, 1995). Large power law exponents are often 76 taken as evidence for a brittle rheology. Currently, however, there is no theory that can reliably 77 predict the spacing of detachment folds in the case of a brittle overburden.

There is thus presently no satisfactory explanation for (1) why deformation in the Zagros Fold Belt is dominated by folding and not by thrusting and for (2) what controls the spacing of folds and how it is linked to crustal rheology. In order to address this, we performed thermomechanical numerical simulations to study the dynamics of detachment folding in the presence of a brittle sedimentary cover.

83 NUMERICAL MODEL

To study the effect of using visco-elasto-plastic rheologies on crustal dynamics, we have performed a series of numerical experiments using the finite element code MILAMIN_VEP (e.g., Kaus, 2010 and GSA Data Repository DR1). The viscosities of the weak layers are assumed to be linear and constant, which is a reasonable approximation for the rheology of salt.

88	The brittle layers have a temperature-dependent rheology of limestone (see DR1), which
89	correspond to the majority of rocks within the sedimentary cover (Fig. 1C, Mouthereau et al.,
90	2007). A linear geotherm of 25 °C.km ⁻¹ is initially applied (see DR1). For the low-temperature
91	conditions of the Zagros Fold Belt, stresses are such that the rocks effectively deform in the
92	brittle regime. Our model domain is initially 200x7.225 km in size (see DR1). The top boundary
93	is a free surface with no erosion (see DR1) and a constant background strain rate of 10^{-15} s ⁻¹ is
94	applied at the right of the model box, which results in 15% shortening after 5.5 Myrs consistent
95	with geological constraints (see DR1). All other sides of the model have free slip conditions.
96	Finally, to initiate the folding, the interface between the salt and the overburden rocks has
97	random noise with maximum amplitude of 100 m. Model simulations are performed for 5.5
98	Myrs, after which results are interpreted.

99 RESULTS FROM NUMERICAL SIMULATIONS

100 With a 1.5 km-thick single basal detachment layer underlying a homogeneous brittle 101 sedimentary cover, the models develop faults rather than folds (Fig. 2B). Such faults develop at 102 early stages with a spacing that is approximately twice the brittle layer thickness. Subsequent 103 deformation is locked around these folds that have a box-fold geometry. Compared to the Zagros 104 Fold Belt, we thus obtain a too large wavelength and an incorrect deformation style. Additional 105 simulations where we varied the frictional parameters of the crust, or the viscosity of the salt 106 layer gave similar results (see Fig. DR1). We thus infer that it is impossible to reproduce the 107 observed finite wavelength of Zagros Fold Belt folds (Fig. 1) by considering only one weak basal 108 décollement layer, unless this layer has an unrealistically large thickness.

109 A detailed look at the stratigraphic column, however, reveals that the sedimentary cover 110 is not rheologically homogeneous. Instead there are several layers that are composed of relatively

111	weak rocks such as evaporites or shales (Fig. 1B, C, see detailed descriptions in McQuarrie,
112	2004; Sherkati et al., 2006 and Mouthereau et al., 2007). A second set of simulations took this
113	fine-scale rheological structure into account (Fig. 3). The results are remarkably different from
114	the previous experiments: rather than being fault-dominated, deformation is now achieved by
115	folding, with a final wavelength similar to the one observed in the Zagros Fold Belt (Fig. 3). An
116	analysis of the simulation shows that the spacing of the folds is fixed at a very early stage, after
117	which the individual structures grow without clear geometric pattern, in accordance with field
118	constraints (Mouthereau et al., 2007). The initial fine-scale rheological stratification of the
119	sediment cover of the Zagros Fold Belt thus has a first-order effect on the development of upper
120	crustal-scale structures. These results are in full agreement with a recent study of active
121	seismicity in the Zagros Fold Belt which showed that both the Hormuz salt layer and the
122	intermediate layers are mechanically-weak zones that form barriers to rupture for active faults

123 (Nissen et al., 2010).

124 CONSTRAINTS ON CRUSTAL RHEOLOGY

125 The simulations presented in this study highlight the different modes of deformation that 126 might occur in fold-and-thrust belts. However, they give limited insights into the underlying 127 physics, as it remains unclear how sensitive the spacing of structures is to the rheology of the 128 crust. For this reason, we developed a semi-analytical methodology drastically reducing the 129 computational requirements that allows us to predict the outcome of numerical simulations in a 130 large parameter space (see details in DR2). The resulting wavelength versus growth rate 131 diagrams have a single maximum as a function of non-dimensional wavelength (Fig. 4A). 132 Rigorously, these semi-analytical results are only valid for very small deformations. Yet, a 133 comparison with numerical simulations reveals that they correctly predict the spacing of folds

134	even after 5.5 Myrs, which confirms that fold-spacing is selected at a very early stage in the
135	evolution of a fold and thrust belt (Fig. 4A).
136	Results for a homogeneous and brittle sedimentary cover reveal that the dominant growth
137	rate is smaller than 20, which essentially means that folding will not be able to overcome
138	background pure-shear thickening. Indeed, our numerical simulations indicate that this leads to
139	fault-dominated deformation rather than folding (box folds, Fig. 2). If, on the other hand, weak
140	layers are taken into account in the sedimentary sequence, the growth rate is significantly larger
141	and the dominant wavelength is reduced (Fig. 4B). The addition of a single weak layer is
142	sufficient to switch deformation styles from fault- to fold- dominated, and elasticity has a minor
143	effect only.
144	Using the same semi-analytical methodology, we performed a large number of
145	simulations and found that the two most important parameters are the viscosity of the salt/weak
146	layers and the friction angle of the crust, whereas rock density plays little to no role. Plots of
147	dominant wavelength and growth rate versus those two parameters show an approximate equal
148	dependence on the two parameters (Fig. 4). The results also show that weak layers in all cases
149	yields growth rates that are sufficiently large for the folding instability to dominate faulting.
150	In the case of Zagros Fold Belt, the effective viscosity of salt has been determined to be
151	close to 10 ¹⁸ Pa.s, a value consistent with scaled laboratory-derived values (Spiers et al., 1990)
152	and other modeling studies (Van Keken et al., 1993; Mouthereau et al., 2006). If we combine this
153	with our method, we estimate that the effective friction angle for the crust in the Zagros Fold
154	Belt on geological timescales is around $5^{\circ}+/-5^{\circ}$ (Fig. 4B).
155	DISCUSSIONS AND CONCLUSIONS

156	Contrary to the common view of fold belts that often consider a single major basal
157	décollement only, we demonstrate through the example of the Zagros Mountains that the whole
158	stratigraphic sequence might influence the dynamics of the belt. Heterogeneities within the
159	sedimentary cover, and weak layers in particular, control whether deformation is dominated by
160	crustal-scale folds or by thrusts. The stratigraphy of a fold belt plays a much larger role than
161	previously appreciated and should thus be taken into account if one wishes to reconcile field
162	observations with physically consistent models of geological processes.
163	Balancing geological cross-sections in fold-thrust belts is a difficult exercise that aims at
164	providing a consistent structural and kinematic interpretation of usually independent structural
165	data. Our method paves the way for developing future generations of 2D and 3D dynamic
166	reconstruction models for fold and thrust belts (e.g., Lechmann et al. 2010).
167	Moreover, we show that the regular spacing of folds puts constraints on the rheology of
168	the crust on geological timescales. In the case of Zagros Fold Belt, the value for the friction
169	angle we obtained in this manner is small (<10°), which indicates that the crust was rather weak,
170	potentially due to large fluid pressures (e.g., Huismans et al. 2005).

171 ACKNOWLEDGMENTS

This work benefits from fruitful discussions with L. Le Pourhiet, S. Schmalholz and
Y. Podladchikov. Constructive reviews from D. van Hinsbergen and 2 anonymous reviewers
have also largely contributed to improve the manuscript. BJP Kaus was partly supported by
ERC Starting Grant 258830.

176 **REFERENCES CITED**

Alavi, M., 2004, Regional stratigraphy of the Zagros fold-thrust belt of Iran and its proforeland
evolution: American Journal of Science, v. 304, p. 1–20, doi:10.2475/ajs.304.1.1.

179 Burg, J.-P., Kaus, B.J.P., and Podladchikov, Y.Y., 2004, Dome structures in collision orogens: 180 Mechanical investigation of the gravity/compression interplay, in Whitney, D.L., et al., eds., 181 Gneiss Domes in Orogeny: Geological Society of America Special Paper 380, p. 47–66. 182 Hatzfeld, D., and Molnar, P., 2010, Comparisons of the kinematics and deep structures of the 183 Zagros and Himalaya and of the Iranian and Tibetan plateaus and geodynamic implications: 184 Review of Geophysics, v. 48, no. RG2005, doi:10.1029/2009RG000304. 185 Huismans, R. S., S. J. H. Buiter, and C. Beaumont (2005), Effect of plastic-viscous layering and 186 strain softening on mode selection during lithospheric extension, J. Geophys. Res., 110, B02406, 187 doi:10.1029/2004JB003114. 188 Jackson, M.P.A., Cornelius, R.R., Craig, C.H., Gansser, A., Stöcklin, J., and Talbot, C.J., 1990, 189 Salt diapirs of the Great Kavir, central Iran: Geological Society of America Memoir 177, 190 139 p. 191 Jahani, S., Callot, J.P., Letouzey, J., and Frizon de Lamotte, D., 2009, The eastern termination of 192 the Zagros Fold-and-Thrust Belt, Iran: Structures, evolution, and relationships between salt 193 plugs, folding, and faulting: Tectonics, v. 28, p. TC6004, doi:10.1029/2008TC002418. 194 Kaus, B.J.P., 2010, Factors that control the angle of shear bands in geodynamic numerical 195 models of brittle deformation: Tectonophysics, v. 484, p. 36-47, 196 doi:10.1016/j.tecto.2009.08.042. 197 Kaus, B.J.P., Steedman, C., and Becker, T.W., 2008, From passive continental margin to 198 mountain belt: Insights from analytical and numerical models and application to Taiwan: 199 Physics of the Earth and Planetary Interiors, v. 171, p. 235–251, 200 doi:10.1016/j.pepi.2008.06.015.

- 201 Lechmann, S.M., Schmalholz, S.M., Burg, J.-P., Marques, F.O., 2010. Dynamic unfolding of
- 202 multilayers: 2D numerical approach and application to turbidites in SW Portugal.
- 203 Tectonophysics 494 (1-2), p. 64-74.
- 204 McQuarrie, N., 2004, Crustal scale geometry of the Zagros fold-thrust belt, Iran: Journal of
- 205 Structural Geology, v. 26, p. 519–535, doi:10.1016/j.jsg.2003.08.009.
- 206 Mouthereau, F., Lacombe, O., and Meyer, B., 2006, The Zagros folded belt (Fars Iran):
- 207 Constraints from topography and critical wedge modelling: Geophysical Journal
- 208 International, v. 165, p. 336–356, doi:10.1111/j.1365-246X.2006.02855.x.
- 209 Mouthereau, F., Tensi, J., Bellahsen, N., Lacombe, O., De Boisgrollier, T., and Kargar, S., 2007,
- 210 Tertiary sequence of deformation in a thin-skinned/thick skinned collision belt: The Zagros
- 211 Folded Belt (Fars, Iran): Tectonics, v. 26, p. TC5006, doi:10.1029/2007TC002098.
- 212 Nissen, E., Yamini-Fard, F., Tatar, M., Gholamzadeh, A., Bergman, E., Elliott, J.R., Jackson,
- J.A., and Parsons, B., 2010, The vertical separation of mainshock rupture and
- 214 microseismicity at Qeshm island in the Zagros fold-and-thrust belt, Iran: Earth and Planetary
- 215 Science Letters, v. 296, p. 181–194, doi:10.1016/j.epsl.2010.04.049.
- 216 Ranalli, G., 1995, Rheology of the Earth, 2nd ed.: London, Chapman and Hall.
- 217 Schmalholz, S.M., 2006, Scaled amplification equation: A key to the folding history of buckled
- 218 viscous single-layers: Tectonophysics, v. 419, p. 41–53, doi:10.1016/j.tecto.2006.03.008.
- 219 Schmalholz, S.M., Podladchikov, Y., and Burg, J.-P., 2002, Control of folding by gravity and
- 220 matrix thickness: Implications for large-scale folding: Journal of Geophysical Research,
- 221 v. 107, 2005, doi:10.1029/2001JB000355.

- 222 Sherkati, S., and Letouzey, J., 2004, Variation of structural style and basin evolution in the
- 223 central Zagros (Izeh zone and Dezful embayment): Iran: Marine and Petroleum Geology,
- v. 21, p. 535–554, doi:10.1016/j.marpetgeo.2004.01.007.
- 225 Sherkati, S., Letouzey, J., and Frizon de Lamotte, D., 2006, Central Zagros fold-thrust belt
- 226 (Iran): New insights from seismic data, field observation, and sandbox modelling: Tectonics,
- 227 v. 25, p. TC4007, doi:10.1029/2004TC001766.
- 228 Simpson, G.D.H., 2009, Mechanical modeling of folding versus faulting in brittle-ductile
- 229 wedges: Journal of Structural Geology, v. 31, p. 369–381, doi:10.1016/j.jsg.2009.01.011.
- 230 Spiers, C.J., Schutjens, P.M.T.M., Brzesowsky, R.H., Peach, C.J., Liezenberg, J.L., and Zwart,
- H.J., 1990, Experimental determination of constitutive parameters governing creep of
- rocksalt by pressure solution: Geological Society of London Special Publication 54, p. 215–
 227.
- Stocklin, J., 1968, Structural history and tectonics of Iran; A review: The American Association
 of Petroleum Geologists Bulletin, v. 52, p. 1229–1258.
- 236 Talebian, M., and Jackson, J.A., 2004, Reappraisal of earthquake focal mechanisms and active
- shortening in the Zagros mountains of Iran: Geophysical Journal International, v. 156,
- 238 p. 506–526, doi:10.1111/j.1365-246X.2004.02092.x.
- 239 Van Keken, P.E., Spiers, C.J., Van den Berg, A.P., and Muyzert, E.J., 1993, The effective
- viscosity of rocksalt: Implementation of steady state creep laws in numerical models of salt
- 241 diapirism: Tectonophysics, v. 225, p. 457–476, doi:10.1016/0040-1951(93)90310-G.

242 FIGURE CAPTIONS

- 243 Figure 1. Field constraints for the Zagros folded belt. A: Topography illustrates the regular
- spacing of folds with amplitude \sim 500–1000 m over on an area of \sim 80 000 km². Fold crest length

245	are of \sim 50 km in average. Inset shows the distribution of fold wavelengths measured for 88
246	anticlines from the Zagros Folded Belt. B: Cross-section (aa') across the Zagros Fold Belt based
247	on field measurement (Mouthereau et al., 2007). λ corresponds to the average wavelength of the
248	folds. This value is slightly smaller than the 15.8 \pm – 5.3 km from Mouthereau et al. (2007) that
249	took into account the folds from the whole Fars area. MFF and SF correspond to the seismogenic
250	Mountain Front Fault and the Surmeh Fault, respectively, associated with basement faulting.
251	Vertical fold velocity is 0.3–0.6 mm.yr ⁻¹ . C: Synthetic stratigraphic log where WL1, WL2 and
252	WL3 correspond to the weak layers in the sedimentary sequence (Fm: Formation).
253	Figure 2. Simulation with a basal décollement layer only. A: Initial setup with a sedimentary
254	thickness of 7.225 km. All rocks above the basal salt layer are homogeneous and have a friction
255	angle of 5° and a cohesion of 20 MPa. A background strain rate of 10^{-15} s ⁻¹ is imposed at the
256	right model boundary. B: Geometry, strain rate, and vertical velocities after 1.5 Myrs and 5.5
257	Myrs respectively. Deformation is localized along crustal-scale plastic shear zones and
258	deformation structures are fault-dominated.
259	Figure 3. Simulation with intermediate crustal detachment layers. A: Initial setup as in figure 2,
260	but with three additional weak crustal detachment layers with 10 ¹⁸ Pa.s. B: Snapshots of
261	geometry, strain rate and vertical velocities at different times, which illustrate that crustal-scale
262	folds rather than faults dominate the deformation pattern. Note that folds do not grow
263	continuously with time, but rather grow to certain amplitude after which activity switches to a
264	different fold.
265	Figure 4. Influence of multiple weak layers and elasticity on folding. λ , H, q and $\dot{\epsilon}$ corresponds
266	to the wavelength, the total thickness, the growth rate and the background strain rate,

267 respectively. VP and VEP correspond to visco-plastic and visco-elasto-plastic simulations,

268	respectively. This diagram was produced using the semi-analytical approach described in DR2.
269	A: Growth rate values obtained for given values of λ/H for 0, 1, 2 and 3 weak layers. For each
270	case, the characteristic wavelength value corresponds to the highest value of growth rate (e.g.,
271	white star for the case with 3 weak layers). Insets show results of numerical simulations after 5.5
272	Myrs, which develop folds with a spacing that is in excellent agreement with the predicted
273	characteristic wavelength. A single basalt décollement layer results in small folding growth rates
274	and in thrust-dominated deformation. Addition of one or more weak layers to the brittle
275	sedimentary cover results increases the growth rate significantly and leads to folding-dominated
276	deformation. The ZFB brown area corresponds to the λ /H ratio of the Zagros Fold Belt. B:
277	diagrams of characteristic wavelength (left) and corresponding growth rate (right) versus
278	viscosity of the weak layers and friction angle of the crust. Thick white lines show the
279	constraints for Zagros Fold Belt (average +/- 1 standard deviation). As in the Zagros Fold Belt
280	salt viscosity is constrained independently, the best-fit friction angle for the crust is $5+/-5^{\circ}$. The
281	white star corresponds to the simulation of figure 3.
282	¹ GSA Data Repository item 2011xxx, xxxxxxx, is available online at
283	www.geosociety.org/pubs/ft2011.htm, or on request from editing@geosociety.org or Documents
284	Secretary, GSA, P.O. Box 9140, Boulder, CO 80301, USA.







