



# New insights in the use of carbon isotopes as tracers of DOC sources and water pathways in headwater catchments

Thibault Lambert, Anne-Catherine Pierson-Wickmann, Gérard Gruau, Patrice Petitjean, Jean-Noël Thibault, Laurent Jeanneau

## ► To cite this version:

Thibault Lambert, Anne-Catherine Pierson-Wickmann, Gérard Gruau, Patrice Petitjean, Jean-Noël Thibault, et al.. New insights in the use of carbon isotopes as tracers of DOC sources and water pathways in headwater catchments. EGU General Assembly 2013, Apr 2013, Vienne, Austria. pp. EGU2013-9595. insu-00853603

**HAL Id: insu-00853603**

**<https://insu.hal.science/insu-00853603>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## **New insights in the use of carbon isotopes as tracers of DOC sources and water pathways in headwater catchments**

Thibault Lambert (1), Anne-Catherine Pierson-Wickmann (1), Gérard Gruau (1), Patrice Petitjean (1), Jean-Noël Thibault (2), and Laurent Jeanneau (1)

(1) OSUR, Université de Rennes 1, France, (2) INRA, St Gilles, France

Despite the significant importance of dissolved organic carbon (DOC) in aquatic ecosystems, the processes controlling DOC delivery to stream waters at the catchment scale are still poorly understood, in particular with regards to the relative importance of riparian versus upland soils as DOC sources. In this respect, the stable carbon isotopic composition of DOC ( $\delta^{13}\text{C}_{\text{DOC}}$ ) appears to be a promising tool as different  $\delta^{13}\text{C}_{\text{DOC}}$  values are anticipated between wetland and upland soil DOC, because of differences in soil oxygenation, soil humidity and soil organic matter degradation scheme. However, care must be exercised because of the possible occurrence of seasonal variations in the  $\delta^{13}\text{C}_{\text{DOC}}$  values of both riparian and upland DOC, and because also of the possible mixing of DOC coming from spatially distinct sources.

The markedly different isotopic patterns obtained during high resolution monitoring (<hourly basis) of six successive storm events occurring between November 2010 and February 2011 in a small headwater catchment from western France illustrate this difficulty. Survey of the  $\delta^{13}\text{C}_{\text{DOC}}$  values of the DOC during these storm events showed that the variability in intra-storm DOC isotopic composition markedly evolved through time, with some storms exhibiting strongly variable  $\delta^{13}\text{C}_{\text{DOC}}$  values ( $> 2 \delta$  units), while others showed no, or much more restricted isotopic variations. A comparison of these results with previously published data revealed that this temporal variability of intra-storm  $\delta^{13}\text{C}_{\text{DOC}}$  values is the exact transposition of the temporal variability of  $\delta^{13}\text{C}_{\text{DOC}}$  values that was found in the riparian soil waters of this catchment during the same period. The latter variability has been shown to arise from the combined effect of changes in the production mechanisms and ultimate sources of riparian DOC and of the lateral input in the riparian domains of an isotopically heavier DOC component coming from more upland areas.

Overall, results from this study confirm that upland domains may be significant contributors of stream DOC flux in headwater catchments. They also show that upland soils behave as a size-limited reservoir with respect to DOC production, whereas more highly productive soils in the wetland domains act as a near-infinite reservoir. Through this study, we show that the isotopic composition of DOC is an extremely powerful tool for tracing DOC sources and DOC transport mechanisms in headwater catchments, demonstrating in the meantime that the use of this tool requires that the temporal and spatial variability of the isotopic signatures of all potential DOC sources in the catchment is known accurately. Providing that this condition is fulfilled, the isotopic tool can allow up to quantify the proportions of DOC - and of corresponding water flows - coming from different contributing areas which may be of great importance for better understanding and better modeling of DOC transfer and water routing through the landscape.