

## Experimental warming interacts with soil moisture to discriminate plant responses in an ombrotrophic peatland

Alexandre Buttler, B.J.M. Robroek, Fatima Laggoun-Défarge, Vincent E. J. Jassey, Cédric Pochelon, Gregory Bernard, Frédéric Delarue, Sébastien Gogo, Pierre Mariotte, Edward A. D. Mitchell, et al.

### ► To cite this version:

Alexandre Buttler, B.J.M. Robroek, Fatima Laggoun-Défarge, Vincent E. J. Jassey, Cédric Pochelon, et al.. Experimental warming interacts with soil moisture to discriminate plant responses in an ombrotrophic peatland. Journal of Vegetation Science, 2015, 26 (5), pp.964-974. 10.1111/jvs.12296. insu-01163362

## HAL Id: insu-01163362 https://insu.hal.science/insu-01163362

Submitted on 12 Jun 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Experimental warming interacts with soil moisture to discriminate plant responses in an ombrotrophic peatland

Journal:	Journal of Vegetation Science
Manuscript ID:	JVS-RA-02979.R1
Manuscript Type:	Research article
Date Submitted by the Author:	n/a
Complete List of Authors:	Buttler, Alexandre; EPFL-WSL, Laboratory of Ecological Systems - ECOS Robroek, Bjorn; Utrecht University, Institute of Environmental Biology; EPFL-WSL, Laboratory of Ecological Systems - ECOS Laggoun, Fatima; CNRS/INSU, ISTO, UMR 7327 Jassey, Vincent; EPFL-WSL, Laboratory of Ecological Systems - ECOS Pochelon, Cédric; EPFL-WSL, Laboratory of Ecological Systems - ECOS Bernard, Gregory; Fédération des Conservatoires d'espaces naturels, Pôle- Relais Tourbières Delarue, Frédéric; Muséum National d'Histoire Naturelle, Laboratoire de Minéralogie et de Cosmochimie du Muséum, UMR 7202 CNRS Gogo, Sébastien; CNRS/INSU, ISTO, UMR 7327 Mariotte, Pierre; EPFL-WSL, Laboratory of Ecological Systems - ECOS Mitchel, Edward; Université de Neuchâtel, Laboratory of Soil Biology, Institute of Biology Bragazza, Luca; EPFL-WSL, Laboratory of Ecological Systems - ECOS
Keywords:	climate change, vegetation, Sphagnum, OTC, temperature, drought, NDVI, point-intercept survey

SCHOLARONE<sup>™</sup> Manuscripts

1	Research article
2	Estimated size: 6617 words (8 pages), figures 2 pages, total 10 pages
3	
4	Experimental warming interacts with soil moisture to discriminate plant responses in an
5	ombrotrophic peatland
6	
7	Running head: Experimental warming and plant responses
8	
9	Alexandre Buttler, Bjorn J.M. Robroek, Fatima Laggoun-Défarge, Vincent E.J. Jassey, Cédric
10	Pochelon, Gregory Bernard, Frédéric Delarue, Sébastien Gogo, Pierre Mariotte, Edward A.D.
11	Mitchell & Luca Bragazza
12	
13	Buttler, A. (Corresponding author, alexandre.buttler@epfl.ch) <sup>1, 2, 3, 12</sup>
14	Robroek, B.J.M. (bjorn.robroek@epfl.ch) <sup>1, 2, 11</sup>
15	Laggoun-Defarge, F. (Fatima.Laggoun-Defarge@univ-orleans.fr) <sup>4,5,6</sup>
16	Jassey, V.E.J. (vincent.jassey@epfl.ch) <sup>1, 2, 3</sup>
17	Pochelon, C. (cedric.pochelon@bluewin.ch) <sup>1,2</sup>
18	Bernard, G. (gregory.bernard@reseau-cen.org) <sup>1, 2, 14</sup>
19	Delarue, F. (fdelarue@mnhn.fr) <sup>4, 5, 6, 7</sup>
20	Gogo, S. (Sebastien.Gogo@univ-orleans.fr) <sup>4, 5, 6</sup>
21	Mariotte, P. (pierre.mariotte@hotmail.com) <sup>1, 2, 12, 13</sup>
22	Mitchell, E.A.D. (edward.mitchell@unine.ch) <sup>9,10</sup>
23	Bragazza, L. ( <u>luca.bragazza@wsl.ch</u> ) <sup>1, 2, 8</sup>
24	

- 25 Author contributions: A.B. wrote the manuscript, A.B., G.B and C.P. collected the vegetation data,
- A.B., F.L.D, S.G. and E.M. conceived the experiment, B.R., V.J. and F.D. helped for vegetation and
- 27 micrometeorological data analysis and all contributed to the interpretation of results and commented
- 28 on the manuscript.
- 29

30 Addresses:

- <sup>1</sup>Ecole Polytechnique Fédérale de Lausanne EPFL, School of Architecture, Civil and Environmental
- 32 Engineering ENAC, Laboratory of ecological systems ECOS, Station 2, 1015 Lausanne,
- 33 Switzerland.
- <sup>2</sup>Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Site Lausanne, Station 2,
- 35 1015 Lausanne, Switzerland.
- <sup>3</sup>Laboratoire de Chrono-Environnement, UMR CNRS 6249, UFR des Sciences et Techniques, 16
- 37 route de Gray, Université de Franche-Comté, F-25030 Besançon, France.
- <sup>4</sup>Université d'Orléans, ISTO, UMR 7327, 45071 Orléans, France.
- <sup>5</sup>CNRS/INSU, ISTO, UMR 7327, 45071 Orléans, France.
- 40 <sup>6</sup>BRGM, ISTO, UMR 7327, BP 36009, 45060 Orléans, France.

<sup>41</sup> <sup>7</sup>Laboratoire de Minéralogie et de Cosmochimie du Muséum, UMR 7202 CNRS, Muséum National

- 42 d'Histoire Naturelle, 57 rue Cuvier, 75231 Paris Cedex 05, France.
- <sup>43</sup><sup>8</sup>University of Ferrara, Department of Life Science and Biotechnologies, Corso Ercole I d'Este 32,
- 44 I-44121 Ferrara, Italy.
- <sup>45</sup> <sup>9</sup>Laboratory of Soil Biology, Institute of Biology, University of Neuchâtel, CH-2000 Neuchâtel,
- 46 Switzerland.
- <sup>47</sup> <sup>10</sup>Jardin Botanique de Neuchâtel, Chemin du Perthuis-du-Sault 58, 2000 Neuchâtel, Switzerland.
- 48 <sup>11</sup>Ecology and Biodiversity, Institute of Environmental Biology, Utrecht University, Padualaan 8,
- 49 3584 CH Utrecht, The Netherlands.
- <sup>12</sup>Department of Environmental Science, Policy and Management, Suding Lab, University of
- 51 California Berkeley, 147 Hilgard Hall, Berkeley, CA 94720.
- <sup>13</sup>University of Sydney, Centre for Carbon, Water and Food Biogeochemistry Group 380
- 53 Werombi Road, Camden NSW 2570.
- <sup>14</sup>Pôle-relais Tourbières, Fédération des Conservatoires d'espaces naturelsK7, rue Voirin, 25000
- 55 Besançon, France.

57 Keywords: climate change; vegetation, *Sphagnum*; OTC; temperature; drought; NDVI; point-58 intercept survey.

59

60

#### 61 ABSTRACT

62

Question: A better understanding of the response of *Sphagnum* mosses and associated vascular plants to climate warming is relevant for predicting the carbon balance of peatlands in a warmer world. Open-top chambers (OTCs) have been used to investigate the effect on soil biogeochemical processes in peatlands, but little information is available on the effects of OTCs on microclimate conditions and the associated response of plant community. We aimed to understand how simulated warming and differences in soil moisture affect plant species cover.

Methods and location: We used OTCs to measure the effect of a near-ground temperature increase (+ 1.5 °C on average) on vegetation dynamics over five growing seasons (2008-2012) in a *Sphagnum*-dominated peatland (French Jura), in two adjacent microhabitats with different hydrological conditions – wet and dry. Microclimatic conditions and plant species abundance were monitored at peak biomass in years 1, 2, 3 and 5 and monthly during the plant-growing season in year 5.

Results: The response of plants to warming differed between vascular plants and bryophytes as well as among species within these groups and also varied in relation to soil moisture. *Andromeda polifolia* abundance responded positively to warming, while *Vaccinium oxycoccus* responded negatively and *Eriophorum vaginatum* showed a high resistance. 79 Conclusion: Depth of rooting of vascular plants appeared to control the response in plant 80 abundance, while moss abundance depended on various other interacting factors, such as shading 81 by the vascular plant community, precipitation and soil moisture.

- 82
- 83
- 84

#### 85 INTRODUCTION

Peatlands are long-term sinks of atmospheric carbon (C) due to the accumulation of peat (Yu 2011). These ecosystems develop in areas that are hydrologically and climatically characterized by a net surplus of water. As a consequence, prevailing soil anoxic conditions hamper microbial decomposition of organic matter (Holden 2005).

90 In northern hemisphere ombrotrophic (= rain fed) peatlands, bryophytes of the genus Sphagnum 91 (peat mosses) are generally the dominant plant species. Peat mosses produce litter with 92 antimicrobial properties, which hampers microbial decomposition (Hajek et al. 2011), leading to the 93 build-up of peat. Further, Sphagnum mosses can affect the abundance and the performance of 94 associated vascular plants (Dorrepaal et al. 2006) by regulating soil moisture, energy balance, soil 95 temperature, and nutrient availability (Turetsky et al. 2012). Ultimately, the inter-specific 96 competition between Sphagnum mosses and vascular plants controls peat accumulation because an 97 increase of vascular plant abundance and in particular their shading effect can reduce Sphagnum 98 growth (Heijmans et al. 2002; Bragazza et al. 2013). Changes in the competitive balance between 99 plant species of different growth forms has been shown to alter the C balance in peatlands (Kuiper 100 et al. 2014), potentially by changes in the litter quality, which ultimately affects microbial 101 decomposition (Bragazza et al. 2007; Dorrepaal et al. 2007; Gogo et al. 2011).

102 Climate warming and associated extreme climatic events can be detrimental to *Sphagnum* 103 mosses as, like all terrestrial bryophytes, these plants cannot actively control their water balance and

104 therefore rely on atmospheric precipitation to maintain suitable hydrological conditions (Robroek et 105 al. 2007, Bragazza 2008; Nijp et al. 2014). Increased air temperature can increase 106 evapotranspiration with direct negative effects on Sphagnum productivity (Gerdol 1995; Weltzin et 107 al. 2001), but with positive effects on vascular plant growth by drying out the top soil and 108 enhancing soil oxygenation (Weltzin et al. 2000; Bragazza et al. 2013), to which plants of different 109 functional types might respond differently (Kuiper et al. 2014). A better understanding of the 110 response of Sphagnum mosses and associated vascular plants to climate warming is then relevant 111 for predicting the C balance of peatlands in a warmer world. Indeed, a warmer climate has been 112 reported to alter plant species abundance in peatlands, in particular favoring ericaceous dwarf 113 shrubs (Weltzin et al. 2003, Breeuver et al. 2009), which will affect soil biogeochemical processes 114 (Bragazza et al. 2013).

115 Under field conditions, simulation of climate warming can be obtained by using passive 116 warming open-top chambers (OTCs) (Marion 1997). Although OTCs are recognized to increase 117 temperature consistent with general circulation models, they also affect a set of microclimatological 118 variables such as moisture, light, snow accumulation and wind speed (Bokhorst et al. 2013). In 119 peatland studies, OTCs have been reliably used to investigate the effect on soil biogeochemical 120 processes (Dorrepaal et al. 2009), but little information is available on the effects of OTCs on 121 microclimate conditions and the associated response of plant species (Weltzin et al. 2003, Keuper et 122 al. 2011; Delarue et al. 2014; Jassey et al. 2013).

In the present study we used OTCs to simulate increased soil and air temperature, and we monitored changes in vegetation cover in a *Sphagnum*-dominated peatland over five years of experimental warming. We aimed to understand how species abundance and plant community composition responded to simulated warming and associated changes in soil moisture. We hypothesized that plant responses will vary among functional types; and more specifically that 1) vascular plants will respond to warming according to their rooting depth, and 2) *Sphagnum* mosses will have a more complex response depending on the interaction between micrometeorologicalconditions and abundance of vascular plants.

131

#### 132 MATERIAL AND METHODS

133

134 *Study site* 

The study was performed on an undisturbed, ombrotrophic *Sphagnum*-dominated peatland situated in the Jura Mountains (The Forbonnet peatland, France, 46°49'35''N, 6°10'20''E, 840 m a.s.l). Climate is characterized by relatively cold winters (average temperature -1.4°C) and mild summers (average temperature 14.6°C), with annual mean temperature of 7.5°C and annual mean precipitation between 1300 and 1500 mm (Laggoun-Défarge et al. 2008).

140 Two areas (microhabitats) were selected based on their hydrological conditions. The first 141 microhabitat (hereafter called 'wet') was a Sphagnum-dominated relatively flat area bordering a 142 transitional poor fen, characterized by a moss cover dominated by Sphagnum fallax. Vascular plants 143 such as Eriophorum vaginatum and Andromeda polifolia were recorded with low abundance, 144 together with Scheuchzeria palustris, Drosera rotundifolia and Vaccinium oxycoccus. The second 145 microhabitat (hereafter called 'dry') was adjacent to the wet microhabitat, but in a area with 146 generally deeper water tables. The dry microhabitat was characterized by a pattern of hummocks 147 dominated by Sphagnum magellanicum, V. oxycoccus, E. vaginatum and Calluna vulgaris, and 148 lawns with S. fallax, Carex rostrata and A. polifolia. Tree (Pinus rotondata) encroachment occurs 149 in this dry microhabitat.

150

151 Experimental design

152 In April 2008, in both the dry and the wet microhabitats, six representative plots  $(3 \times 3 \text{ m})$  were 153 selected, which were then randomly allocated to either the control or the warming treatment

(Laggoun-Défarge et al., 2008). Warming treatments were obtained by using open-top chambers (OTCs),-which consisted of transparent polycarbonate hexagonal chambers (50 cm high, 1.7 m topwidth, 2.4 m base-width). OTCs allow quasi-natural transmittance of visible wavelengths, and minimize the transmittance of re-radiated infrared wavelengths (Marion et al., 1997). All OTCs were raised 10 cm above the peat surface to allow air circulation. The minimal (adjacent plots) and maximal (distant plots) distance between plots were *c*. 5 and 30 m, respectively.

160

#### 161 Environmental monitoring

162 From November 2008 to November 2012, peat temperature (7 cm below the moss carpet) and air 163 temperature (at 10 cm above the moss carpet) were monitored at 30-minute intervals using 164 thermocouple probes connected to a data logger (CR-1000 Campbell). Sensors were repositioned as 165 necessary to keep height and depth of measurement constant despite moss carpet accumulation. In 166 2012, these measurements were augmented with combined soil temperature/moisture sensors. In 167 each microhabitat (wet/dry) and treatment (OTC/control) a randomly chosen plot was equipped 168 with soil temperature-moisture probes (Decagon 5TM) at 3 cm and 5 cm depth. Furthermore, 169 precipitation at the study site and water table depth in randomly chosen plot were continuously 170 monitored by a meteorological station and sensors connected to the CR-1000 data logger.

171

#### 172 Plant species abundance

At peak biomass (July or August, depending on the year) in 2008, 2009, 2010 and 2012, plant species abundance was measured using the point-intercept frequency method (Buttler 1992) in 50 × 50 cm subplots located in flat lawns. We used a Plexiglas frame with adjustable legs placed above a permanently marked subplot. A 20 holes ruler was moved along 20 different positions so as to obtain 400 measuring points on a regular grid. A metal pin (1 mm diameter) was lowered through each hole in the ruler and each contact of the pin with living vegetation and litter was recorded by

species until the pin reached the *Sphagnum* surface. This field technique could account for threedimensional biomass distribution because, at each point, each plant that was intercepted was recorded. The relative frequency of each species was then calculated.

182 To allow analyses on seasonal dynamics of plant species cover, during the growing season 2012 183 (April - September), monthly pictures were taken from aforementioned subplots. In order to obtain 184 high-resolution images, the subplots were divided into four  $25 \times 25$  cm quadrats. For the digital 185 image analysis, a grid of 100 points was laid on each  $25 \times 25$  cm picture, after which the four grids 186 were merged into a single 400 points grid-picture. Species overlaying the grid intersects were 187 identified at 200% digital magnification. The relative frequency of each species was then calculated 188 for each sampling date, as for the point-intercept method. Because we could not reliably distinguish 189 S. fallax and S. rubellum in our point-intercept field measures and in the frequency analysis on 190 photographs, we pooled these two species.

191

#### 192 Sphagnum height growth

The cranked wire method (Clymo 1970) was used to measure *Sphagnum* growth. At the start of the 2011 and 2012 growing seasons (early April, after snow melt) cranked wires (5-10) were placed in each subplot. At the beginning of November, before snowfall, the length of the cranked wire above the moss surface was measured with a graduated pipe sliding down along the cranked wire and the average height of the *Sphagnum* capitula around the wire was calculated.

198

199 Seasonality of plant community

The normalized difference vegetation index (NDVI) was used as a phenological proxy and measured every fortnight between March and November 2012 using a portable spectroradiometer sensor. This measure is also used as an indirect measure of bulk biomass (Soudani et al. 2012). The sensor was maintained at 50 cm above the moss carpet, which allowed obtaining a NDVI signal of

the whole plant community, including the highest plants, while covering only the subplot surface. In each  $50 \times 50$  cm subplot and for each campaign of sampling, 10 consecutive measures were taken according to a grid sampling design and then averaged to have a single NDVI value per plot. The NDVI measurements were performed under direct sunlight and with an umbrella to shadow the subplot.

209

210 Statistical analysis

211 Differences in vegetation between years (i.e. 2008, 2009, 2010, 2012) were analyzed using 212 relative species frequency data obtained from the point-intercept method in the field, whereas 213 seasonal trend of plant species cover during the 2012 growing season was analyzed with species 214 relative frequency data obtained from the photographic image analysis. Analysis at species level 215 was done with Linear Mixed Models, with treatment (OTC vs. control) and years (or months for the 216 seasonal data set in 2012) as fixed factors, and years (or months) nested in plots and blocks (wet 217 and dry microhabitats) as random factors. The same model was also used to test the block effect. 218 Sphagnum growth was analyzed with Linear Mixed Models, with treatment and block as fixed 219 factors, and location of growth measures within plots given as random nested factors. The model 220 was also tested for each block separately. NDVI was also analyzed with Linear Mixed Model, with 221 treatment and block as fixed factors, and dates nested in plots as random factors. Random effects on 222 the intercept were used to correct the inflation of the residual degrees of freedom that would occur 223 if repeated measurements within sites would have been used as true replicates (Pinheiro & Bates 224 2000). In addition, for NDVI, separate ANOVA's were performed for each block and date. 225 Pearson's correlation coefficients were calculated and tested between NDVI signals and species 226 frequencies. Differences in monthly means of differences in daily mean air and soil temperature 227 were assessed by ANOVA's. Comparisons of mean water level depth between blocks were done 228 with Student t-tests. Data were log transformed prior to analysis. All statistical analyses were 229 performed in R (R Core Team, 2014).

230 Results

231 232

233

*Micrometeorology* 

234

235 In general, over the period 2009-2012, monthly air temperature in the OTCs was 0.2-1.5 °C

236 higher than in the control plots (p < 0.05). Interestingly, OTCs rarely affected soil temperature, and 237 when they did (p < 0.05), the pattern was less clear, varying between a warming effect up to 0.3 °C 238 in some cases, and a cooling effect up to -1 °C in some others (see also Delarue et al. 2011, 2014, 239 2015, Jassey et al. 2011). These effects of OTC on soil temperature were not related to the season, 240 neither to the microhabitat. Soil moisture and temperature measured in 2012 with Decagon sensors 241 in a subset of 4 plots (dry control, dry OTC, wet control and wet OTC) indicate that soil 242 temperature was slightly higher in OTCs this particular year (mean daily differences between OTC 243 and control plots for soil temperature was +0.6 °C in both microhabitat), and that moisture was 244 slightly lower in OTC plots (mean daily differences between OTC and control plots for soil 245 moisture were -0.011 and -0.067 (VWC) in dry and wet microhabitat, respectively).

246 Monthly cumulative rainfall during the period of *Sphagnum* growth measurements was higher in 247 2012 (2108 mm) than in 2011 (1376 mm). Depth of water level fluctuated over time, but highest 248 water levels were recorded during spring 2012. Mean water level during the measurement period of 249 Sphagnum growth was significantly higher in 2012 than in 2011 (Student t-test; p=0.002 in dry, 250 p=0.004 in wet). Mean water level was also higher in the wet than the dry microhabitat (p=0.004).

251

252

254

253 Annual change in species abundance

255 Considering all vascular plants, abundance (relative frequency) increased significantly with the 256 OTC treatment (Figure 1). At species level, it appeared that with warming A. polifolia increased its

257 abundance consistently in wet and dry microhabitats, whereas the abundance of V. oxycoccus 258 decreased. D. rotundifolia abundance responded marginally to the OTC treatment whereas E. 259 vaginatum did not respond. Some other species responded contrastingly to the OTC treatment in the 260 two microhabitats, for example *P. strictum* and *S. fallax+rubellum* increased their abundance in the 261 dry microhabitat, whereas their abundance decreased or did not change in the wet microhabitat 262 (treatment x block p=0.0367 and 0.0767, respectively). Litter increased in the wet microhabitat, and 263 decreased in the dry microhabitat (treatment x block p=0.0001). Significant changes occurred also 264 over time, with a decrease of abundance for V. oxycoccus and S. fallax+rubellum, and an increase 265 for *E. vaginatum* and litter.

There was also a strong microhabitat effect related to the dry and wet situations (p<0.05), except for *Drosera rotundifolia* for which the microhabitat effect was marginally significant, and for *Carex rostrata*, *C. limosa*, *Calluna vulgaris* and *Pinus rotundata* for which there was no significant microhabitat effect. *Vaccinium oxycoccus*, *D. rotundifolia*, *Scheuchzeria palustris*, and *Sphagnum fallax+rubellum* were more abundant in the wet microhabitat, whereas species such as *Andromeda polifolia*, *Eriophorum vaginatum* and *Polytrichum strictum* were more frequent in the dry microhabitat, or exclusively found in the latter one (*S. magellanicum*).

273

#### 274 *Monthly change in species abundance*

During the 2012 season, total vascular plant abundance and, more specifically *Andromeda polifolia* frequency increased significantly in the OTCs (Figure 2). *Vaccinium oxycoccus* abundance decreased in the wet microhabitat (treatment x block p=0.0034) whereas that of *Sphagnum fallax+rubellum* declined in the wet microhabitat, but increased in the dry microhabitat (treatment x block p=0.008). Litter increased in the wet microhabitat and decreased in the dry microhabitat (treatment x block p=0.0001). Significant changes occurred also over the monthly course of 2012, 281 with a decrease in abundance for S. magellanicum (present only in the dry microhabitat) and litter,

282 and an increase for V. oxycoccus and total vascular plants.

283

284 Sphagnum growth

285

286 The length increment of Sphagnum species differed between years, being higher in 2012 as 287 compared to 2011 (Figure 3). In 2011, Sphagnum growth was significantly lower in the wet 288 microhabitat, but warming did not affect the length increment. In 2012, there were significant 289 microhabitat and treatment effects as well as a cross effect, with increased moss growth in the 290 warmed plots of the dry microhabitat (p=0.017), whereas no-significant changes were observed in 291 the wet microhabitat.

292

### ndex Normalized Difference Vegetation Index 293

294

NDVI values taken under direct sunlight and in the shade were strongly correlated ( $R^2=0.83$ ), yet 295 296 measurements at the start of the season (early May) were less correlated ( $R^2=0.49$ ) than 297 measurements close to the peak of biomass (mid-June;  $R^2=0.82$ ). The precision of the 298 measurements was calculated as the mean difference between each pairwise measurement. It gives 299 an error of 1.6% under sun and 2.0% in the shade. As such, we only used NDVI data from sun 300 measurements in further analyses.

301 Throughout the season, OTCs increased the vegetation's NDVI signal (p=0.001), yet the effect 302 of OTCs differed between wet and dry microhabitats (block x treatment p=0.002). In the dry 303 microhabitat, the NDVI signal in the OTC exceeded the signal in control plots throughout the 304 season (Figure 4a). Such difference was mainly significant at the beginning of the growing season, 305 and marginally different in summer. In the wet microhabitat, however, the difference between OTC 306 and control plots was less obvious, and occurred during a more limited time in summer (Figure 4b).

NDVI was positively correlated to vascular plant abundance (r=0.85, p<0.001), and negatively with litter (r=-0.60, p<0.001). The correlation between NDVI and *Sphagnum* cover was negative (r=-0.24, p<0.045).

#### 310 Discussion

#### 311 *Effect of OTC on microclimate*

312 The observed OTC warming effect on near-ground air temperature was in the range of those 313 found in other studies (e.g., Marion et al., 1997; Hollister & Webber, 2000; Dorrepaal et al. 2003; Sullivan et al., 2005, Walker et al. 2015). Yet, in accordance with Dabros et al. (2010), we observed 314 315 that OTCs can reduce soil temperature, as a result of increased evapotranspiration (Delarue et al. 316 2011). Daily temperature fluctuations, minimum and maximum values and day or night means have 317 all been shown to be affected by OTC warming (e.g. Marion et al. 1997; Dabros et al. 2010). A 318 previous study at the same site revealed that daily soil temperature amplitudes were smaller but 319 often significantly different between OTC's and control plots in winter months (Delarue et al. 320 2011). In summer months, amplitudes were larger, and even more so in OTCs as compared to 321 control plots in April, May and June in the dry microhabitat (but differences were marginally or not 322 significant), whilst an opposite trend appeared in the wet microhabitat. The additional Decagon 323 sensors installed in 2012, which integrate more soil volume, might be more adapted to the very 324 porous peat soil than tiny thermocouples. Despite the absence of replication, it is reasonable to 325 recognize that our OTCs had a slight warming effect on soils for the largest part of time, yet this 326 effect coincided with a slight reduction in soil volumetric water content as previously reported 327 (Jassey et al. 2011)

328

329 *Effect of OTCs on vegetation* 

330 We showed that experimental warming differently influenced plant abundance (or frequency),

331 depending on the underlying hydrological conditions and on individual species identity. A. polifolia 332 increased in abundance while V. oxycoccus decreased, and both S. fallax+rubellum and P. strictum 333 responded positively to the OTC treatment in the dry microhabitat, and decreased or remained 334 constant in the wet microhabitat. Among these species, the response of A. polifolia is interesting. 335 This species has been shown to be sensitive to changes in environmental conditions such as 336 snow cover and spring/summer warming (Aerts et al. 2006; Jassey et al. 2013; Robroek et al. 337 2013). It allocates about 98% of the total biomass belowground, concentrated in the 0-15 cm 338 peat layer, with the bulk at 2.5-7.5 cm, and a continuous rooting system from recent vegetation 339 down into the peat (Wallén 1986). Roots of A. polifolia may reach a depth of 45 cm (Jaquemart 340 1998) and fine roots form up to 24% of the total below ground biomass (Wallen 1986). A. polifolia 341 might therefore be expected to have a somewhat deeper root system as compared to several other 342 species enabling the species to sustain growth when soil dries out at the surface. Differently, V. 343 oxycoccus has a shallow root system hardly penetrating more than a few centimeters below the 344 living parts of the moss layer (Jacquemart 1998). With its shallow roots, this species relies on the 345 water-conducting capacity of the Sphagnum mosses for its water supply (Malmer et al. 1994). 346 Indeed, a decrease of Sphagnum water content in the OTC treatment has been demonstrated by 347 Jassey et al. (2011). A subsequent lowering of water table will also reduce the growth rate of V. 348 oxycoccus through water stress (Malmer et al. 1994, Jacquemart 1998). Coriaceous leaves of A. 349 *polifolia*, their silvery-glaucous lower surface and their revolute margins, could indicate drought 350 resistance. But anatomical features of V. oxycoccus such as thick epicuticular waxes and stomata on 351 the abaxial surface only are considered similar to those of desert plant characteristic to a high ability 352 to avoid drought (Jacquemart 1997). Drier soil might also promote ericoid mycorrhizal infection 353 (Cullings 1996) and thus better access to nutrients in the soil, but this might equally apply to other 354 Ericaceous present in the bog, such as V. oxycoccus.

355 An alternative explanation for the decrease of V. oxycoccus with OTC warming may come from 356 direct nutrient resource competition (Malmer et al. 1994), in particular with A. polifolia. Because, 357 however, OTC warming coincided with a decrease in moisture content, a direct effect of moisture 358 limitation on the shallow rooting V. oxycoccus cannot be ruled out. The evidence of moisture 359 limitation is supported by the lack of response to the OTC by *E. vaginatum*. This species is known 360 to root deeply into the peat, up to 1.5 meters, with roots having highest elongation rates at low soil 361 temperatures (Shaver & Billings 1977). Contrary to A. polifolia and V. oxycoccus, no mycorrhizal 362 association has been reported for *E. vaginatum*. In warming experiment, no significant differences 363 were reported for *E. vaginatum* root production between OTC's and control plots, albeit a tendency 364 to higher production was observed (Sullivan & Welker 2005). We believe that the resistance of E. 365 *vaginatum* is higher as compared to more shallow rooted species, and that an effect of warming has 366 to be expected on a longer term, as soil warms up at lower depth. Weltzin et al. (2003) found that 367 within bog and fen communities, different life forms and species can respond differently to 368 warming and water table manipulation. They showed that warming may lead to an increase in 369 abundance of woody plants, as does lowering water table as a result of changes in rates of 370 evapotranspiration. In the peatlands studied by Weltzin et al. (2003), Andromeda glaucophylla 371 responded mostly to the heat treatment, while other dominant shrubs (such as *Vaccinium oxycoccus*) 372 did not, whereas graminoids decreased their cover. As observed by Weltzin et al. (2000), change in 373 belowground biomass can be even more important. Our wet and dry sites showed similar trends for 374 vascular plants, but not for mosses. The better growth of some mosses in warmed plots might be 375 transient since, on the long run, mosses will be out-competed by the concomitant increase of 376 Andromeda shrubs. Overall, these observations could point to a different evolution of such 377 contrasted dry and wet situations, as for "bog" and "fen" like facies (Weltzin et al. 2003).

The increase of litter in the wet microhabitat and its decrease in the dry microhabitat under the OTC treatment does not follow the pattern of the abundance of total vascular plants, which could

indicate that litter abundance is mainly driven by the differential decomposition rate in the dry and wet microhabitats. Nevertheless, this relationship between warming and decomposition is not straightforward (Aerts 2006) and would deserve a proper litter decomposition experiment.

383 Moss length increment was also higher in the dry microhabitat in 2011, and was promoted by 384 OTC warming in 2012 (Figure 3). In the dryer 2011 year, the higher cover of vascular plants, i.e. 385 *Eriophorum* leaves in the dry microhabitat is the determinant factor for moss growth and might 386 have had a facilitating effect on S. fallax+rubellum in limiting the drying out of the moss carpet 387 (Grosvernier et al. 1994, Buttler et al. 1998, Walker et al. 2015). Instead, in the wetter year 2012, 388 when water limitation was not critical and engendered an overall higher length increment, the 389 warming effect by OTC's could promote further the moss growth in the dry microhabitat. Adaptive 390 seasonal acclimation has been shown for S. fallax, as reflected by the maintenance of relatively high 391 net photosynthetic rates to lower water contents and no response differences when plants were 392 collected from different heights above the water table (Titus et al. 1983). In our study, we pooled S. 393 fallax and S. rubellum, because we could not reliably distinguish these two species in our point-394 intercept field measures and in the frequency analysis on photographs. Nevertheless, some field 395 sampling in the plots showed that S. fallax is by far the most frequent species and that S. rubellum 396 occurs only on some very limited patches. Similarly to S. fallax, S. rubellum has been shown to 397 adapt and maintain similar height increment at different experimental water table depths (Robroek 398 et al. 2007). Therefore, shading may be critical for the development of these mosses, mediating soil 399 moisture and near ground temperature, which both are direct drivers of growth. Our results show 400 that shading can facilitate growth in dry years, but when wetter years are combined with warmer 401 conditions, in our case the OTC effect, this shading effect on growth is out-ruled by temperature. 402 Nevertheless, a better length increment of Sphagnum might also, on the long run, change the 403 structure of the moss carpet and consequently its moisture holding and transport capacity 404 (Dorrepaal et al. 2003). Therefore, when analyzing long term-effect of global change, not only the

405 change in species composition should be considered (Heijmans et al. 2008), but also the change of 406 vegetation structure might be critical. In their work, Walker et al. (2015) mentioned a mismatch in 407 the growth responses of different plant functional types to simultaneous climate and vegetation 408 change. This was due to the complexity of interacting factors between dwarf-shrubs, bryophytes 409 and graminoids, suggesting that some functional types reflecting a direct response to increased air 410 temperature, some others an indirect one.

The evolution of aboveground biomass over the seasons was well reflected in the NDVI, which was also higher in OTC plots. Interestingly, NDVI was also better correlated to vascular plants than to mosses, which explains the more obvious difference in the dry plot, where vascular plants are more abundant. Indeed, when mosses dry out temporarily in dry periods, which can be observed in the wet microhabitat where the moss carpet is less protected from wind, reflectance is lowered. This differentiation between mosses and higher plants is critical in studying the carbon cycle using airborne remote sensing (Yan et al. 2014).

418

#### 419 Conclusion

Our study on the response of plant cover to experimental warming in two contrasting hydrological peatland microhabitats – wet and dry – shows that experimental warming by OTC interacts with soil wetness, resulting in divergent responses of vascular plants and bryophytes. For vascular plants, the depth of rooting seems the key in determining the response of plant cover, whereas for mosses, the growth is under the dependence of various other interacting factors, such as shading by shrubs and large statured vascular plants, which contributes further to regulate the microclimate conditions in the moss carpet.

427

#### 428 Acknowledgments

429 This work was funded as part of the PEATWARM initiative through an ANR (French National 430 Agency for Research) grant (ANR-07-VUL-010). We acknowledge support from Labex 431 VOLTAIRE (ANR-10-LABX-100-01) and grant PSPB-013/2010 from Switzerland through the 432 Swiss Contribution to the enlarged European Union. P.M. acknowledges the support of the Swiss National Science Foundation (n<sup>0</sup> PBELP3 146538). The authors are also indebted to the Regional 433 434 Scientific Council of Natural Heritage of the Franche-Comté Region for permitting access to Le 435 Forbonnet site. We also thank M.-L- Toussaint for providing the meteorological data, E. Farrer and 436 L. M. Hallet for statistical advices, and B. Corboz for technical assistance.

#### 437 **References**

- Aerts, R. 2006. The freezer defrosting: global warming and litter decomposition rates in cold
  biomes. *Journal of Ecology* 94: 713-724.
- Aerts, R, Cornelissen, H. & Dorrepaal, E. 2006. Plant performance in a warmer world: general
  responses of plants from cold, northern biomes and the importance of winter and spring
  events. *Plant Ecology* 182: 65-77.
- Bragazza, L., Siffi, C., Iacumin, P. & Gerdol, R. 2007. Mass loss and nutrient release during litter
  decay in peatland: the role of microbial adaptability to litter chemistry. *Soil Biology & Biochemistry* 39: 257-267.
- Bragazza, L. 2008. A climatic threshold triggers the die-off of peat mosses during an extreme
  heatwave. *Global Change Biology* 14: 2688-2695.
- Bragazza, L., Parisod, J., Buttler, A. & Bardgett, R.D. 2013. Biogeochemical plant-soil microbe
  feedback in response to climate warming. *Nature Climate Change* 3: 273-277.
- Breeuwer, A., Robroek, B.J.M., Limpens, J., Heijmans, M.M.P.D., Schouten, M.G.C. & Berendse,
  F. 2009. Decreased summer water table depth affects peatland vegetation. *Basic and Applied Ecology* 10: 330–339.
- Bokhorst, S., Huiskes, A., Aerts, R., Convey, P., Cooper, E.J., Dalen, L., Erschbamer, B.,
  Gudmundsson, J., Hofgaard, A., Hollister, R.D., Johnstone, J. et al. 2013. Variable
  temperature effects of open top chamber at polar and alpine sites explained by irradiance
  and snow depth. *Global Change Biology* 19: 64-74.

- Buttler, A., Grosvernier, Ph. & Matthey, Y. 1998. Development of *Sphagnum fallax* diaspores on
  bare peat with implications for the restoration of cut-over bogs. *Journal of Applied Ecology*35: 800-810.
- 460 Buttler, A. 1992. Permanent plot research in wet meadows and cutting experiment. *Vegetatio*461 103(2): 113-124.
- 462 Clymo, R.S. 1970. The Growth of Sphagnum: Methods of Measurement. *Journal of Ecology* 58(1):
  463 13-49.
- 464 Cullings, K.W. 1996. Single phylogenetic origin of ericoid mycorrhizae within the Ericaceae.
   465 *Canadian Journal of Botany* 74: 1896-1909.
- 466 Dabros, A., Fyles, J.W. & Strachan, I.B. 2010. Effects of open-top chambers on physical properties
  467 of air and soil at post-disturbance sites in northwestern Quebec. *Plant Soil* 333 : 203-218.
- Delarue, F., Laggoun-Défarge, F., Buttler, A., Gogo, S., Jassey, V.E.J & Disnar, J.R. 2011. Effect
  of short-term ecosystem experimental warming on water extractable organic matter in an
  ombrotrophic *Sphagnum* peatland (Le Forbonnet, France). *Organic Geochemistry* 42: 10161024.
- 472 Delarue, F., Gogo, S., Buttler, A., Bragazza, L., Jassey, V., Bernard, G. & Laggoun-Défarge, F.
  473 2014. Indirect effects of experimental warming on dissolved organic carbon content in
  474 subsurface peat. J. Soils Sediments 14(11): 1800-1805.
- Delarue, F., Buttler, A., Bragazza, L., Grasset, L., Jassey, V.E.J., Gogo, S. & Laggoun, F. 2015.
  Experimental warming differentially affects microbial structure and activity in two
  contrasted moisture sites in a *Sphagnum*-dominated peatland. *Geoderma* (in press).
- 478 Dorrepaal, E., Aerts, R., Cornelissen, J.H.C. & Van Logtestijn, R.S.P. 2003. Summer warming and
  479 increased winter snow cover affect *Sphagnum fuscum* growth, structure and production in a
  480 sub-arctic bog. *Global Change Biology* 10: 93-104.
- 481 Dorrepaal, E., Cornelissen, J. H. C. & Aerts, R. 2007. Changing leaf litter feedbacks on plant
  482 production across contrasting sub-arctic peatland species and growth forms. *Oecologia* 151:
  483 251–261.
- 484 Dorrepaal, E., Aerts, R., Cornelissen, J.H.C., Van Logtestijn, R.S.P. & Callaghna, T.V. 2006.
   485 Sphagnum modifies climate-change impacts on subarctic vascular bog plants. Functional
   486 Ecology 20: 31–41.
- 487 Dorrepaal. E., Toet, S., van Logtestijn, R.S.P., Swart, E., van de Weg, M.J. Callaghan, T.V. &
  488 Aerts, R. 2009. Carbon respiration from subsurface peat accelerated by climate warming in
  489 the subarctic. *Nature* 460: 616-619.

- Gogo S., Laggoun-Défarge F., Delarue F. & Lottier N., 2011. Invasion of a Sphagnum-peatland by
   Betula spp and Molinia caerulea impacts organic matter biochemistry. Implications for
   carbon and nutrient cycling. *Biogeochemistry* 106: 1, 53-69
- Grosvernier, Ph., Matthey, Y. & Buttler, A. 1994 *Microclimate and physical properties of peat: new clues to the understanding of bog regeneration processes*. In: Restoration of Temperate
  Wetlands (Edited by B. D. Wheeler, S. C. Shaw, W. S. Fojt and R. A. Robertson), John
  Wiley & Sons, 435-450.
- Gerdol, R., 1995. The growth dynamics of sphagnum based on field-measurements in a temperate
  bog and on laboratory cultures. *Journal of Ecology* 83: 431-437.
- 499
- Hajek, T., Balance, S., Limpens, J., Zijlstra, M. & Verhoeven, J. T. A. 2011. Cell-wall
   polysaccharides play an important role in decay resistance of *Sphagnum* and actively
   depressed decomposition in vitro. *Biogeochemistry* 103: 45-57.
- Heijmans, M.M.P.D., Klees, H. & Berendse, F. 2002. Competition between Sphagnum
   magellanicum and Eriophorum angustifolium as affected by raised CO2 and increased N
   deposition. *Oikos* 97: 415–425.
- Heijmans, M.M.P.D., Mauquoy, D, van Geel, B. and Berendse, F. 2008. Long-term effects of
  climate change on vegetation and carbon dynamics in peat bogs. *Journal of Vegetation Science* 19: 307-320.
- Holden, J. 2005. Peatland hydrology and carbon release: why small-scale process matters. *Phil. Trans. R. Soc. A* 363: 2891–2913.
- Hollister, R.D. & Webber, P.J. 2000. Biotic validation of small open-top chambers in a tundra
  ecosystem. *Global Change Biology* 6: 835-842.
- Jacquemart, A.L. 1997. Vaccinium oxycoccus L. (Oxycoccus palustris Pers.) and Vaccinium
   microcarpum (Turcz. ex Rupr.) Schmalh. (Oxycoccus microcarpus Turcz. ex Rupr.).
   *Journal of Ecology* 85: 381–396.
- Jacquemart, A.L. 1998. Biological Flora of the British Isles Andromeda polifolia L. *Journal of Ecology* 86: 527-541.
- Jassey, V.E.J., Chiapusio, G., Gilbert, D., Buttler, A. Toussaint, M-L., Binet, P. 2011. Experimental
  climate effect on seasonal variability of polyphenols/phenoloxidase interplay along a narrow
  fen-bog ecological gradient in *Sphagnum fallax*. *Global Change Biology* 17(9): 2945-2957.
- Jassey V.E., Chiapusio G., Binet P., Buttler, A., Laggoun-Defarge, F., Delarue, F., Bernard, N.,
  Mitchell, E.A.D., Toussaint, M.L., Francez, A.J. & Gilbert, D. 2013. Above- and

belowground linkages in Sphagnum peatland: climate warming affects plant-microbial
 interactions. *Global Change Biology*, 19, 811–823.

- Keuper, F., Dorrepaal, E., Van Bodegom, P.M., Aerts, R., Van Logtestijn, R.S.P., Callaghan, T.V.
  & Cornelissen, J.H.C. 2011. A Race for Space? How Sphagnum fuscum stabilizes
  vegetation composition during long-term climate manipulations. *Global Change Biology*17(6): 2162-2171.
- Kuiper, J. J., Mooij, W. M., Bragazza, L. & Robroek, B.J.M. 2014. Plant functional types define
   magnitude of drought response in peatland CO2 exchange. *Ecology* 95: 123–131.
- Laggoun-Défarge, F., Gilbert, D., Buttler, A., Epron, D., Francez, A.-J., Grasset, L., Guimbaud, C.,
   Mitchell, E.A.D & Roy, J.-C. 2008. Effects of experimental warming on carbon sink
   function of a temperate pristine mire : the *PEATWARM* project. *Proceedings of the 13th International Peat Congress*, 599-602.
- Malmer, N., Svensson, B. & Wallen, B. 1994. Interactions between Sphagnum mosses and field
   layer vascular plants in the development of peat-forming systems. *Folia Geobotanica et Phytotaxonomica Praha* 29: 483-496
- Marion, G.M., Henry, G.H.R., Freckman, D.W., Johnstone, J., Jones, G., Jones, M.H., Levesque, E.,
  Molau, U., Mølgaard, P., Parsons, A.N., Svoboda, J. & Virginia, R.A. 1997. Open-top
  designs for manipulating field temperature in high-latitude ecosystems. *Global Change Biology* 3: 20–32.
- Nijp, J.J., Limpens, J., Metselaar, K., van der Zee, S.E., Berendse, F. & Robroek, B.J. 2014. Can
  frequent precipitation moderate the impact of drought on peatmoss carbon uptake in
  northern peatlands? *New Phytologist* 203:70-80.
- 545 Pinheiro, J.C. & Bates, D.M. 2000. *Linear mixed-effects models: basic concepts and examples*.
  546 Springer ISBN 978-1-4419-0318-1.
- R Core Team 2014. R: *A language and environment for statistical computing*. R Foundation for
  Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL http://www.Rproject.org/.
- Robroek, B.J.M., Limpens, J., Breeuwer, A. & Schouten, M.G.C. 2007. Effect of water level and
   temperature on performance of four *Sphagnum* species. *Plant Ecology* 190: 97-107.
- Robroek, B.J.M., Heijboer, A., Jassey, V.E.J., Hefting, M.M., Rouwenhorst, T.G., Buttler, A. &
  Bragazza, L. 2013. Snow cover manipulation effects on microbial community structure and
  soil chemistry in a mountain bog. *Plant Soil* 369 (1-2) : 151-164.

- Shaver, G.R. and Billings, W.D. 1977. Effect of day-length and temperature on root elongation in
   tundra graminoids. *Oecologia* 28: 57-65.
- Soudani, K, Hmimina, G., Delpierre, N., Pontailler J.-Y., Aubinet, M., Bonal, D., Caquet, B., de
   Grandcourt, A., Burban. B., Flechard. C., Guyon D. et al. 2012. Ground-based Network of
   NDVI measurements for tracking temporal dynamics of canopy structure and vegetation
   phenology in different biomes. *Remote Sensing of Environment* 123: 234–245.
- Sullivan, P.F. & Welker, J.M. 2005. Warming chambers stimulate early season growth of an arctic
   sedge: results of a minirhizotron field study. *Oecologia* 142: 616-626.
- Titus, J.E., Wagner, D.J. & Stephens, M.D. 1983. Contrasting water relations of photosynthesis for
   two Sphagnum mosses. Ecology 64 (5): 1109-1115.
- Turetsky, M.R., Bond-Lamberty, B., Euskirchen, E., Talbot, J., Frolking, S., McGuire, A.D. &
  Tuittila, E.S. 2012. The resilience and functional role of moss in boreal and arctic
  ecosystems. *New Phytologist* 196: 49-67.
- Wallen, B. 1986. Above and below ground dry mass of the three main vascular plants on
  hummocks on a subarcticp peat bog. *Oikos* 46: 51-56.
- Walker, T.N., Ward, S.E., Ostle, N.J. & Bardgett, R.D. 2015. Contrasting growth responses of
  dominant peatland plants to warming and vegetation composition. Oecologia, DOI
  10.1007/s00442-015-3254-1.
- Weltzin, J. F., Pastor, J., Harth, C., Bridgham, S. D., Updegraff, K., & Chapin, C. T. 2000.
  Response of bog and fen plant communities to warming and water-table manipulations. *Ecology* 81: 3464–3478.
- Weltzin, J. F., Harth, C., Bridgham, S. D., Pastor, J. & Vonderharr, M. 2001. Production and
   microtopography of bog bryophytes: response to warming and water-table manipulations.
   *Oecologia* 128: 557–565.
- Weltzin, J.F., Brigham, S.D., Pastor, J., Chen, J. & Harth, C. 2003. Potential effects of warming and
  drying on peatland plant community composition. *Global Change Biology*, 9, 141-151.
- Yu, Z.C. 2011. Holocene carbon flux histories of the world's peatlands: global carbon-cycle
   implications. *Holocene* 21: 761-774.
- Yan, W et al. 2014. Differentiating moss from higher plants is critical in studying the carbon cycle
  of the boreal biome. *Nature Communications* 5: 4270. DOI: 10.1038/ncomms5270
  |www.nature.com/naturecommunications.
- 586
- 587

588	
589	

#### 590 **Captions to figures**

591

592 Figure 1: Changes of the relative frequencies of dominant plant species from 2008 to 2012 (no 593 records in 2011) in the Forbonnet peatland (French Jura). Mean values (n=3) and standard error 594 bars are given for dry (solid line) and wet (broken line) microhabitats, and for control (black line) 595 and OTC (grey lines) plots. Significance of Linear Mixed Model is given for treatment effect (T) 596 and for year effect (Y). Species that are not illustrated: *Carex pauciflora* (appeared in 2009 and then 597 was only seen in one plot the years after), Carex limosa (appeared only in 3 plots in 2009), Calluna 598 vulgaris (only in 2 different plots in different years), *Pinus rotundata* (only in one plot in one year). 599 600 Figure 2: Changes of the relative frequencies of species from April to September 2012 in the 601 Forbonnet peatland (French Jura). Mean values (n=3) and standard error bars are given for dry

(solid line) and wet (broken line) microhabitats, and for control (black line) and OTC (grey lines)
plots. Significance of Linear Mixed Model is given for treatment (T) and month effect (M). Species
that are not illustrated; with no significant differences: *Scheuchzeria palustris, Eriophorum vaginatum, Carex pauciflora, Carex rostrata, Drosera rotundifolia, Polytrichum strictum*; present
only in one plot: *Calluna vulgaris, Carex pauciflora* and *Polytrichum strictum*.

607

 $\begin{array}{ll} 608 \quad Figure \ 3: \ Length \ increment \ (mm) \ of \ Sphagnum \ mosses \ in the \ different \ treatments \ (OTC \ vs \ control) \\ only \ and \ microhabitats \ (dry \ vs \ wet) \ in \ 2011 \ (a: \ May \ 13 - November \ 27, n=60) \ and \ 2012 \ (b: \ March \ 30 - 610 \ November \ 6, \ n=120) \ in \ the \ Forbonnet \ peatland \ (French \ Jura). \ Vertical \ bars \ represent \ standard \\ errors. \ Scales \ of \ both \ graphs \ are \ different. \ Significance \ levels \ of \ Linear \ Mixed \ Model \ are \ given \ for \\ 612 \ microhabitat \ (B) \ and \ treatment \ effect \ (T), \ and \ for \ their \ interaction \ (B \ x \ T). \end{array}$ 

613

*Figure 4:* Normalised Difference Vegetation Index (NDVI) at plot scale in the dry (a) and wet (b) microhabitats in year 2012 in the Forbonnet peatland (French Jura). Curves represent mean values (n=3) for control (black) and OTC (grey) plots. Significance codes for ANOVA's are (\*\*\*) p<0.001; (\*\*) p<0.01; (\*) p<0.05; (.) p<0.1.

- 618
- 619

620 Figure 1

- 621
  622
  623
  624
  625
- 626 Figure 2
- 627

628	
629	Figure 3
630	
631	

632			
633			
634			
635	Figure 4		
636			

637

638 Electronic appendices

639

Supporting Information to the paper Buttler et al. 2015. Experimental warming interacts with soil moisture to discriminate plant responses in an ombrotrophic peatland. *Journal of Vegetation Science*. Appendix S1. Relationship between point-intercept frequency measures of vegetation and picture frequency analysis from the Forbonnet peatland (French Jura) for a: total vascular plants (Pearson r: 0.733, p<0.01), b: litter (Pearson r=0.873, p<0.001) and c: total *Sphagnum* and *Polytrichum strictum* (Pearson r=0.942, p<0.001).

646

The correlation between frequencies obtained by means of field point-intercepts and photographic analyses gives a good linear relation with vascular plants but shows an underestimation of mosses with the picture analysis. This is because leaves of vascular plants might hide the understory of mosses which than cannot be recorded in the one-layer photographic method, whereas in the field, the needle can hit more than one species or individual. This underestimation is mostly noticeable at intermediate moss cover, when vascular plants are most abundant. Conversely, the picture analysis overestimates litter when it is dense.

655

656

657





**Journal of Vegetation Science** 





