



Multiplication of microbes below 0.690 water activity: implications for terrestrial and extraterrestrial life

Andrew Stevenson, Jürgen Burkhardt, Charles S. Cockell, Jonathan A. Cray,
Jan Dijksterhuis, Mark Fox-Powell, Terence P. Kee, Gerhard Kminek, Terry J.
Mcgenity, Kenneth N. Timmis, et al.

► To cite this version:

Andrew Stevenson, Jürgen Burkhardt, Charles S. Cockell, Jonathan A. Cray, Jan Dijksterhuis, et al..
Multiplication of microbes below 0.690 water activity: implications for terrestrial and extraterrestrial
life. *Applied and Environmental Microbiology*, 2015, 17 (2), pp.257-277. 10.1111/1462-2920.12598 .
insu-01206203

HAL Id: insu-01206203

<https://insu.hal.science/insu-01206203>

Submitted on 7 Mar 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



UNIVERSITY OF LEEDS

This is a repository copy of *Multiplication of microbes below 0.690 water activity: implications for terrestrial and extraterrestrial life*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/121554/>

Version: Accepted Version

Article:

Stevenson, A, Burkhardt, J, Cockell, CS et al. (12 more authors) (2015) Multiplication of microbes below 0.690 water activity: implications for terrestrial and extraterrestrial life. *Environmental Microbiology*, 17 (2). pp. 257-277. ISSN 1462-2912

<https://doi.org/10.1111/1462-2920.12598>

© 2014 Society for Applied Microbiology and John Wiley & Sons Ltd. This is the peer reviewed version of the following article: Stevenson, A., Burkhardt, J., Cockell, C. S., Cray, J. A., Dijksterhuis, J., Fox-Powell, M., Kee, T. P., Kminek, G., McGenity, T. J., Timmis, K. N., Timson, D. J., Voytek, M. A., Westall, F., Yakimov, M. M. and Hallsworth, J. E. (2015), Multiplication of microbes below 0.690 water activity: implications for terrestrial and extraterrestrial life. *Environ Microbiol*, 17: 257–277. doi:10.1111/1462-2920.12598, which has been published in final form at <https://doi.org/10.1111/1462-2920.12598>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Multiplication of microbes below 0.690 water activity: implications for terrestrial and extraterrestrial life

Andrew Stevenson¹, Jürgen Burkhardt², Charles S. Cockell³, Jonathan A. Cray¹, Jan Dijksterhuis⁴, Mark Fox-Powell³, Terence P. Kee⁵, Gerhard Kminek⁶, Terry J. McGenity⁷, Kenneth N. Timmis⁸, David J. Timson¹, Mary A. Voytek⁹, Frances Westall¹⁰, Michail M.Yakimov¹¹, John E. Hallsworth¹

¹Institute for Global Food Security, School of Biological Sciences, MBC, Queen's University Belfast, Belfast, BT9 7BL, Northern Ireland. ²Plant Nutrition Group, Institute of Crop Science and Resource Conservation, University of Bonn, Karlrobert-Kreiten-Str. 13, D-53115 Bonn, Germany. ³UK Centre for Astrobiology, School of Physics and Astronomy, University of Edinburgh, Edinburgh EH9 3JZ, UK. ⁴CBS Fungal Biodiversity Centre, Uppsalalaan 8, NL-3584, CT Utrecht, Netherlands. ⁵University of Leeds, School of Chemistry, Leeds, LS2 9JT, West Yorkshire, UK. ⁶ESA-ESTEC, Keplerlaan 1, 2200 Noordwijk, The Netherlands. ⁷University of Essex, School of Biological Sciences, Colchester, CO4 3SQ, Essex, UK. ⁸Institute of Microbiology, Technical University Braunschweig, Spielmannstrasse 7, D-38106 Braunschweig, Germany. ⁹NASA Headquarters, Washington, DC 20546-0001, USA. ¹⁰Centre de Biophysique Moléculaire, CNRS, Rue Charles Sadron, and Centre de Recherches sur les Matériaux à Haute Température, 1D, avenue de la recherche scientifique, 45071 Orléans Cedex 2, France. ¹¹Istituto per l'Ambiente Marino Costiero, CNR, 98122 Messina, Italy.

*For correspondence. E-mail j.hallsworth@qub.ac.uk; Tel: (+44) 289097 2314; Fax: (+44) 289097 5877.

Running title: Multiplication of microbes at low water-activity

Keywords: Aspergillus, habitability, halophilic Archaea and Bacteria, limits for life, Mars, xerophilic fungi, water activity

33 Summary

34

35 Since a key requirement of known life-forms is available water (water-activity; a_w), searches
36 for signatures of past life in terrestrial and extraterrestrial environments have recently
37 targeted places known to have contained significant quantities of biologically available water.
38 The lower limit of water activity that enables cell division is ~ 0.605 which, until now, was only
39 known to be exhibited by a single eukaryote; the sugar-tolerant, fungal xerophile *Xeromyces*
40 *bisporus*. The first forms of life on Earth were, however, prokaryotic. Furthermore, early life
41 on Earth inhabited high-salt environments, suggesting an ability to withstand low water-
42 activity. Recent evidence indicates that some halophilic Archaea and Bacteria have water-
43 activity limits more or less equal to those of *X. bisporus*. Regardless of species, cellular
44 systems are sensitive to minute differences in water activity (of < 0.005 a_w -units) so there is a
45 need to determine water-activity values to three decimal places. We discuss water activity in
46 relation to the limits of Earth's present-day biosphere; the possibility of microbial
47 multiplication by utilizing water from thin, aqueous films or non-liquid sources; whether
48 prokaryotes were the first organisms able to multiply at the 0.605 - a_w limit; and whether
49 extraterrestrial aqueous milieu of ≥ 0.605 a_w can resemble fertile microbial habitats found on
50 Earth.

51

52

53

54 Introduction

55 Given the fact that water is one of the principal ingredients of cellular life (Daniel et al.,
56 2004), insights into the minimum water requirements of cells are imperative to understanding
57 the functionality of living-systems at every level (from biomacromolecule to biosphere), as
58 well as the origins of life, in an environmental context. The generally held opinion is that life
59 appeared independently on Earth and, possibly, elsewhere in the Solar System (Clancy et
60 al., 2005); though one other explanation for the presence of life on Earth is that it appeared
61 on another planet and was transported here in the form of prokaryotes or their ancestors (an
62 idea known as panspermia; Thomson, 1871). Until recently, eukaryotic microbes have held
63 the record for life under water-constrained conditions, as some species are capable of cell
64 division down to a water activity (a_w)¹ of 0.605 at high sugar concentrations (Pitt and
65 Christian, 1968; Williams and Hallsworth, 2009). Whereas such data have formed the basis
66 of international policy for planetary protection in relation to space-exploration missions (see

¹ Water activity, the mole fraction of water, is defined by an equation (water activity = vapour pressure of the solution/vapour pressure of the water) which is derived from Raoult's Law; this parameter and its derivation are discussed in detail by Brown (1990) and Grant (2004).

below), sugar-rich substrates have very limited applicability to those extraterrestrial habitats with which we are familiar. Historically, the accepted limit for cell division of prokaryotic microbes has been 0.755 a_w ; this applied to a small fraction of halophilic species at high salt concentrations (for references, see Grant, 2004). However, both culture-based and culture-independent studies provide evidence for multiplication and metabolic activity of halophilic Archaea and Bacteria in the range 0.680 to 0.605 a_w , both in their natural habitats in situ, and in vitro (Javor, 1984; Yakimov et al., 2014; A. Stevenson et al., submitted). Other studies have shown that, whereas the vast majority of yeasts and fungi are active somewhere within the range 1 to 0.720 a_w (Pitt, 1975; Brown, 1976), only ~12 species have been observed to grow and/or germinate at < 0.700 a_w (Williams and Hallsworth, 2009; A. Stevenson et al., submitted). Here, we discuss the evidence for microbial activity below at or below 0.690, which represents the very edge of the functional biosphere on Earth. Low water activity is also discussed in relation to early life on Earth, the plausibility of cell division in habitable extraterrestrial environments in which biologically available water is present, and a series of unanswered scientific questions.

82

83 Water-activity at the outer edges of the microbial biosphere

84

85 The primary physical determinants of the habitable space on Earth are temperature and
86 water activity; these parameters are also used to designate the ‘Special Regions’ of Mars in
87 which microbial cell-division might feasibly take place (Beaty et al., 2006; Kminek et al.,
88 2010; J. D. Rummel et al., unpublished)². The temperature window over which microbes are,
89 collectively, capable of cell division (i.e. from -18 to +122°C; Takai et al., 2008; Chin et al.,
90 2010) spans ≤ 40% of the entire range of temperatures to which life-systems on Earth can
91 be exposed; i.e. from approximately -90°C to ≥ 250°C (for some hydrothermal vents; Fig.
92 1a). By contrast, environmental water-activity values range from 1 to 0 and most cellular
93 systems of known life-forms on Earth are only active in the range, or a segment of the range,
94 1 to 0.900 a_w (Fig. 1b; Brown, 1976; Grant, 2004). For example, there is a drop-off in
95 measurable metabolic activity in many soils at ≤ 0.890 a_w (Moyano et al., 2012; 2013;
96 Stevenson and Hallsworth, 2014). However, metabolic activity and cell-division has been

² Planetary protection in relation to space missions aims to protect those planets where spacecraft are landed, as well as Earth, from accidental contamination with non-native life-forms (Kminek et al., 2010; 2014). Mars Special Regions have been defined according to the activities of the NASA Mars Exploration Program Analysis Group (MEPAG), Special Regions-Scientific Analysis Group 1 (SR-SAG1) and the Committee on Space Research (COSPAR), which is part of the International Council for Science. Both these committees conservatively recommended 0.500 a_w as the limit beyond which no known terrestrial microorganism is capable of multiplication; implying that any environment of Mars with a water activity of > 0.500 may potentially enable proliferation of xerophilic microbes if they happened to arrive as accidental passengers on spacecraft sent from Earth (Fig. 1; Beaty et al., 2006; Kminek et al., 2010). A revised analysis of Mars Special Regions is currently underway by the MEPAG SR-SAG2 (J. D. Rummel et al., unpublished).

97 reported below 0.900 a_w for a great number of xerotolerant/philic and halotolerant/philic
98 microbes (Brown, 1976; Grant, 2004), and even below 0.755 a_w for both eukaryotic and
99 prokaryotic species (Javor, 1984; Williams and Hallsworth, 2009; Yakimov et al., 2014; A.
100 Stevenson et al., submitted). Of the microbes known to multiply below 0.720, the majority
101 (unlike *X. bisporus*) are not obligate osmophiles that must inhabit sugar-rich substrates;
102 these include halophilic prokaryotes and xerophilic fungi such as *Aspergillus penicillioides*
103 and *Eurotium herbariorum* (Samson and Lustgraaf, 1978; Williams and Hallsworth, 2009;
104 Yakimov et al., 2014; A. Stevenson et al., submitted)³. Even for the most xerophilic microbes
105 thus far characterized (see Pitt, 1975; Javor, 1984; Williams and Hallsworth, 2009; A.
106 Stevenson et al., submitted), rates of cell division typically decrease by an order of
107 magnitude between 0.870 and 0.770 a_w , and by a further order of magnitude between 0.770
108 and 0.670 a_w (Stevenson and Hallsworth, 2014; A. Stevenson et al., submitted). There are
109 only reports of cell division for between 20 and 30 microbial species or communities at $\leq$
110 0.690 a_w (see Javor, 1984; Yakimov et al., 2014; A. Stevenson et al., submitted). Whereas
111 all of these species are extreme, obligately xerophilic eukaryotes or extreme, obligately
112 halophilic prokaryotes which have low rates of cell division - or are incapable of growth -
113 close to 1 a_w , the ultimate limit for multiplication of even the most resilient strains appears to
114 be ~ 0.61 a_w (Pitt and Christian, 1968; A. Stevenson et al., submitted). For microbes on
115 Earth, therefore, biotic activity spans approximately 40% of the available water-activity
116 range, thus emphasizing the potency of water as a determinant of the functional biosphere.
117 The overwhelming majority of microbial systems are metabolically active somewhere within
118 the ranges 5 to 40°C, and 1 to 0.900 a_w , which represent even smaller portions of the
119 environmentally pertinent temperature and water-activity ranges; i.e. only 10% in each case
120 (Fig. 1). Of the microbial systems characterized thus far, the 20 to 30 known to be active at $\leq$
121 0.690 a_w (Fig. 1; Javor, 1984; Yakimov et al., 2014; A. Stevenson et al., submitted) represent
122 the most extreme forms of life to have penetrated these kinds of hostile environment⁴.

123 Some reports have alluded to the possibility of microbial growth and metabolism at
124 the otherwise unprecedented water-activity values of 0.382 (for deep-sea halophiles in
125 $MgCl_2$ -saturated brine; van der Wielen et al., 2005), < 0.450 (for halophiles in the $CaCl_2$ -rich,
126 Antarctic Don Juan Pond; Siegel, 1979), 0.500 (Actinobacteria isolated from algal mats and
127 cultured in soil-based substrates; Doroshenko et al., 2005; 2006; Zvyagintsev et al., 2009;

³ This has implications for preventing contamination of other planetary bodies (see above) which, as far as we know, lack sugar-rich environments, during space exploration missions.

⁴ Habitats which have sufficiently low water-activity to exclude almost all forms of life on Earth and, therefore, have a characteristically low biodiversity (especially those of < 0.690 a_w) are fertile habitats for those extremophiles which thrive there due to minimal competition and, frequently, a lack of grazers and predators (for references, see Cray et al., 2013b). Such low-water activity habitats are, however, typically too biologically hostile and insufficiently biodiverse to act as open habitats for microorganisms (Cray et al., 2013b; Lievens et al., 2014; Oren and Hallsworth, 2014).

2012), 0.570 (for halophiles in acidic saline lakes; Mormile et al., 2009), 0.600 (for germination of *Wallemia sebi* [a xerophilic basidiomycete] on high-sugar substrates; Frank and Hess, 1941) and 0.600 (reported value for optimum growth of halophiles (Jaenicke and Bohm, 1998), and biotic activity in salt lakes; Cobucci-Ponzano et al., 2006). Some of these values were hypothetical (see below), and the other claims have not been accepted or have been refuted by authors of a number of subsequent studies (Pitt and Christian, 1968; Wynn-Williams, 1996; Beaty et al., 2006; Hallsworth et al., 2007; Kminek et al., 2010; Oren, 2011; Stevenson and Hallsworth, 2014; A. Stevenson et al., submitted; J. D. Rummel et al., unpublished). The Don Juan Pond (located within the McMurdo Dry Valleys, Antarctica) is a CaCl_2 -saturated brine-pool situated in a closed basin and fed by seasonal melt-water streams and deliquescent seepages, both of which are thought to deliver CaCl_2 to the lake (Dickson, 2013). Its volume fluctuates but is typically $\sim 3000 \text{ m}^3$ (slightly larger than an Olympic swimming pool), and it is amongst the most saline large-scale bodies of water known on Earth. This pond rarely, if ever, freezes despite winter temperatures of $\leq -51^\circ\text{C}$ (Siegel 1979; Marion 1997; Grant, 2004). While annual temperatures of the pond's water and the surrounding sediments are occasionally above 0°C , they remain below -20°C for the majority of the year (Samarkin et al., 2010) so it is highly unlikely that microbial life could multiply there (for references, see Chin et al., 2010; Kminek et al., 2010; J. D. Rummel et al., unpublished). Saturated solutions of CaCl_2 , as found in the Don Juan Pond, are highly chaotropic and are therefore likely to prevent microbial growth (and may even be sterile environments; Duda et al., 2004; 2005; Hallsworth et al., 2007; Cray et al., 2013a; 2013b; Oren, 2013; Yakimov et al., 2014). Nitrous oxide emissions recorded from the surrounding sediments, frequently attributed to the biological transformation of nitrogenous compounds, are apparently the result of abiotic reactions between brine nitrates and Fe^{II} -bearing minerals (Samarkin et al., 2010). The water activity of the MgCl_2 -dominated, deep-sea hypersaline brine studied by van der Wielen et al. (2005) is ~ 0.382 at the in situ temperature of 14.5°C (Winston and Bates, 1960; Hallsworth et al., 2007). Culture-dependent and culture-independent studies of this brine, and investigations into the biophysics of macromolecular interactions, indicate that both its potent chaotropicity (even at water-activity values which would otherwise be permissive for cell division) and exceptionally low water-activity prohibit life processes (Hallsworth et al., 2007; Yakimov et al., 2014), as these parameters do for solutions of comparable salts (Winston and Bates, 1960; Duda et al., 2004; Hallsworth et al., 2003a; Kminek et al., 2010; Oren, 2011; Cray et al., 2013a; 2013b). Speculations that microbial metabolism and cell division occur at $\sim 5 \text{ M}$ MgCl_2 are inconsistent with the virtual sterility of the Dead Sea when MgCl_2 concentrations become elevated (Oren, 1999; 2010; Oren, 2013) or the CaCl_2 -dominated Don Juan Pond (Siegel et al., 1983; Samarkin et al.,

2010; Oren, 2013) where concentrations of divalent chloride salts reach critical concentrations which are prohibitive for all life processes (Hallsworth et al., 2007; Cray et al., 2013a; Oren, 2013; Yakimov et al., 2014). Although there is a theoretical possibility that some microbes have evolved specialised structures which isolate cells from such hostile habitats whilst permitting biotic activity, to our knowledge no such structures have yet been reported for any microbial species in any type of extremely chaotropic (e.g. Hallsworth et al., 2007; Yakimov et al., 2014) or low water-activity (≤ 0.600) environment.

Reports of germination and subsequent cell division during germ-tube formation of several Actinobacteria (i.e. *Streptomyces albidoflavus* [syn. *Streptomyces odorifer*], *Streptomyces rectiviolaceus*, and a *Micromonospora* strain) at 0.500 a_w (which were carried out by one group: Doroshenko et al., 2005; 2006; Zvyagintsev et al., 2009; 2012) are apparently erroneous (see Stevenson and Hallsworth, 2014). Independent studies have demonstrated that none of these species was capable of growth below 0.895 a_w , and the theoretical water-activity minimum for the most xerotolerant (a strain of *Streptomyces albidoflavus*) was ~ 0.870 (Stevenson and Hallsworth, 2014). Proposed limits of 0.570 or 0.600 a_w for biotic activity of halophiles were speculative (i.e. not derived from determinations of water-activity; Jaenicke and Bohm, 1998; Mormile et al., 2009; Cobucci-Ponzano et al., 2006), and sources of experimental error in studies of *W. sebi* germination have been discussed previously (Pitt and Christian, 1968). Furthermore, apparent microbial growth within terrestrial brine lakes which can reach values of ≤ 0.600 a_w may have actually occurred at higher water-activity values given the seasonal and other temporal fluctuations of the in situ salt concentrations (Oren, 1988; 1993; Cobucci-Ponzano et al., 2006; Mormile et al., 2009).

Although the established temperature minima for multiplication of the most psychrophilic microbes are in the region of -15 to -18°C (for references, see Chin et al., 2010; Kminek et al., 2010), there are numerous sources of evidence for metabolic activity considerably below this range (Kminek et al., 2010; J. D. Rummel et al., unpublished). By contrast, there is a paucity of data to demonstrate metabolic activity below the accepted water-activity minimum for microbial cell division (i.e. 0.605; Kminek et al., 2010; Yakimov et al., 2014; A. Stevenson et al., unpublished; J. D. Rummel et al., unpublished). In relation to the water-activity limit for life, it is noteworthy that trehalose, a hygroscopic substance which accumulates in desiccated microbial cells and may facilitate the acquisition and retention of water, cannot efficiently absorb water from the vapour phase at equilibrium relative humidities of less than $\sim 50\%$, equivalent to 0.500 a_w (Fakes et al., 2000). Whereas some enzymes can remain catalytic at water activities of < 0.500 (Kurkal et al., 2005; Lopez et al., 2010), there is evidence that DNA becomes disordered, and is therefore no longer

transcribable, below a water activity of 0.550 (Falk et al., 1963). Furthermore, strand breaks have been recorded at 0.530 a_w in bacterial cells (Asada et al., 1979). It has, therefore, long-been considered unlikely that cellular systems could function at water activities substantially lower than 0.600 (e.g. Pitt, 1975; Brown, 1976; 1990; Sutton and Hildebrand, 1985; J. D. Rummel et al., unpublished). However, interactions between the various factors which determine the biophysical limits for cellular integrity and biotic activity at low water-activity are complex and have yet to be fully elucidated. Macromolecular integrity and functionality can depend on the net effect of prevailing conditions such as temperature, chaos/kosmotropicity, pressure and water activity (Hallsworth, 1998; Hallsworth et al., 2007; Williams and Hallsworth, 2009; Bhaganna et al., 2010; Chin et al., 2010; Yakimov et al., 2014) and it may be possible that, in some as-yet-undiscovered environments, cells are capable of metabolism at $< 0.600 a_w$.

Microbial cell division via utilization of water which is not in the bulk liquid-phase

Water is more or less ubiquitous on Earth and in other parts of the Solar System (Bradley et al., 2014; Küppers et al., 2014); it may be present within the atmospheres, subsurface, rocks and regolith, polar ice-sheets, glaciers, and/or subsurface oceans of planetary bodies, in vapour plumes extruded into space, and – indeed – within space itself⁵. Whereas here on Earth, we tend to be most familiar with water in its bulk-liquid phase, in both terrestrial and extraterrestrial environments, it can also be present in a variety of forms. In addition to ice and vapour, these include thin aqueous films on/at various types of surfaces and interfaces, or as molecules hydrating mineral, organic, and other substances (Kminek et al., 2010; Toner et al., 2014; J. D. Rummel et al., unpublished). Liquidity of water is determined by temperature, pressure, the presence of solutes and/or gases, and molecular interactions between other materials or substances and water molecules - as well as processes such as salt deliquescence, sublimation of ice, frost formation, condensation or dew-formation on surfaces or within the gaseous phase, aerosol formation, and precipitation (Watanabe and Mizoguchi, 2002; Jepsen et al., 2007; Möhlmann; 2008; 2009; 2012; Argyris et al., 2008; Chin et al., 2010; Pavlov et al., 2010; Bing and Ma, 2011).

Thin aqueous films can exist on various surfaces including those of ice and biological and mineral structures, and the water within these films can remain in the liquid phase under a wide range of conditions (Pearson and Derbyshire, 1974; Raviv et al., 2001; Wolfe et al., 2002; Jepsen et al., 2007; Möhlmann; 2004; 2008; 2009; 2011; 2012; J. D. Rummel et al.,

⁵ See Waite et al. (2006); Nimmo et al. (2007); Tosca et al. (2008); Campins et al. (2010); Sohl et al. (2010); Carter et al. (2013); Martínez and Renno (2013); and Bradley et al. (2014).

unpublished). The depth of thin films can range from > 1 mm to a monolayer of water molecules (~0.3 nm; Möhlmann, 2004; 2005), and they can be stable (Möhlmann, 2012) or highly ephemeral (Burkhardt and Hunsche, 2013). At the temperatures and pressures which typically prevail in Earth's biosphere, aqueous films of ~1 mm are primarily made up of water which is biologically available (e.g. Qvit-Raz et al., 2008, Burch et al., 2013). Whereas we speculate that single-monolayer films do not provide water that can be accessed by cellular systems. It has, however, been suggested that microbes can utilize fluid films with a mean thickness equivalent to that of three water molecules (Harris, 1981; Beaty et al., 2006); a hypothesis that may be inconsistent with the lack of solute diffusion in very thin films (Derjaguin and Churaev, 1986; Hu and Wang, 2003) which indicate that the water in films as thin as this is not in the liquid phase⁶. Despite the circumstantial evidence (see also Rivkina et al., 2000), there is a paucity of data thus far available which convincingly demonstrate that water in thin films that are equivalent to between one and three water molecules in depth is biologically available.

There are three possible sources of liquid water in otherwise desiccated and cold areas such as those which are characteristic of Mars: (1) interfacial water as a thin film (several water-molecular in depth) forming on mineral surfaces by adsorption or, on ice, as pre-melted ice (Dash et al. 2006, Möhlmann, 2011); (2) brines forming on salt crystals via deliquescence; and (3) subsurface melt-water below an ice covering due to a solid-state 'greenhouse' effect (Möhlmann, 2011). Process (2) is a particularly effective mechanism by which liquid water can be generated on Earth and, almost certainly, in extraterrestrial locations (Möhlmann, 2011). The condensing water vapour can potentially reach the dry weight of the deliquescent salt, and will exceed it if the humidity exceeds the deliquescence relative humidity. Deliquescence of NaCl, as equilibrium relative humidity increases from 65 to 80%, can be observed in Movie S1. Most salts (and, indeed, many organic substances) are hygroscopic and will attract water to their surface at equilibrium relative humidities of ≤ 100%. Each salt becomes deliquescent at a specific relative humidity, thereby dissolving as the water vapour condenses. The deliquescence relative humidity for a given salt and its (usually slight) temperature-dependence quantitatively correspond to both the water activity values of, and equilibrium relative humidity values for, saturated solutions of a given salt (Winston and Bates, 1960). If the equilibrium relative humidity is higher than a salt's deliquescence relative humidity, the water activity of the salt solution will equilibrate with the relative humidity of the atmosphere, so the salt solution will become more dilute. Mixtures of substances (e.g. mixtures of different salts or salts plus sugars) will have a deliquescence

⁶ This inconsistency also raises the possibility that the high water-activity values associated with very thin films (Harris, 1981; Papendick and Campbell, 1981) could be a consequence of methodological error.

relative humidity below that of each individual component (Mauer and Taylor, 2010). In addition to the reduced water activity, salts also reduce the freezing point, and cryobrines may be stable far below the melting point of water, e.g. under Martian conditions (Möhlmann, 2011, Martínez and Renno, 2013).

Within the Earth's biosphere, brine formation may play a role for diverse microbial species – especially those that are halotolerant or halophilic – which are located within bioaerosols, or on mineral or biological surfaces (e.g. leaf surfaces) and are exposed to humid air (Potts, 1994). For example, adapted species can reproduce within the phyllosphere of salt-exuding desert plants (Qvit-Raz et al., 2008, Burch et al., 2013) and, at subzero temperatures, in supercooled water in the atmosphere (Sattler et al., 2001). *Pseudomonas syringae*, which is not haliophilic, is a species widely transported within bioaerosols and its cells are highly effective as ice nuclei because they have protein coatings that cause water to freeze at relatively warm temperatures (Christner et al., 2008; Morris et al., 2014). Being surrounded by ice, they may benefit from the solid-state greenhouse effect which involves the internal formation of thin films due to the penetration and retention of shortwave radiation within the ice.

Microbes can obtain water from the vapour-phase, a process which has been observed in lichens (Lange et al., 2006; Pintado and Sancho, 2002) as well as the propagules of various species (Waldham and Halvorson, 1954; Pasanen et al., 1991; Reponen et al., 1996). Other studies have demonstrated that microbial cells also generate considerable quantities of water via their metabolic activity (Oriol et al., 1988; Nagel et al., 2001; Marcano et al., 2002; Kreuzer-Martin et al., 2005; 2006; de Goffau et al., 2011), up to 70% of the cell's water according to radio-labelled gas uptake experiments (Kreuzer-Martin et al. 2005; 2006). Spore germination of powdery mildews, such as by the *Erysiphe* and *Uncinula* species, has been observed at low equilibrium relative humidities (0 to 10%) without a visible extracellular source of liquid water (Brodie and Neufield, 1942; Manners and Hossain, 1963; Carroll and Wilcox, 2003); although it is not clear whether condensation processes and/or thin films might act to shuttle water to the cell. Desiccated lichens are able to absorb water at an equilibrium relative humidity of $\geq 82\%$ and thereby commence photosynthesis (Pintado and Sancho, 2002; Lange et al. 2006). Various lines of evidence suggest that microorganisms may be capable of cell division without an extracellular supply of liquid water (see also Miller and Chibnall, 1932; Yarwood, 1950; Peterson and Cowling, 1972; Lange et al., 1986; 1994). However, there is a paucity of convincing data to irrefutably affirm this hypothesis. Furthermore, systematic studies of water-activity limits for cell division of phylogenetically diverse extremotolerant and extremophilic microbes suggest that cell division would be implausible at values much below 0.600 a_w (i.e. 60% equilibrium relative

304 humidity) (Pitt and Christian, 1968; Brown, 1976; Williams and Hallsworth, 2009; A.
305 Stevenson et al., unpublished). This question is equally pertinent to life on Earth, and the
306 aqueous milieu found elsewhere in the Solar System (not least in relation to planetary
307 protection; see above).

308
309 Implications for the evolution of microbial life on Earth
310

311 The most solute-tolerant Bacteria and Archaea (i.e. extreme, obligate halophiles) are only
312 able to grow at their water-activity minima under hypersaline conditions. Some of these
313 organisms thrive under conditions which resemble those that would have been available on
314 the early Earth; indeed, many of the extreme halophiles thus far studied exhibit their water-
315 activity minimum for cell division at elevated temperatures (Robinson et al., 2005). There is
316 some debate regarding the temperature of the early seas; earlier estimates of 70-80°C
317 (Knauth and Lowe, 2003) are now considered to be too high (the $\delta^{18}\text{O}$ values on which the
318 calculations were based were skewed due to increased seawater temperatures which
319 resulted from inputs of hydrothermal fluids from the crust). More recent estimates based on
320 analysis of oxygen and hydrogen isotopes (i.e. $\delta^{18}\text{O}$ and δD , respectively) are about 40°C
321 (Blake et al., 2010). However, the high mantle heat flow on the early Earth drove a highly
322 active hydrothermal circulatory system that contributed hot, salty (de Ronde et al., 1997),
323 silica-rich fluids to the local environment (Westall, 2012). It has been proposed that
324 primordial life may have first occurred within saline environments on early Earth (Dundas,
325 1998), and recent evidence suggests that the abiotic formation of primitive proteins can
326 indeed occur under saline conditions (Longo et al., 2013; Longo and Blaber, 2014).
327 Understanding the way in which water-condensing chemical reactions could have led to the
328 emergence of key biomolecules (eg: peptides and nucleic acids) is essential to
329 understanding the origins of life (da Silva and Holm 2014 and references therein).
330 Prokaryote life (anaerobic) was relatively abundant in these early environments and left
331 behind numerous signatures of its presence (Westall, 2012). There are stratified salt
332 deposits of various ages across large regions of the Earth, indicating that concentrated salt-
333 waters/brines have existed across the planet's geologic history (Warren, 2010). Direct
334 association of an early photosynthetic microbial community with evaporitic conditions is
335 documented in 3.33 billion-year-old volcanic sands from the Barberton greenstone belt,
336 South Africa (Figure 2; Westall et al., 2006, 2011). The uppermost layers of a desiccated
337 biofilm, formed on sediments deposited in shallow waters that were partially exposed to air,
338 are interlayered with tiny evaporate crystals (microns in size and including aragonite,
339 gypsum, halite and magnesium calcite; Figure 2). Evaporitic precipitates have been

described from other formations on the early Earth, including the 3.42 billion-year-old Buck Reef Chert in Barberton (Lowe and Fisher-Worrell, 1999) and the 3.43 billion-year-old Strelley Pool Chert of the Pilbara in Australia (Allwood et al., 2007). The early terrestrial phototrophs were quite advanced on the evolutionary scale compared to chemotrophs. Although, to date, no direct association of chemotrophic biosignatures with the early evaporitic deposits has been identified, these more primitive organisms were nevertheless also common (Westall, 2012; Westall et al., 2013). If primitive cells did reach the early Earth through panspermia, experiments simulating the entry of meteorites containing microorganisms into the Earth's atmosphere have shown (1) that phototrophs could not have been transported to Earth by these means (Cockell et al., 2007) and (2) that, if resilient forms of life were hidden in meteorites, they would need to be buried at depths of at least 5 cm in cracks within the meteorite in order to withstand the heat of entry (Foucher et al., 2010).

Regardless of how (and where) life originated, it seems most likely that it was prokaryotes (known to have preceded eukaryotes by ~2 billions years), in saline environments, which first reached the 0.605- a_w limit. Some of the oldest known fossils are those of prokaryotic cells (dating from ~3.5 billion years ago [Frances, is this the same environment as the 3.33 billion above? And if so, do we need to use the same number?....also is there any repetition between these sentences concerning the Barberton work here and those in the paragraph above? If not then that's fine.]) which apparently lived in salt-rich environments, as evaporite minerals such as magnesium calcite and halite were found embedded in the biofilm of an extant [Frances, was this mat fossilized or alive?] microbial mat, discovered in the Barberton greenstone belt, South Africa [Frances, given that the Barberton greenstone belt and it's location were mentioned above does this need repeating here?] (Westall et al., 2001; 2006), and similarly within ancient stromatolite columns from the Pilbara Craton, Western Australia (Allwood et al., 2007). Intriguingly, molecular analysis of modern stromatolite communities revealed that 74% of archaeal clones were closely related to the Halobacteria (Burns et al., 2004), which frequently dominate hypersaline environments (Oren, 2002). These prokaryotic halophiles were exposed to, and presumably inhabited [Frances should there be a comma here...] evaporitic environments containing [...or a comma here?] elevated concentrations of magnesium and characterized by water activities of considerably less than 0.755 (and can, indeed, be considerably below 0.600 a_w , depending on salt concentrations; Winston and Bates, 1960; Hallsworth et al., 2007; Yakimov et al., 2014; A. Stevenson et al., submitted). Indeed, the signatures of past life forms, including stromatolites, can be common in evaporitic deposits (Rothschild and Mancinelli, 2001).

Much later, and presumably in terrestrial locations, the Eukarya must have developed a similar resilience during growth at high concentrations of solutes which are produced via biogenic activity; namely sugars and polyols. Indeed, extremophilic Eukarya are considerably less salt-tolerant than their bacterial and archaeal counterparts, and it may be that the prokaryotes are yet to evolve an ability to grow at low water-activity in non-saline substrates (their current record is in the range 0.850 to 0.800; Lievens et al., 2014; R. Santos et al., submitted; A. Stevenson et al., submitted). Microbial, and indeed all biological, cells are not pure-water reactors with water activity of 1 (Trevors and Pollack, 2005), but consist of gels within which modulation of water activity along with speciation as a result of the solute-exclusion principle are central to effective cellular function. Indeed, a metabolic ability to maintain the cellular system at this level is one of the fundamental, defining characteristics of life itself.

Extraterrestrial, aqueous milieu which resemble fertile habitats on Earth

Liquid water was, and may still be, present in numerous locations in the Solar System. On Mars, for example, there is abundant geomorphological evidence for the presence of liquid water on the planet in the past (Carr, 2006) and possibly even, ephemerally, in the present (Möhlmann, 2011; McEwan et al., 2014; J. D. Rummel et al., unpublished). Such evidence includes the formation of secondary minerals through the aqueous alteration of the basaltic rocks that cover the surface of the planet (e.g. Carter et al., 2013; Martínez and Renno, 2013). It has been calculated that the water activities of evaporite deposits and bodies of saline water on early Mars were as high as 0.780 to 0.860 (Tosca et al., 2008), which is well within the ranges for microbial species from each Domain-of-life (Javor, 1984; Grant, 2004; Williams and Hallsworth, 2009; Stevenson et al., submitted).

The various brines on Jupiter's moon Europa are composed primarily of water and salts such as MgSO_4 , Na_2SO_4 , and/or Na_2CO_3 (and, in some cases also contain sulfuric acid; Muñoz-Iglesias et al., 2013). Saturated solutions of these salts have water-activity values of 0.900, 0.930 and 0.920 respectively (at 20°C, 1 atm; Winston and Bates, 1960), although it is currently unclear what the values would be under the prevailing conditions on Europa. At the lower temperatures, and the in situ pressures, on Europa the solubility of ions and, conversely, the precipitation of salts can also vary leading to increases in water activity (Marion et al., 2003; 2005), the water activity of a saturated Na_2CO_3 solution at 10°C, for example, is 0.990 (Winston and Bates, 1960). Whereas water-activity values for individual brines will vary according to their ionic composition (and pH, which also influences solubilities of some salts), it seems likely that the in-situ water activities are sufficiently high

to span the entire range for known life (Javor, 1984; Williams and Hallsworth, 2009; Stevenson et al., submitted).

Water has also been identified in asteroidal materials, for example the Monahans (1998) H5 chondrite which contained hypersaline fluid inclusions composed predominantly of saturated NaCl (Zolensky et al., 1999) having a water activity of 0.760 at 20°C and 0.750 at 2°C at 1 atm (Winston and Bates, 1960), although these values will vary with pressure. Fluid inclusions have been identified in an increasing number of asteroidal specimens including the Zag (1998) meteorite (Rubin et al., 2002). Furthermore, organic molecules have been detected in the fluid inclusions of some of these asteroidal bodies (e.g. Fries et al., 2012), so the composition of these asteroidal fluids can be close to those of the media and substrates in which halophiles occur. For instance, halophiles in hypersaline fluid inclusions of salt crystals from evaporite deposits contain Archaea, Bacteria, and algae (*Dunaliella* species)⁷. Many NaCl-saturated habitats contain a remarkably high microbial biomass and are characterised by intense competition (Antón et al., 2002; Daffonchio et al., 2006; Baati et al., 2008; Elevi Bardavid et al., 2008; Khemakhem et al., 2010) during which some species - which are known as “microbial weeds” (Cray et al., 2013b; Oren and Hallsworth, 2014) - achieve dominance of the communities including Archaea, Bacteria and Eukarya (e.g. *Haloquadratum walsbyi*, *Salinibacter ruber*, and *Dunaliella salina*; for references see Cray et al., 2013b; Oren and Hallsworth, 2014). The microbes that dominate and/or are most frequently isolated from the fluid inclusions of salt crystals found in evaporite deposits include a number of species known to be capable of cell division in the range 0.710 to 0.605 (or their close relations, such as *Dunaliella*, *Halocarcula*, *Halobacterium*, *Halococcus*, *Halorubrum* and *Natrinema* spp.: Stan-Lotter et al., 2000; Schubert et al., 2009b; Lowenstein et al., 2011; Gramain et al., 2011; A. Stevenson et al., submitted). In relation to water activity, the biotic activity of microorganisms - including halophiles – is plausible for some of the aqueous milieu found in extraterrestrial environments. Indeed, some of these locations resemble highly fertile habitats for known halophiles (see also A. Stevenson et al., submitted).

Planets which are neither too close to nor too far from a star and could, theoretically at least, accommodate active biological systems are said to be in the Circumstellar Habitable Zone or Goldilocks Zone of their respective solar system (Strughold, 1953). This designation is based on criteria, such as size of the planet and its absolute distance from the star it orbits, whether illuminosity could permit photosynthesis, having surface temperatures which

⁷ See McGenity et al. (2000); D'Hondt et al. (2002); Schubert et al. (2009a); Gramain et al. (2011); Lowenstein et al. (2011); Lomstein et al. (2012); Valentine (2013). Cyanobacteria are known to be metabolically active in evaporite deposits (the in situ water-activity limit for this activity has yet to be determined; Rothschild et al., 1994).

are biologically permissive for at least some of the time (variously defined as 0 to 100°C, or -25 to +122°C; Franck et al. 2007; Takai et al., 2008; Kminek et al., 2010; Harrison et al., 2013), and/or whether they have liquid water (Rampino and Caldeira, 1994; von Bloh et al., 2011). However, these criteria (and indeed the habitable-zone concept) have limited applicability or validity for a variety of reasons. Ecosystems exist on Earth which do not depend on photosynthetic activity (Chivian et al., 2008; Teixeira et al., 2013) and, indeed, the earliest forms of life were not photosynthetic (Westall, 2012); furthermore, there is circumstantial evidence that an extracellular source of liquid water is not obligatory for microbial life (see above). What is more, biologically permissive conditions may prevail in specific environments or substrates on otherwise hostile planetary bodies (for examples in relation to moons of Saturn, see Raulin, 2006; Nimmo et al., 2007; Parkinson et al., 2008). And finally, various activities of solutes can both prevent freezing of water and expand biotic windows of microbes and may do so to a degree yet to be determined (see below; Chin et al., 2010; J. D. Rummel et al., unpublished).

Water can remain liquid at temperatures far lower than those known to permit microbial cell-division (i.e. approximately -18°C; see references in Chin et al., 2010). Liquid water (in various forms, from thin films to underground oceans) may be found in many environments on Mars as well as planetary moons (Europa, Ganymede, Enceladus, etc). Diverse lines of evidence suggest that both photosynthetic and non-photosynthetic microbes may be capable of metabolism and cell division by hygroscopic absorption of water vapour and/or acquiring water from their substratum (as a sole extracellular source of water) both in vitro and in their natural habitats on Earth⁸, and utilize a variety of mechanisms for the acquisition and retention of water (e.g. production and accumulation of trehalose and other hygroscopic substances which optimize the acquisition and retention of water, morphological changes which minimize water loss, hydrotactic responses, inhabiting high-humidity niches, and construction of soil features to enhance water capture and retention; Garcia-Pichel and Pringault, 2001; Garvie et al., 2008; de Goffau et al. 2011; Williams et al., 2012; Rajeev et al., 2013; Zakharova et al., 2013). Furthermore, as noted above, some microbial cells can generate vast quantities of water via their metabolic activities (Miller, 1932; Peterson and Cowling, 1973; Oriol et al., 1988; Nagel et al., 2001; Marcano et al., 2002; Hocking, 2003; Kreuzer-Martin et al., 2005; 2006). Indeed, studies of bacterial cells demonstrate that up to 70% of intracellular water can be derived in this way (Kreuzer-Martin et al., 2005; 2006) and other studies demonstrate that cells can maintain higher intracellular water-activity than that of the environment; de Goffau et al. (2011).

⁸ E.g. fungi, lichens and cyanobacteria (Snow, 1949; Armolick and Dickson, 1956; Pitt and Christian, 1968; Ayerst, 1969; Bootsma et al., 1973; Drewello and Weissmann, 1997; Shomari and Kennedy, 1999; Lange et al., 2006; Wierzchos et al., 2011; Zakharova et al., 2013).

478 The atmosphere of Saturn's moon Enceladus can contain $\geq 90\%$ water vapour (Waite
479 et al., 2006) and, whereas its terrestrial surface is approximately -200°C (Brown et al., 2006),
480 plumes of water vapour and ice which are released into space are thought to originate in
481 subsurface oceans that have temperatures in the range -23 to -3°C (Nimmo et al., 2007;
482 Parkinson et al., 2008); i.e. temperatures which are permissive for the metabolic activity of
483 psychrotolerant and psychrophilic microbes (Collins and Buick, 1989; Chin et al., 2010,
484 Kminek et al., 2010, Mykytczuk et al., 2013). Various salts, nitrogenous compounds, and
485 organic substances have been identified in the atmosphere of Enceladus and E-ring ice
486 grains of Saturn (which may originate from Enceladus) including NaCl , NaHCO_3 , NaCO_3 , N_2 ,
487 ammonia, hydrogen cyanide, CO and CO_2 , methane, acetylene, and propane (Matson et al.,
488 2007; Postberg et al., 2009; 2011). Under conditions prevalent on Earth, bioaerosols can be
489 fertile habitats characterized by high levels of microbial diversity, biomass, and metabolic
490 activity (Fahlgren et al., 2010; Womack et al., 2010; 2012). In relation to the atmosphere of
491 Enceladus and/or the watery plumes which it emits into space, it is intriguing to speculate
492 what the water activity of liquid droplets in, or the humidity of, the gaseous phase might be
493 (presumably close to 100%) and whether the temperatures within these plumes can ever be
494 considerably higher than -200°C . It should be noted that, whereas definitive evidence from
495 culture-based studies of microbial systems on Earth indicate limits for cell division of
496 approximately $+122^{\circ}\text{C}$ or -18°C (Collins and Buick, 1989; Takai et al., 2008; Chin et al.,
497 2010; Harrison et al., 2013), circumstantial evidence from other biochemical or geochemical
498 data suggest biotic activity under more extreme conditions (down to about -40°C , and up to
499 approximately $+140^{\circ}\text{C}$; Parkes et al., 2000; Kminek et al., 2010; J. D. Rummel et al.,
500 unpublished).

501 Although the Earth is located within the region allocated as the Goldilocks Zone of our
502 own Solar System, it hosts many environments which do not permit life process and are
503 therefore essentially sterile due to, for example, low water activity, high chaotropicity,
504 excessively high or low temperatures, pH of > 12 , plus combinations of conditions such as
505 high salt and low pH or high temperature and high pH (e.g. Brown, 1990; Hallsworth, 1998;
506 Parkes et al., 2000; Grant, 2004; Hallsworth et al., 2007; Harrison et al., 2013; Yakimov et
507 al., 2014). Under all these conditions cells also need adequate energy sources and nutrients
508 for maintenance and growth which may require electron donors and acceptors for
509 respiration etc. Some combinations of conditions can slightly extend extremes for growth,
510 such as high pressure and temperatures; furthermore survival can occur under conditions

where growth cannot⁹. Conversely, planetary bodies which are basically hostile to life may nevertheless harbour small-scale, biologically permissive domains (Kminek et al., 2010; J. D. Rummel et al., unpublished). Solute activities represent one of the determinants for potential habitability on Earth; for example, chaotropicity can enable cellular function at low temperatures and kosmotropicity may enable cellular function in high-temperature environments or those dominated by chaotropic substances¹⁰. The ways in which water activity and other solute activities can interact to determine the physicochemical limits for life (e.g. Williams and Hallsworth, 2009; Chin et al., 2010) have yet to be fully characterized. Furthermore, there is little information on the way in which availability of nutrients and other resources can determine tolerance limits to physicochemical stress parameters (e.g. Daffonchio et al., 2006; J. P. Harrison et al., submitted). Once the interactions between such factors are better understood, the currently accepted criteria for habitability will require revision (Beaty et al., 2006; Marion et al., 2003; Marion and Kargel, 2008; Tosca et al., 2008; Kminek et al., 2010; Harrison et al., 2013; J. D. Rummel et al., unpublished).

How sensitive are cells to minute changes in water activity? And other unanswered questions

In their environmental context, microbes are exposed to complexity at multiple levels; in relation to (i) the dynamics of physical and chemical parameters, (ii) the antimicrobials and other substances produced by other cells in the vicinity, (iii) varying availability of resources, and countless other factors. Water activity, in particular, can oscillate (Cray et al., 2013b; Lievens et al., 2014), and may do so across a range of timescales from a fraction of a second, for example to days or longer. The majority of stress-biology studies which quantify water activity do so to either one or two decimal places. We propose here that water activity ought to be determined to an accuracy of three decimal places (Winston and Bates, 1960; Williams and Hallsworth, 2009; A. Stevenson et al., submitted) as this is more closely aligned with the sensitivity of cellular systems. All technologies used to quantify the water activity of undefined substrates are associated with some degree of error (see Winston and Bates, 1960, Greenspan, 1977, Hallsworth and Nomura, 1999, Yu et al., 2009). Commercially available apparatus for water-activity determination are associated with a net

⁹ The propagules/cells of many microbes are highly resilient to exposure to extremes of temperature, uv, pH, chaotropicity, desiccation and other stresses (e.g. Wyatt et al., 2014; R. Santos et al., submitted), even over long timescales, and so are capable of surviving conditions found in extraterrestrial locations (see above).

¹⁰ See Hallsworth (1998a); Hallsworth et al. (1998b; 2003a; 2003b; 2007); Williams and Hallsworth (2009) Bhaganna et al. (2010); Chin et al. (2010); McCammick et al. (2010); Bell et al. (2013); Cray et al. (2013a; 2013b); Lievens et al. (2014); Yakimov et al. (2014). Whereas chaotropic substances are typically less polar than water and disorder biomacromolecules, kosmotropic substances are usually more polar than water and thereby structure or rigidify macromolecular systems (see Cray et al., 2013a, and references therein).

variation (accounting for both accuracy and repeatability) of ± 0.010 to 0.020 water-activity units (A. Stevenson et al., submitted). At 0.600 water activity, this is equivalent to variations of water potential between ± -2.3 and -4.5 MPa respectively). For the purposes of biological and food-related research it has been suggested, that levels of accuracy of ± 0.010 (Labuza et al., 1976; Roa and Tapia, 1998), ± 0.020 (Troller and Christian, 1978; Sereno et al., 2001), ± 0.005 (Ferro Fontán and Chirife, 1981; Hallsworth and Nomura, 1999), or ± 0.001 a_w are appropriate (Winston and Bates, 1960). Our earlier studies (Williams and Hallsworth, 2009; A. Stevenson et al., submitted) suggest that microbial cells can be sensitive to differences/changes of < 0.010 water activity. For example, water-activity differences of < 0.005 units have impacted growth rates for diverse strains of xerophilic fungi by between 40 and 80% (A. Stevenson et al., submitted) which, in turn, implies fundamental differences at every level of the cellular system, from gene expression to physiological and developmental processes. On glycerol-supplemented media at water activities of 0.799 and 0.795 growth-rates for *A. penicillioides* varied between 1.13 and 0.642 mm d⁻¹ for strain JH06THH and between 1.20 and 0.732 mm d⁻¹ for strain JH06THJ; and on MgCl₂-supplemented media at water activities of 0.915 and 0.907 rates for *X. bisporus* varied between 3.96 and 1.43 mm d⁻¹ for strain FRR 0025, 2.55 and 0.533 mm d⁻¹ for strain FRR 2347, and 2.13 and 0.800 mm d⁻¹ for strain FRR 3443 (A. Stevenson et al., submitted). These data raise the tantalizing question of whether microbial cells are sensitive to water-activity differences down to the fourth, or even fifth, decimal place¹¹. It is noteworthy that, for a hypothetical microbial species which has a temperature window for cell division spanning from 5 to 40°C (i.e. a 35°C range), a temperature change of 10 , 1 or 0.1°C would represent a $1/3.5$, $1/35$ and $1/350$ fraction of this window, respectively. If the water-activity window for this microbe spanned from 1 to 0.900 a_w (i.e. 0.100 a_w -units in total), $1/3.5$, $1/35$ - and $1/350$ -portions of this window would correspond to 0.02857 , 0.00286 and 0.00029 a_w units, respectively. This underlines the fact that water-activity determinations to one decimal place (equivalent, in this example, to $\sim 29^\circ\text{C}$) can lack biological meaning, and those made to two decimal places (equivalent to an accuracy level of up to 2.9°C) are far less accurate than we would accept for biological studies of temperature or other environmental parameters. In relation to microbial multiplication on Earth, the water-activity and temperature windows for life span 0.400 a_w -units and 140°C , respectively (Fig. 1). In the context of stress biology, and at the scale of the

¹¹ Based on the use of Novasina technology (Axair Ltd., Pfäffikon, Switzerland) and a protocol incorporating a range of precautionary measures we achieve an accuracy of ± 0.001 water-activity units (A. Stevenson et al., submitted). Whereas calculations can be carried out to enable the expression of water-activity values to the fourth decimal place, these have been based on a number of assumptions which, collectively, result in unacceptable levels of uncertainty (Greenspan, 1977; Yu et al., 2009). Such a level of accuracy would be highly desirable in many spheres of biological research but empirical determinations of water activity to the fourth decimal place are currently unattainable.

biosphere, the expression of water activity to decimal place leads to an unacceptable level of accuracy, as 0.100 a_w units equates to a temperature of 35°C. Even water-activity determinations to three decimal places (equivalent to an accuracy level of ~0.3°C) are imposed by technological limitations rather than being dictated by the sensitivity level of the cell.

It remains unclear whether microorganisms are capable of subsistence without an extracellular supply of liquid water, and the biological availability of water in various types of aqueous film has also yet to be quantified. Cells may be able to acquire and retain water (de Goffau et al., 2011) which can be utilized when water activity falls below biologically permissive levels (for instance, see the studies of powdery mildew cited above) but there is no definitive evidence that this does indeed occur (and, if so, what mechanisms are involved) at present (J. D. Rummel et al., unpublished). Culture-independent studies are needed for high-solute, and other low-water activity, habitats to establish whether metabolic activity below the threshold for cell division (0.605 a_w) is commonplace at different locations within the microbial biosphere. In contrast with the increasing understanding of molecular-level adaptations in many other forms of extremophile, there is a paucity of information in relation to physiological, biochemical, and genetic mechanisms which facilitate halophile/xerophile function at < 0.690 a_w ¹². Further work is also needed to elucidate the roles that low water-activity substrates have played, and continue to play, in the evolution of both prokaryotic and eukaryotic systems. In the context of habitability, work is also needed to elucidate the interactions between type and concentration of ions, chao-/kosmotropicity, and water activity in relation to complex brines such as current those found in various locations on Earth (Siegel et al., 1983; Oren, 1988; Hallsworth et al., 2007; Yakimov et al., 2014) and those likely to have existed on early Earth or ancient Mars (Tosca et al., 2008). For ecosystems located in extremely hostile habitats, some reports hint that microbial life can be discontinuous and fragmented (Hopkins et al., 2005). In some low water-activity habitats, it may be that active cells can be located in otherwise biologically non-permissive zones, and pockets of sterility exist within otherwise inhabited zones. Furthermore, in some locations microbes may be inactive for most of the time and yet functional for short periods. It has yet to be determined, for example, whether slow cell divisions (over 100s or 1000s years) can occur in microbial communities which may subsist in nature at water activities below the known 0.605 a_w limit. In relation the water-activity limits for microbial life, we know much about the outer edges of Earth's biosphere yet, in the relation to the in situ conditions of microbial habitats, we still know relatively little.

¹² This also acts as a barrier to the biotechnological exploitation of these extremophiles and the macromolecular systems derived from them.

608 **Acknowledgements**

609 We are grateful to Dave W. Beaty (Jet Propulsion Laboratory, California Institute of
610 Technology, USA), Kathleen C. Benison (West Virginia University, USA), Ben Clark (Space
611 Science Institute, USA), Peter N. Golyshin (Bangor University, Wales), Jesse P. Harrison
612 (UK Centre for Astrobiology, The University of Edinburgh, UK), Ailsa D. Hocking (CSIRO
613 Division of Food and Nutritional Sciences, Australia), Barbara J. Javor (Southwest Fisheries
614 Science Center, USA), Tom L. Kieft (New Mexico Tech., USA), Chris R. Omelon (University
615 of Texas at Austin, USA), Aharon Oren (The Hebrew University of Jerusalem, Israel),
616 R. John Parkes (Cardiff University, Wales), John D. Rummel (East Carolina University,
617 USA), and Andrew Steele (Carnegie Institution of Washington, USA) for fruitful
618 discussions. Invaluable technical and logistical assistance was provided by Kalpa J.
619 Hallsworth (Bangor, UK) and Knut Wichterich (University of Bonn, Germany).

620

621 **Funding Information** Funding was received from the Department of Agriculture and Rural
622 Development and the Research (Northern Ireland) and Enterprise Directorate of Queen's
623 University Belfast.

624

625

626

627

References

- Allwood, A.C., Walter, M.R., Burch, I.W., and Kamber, B.S. (2007) 3.43 billion-year-old stromatolite reef from the Pilbara Craton of Western Australia: ecosystem-scale insights to early life on Earth. *Precambrian Res* **158**: 198–227.
- Antón, J., Oren, A., Benlloch, S., Rodríguez-Valera, F., Amann, R., and Rosselló-Mora, R. (2002) *Salinibacter ruber* gen. nov., sp. nov., a novel, extremely halophilic member of the Bacteria from saltern crystallizer ponds. *Int J Syst Evol Microbiol* **52**: 485–491.
- Argyris, D., Tummala, N.R., Striolo, A., and Cole, D.R. (2008) Molecular structure and dynamics in thin water films at the silica and graphite surfaces *J Phys Chem C* **112**: 13587–13599.
- Armolick, N., and Dickson, J.G. (1956) Minimum humidity requirement for germination of conidia of fungi associated with storage of grains. *Phytopathology* **46**: 462–465.
- Asada, S., Takano, M., and Shibasaki, I. (1979) Deoxyribonucleic acid strand breaks during drying of *Escherichia coli* on a hydrophobic filter membrane. *Appl Environ Microbiol* **37**: 266–273.
- Ayerst, G. (1969) The effects of moisture and temperature on growth and spore germination in some fungi. *J Stored Prod Res* **5**: 127–141.
- Baati, H., Guermazi, S., Amdouni, R., Gharsallah, N., Sghir, A., and Ammar, E. (2008) Prokaryotic diversity of a Tunisian multipond solar saltern. *Extremophiles* **12**: 505–518.
- Beaty, D.W., Buxbaum, K.L., Meyer, M.A., Barlow, N.G., Boynton, W.V., Clark, B.C., et al. (2006) Findings of the special regions science analysis group. *Astrobiology* **6**: 677–732.
- Bell, A.N.W., Magill, E., Hallsworth, J.E., and Timson, D.T. (2013) Effects of Alcohols and Compatible Solutes on the Activity of beta-Galactosidase. *Appl Biochem Biotech* **169**: 786–796.
- Bhaganna, P., Volkers, R.J.M., Bell, A.N.W., Kluge, K. Timson, D.J., McGrath, J.W., et al. (2010) Hydrophobic substances induce water stress in microbial cells. *Microb Biotechnol* **3**: 701–716.
- Bing, H., and Ma, W. (2011) Laboratory investigation of the freezing point of saline soil. *Cold Reg Sci Technol* **67**: 79–88.
- Blake, R.E., Chang, S.J. and Lepland, A. (2010) Phosphate oxygen isotope evidence for a temperate and biologically active Archean ocean. *Nature* **464**: 1029–1033.
- Bootsma, A., Gillespie, T.J., and Sutton, J.C. (1973) Germination of *Phyllosticta maydis* conidia in an incubation chamber with control of high relative humidities. *Phytopathology* **63**: 1157–1161.

- Bradley, J.P., Ishii, H.A., Gillis-Davis, J.J, Ciston, J., Nielsen, M.H., Bechtel, H.A., et al. (2014) Detection of solar wind-produced water in irradiated rims on silicate minerals. *Proc Natl Acad Sci* doi: 10.1073/pnas.1320115111
- Brodie, H.J., and Neufeld, C.C. (1942) The development and structure of the conidia of *Erysiphe polygoni* DC and their germination at low humidity. *Can J Res* **20**: 41–61.
- Brown, A.D. (1976) Microbial water stress. *Bacteriol Rev* **40**: 803–846.
- Brown, A.D. (1990) *Microbial Water Stress Physiology. Principles and Perspectives*. John Wiley and Sons, Chichester, United Kingdom.
- Brown, R.H., Clark, R.N., Buratti, B.J., Cruickshank, D.P., Barnes, J.W., Mastrapa, R.M.E., et al. (2006) Composition and physical properties of Enceladus' surface. *Science* **10**: 1425–1428.
- Burch, A.Y., Finkel, O.M., Cho, J.K., Belkin, S. and Lindow, S.E. (2013) Diverse microhabitats experienced by *Halomonas variabilis* on salt-secreting leaves. *Appl Environ Microbiol* **79**: 845–852.
- Burkhardt J. and Hunsche M. (2013) “Breath figures” on leaf surfaces – formation and effects of microscopic leaf wetness. *Front Plant Sci*. **4**: 1–9.
- Burns, B.P., Goh, F., Allen, M., and Neilan, B.A. (2004) Microbial diversity of extant stromatolites in the hypersaline marine environment of Shark Bay, Australia. *Environ Microbiol* **6**: 1096–1101.
- Campins, H., Hargrove, K., Pinilla-Alonso, N., Howell, E.S., Kelley, M.S., Licandro, J., et al. (2010) Water ice and organics on the surface of the asteroid 24 Themis. *Nature* **464**: 1320–1321.
- Carr, M. (2006) *The surface of Mars*. Cambridge University Press.
- Carroll, J.E., and Wilcox, W.F. (2003) Effects of humidity on the development of grapevine powdery mildew. *Phytopathology* **93**: 1137–1144.
- Carter, J., Poulet, F., Bibring, J.P., Mangold, N., and Murchie, S. (2013) Hydrous minerals on Mars as seen by the CRISM and OMEGA imaged spectrometers: updated global view. *J Geophys Res-Planet* **118**: 831–858.
- Chin, J.P., Megaw, J., Magill, C.L., Nowotarski, K., Williams, J.P., Bhaganna, P., et al. (2010) Solutes determine the temperature windows for microbial survival and growth. *Proc Natl Acad Sci USA* **107**: 7835–7840.

720 Chivian, D., Brodie, E.L., Alm, E.J., Culley, D.E., Dehal, P.S., DeSantis, T.Z., et al. (2008)
 721 Environmental genomics reveals a single-species ecosystem deep within Earth. *Science*
 722 **322**: 275–278.

723

724 Christner, B.C., Morris, C.E., Foreman, C.M., Cai, R., and Sands, D.C. (2008) Ubiquity of
 725 Biological Ice Nucleators in Snowfall. *Science* **319**: 1214-1214.

726

727 Clancy, P., Brack, A., Horneck, G. (2005) Looking for life: Searching the Solar System.
 728 Cambridge University Press, Cambridge.

729

730 Cobucci-Ponzano, B., Rossi, M., and Moracci, M. (2006) Interrupted genes in extremophilic
 731 Archaea: mechanisms of gene expression in early organisms. *Orig Life Evol Biosph* **36**:
 732 487–492.

733

734 Cockell, C.S., Brack, A., Wynn-Williams, D.D., Baglioni, P., Brandstätter, F., Demets, R., et
 735 al. (2007) Interplanetary transfer of photosynthesis: an experimental demonstration of a
 736 selective dispersal filter in planetary island biogeography. *Astrobiology* **7**: 1–9.

737

738 Collins, M.A., Buick, and R.K. (1989) Effect of temperature on the spoilage of stored peas by
 739 *Rhodotorula glutinis*. *Food Microbiol* **6**: 135–142

740

741 Cray, J.A., Bell, A.N.W., Bhaganna, P., Mswaka, A.Y., Timson, D.J., and Hallsworth,
 742 J.E. (2013a) The biology of habitat dominance; can microbes behave as weeds? *Microb*
 743 *Biotechnol* **6**: 453–492.

744

745 Cray, J.A., Russell, J.T., Timson, D.J., Singhal, R.S., and Hallsworth, J.E. (2013b) A
 746 universal measure of chaotropy and kosmotropy. *Environ Microbiol* **15**: 287–296.

747

748 Da Silva, J.A.L. and Holm, N.G. (2014) Borophosphates and silicophosphates as plausible
 749 contributors to the emergence of life. *J Coll Interface Sci*. DOI: 10.1016/j.jcis.2014.02.034.

750

751 Daffonchio, D., Borin, S., Brusa, T., Brusetti, L., van der Wielen, P.W.J.J., Bolhuis, H., et al.
 752 (2006) Stratified prokaryote network in the oxic–anoxic transition of a deep sea halocline.
 753 *Nature* **440**: 203–207.

754

755 Daniel, R.M., Finney, J.L., and Stoneham, M. (2004) The molecular basis of life: is life
 756 possible without water? A discussion meeting held at the Royal Society, London, UK, 3-4
 757 December 2003. *Philos Trans R Soc Lond B Biol Sci* **359**: 1141–1328.

758

759 Dash, J.G., Rempel, A.W., and Wettlaufer, J.S. (2006) The physics of premelted ice and its
 760 geophysical consequences. *Rev Mod Phys* **78**: 698–741.

761

762 de Goffau, M.C., van Dijk, J.M., and Harmsen, H.J.M. (2011) Microbial growth on the edge of
 763 desiccation. *Environ Microbiol* **13**: 2328–2335.

764

- de Ronde, C.E.J., Channer, D.M.DeR., Faure, K., Bray, C.J., and Spooner, E.T.C. (1997) Fluid chemistry of Archean seafloor hydrothermal vents; implications for the composition of circa 3.2 Ga seawater. *Geochim Cosmochim Acta* **61**: 4025–4042.
- Derjaguin, B.V., and Churaev, N.V. (1986) Properties of water layers adjacent to interfaces. In: *Fluid Interfacial Phenomena*, Croxton C.A. (Ed.). John Wiley & Sons, New York, pp. 663–738.
- D'Hondt, S., Rutherford, S., and Spivack, A.J. (2002) Metabolic activity of subsurface life in deep-sea sediments. *Science* **295**: 2067–2070.
- Dickson, J.L., Head, J.W., Levy, J.S., and Marchant, D.R. (2013) Don Juan Pond, Antarctica: near-surface CaCl₂-brine feeding Earth's most saline lake and implications for Mars. *Sci Rep* **3**: 1166.
- Doroshenko, E.A., Zenova, G.M., Zvyagintsev, D.G., and Sudnitsyn, I.I. (2005) Spore germination and mycelial growth of streptomycetes at different humidity levels. *Mikrobiologiya* **74**: 690–694.
- Doroshenko, E.A., Zenova, G.M., Sudnicin, I.I., and Zvyagintsev, D.G. (2006) Influence of humidity on soil mycelial bacteria. *Vestn Mosk U Poch* **1**: 45–48.
- Drewello R., and Weissmann, R. (1997) Microbially influenced corrosion of glass. *Appl Microbiol Biotechnol* **47**: 337–346
- Duda, V.I., Danilevich, V.N., Suzina, N.E., Shorokhova, A.P., Dmitriev, V.V., Mokhova, O.N., et al. (2004) Changes in the fine structure of microbial cells induced by chaotropic salts. *Mikrobiologiya* **73**: 341–349.
- Duda, V.I., Danilevich, V.N., Akimov, V.N., Suzina, N.E., Dmitriev, V.V., and Shorokhova, A.P. (2005) Fluorescence microscopic study of microorganisms treated with chaotropic agents. *Mikrobiologiya* **74**: 434–439.
- Dundas, I. (1998) Was the environment for primordial life hypersaline? *Extremophiles* **2**: 375–377.
- Elevi Bardavid, R., Khristo, P., and Oren, A. (2008) Interrelationships between *Dunaliella* and halophilic prokaryotes in saltern crystallizer ponds. *Extremophiles* **12**: 5–14.
- Fahlgren, C., Hagström, A., Nilsson, D., and Zweifel, U.L. (2010) Annual variations in the diversity, viability, and origin of airborne bacteria. *Appl Environ Microbiol* **76**: 3015–3025.
- Fakes, M.G., Dali, M.V., Haby, T.A., Morris, K.R., Varia, S.A., and Serajuddin, A.T.M. (2000) Moisture sorption behavior of selected bulking agents used in lyophilized products. *PDA J Pharm Sci Technol* **54**: 144–149.

811 Falk, M., Hartman, K.A., and Lord, R.C. (1963) Hydration of deoxyribonucleic acid. II. An
812 infrared study. *J Am Chem Soc* **85**: 387–391.

813

814 Ferro Fontán, C., and Chirife, J. (1981) The evaluation of water activity in aqueous-solutions
815 from freezing-point depression. *J Food Technol* **16**: 21–30.

816

817 Foucher, F., Westall, F., Brandstatter, F., Demets, R., Parnell, J., Cockell, C.S., et al. (2010)
818 Testing the survival of microfossils in artificial martian sedimentary meteorites during entry
819 into Earth's atmosphere: the STONE 6 experiment. *Icarus* **207**: 616–630.

820

821 Franck, S., von Bloh, W., and Bounama, C. (2007) Maximum number of habitable planets at
822 the time of Earth's origin: new hints for panspermia and the mediocrity principle. *Int J*
823 *Astrobiol* **6**: 153–157.

824

825 Frank, M., and Hess, E. (1941) Studies on salt fish: V. Studies on *Sporendonema epizoum*
826 from "Dun" salt fish. *Can J Fish Aquat Sci* **5b**: 276–286.

827

828 Fries, M.D., Steele, A., and Zolensky, M. (2012) Halogen-Substituted Methane in Monahans
829 Halite. *Meteorit Planet Sci Supplement* **75**: 5381.

830

831 Garcia-Pichel, F., and Pringault, O. (2001) Cyanobacteria track water in desert soil.
832 *Nature* **413**: 380–381.

833

834 Garvie, L.A., Knauth, L.P., Bungartz, F., Klonowski, S., and Nash, T.H. 3rd. (2008) Life in
835 extreme environments: survival strategy of the endolithic desert lichen *Verrucaria*
836 *rubrocincta*. *Naturwissenschaften* **95**: 705–712.

837

838 Gramain, A., Díaz, G.C., Demergasso, C., Lowenstein, T.K., and McGenity, T.J. (2011)
839 Achaean diversity along a subterranean salt core from the Salar Grande (Chile). *Environ*
840 *Microbiol* **13**: 2105–2121.

841

842 Grant, W.D. (2004) Life at low water activity. *Philos Trans R Soc Lond B Biol Sci* **359**: 1249–
843 1266.

844

845 Greenspan L. (1977) Humidity fixed points of binary saturated aqueous solutions. *J Res Nat*
846 *Bur Stand-A Phys Chem* **81A**: 89–96.

847

848 Hallsworth, J.E. (1998) Ethanol-induced water stress in yeast. *J Ferment Bioeng* **85**: 125–
849 137.

850

851 Hallsworth, J.E., Nomura, Y., and Iwahara, M. (1998) Ethanol-induced water stress and
852 fungal growth. *J Ferment Bioeng* **86**: 451–456.

853

854 Hallsworth, J.E., and Nomura, Y. (1999) A simple method to determine the water activity of
855 ethanol-containing samples. *Biotechnol Bioeng* **62**: 242–245.

- Hallsworth, J.E., Heim, S. and Timmis, K.N. (2003a) Chaotropic solutes cause water stress in *Pseudomonas putida*. *Environ Microbiol* **5**: 1270–1280.
- Hallsworth, J.E., Prior, B.A., Nomura, Y., Iwahara, M., and Timmis, K.N. (2003b) Compatible solutes protect against chaotrope (ethanol)-induced, nonosmotic water stress. *Appl Environ Microbiol* **69**: 7032–7034.
- Hallsworth, J.E., Yakimov, M.M., Golyshin, P.N., Gillion, J.L.M., D'Auria, G., Alves, F.L., et al. (2007) Limits of life in MgCl₂-containing environments: chaotropicity defines the window. *Environ Microbiol* **9**: 803–813.
- Harris, R.F. (1981). The effect of water potential on microbial growth and activity. *Water Potential Relations in Soil Microbiology*, J.F. Parr et al., (Eds.). Soil Science Society of America, Madison, WI, pp. 23-95.
- Harrison, J.P., Gheeraert, N., Tsigelnitskiy, D., and Cockell, C.S. (2013) The limits for life under multiple extremes. *Trends Microbiol* **21**: 204–212.
- Harrison, J. P., Hallsworth, J. E. and C. S. Cockell (2014) Resource limitation redefines the limits for life on Earth. Under review for *Environmental Microbiology Special Issue*.
- Hocking, A.D. (2003) Microbiological facts and fictions in grain storage. In *Stored Grain in Australia*. Wright, E.J., Webb, M.C., and Highley, E., (Eds.). Proceedings of the Australian Postharvest Technical Conference, Canberra. CSIRO: 55-58.
- Hopkins, B., Elberling, B., Greenfield, L.G., Gregorich, E.G., Novis, P., O'Donnell, A.G., et al. (2005) Soil microorganisms in Antarctic dry valleys: resource supply and utilization. In *Microorganisms and Earth Systems-Advances in Geomicrobiology*, Gadd, G., Semple, K., and Lappin-Scott, H. (Eds.). Cambridge University Press, Cambridge, UK, pp. 71–84.
- Hu, Q., and Wang, J.S.Y. (2003) Aqueous-phase diffusion in unsaturated geologic media: a review. *Crit Rev Environ Sci Technol* **33**: 275–297.
- Jaenicke, R., and Bohm, G. (1998) The stability of proteins in extreme environments. *Curr Opin Struct Biol* **8**: 738–748.
- Javor, B.J. (1984) Growth potential of halophilic bacteria isolated from solar salt environments: carbon sources and salt requirements. *Appl Environ Microbiol* **48**: 352–360.
- Jepsen, S.M., Priscu, J.C., Grimm, R.E., and Bullock, M.A. (2007) The potential for lithoautotrophic life on Mars: application to shallow interfacial water environments. *Astrobiology* **7**: 342–354.
- Khemakhem, H., Elloumi, J., Moussa, M., Aleya, L., and Ayadi, H. (2010) The concept of ecological succession applied to phytoplankton over four consecutive years in five ponds featuring a salinity gradient. *Estuar Coast Shelf Sci* **88**: 33–44.

903 Kminek, G., Rummel, J. D., Cockell, C. S., Atlas, R., Barlow, N., Beaty, D., et al. (2010)
 904 Report of the COSPAR Mars special regions colloquium. *Adv Space Res* **46**: 811–829.
 905

906 Kminek, G., Conley, C., Allen, C.C., Bartlett, D.H., Beaty, D. W., Benning, D.G., et al. (2014)
 907 Report of the workshop for life detection on samples from Mars. *Life Sci Space Res* **2**: 1–5.
 908

909 Knauth, L.P., and Lowe, D.R. (2003) High Archean climatic temperature inferred from
 910 oxygen isotope geochemistry of cherts in the 3.5 Ga Swaziland Supergroup, South Africa.
 911 *Geol Soc Am Bull* **115**: 566–580.

912 Kreuzer-Martin, H.W., Ehrlinger, J.R., and Hegg, E.L. (2005) Oxygen isotopes indicate most
 913 intracellular water in log-phase *Escherichia coli* is derived from metabolism. *Proc Natl Acad*
 914 *Sci* **102**: 17337–17341.

915 Kreuzer-Martin, H.W., Lott, M.J., Ehleringer, J.R., and Hegg, E.L. (2006) Metabolic
 916 processes account for the majority of the intracellular water in log-phase *Escherichia coli*
 917 cells as revealed by hydrogen isotopes. *Biochemistry* **45**: 13622–13630.

918 Kurkal, V., Daniel, R.M., Finney, J.L., Tehei, M., Dunn, R.V., and Smith, J.C. (2005) Enzyme
 919 activity and flexibility at very low hydration. *Biophys J* **89**: 1282–1287.

920 Küppers, M., O'Rourke, L., Bockelée-Morvan, D., Zakharov, V., Lee, S., von Allmen, P., et
 921 al. (2014) Localized sources of water vapour on the dwarf planet (1) Ceres. *Nature* **505**:
 922 525–527.
 923

924 Labuza, T.P., Acott, K., Tatini, S.R., Lee, R.Y., Flink, J., and McCall, W. (1976) Water activity
 925 determination - collaborative study of different methods. *J Food Sci* **41**: 910–917.
 926

927 Lange, O.L., Kilian, E., and Ziegler, H. (1986) Water vapour uptake and photosynthesis of
 928 lichens: performance differences in species with green and blue-green algae as
 929 phycobionts. *Oecologia* **71**: 104–110.
 930

931 Lange, O.L., Meyer, A., Zellner, H., and Heber, U. (1994) Photosynthesis and water relations
 932 of lichen soil crusts: field measurements in the coastal fog zone of the Namib Desert. *Funct*
 933 *Ecol* **8**: 253–264.
 934

935 Lange, O.L., Allan Greene, T.G., Melzer, B., Meyer, A., and Zelinier, H. (2006) Water
 936 relations and CO₂ exchange of the terrestrial lichen *Teloschistes capensis* in the Namib fog
 937 desert: measurements during two seasons in the field and under controlled conditions. *Flora*
 938 **201**: 268–280.
 939

940 Lievens, B., Hallsworth J.E., Belgacem, Z.B., Pozo, M.I., Stevenson, A., Willems, K.A., et al.
 941 (2014) Microbiology of sugar-rich environments: diversity, ecology, and system constraints.
 942 *Environ Microbiol.* DOI: 10.1111/1462-2920.12570
 943

- Lomstein, B.A., Langerhuus, A.T., D'Hondt, S., Jorgensen, B.B., and Spivack, A.J (2012) Endospore abundance, microbial growth and necromass turnover in deep sub-seafloor sediment. *Nature* **484**: 101–104.
- Longo, L.M., Lee, J., and Blaber, M. (2013) Simplified protein design biased for prebiotic amino acids yields a foldable, halophilic protein. *Proc Natl Acad Sci USA* **110**: 2135–2139.
- Longo, L.M., and Blaber, M. (2014) Prebiotic protein design supports a halophile origin of foldable proteins. *Front Microbiol* **4**: 418.
- Lopez, M., Kurkal-Siebert, V., Dunn, R.V., Tehei, M., Finney, J.L., Smith, J.C. et al. Activity and dynamics of an enzyme, pig liver esterase, in near-anhydrous conditions. *Biophys J* **99**: L62–64.
- Lowe, D.R., and Fisher-Worrell, G., (1999) Sedimentology, mineralogy, and implications of silicified evaporites in the Kromberg Formation, Barberton greenstone belt, South Africa. In *Geologic evolution of the Barberton greenstone belt, South Africa*, Lowe, D.R., and Byerly, G.R., (Eds.). *Geol Soc Am Spec Pap* **329**: 167–188.
- Lowenstein, T.K., Schubert, B.A., and Timofeeff, M.N. (2011) Microbial communities in fluid inclusions and long-term survival in halite. *GSA Today* **21**: 4–9.
- Marcano, V., Benitez, P., and Palacios-Prü, E. (2002) Growth of a lower eukaryote in non-aromatic hydrocarbon media C₁₂ and its exobiological significance. *Planet Space Sci* **50**: 693–709.
- Manners, J.G., and Hossain, S.M.M. (1963) Effects of temperature and humidity on conidial germination in *Erysiphe graminis*. *T Brit Mycol Soc* **46**: 225–234.
- Marion, G.M. (1997) A theoretical evaluation of mineral stability in Don Juan Pond, Wright Valley, Victoria Land. *Antarct Sci* **9**: 92–99.
- Marion, G.M., Fritsen, C.H., Eicken, H., and Payne, M.C. (2003) The search for life on Europa: limiting environmental factors, potential habitats, and Earth analogues. *Astrobiology* **3**: 785–811.
- Marion, G.M., Kargen, J.S., Catling, D.C., and Jakubowski, S.D. (2005) Effects of pressure on aqueous chemical equilibria at subzero temperatures with applications to Europa. *Geochim Cosmochim Acta* **69**: 259–274.
- Marion, G.M., and Kargel, J.S. (2008) *Cold Aqueous Planetary Geochemistry with FREZCHEM: From Modeling to the Search for Life at the Limits*. Springer-Verlag, Berlin, Germany.
- Martínez, G.M., and Renno, N.O. (2013) Water and brines on Mars: current evidence and implications for MSL. *Space Sci Rev* **175**: 29–51.

990 Matson D.L, Castillo, J.C., and Lunine, J. (2007) Enceladus' plume: compositional evidence
991 for a hot interior. *Icarus* **187**: 569–573.

992

993 Mauer, L.J., and Taylor, L.S. (2010) Water-Solids Interactions: deliquescence. *Annu Rev*
994 *Food Sci Tech* **1**: 41–63.

995

996 McCammick, E.M., Gomase, V.S., Timson, D.J., McGenity, T.J., and Hallsworth, J.E. (2009)
997 Water-hydrophobic compound interactions with the microbial cell. In *Handbook of*
998 *Hydrocarbon and Lipid Microbiology – Hydrocarbons, Oils and Lipids: Diversity, Properties*
999 *and Formation*, Timmis, K.N. (Ed.). Springer, New York, USA, Vol. **2**, pp. 1451– 1466.

1000

1001 McEwen, A.S., Dundas, C.M., Mattson, S.S., Toigo, A.D., Ojha, L., Wray, J.J., et al.. (2014)
1002 Recurring slope lineae in equatorial regions of Mars. *Nature Geoscience* **7**: 53–58.

1003

1004 McGenity, T.J., Gemmell, R.T., Grant, W.D., and Stan-Lotter, H. (2000) Origins of halophilic
1005 microorganisms in ancient salt deposits. *Environ Microbiol* **2**: 243–250.

1006

1007 Miller, E.J., and Chibnall, A. C. (1932) The proteins of grasses. *Biochem J* **26**: 392–402.

1008

1009 Möhlmann, D.T.F. (2004) Water in the upper Martian surface at mid- and low-latitudes:
1010 presence, state, and consequences. *Icarus* **168**: 318–323.

1011

1012 Möhlmann, D. (2005) Adsorption water-related potential chemical and biological processes
1013 in the upper Martian surface. *Astrobiology* **5**: 770–777.

1014

1015 Möhlmann, D.T.F. (2008) The influence of van der Waals forces on the state of water in the
1016 shallow subsurface of Mars. *Icarus* **195**: 131–139.

1017

1018 Möhlmann, D. (2009) Are nanometric films of liquid undercooled interfacial water bio-
1019 relevant? *Cryobiology* **58**: 256–261.

1020

1021 Möhlmann, D. (2011) Three types of liquid water in icy surfaces of celestial bodies. *Planet*
1022 *Space Sci* **59**: 1082–1086.

1023

1024 Möhlmann, D. (2012) Widen the Belt of Habitability! *Orig Life Evol Biosph* **42**: 93–100.

1025

1026 Mormile, M.R., Hong, B.Y., and Benison, K.C. (2009) Molecular analysis of the microbial
1027 communities of Mars analog lakes in Western Australia. *Astrobiology* **9**: 919–930.

1028

1029 Morris, C.E., Conen, F., Alex Huffman, J. Phillips, V., Pöschl, U., and Sands, D.C. (2014)
1030 Bioprecipitation: a feedback cycle linking Earth history, ecosystem dynamics and land use
1031 through biological ice nucleators in the atmosphere. *Glob Change Biol* **20**: 341–351.

1032

1033 Moyano, F.E., Vasilyeva, N., Bouckaert, L., Cook, F., Craine, J., Curiel Yuste, J., et al.
1034 (2012) The moisture response of soil heterotrophic respiration: interaction with soil
1035 properties. *Biogeosciences* **9**: 1173–1182.

1036

1037 Moyano, F.E., Manzoni, S., and Chenu, C. (2013) Responses of soil heterotrophic
 1038 respiration to moisture availability: An exploration of processes and models. *Soil Biol*
 1039 *Biochem* **59**: 72–85.

1040

1041 Muñoz-Iglesias, V., Bonales, L.J., and Prieto-Ballesteros, O. (2013) pH and salinity evolution
 1042 of Eurpa's brines: raman spectroscopy study of fractional precipitation at 1 and 300 bar.
 1043 *Astrobiology* **13**: 693–702.

1044

1045 Mykytczuk, N.C.S., Foote, S.J., Omelon, C.R., Southam, G., Greer, C.W., and Whyte, L.G.
 1046 (2013). Bacterial growth at -15C; molecular insights from the permafrost bacterium
 1047 *Planococcus halocryophilus* Or1. *ISME J* **7**: 1211–1226.

1048

1049 Nagel, F.J.J.I., Tramper, J., Bakker, M.S.N., and Rinzema, A. (2001) Model for on-line
 1050 moisture-content control during solid-state fermentation. *Biotechnol Bioeng* **76**: 291–302.

1051

1052 Nimmo, F., Spencer, J.R., Pappalardo, R.T., and Mullen, M.E. (2007) Shear heating as the
 1053 origin of the plumes and heat flux on Enceladus. *Nature* **447**: 289–291.

1054

1055 Oren, A. (1988) The microbial ecology of the Dead Sea. In *Advances in microbial ecology*,
 1056 Marshall, K.C. (Ed.). Plenum Press, New York, pp. 193–229.

1057

1058 Oren, A. (1993) The Dead Sea–alive again. *Experientia* **49**: 518–522.

1059

1060 Oren, A. (1999) Bioenergetic aspects of halophilism. *Microbiol Mol Biol Rev* **63**: 334–348.

1061

1062 Oren, A. (2002) Diversity of halophilic microorganisms: environments, phylogeny,
 1063 physiology, and applications. *J Ind Microbiol Biotechnol* **28**: 56–63.

1064

1065 Oren, A. (2010) The dying Dead Sea: the microbiology of an increasingly extreme
 1066 environment. *Lakes Reser: Res Manag* **15**: 215–222.

1067

1068 Oren, A. (2011) Thermodynamic limits to microbial life at high salt concentrations. *Environ*
 1069 *Microbiol* **13**: 1908–1923.

1070

1071 Oren, A. (2013) Life in magnesium- and calcium-rich hypersaline environments: salt stress
 1072 by chaotropic ions. In *Polyextremophiles: Life Under Multiple Forms of Stress. Cellular*
 1073 *Origin, Life in Extreme Habitats and Astrobiology*. Seckbach, J., Oren, A., and Stan-Lotter,
 1074 H. (Eds.) Springer Science and Business Media, Dordrecht, Netherlands.

1075

1076 Oren, A. and Hallsworth, J.E. (2014) Microbial weeds in hypersaline habitats: the enigma of
 1077 the weed-like *Haloferax mediterranei*. Revised for *FEMS Microbiol Lett*.

1078

1079 Oriol, E., Raimbault, M., Roussos, S., and Viniegra-Gonzales, G. (1988) Water and water
 1080 activity in the solid state fermentation of cassava starch by *Aspergillus niger*. *Appl Microbiol*
 1081 *Biotechnol* **27**: 498–503.

1082

1083 Papendick, R.I. and Campbell, G.S. (1981) Theory and measurement of water potential.
1084 In *Water Potential Relationships in Soil Microbiology*, Parr, J.F., et al. (Eds.). Soil Science
1085 Society of America Publications, Madison, WI, pp. 1–22.
1086

1087 Parkes, R.J., Cragg, B.A., and Wellsbury, P. (2000) Recent studies on bacterial populations
1088 and processes in subseafloor sediments: a review. *Hydrogeol J* **8**:11–28.
1089

1090 Parkinson, C.D., Liang, M.-C., Yung, Y.L., and Kirshcivnk, J.L. (2008) Habitability of
1091 Enceladus: planetary conditions for life. *Orig Life Evol Biosph* **38**: 355–369.
1092

1093 Pasanen, A.L., Pasanen, P., Jantunen, M.J., and Kalliokoski, P. (1991) Significance of air
1094 humidity and air velocity for fungal spore release into the air. *Atmos Environ* **25**: 459–462.
1095

1096 Pavlov, A.K., Shelegedina, V.N., Vdovina, M.A., and Pavlov, A.A. (2010) Growth of
1097 microorganisms in Martian-like shallow subsurface conditions: laboratory modeling. *Int J*
1098 *Astrobiol* **9**: 51–58.
1099

1100 Pearson, R.T., and Derbyshire, W. (1974) NMR studies of water adsorbed on a number of
1101 silica surfaces. *J Colloid Interf Sci* **46**: 232–248.
1102

1103 Peterson, C.A., and Cowling, E.B. (1973) Influence of various initial moisture contents on
1104 decay of sitka spruce and sweetgum sapwood by *Polyporus versicolor* in the soil-block test.
1105 *Phytopathology* **63**: 235–237.
1106

1107 Pintado, A., and Sancho, L.G. (2002) Ecological significance of net photosynthesis activation
1108 by water vapour uptake in *Ramalina capitata* from rain-protected habitats in central
1109 Spain. *Lichenologist* **34**: 403–413.
1110

1111 Pitt, J.I., and Christian, J.H.B. (1968) Water relations of xerophilic fungi isolated from prunes.
1112 *Appl Environ Microbiol* **16**: 1853–1858.
1113

1114 Pitt, J.I. (1975) Xerophilic fungi and the spoilage of foods of plant origin. In *Water Relations*
1115 *of Foods*, Duckworth, R.B. (Ed.). London, United Kingdom: Academic Press, pp. 273–307.
1116

1117 Postberg, F., Kempf, S., Schmidt, J., Brilliantov, N., Beinsen, A., Abel, et al. (2009) Sodium
1118 salts in E-ring ice grains from an ocean below the surface of Enceladus. *Nature* **459**: 1098–
1119 1101.
1120

1121 Postberg, F., Schmidt, J., hillier, J., Kempf, S., and Srama, R. (2011) A salt-water reservoir as
1122 the source compositionally stratified plume on Enceladus. *Nature* **474**: 620–622.
1123

1124 Potts, M. (1994) Desiccation tolerance of prokaryotes. *Microbiol Rev* **58**: 755–805.
1125

1126 Qvit-Raz, N., Jurkevitch, E., and Belkin, S. (2008) Drop-size soda lakes: transient microbial
1127 habitats on a salt-secreting desert tree. *Genetics* **178**: 1615–1622.
1128

1129 Rajeev, L., Da Rocha, U.N., Klitgord, N., Luning, E.G., Fortney, J., Axen, S.D., et

1130 al. (2013). Dynamic cyanobacterial response to hydration and dehydration in a desert
1131 biological soil crust. *ISME J* **7**: 2178–2191.

1132

1133 Rampino, M.R., and Caldeira, K. (1994) The goldilocks problem: climate evolution and long-
1134 term habitability of terrestrial planets. *Annu Rev Astron Astrophys* **32**: 83-114.

1135

1136 Raulin, F. (2008) Astrobiology and habitability of Titan. *Space Sci Rev* **135**: 37–48.

1137

1138 Raviv, U., Laurat, P., and Klein, J. (2001) Fluidity of water confined to subnanometre films.
1139 *Nature* **413**: 51–54.

1140

1141 Rivkina, E., Friedmann, E., McKay, C., and Gilichinsky, D. (2000) Metabolic activity of
1142 permafrost bacteria below the freezing point. *Appl Environ Microbiol* **66**: 3230–3233.

1143

1144 Roa, V., and Tapia, M.S. (1998) Estimating water activity in systems containing multiples
1145 solutes based on solute properties. *J Food Sci* **63**: 559–564.

1146

1147 Robinson, J.L., Pyzyna, B., Atrasz, R.G., Henderson, C.A., Morrill, K.L., Burd, A.M., et al.
1148 (2005) Growth kinetics of extremely halophilic Archaea (family Halobacteriaceae) as
1149 revealed by Arrhenius plots. *J Bacteriol* **187**: 923–929.

1150

1151 Rothschild, L. J. and Mancinelli, R. L. (2001) Life in Extreme Environments. *Nature* **409**:
1152 1092–1101.

1153

1154 Rubin, A.E, Zolensky, M.E., and Bodnar, R.J. (2002) The halite-bearing Zag and Monahans
1155 (1998) meteorite breccias: Shock metamorphism, thermal metamorphism and aqueous
1156 alteration on the H-chondrite parent body. *Meteorit Planet Sci* **37**: 125–141.

1157

1158 Samarkin, V.A., Madigan, M.T., Bowles, M.W., Casciotti, K.L., Priscu, J.C., McKay, C.P., et
1159 el. (2010) Abiotic nitrous oxide emissions from the hypersaline Don Juan Pond in Antarctica.
1160 *Nature* **3**: 341–344.

1161

1162 Samson, R.A., van der Lustgraaf, B. (1978) *Aspergillus penicilloides* and *Eurotium*
1163 *halophilicum* in association with house-dust mites. *Mycopathologia* **64**: 13–16.

1164

1165 Sattler, B., Puxbaum, H., and Psenner, R., (2001) Bacterial growth in supercooled cloud
1166 droplets. *Geophys Res Lett* **28**: 239-242.

1167

1168 Schubert, B.A., Lowenstein, T.K., Timofeeff, M.N., and Parker, M.A. (2009a) How do
1169 prokaryotes survive in fluid inclusions in halite for 30,000 years? *Geology* **37**: 1059–1062.

1170

1171 Schubert, B.A., Lowenstein, T.K., and Timofeeff, M.N. (2009b), Microscopic identification of
1172 prokaryotes in modern and ancient halite, Saline Valley and Death Valley, California:
1173 *Astrobiology* **9**: 467–482.

1174

1175 Sereno, A.M., Hubinger, M.D., Comesaña, J.F., and Correa, A. (2001) Prediction of water
1176 activity of osmotic solutions. *J Food Eng* **49**: 103–114.

1177
1178 Shomari, S.H., and Kennedy, R. (1999) Survival of *Oidium anacardii* on cashew
1179 (*Anacardium occidentale*) in southern Tanzania. *Plant Path* **48**: 505–513.
1180
1181 Siegel, B.Z., McMurty, G., Siegel, S.M., Chen, J., and LaRock, P. (1979) Life in the calcium
1182 chloride environment of Don Juan Pond, Antarctica. *Nature* **280**: 828–829.
1183
1184 Siegel, B.Z., Siegel, S.M., Chen, J., and LaRock, P. (1983) An extraterrestrial habitat on
1185 Earth: the algal mat of Don Juan Pond. *Adv Space Res* **3**: 39–42
1186
1187 Snow, D. (1949) The germination of mould spores at controlled humidities. *Ann Appl Biol* **36**:
1188 1–13.
1189
1190 Sohl, F., Choukroun, M., Kargel, J., Kimura, J., Pappalardo, R., Vance, S., et al.
1191 (2010) Subsurface water oceans on icy satellites: Chemical composition and exchange
1192 processes, *Space Sci Rev* **153**: 485–510.
1193
1194 Stan-Lotter, H., Radax, C., Gruber, C., McGenity, T.J., Legat, A., Wanner, G., et al. (in
1195 press) The distribution of viable micro-organisms in Permo-Triassic rock salt. In 8th World
1196 Salt Symposium, 2000, Geertman, R.M. (Ed.). Elsevier Science, Amsterdam.
1197
1198 Stevenson, A., and Hallsworth, J.E. (2014) Water and temperature relations of soil
1199 Actinobacteria. Revised for *Environ Microbiol Reports*.
1200
1201 Strughold, H. (1953) *The Green and the Red Planet: a Physiological Study of the Possibility*
1202 *of Life on Mars*. University of New Mexico Press, Albuquerque.
1203
1204 Sutton, J.C., and Hildebrand, P.D. (1985) Environmental water in relation to *Peronospora*
1205 *destructor* and related pathogens. *Can J Plant Pathol* **7**: 323–330.
1206
1207 Takai, K., Nakamura, K., Toki., T., Urumu, T., Miyazaki, M., Miyazaki, J., et al. (2008) Cell
1208 proliferation at 122°C and isotopically heavy CH₄ production by a hyperthermophilic
1209 methanogen under high-pressure cultivation. *Proc Natl Acad Sci* **105**: 10949–10954.
1210
1211 Teixeira, S., Olu K., Decker, C., Cunha, R.L, Fuchs, S., Hourdez, S., et al. (2013) High
1212 connectivity across the fragmented chemosynthetic ecosystems of the deep Atlantic
1213 Equatorial Belt: efficient dispersal mechanisms or questionable endemism? *Mol Ecol* **22**:
1214 4663–4680.
1215
1216 Thomson, W. (1871) Presidential Address to the British Association for the Advancement of
1217 Science. *Nature* **4**: 262.
1218
1219 Toner, J.D., Catling, D.C., and Light, B. (2014) Soluble salts at the Phoenix Lander site,
1220 Mars: a reanalysis of the wet chemistry laboratory data. *Geochim Cosmochim Ac* **136**: 142–
1221 168.
1222

1223 Tosca, N.J., Knoll, A.H., and McLennan, S.M. (2008) Water activity and the challenge for life
1224 on early Mars. *Science* **320**: 1204–1207.

1225

1226 Trevors, J.T. and Pollack, G.H. (2005) Hypothesis: the origin of life in a hydrogel
1227 environment. *Prog Biophys Mol Biol* **89**: 1–8.

1228

1229 Troller, J.A., and Christian, J.H.B. (1978) *Water Activity and Food*. Academic Press, New
1230 York, USA.

1231

1232 Valentine, D.L. (2013) Microbiology: intraterrestrial lifestyles. *Nature* **496**: 176–177.

1233

1234 van der Wielen, P.W.J.J., Bolhuis, H., Borin, S., Daffonchio, D., Corselli, C., Giuliano, L., et
1235 al. (2005) The enigma of prokaryotic life in deep hypersaline anoxic basins. *Science* **307**:
1236 121–123.

1237

1238 von Bloh, W., Cuntz, M., Franck, S., and Bounama, C. (2011) Habitability of the Goldilocks
1239 Planet Gliese 581g: results from Geodynamic Models. *Astron Astrophys* **528**: A133.

1240

1241 Waite, J.H., Combi, M.R., Ip, W.-H., Cravens, T.E., McNutt, R.L., Kasprzak, W., et al. (2006)
1242 Cassini ion and neutral mass spectrometer: Enceladus plume composition and structure.
1243 *Science* **311**: 1419–1422.

1244

1245 Waldham, D.C., and Halvorson, H.O. (1954) Studies on the relationship between equilibrium
1246 vapor pressure and moisture content of bacterial endospores. *Appl Micr Biol* **2**: 333–338

1247

1248 Warren, J.K. (2010) Evaporites through time: tectonic, climatic and eustatic controls in
1249 marine and nonmarine deposits. *Earth Sci Rev* **98**: 217–268.

1250

1251 Watanabe, K., and Mizoguchi, M. (2002) Amount of unfrozen water in frozen porous media
1252 saturated with solution. *Cold Regions Sci Technol* **34**: 103–110.

1253

1254 Westall, F., de Wit, M.J, Dann, J., van der Gaast, S., de Ronde C.E.J, and Gerneke, D.
1255 (2001) Early Archean fossil bacteria and biofilms in hydrothermally-influenced sediments of
1256 the Barberton greenstone belt, South Africa. *Precambrian Res* **106**: 93–116.

1257

1258 Westall, F., de Ronde, C.E.J., Southam, G., Grassineau, N., Colas, M., Cockell, C.S., et al.
1259 (2006) Implications of a 3.472–3.333 Gyr-old subaerial microbial mat from the Barberton
1260 greenstone belt, South Africa for the UV environmental conditions on the early Earth. *Phil
1261 Trans R Soc* **361**: 1857–1875.

1262

1263 Westall, F., Cavalazzi, B., Lemelle, L., Marrocchi, Y., Rouzaud, J.-N., Simionovici, A., et
1264 al. (2011) Implications of in situ calcification for photosynthesis in a ~ 3.3 Ga-old microbial
1265 biofilm from the Barberton greenstone belt, South Africa. *Earth Planet Sci Lett* **310**, 468–479.

1266

1267 Westall, F. (2012) Early Earth. In *Astrobiology*, Lunine, J., et al. (Eds.). Cambridge University
1268 Press, pp. 89–114.

1269

1270 Westall, F., Loiseau, D., Foucher, F., Bost, N., Bertrand, M., Vago, J., et al. (2013)
 1271 Habitability on Mars from a Microbial point of view. *Astrobiology* **13**: 887–897.
 1272

1273 Wierzchos, J., Cámara, B., de los Rios, A., Davila, A.F., Sánchez Almazo, I.M., Arteida, O.,
 1274 et al. (2011) Microbial colonization of Ca-sulfate crusts in the hyperarid core of the Atacama
 1275 Desert: implications for the search for life on Mars. *Geobiology* **9**: 44–60.
 1276

1277 Williams, J.P., and Hallsworth, J.E. (2009) Limits of life in hostile environments; no limits to
 1278 biosphere function? *Environ Microbiol* **11**: 3292–3308.
 1279

1280 Williams, A.J., Buck, B.J., and Beyene, M.A. (2012) Biological soil crusts in the Mojave
 1281 Desert, USA: micromorphology and pedogenesis. *Soil Sci Am J* **76**: 1685–1695.
 1282

1283 Winston, P.W. and Bates, P.S. (1960) Saturated salt solutions for the control of humidity in
 1284 biological research. *Ecology* **41**: 232–237.
 1285

1286 Wolfe, J., Bryant, G., and Koster, K.L. (2002) What is “unfreezable water”, how unfreezable
 1287 is it and how much is there? *CryoLetters* **23**: 157–166.
 1288

1289 Womack, A.M., Bohannan, B.J.M., and Green, J.L. (2010) Biodiversity and biogeography of
 1290 the atmosphere. *Phil Trans R Soc B* **365**: 3645–3653.
 1291

1292 Womack, A.M., Artaxo, P.E., Ishida, F., Jardine, K.J., Saleska, S.R., Wiedemann, K.T., et al.
 1293 (2012) Microbial community composition and gene expression in the atmosphere over the
 1294 Brazilian Amazon. ASM, 112th General Meeting American Society for Microbiology, San
 1295 Francisco, California.
 1296

1297 Wyatt, T.T., Golovina, E.A., van Leeuwen, M.R., Hallsworth, J. E., Wösten, H.A.B. and
 1298 Dijksterhuis, J. (2014) Decreases in bulk water and mannitol and accumulation of trehalose
 1299 and trehalose-based oligosaccharides define a two-stage maturation process towards
 1300 extreme stress-resistance in ascospores of *Neosartorya fischeri* (*Aspergillus fischeri*).
 1301 *Environ Microbiol* In press. DOI: 10.1111/1462-2920.12557.
 1302

1303 Wynn-Williams, D.D. (1996) Antarctic microbial diversity: the basis of polar ecosystem
 1304 processes. *Biodivers Conserv* **5**: 1271–1293.
 1305

1306 Yakimov, M.M., Lo Cono, V., La Spada, G., Bortoluzzi, G., Messina, E., Smedile, F., et al.
 1307 (2014) Microbial community of seawater-brine interface of the deep-sea brine Lake Kryos as
 1308 revealed by recovery of mRNA are active below the chaotropicity limit of life. Revised for
 1309 *Environ Microbiol*
 1310


1311 Yarwood (1950) Water content of fungus spores. *Am J Bot* **37**: 636–639.
 1312

1313 Yu, X., Schmidt, A.R., and Schmidt, S.J. (2009) Uncertainty analysis of hygrometer-obtained
 1314 water activity measurements of saturated salt slurries and food materials. *Food Chem* **115**:
 1315 214–226.
 1316

1317 Zakharova, K., Tesei, D., Marzban, G., Dijksterhuis, J., Wyatt, T., and Sterflinger, K. (2013)
1318 Microcolonial fungi on rocks: A life in constant drought? *Mycopathologia* **175**: 537–547.
1319
1320 Zolensky, M.E. Bodnar, R.J., Gibsion, E.K., Nyquist, L.E., Reese, Y., Shih, C.–Y., et al.
1321 (1998) Asteroidal water within fluid inclusion-bearing halite in an H5 chondrite, Monahans
1322 (1998). *Science* **285**: 1377–1379.
1323
1324 Zvyagintsev, D.G., Zenova, G.M., Sudnitsyn, I.I., Gracheva, T.A., Napol'skaya, K.R., and
1325 Belousova, M.A. (2009) Dynamics of spore germination and mycelial growth of
1326 streptomycetes under low humidity conditions. *Microbiology* **78**: 440–444.
1327
1328 Zvyagintsev, D.G., Zenova, G.M., Sudnitsyn, I.I., Gracheva, T.A., Lapygina, E.E.,
1329 Napol'skaya, K.R., et al. (2012) Development of actinomycetes in brown semidesert soil
1330 under low water pressure. *Eurasian Soil Sci* **45**: 717–725.
1331

1332 **Figure legends**

1333



1334

1335 **Figure 1.** Diagrammatic representation of collective biological activity (compound rates cell
1336 division and metabolic activity) for microbes on Earth in relation to prevailing environmental
1337 (a) temperatures and (b) water activities. Red bars indicate the known range for cell division
1338 of microbes (-18 to +122°C, and 1 to 0.605 a_w), and orange dotted lines indicate for (a) the
1339 established limit for cellular metabolism (33°C), and (b) the known limit for physiological
1340 function of DNA (down to 0.530 a_w). Black bars indicate the range in which the overwhelming
1341 majority of microbial activity takes place, and curves represent collective biotic activity of
1342 microbes on Earth. Yellow bars indicate safety margins used for the designation of 'Special
1343 Regions' on Mars (down to -25°C and 0.500 a_w ; Kminek et al., 2010) in relation to
1344 international policy on planetary protection. Horizontal orange arrows indicate zones in which
1345 cell division may take place over extended timescales (10s to 1000s years) though there is a
1346 paucity of data on this topic; this zone for temperature extends considerably below -33°C
1347 because of the possibility that chaotropic substances may enhance flexibility of
1348 macromolecular systems and thereby reduce the temperature minima for microbial activity
1349 by a further 10 to 20°C (Chin et al., 2010).

1350

1351




Figure 2. Early Archaean microbes and evaporites; example from the 3.33 billion-year-old Josefsdal Chert, Barberton Greenstone Belt: (a) layer of evaporite minerals interbedded with layers of a photosynthetic microbial biofilm, (em) evaporite minerals, and (b) details of the diversity of minerals encrusted on the surface of the biofilm. They include here pseudomorphs (silica replaced) of acicular aragonite and losenge-shaped gypsum. Reproduced from Westall et al. (2006) with permission from The Royal Society Press.




Figure 3. Views of two planetary moons which are known to have an abundance of water, some of which may be present as subsurface oceans: (a) the icy surface of Europa, and (b)

1363 jets composed of water vapour, ice particles and organic compounds released from beneath
1364 the surface of Enceladus. Courtesy NASA/JPL-Caltech.

1365

1366 **Supplemental information**

1367

1368 **Movie S1.** Deliquescence of NaCl crystals on the surface of a pine needle (*Pinus sylvestris*)
1369 as humidity rises from approximately 65 to 80% equilibrium relative humidity. The
1370 deliquescence point of NaCl is approximately 75.0% equilibrium relative humidity at 2°C. An
1371 epistomatal chamber is visible but the guard cells are located below this section and cannot,
1372 therefore, be seen. The recording was made using an environmental scanning electron
1373 microscope and equilibrium relative humidity was controlled experimentally within a chamber
1374 (see Burkhardt and Hunsche, 2013).