



HAL
open science

Ozone variability in the troposphere and the stratosphere from the first 6 years of IASI observations (2008–2013)

Catherine Wespes, Daniel Hurtmans, Louisa K. Emmons, Sarah Safieddine, Cathy Clerbaux, David P. Edwards, Pierre-François Coheur

► To cite this version:

Catherine Wespes, Daniel Hurtmans, Louisa K. Emmons, Sarah Safieddine, Cathy Clerbaux, et al.. Ozone variability in the troposphere and the stratosphere from the first 6 years of IASI observations (2008–2013). *Atmospheric Chemistry and Physics*, 2016, 16 (9), pp.5721-5743. 10.5194/acp-16-5721-2016 . insu-01215625

HAL Id: insu-01215625

<https://insu.hal.science/insu-01215625>

Submitted on 10 May 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Ozone variability in the troposphere and the stratosphere from the first 6 years of IASI observations (2008–2013)

Catherine Wespes¹, Daniel Hurtmans¹, Louisa K. Emmons², Sarah Safieddine^{3,a}, Cathy Clerbaux^{1,3}, David P. Edwards², and Pierre-François Coheur¹

¹Spectroscopie de l'Atmosphère, Service de Chimie Quantique et Photophysique, Université Libre de Bruxelles (U.L.B.), Brussels, Belgium

²National Center for Atmospheric Research, Boulder, CO, USA

³Sorbonne Universités, UPMC Univ. Paris 06, Université Versailles St-Quentin, CNRS/INSU, LATMOS-IPSL, Paris, France

^anow at: Department of Civil and Environmental Engineering, Institute of Technology, Cambridge, Massachusetts, USA

Correspondence to: Catherine Wespes (cwespes@ulb.ac.be)

Received: 14 August 2015 – Published in Atmos. Chem. Phys. Discuss.: 14 October 2015

Revised: 4 March 2016 – Accepted: 16 April 2016 – Published: 10 May 2016

Abstract. In this paper, we assess how daily ozone (O_3) measurements from the Infrared Atmospheric Sounding Interferometer (IASI) on the MetOp-A platform can contribute to the analyses of the processes driving O_3 variability in the troposphere and the stratosphere and, in the future, to the monitoring of long-term trends. The temporal evolution of O_3 during the first 6 years of IASI (2008–2013) operation is investigated with multivariate regressions separately in four different layers (ground–300, 300–150, 150–25, 25–3 hPa), by adjusting to the daily time series averaged in 20° zonal bands, seasonal and linear trend terms along with important geophysical drivers of O_3 variation (e.g. solar flux, quasi-biennial oscillation (QBO)). The regression model is shown to perform generally very well with a strong dominance of the annual harmonic terms and significant contributions from O_3 drivers, in particular in the equatorial region where the QBO and the solar flux contribution dominate. More particularly, despite the short period of the IASI data set available up to now, two noticeable statistically significant apparent trends are inferred from the daily IASI measurements: a positive trend in the upper stratosphere (e.g. 1.74 ± 0.77 DU year⁻¹ between 30 and 50° S), which is consistent with other studies suggesting a turnaround for stratospheric O_3 recovery, and a negative trend in the troposphere at the mid-latitudes and high northern latitudes (e.g. -0.26 ± 0.11 DU year⁻¹ between 30 and 50° N), especially during summer and probably linked to the impact of decreasing ozone precursor emissions. The impact of the high tem-

poral sampling of IASI on the uncertainty in the determination of O_3 trend has been further explored by performing multivariate regressions on IASI monthly averages and on ground-based Fourier transform infrared (FTIR) measurements.

1 Introduction

Global climate change is one of the most important environmental problems of today, and monitoring the behaviour of the atmospheric constituents (radiatively active gases and those involved in their chemical production) is key to understanding the present climate and apprehending future climate changes. Long-term measurements of these gases are necessary to study the evolution of their abundance, changing sources and sinks in the atmosphere.

As a reactive trace gas present simultaneously in the troposphere and in the stratosphere, O_3 plays a significant role in atmospheric radiative forcing, atmospheric chemistry and air quality. In the stratosphere, O_3 is sensitive to changes in (photo-)chemical and dynamical processes and, as a result, undergoes large variations on seasonal and annual timescales. Measurements of O_3 total column have indicated a downward trend in stratospheric ozone over the period from the 1980s to the late 1990s relative to the pre-1980 values, which is due to the growth of the reactive bromine and chlorine species following anthropogenic emissions during that

period (WMO, 2003). In response to the 1987 Montreal Protocol and its amendments, with a reduction of the ozone-depleting substances (ODSs; Newchurch et al., 2003), a recovery of stratospheric ozone concentrations to the pre-1980 values is expected (Hofmann, 1996). While earlier works have debated a probable turnaround for the ozone hole recovery (e.g. Hadjinicolaou et al., 2005; Reinsel et al., 2002; Stolarski and Frith, 2006), WMO already indicated in 2007 that the total ozone in the 2002–2005 period was no longer decreasing, reflecting such a turnaround. Since then, several studies have shown successful identification of ozone recovery over Antarctica and over northern latitudes (e.g. Mäder et al., 2010; Salby et al., 2011; WMO, 2011; Kuttippurath et al., 2013; Knibbe et al., 2014; Shepherd et al., 2014). Nevertheless, the most recent papers as well as the WMO (2014) ozone assessment have warned, because of possible underestimation of the true uncertainties in the ozone trends attributed to decreasing effective equivalent stratospheric chlorine (EESC), against overly optimistic conclusions with regard to a possible increase in Antarctic stratospheric ozone (Kramarova et al., 2014; WMO, 2014; Knibbe et al., 2014; de Laat et al., 2015; Kuttippurath et al., 2015; Varai et al., 2015). The causes of the observed stratospheric O₃ changes are hard to isolate and remain uncertain precisely considering the contribution of dynamical variability to the apparent trend and the limitations of current chemistry–climate models to reproduce the observations. The assessment of ozone trends in the troposphere is even more challenging due to the influence of many simultaneous processes (e.g. emission of precursors, long-range transport, stratosphere–troposphere exchanges, STEs), which are all strongly variable temporally and spatially (e.g. Logan et al., 2012; Hess and Zbinden, 2013; Neu et al., 2014). Overall, today there are still large differences in the value of the O₃ trends determined from independent studies and data sets (mostly from ground-based and satellite observations) in both the stratosphere and the troposphere (e.g. Oltmans et al., 1998, 2006; Randel and Wu, 2007; Gardiner et al., 2008; Vigouroux et al., 2008, 2015; Jiang et al., 2008; Kyrölä et al., 2010). In order to improve on this and because O₃ has been recognised as one of the Global Climate Observing System (GCOS) Essential Climate Variables (ECVs), the scientific community has underlined the need of acquiring high-quality global, long-term and homogenised ozone profile records from satellites (Randel and Wu, 2007; Jones et al., 2009; WMO, 2007, 2011, 2014). This specifically has resulted in the ESA Ozone Climate Change Initiative (O₃-CCI; <http://www.esa-ozone-cci.org/>).

The Infrared Atmospheric Sounding Interferometer (IASI) onboard the polar orbiting MetOp, with its unprecedented spatiotemporal sampling of the globe, its high radiometric stability and the long duration of its program (three successive instruments to cover 15 years), in principle provides an excellent means to contribute to the analyses of the O₃ variability and trends. This is further strengthened by the possibility of using IASI measurements to discriminate O₃ dis-

tributions and variability in the troposphere and the stratosphere, as shown in earlier studies (Boynard et al., 2009; Wespes et al., 2009, 2012; Dufour et al., 2010; Barret et al., 2011; Scannell et al., 2012; Safieddine et al., 2013). Here, we use the first 6 years (2008–2013) of the new O₃ data set provided by IASI on MetOp-A to perform a first analysis of the O₃ time development in the stratosphere and in the troposphere. This is achieved globally by using zonal averages in 20° latitude bands and a multivariate linear regression model which accounts for various natural cycles affecting O₃. We also explore in this paper to which extent the exceptional temporal sampling of IASI can counterbalance the short period of data available for assessing trends in partial columns.

In Sect. 2, we give a short description of IASI and of the O₃ retrieved columns used here. Section 3 details the multivariate regression model used for fitting the time series. In Sect. 4, we evaluate how the ozone natural variability is captured by IASI and we present the time evolution of the retrieved O₃ profiles and of four partial columns (upper stratosphere, UST; middle–low stratosphere, MLST; upper troposphere–lower stratosphere, UTLS; middle–low troposphere, MLT) using 20° latitudinal averages on a daily basis. The apparent dynamical and chemical processes in each latitude band and vertical layer are then analysed on the basis of the multiple regression results using a series of common geophysical variables. The “standard” contributors in the fitted time series, as well as a linear trend term, are analysed in the specified altitude layers. Finally, the trends inferred from IASI are compared against those from Fourier transform infrared (FTIR) spectroscopy for six stations in the Northern Hemisphere.

2 IASI measurements and retrieval method

IASI measures the thermal infrared emission of the Earth’s atmosphere between 645 and 2760 cm⁻¹ with a field of view of 2 × 2 circular pixels on the ground, each of 12 km diameter at nadir. The IASI measurements are taken every 50 km along the track of the satellite at nadir, but also across-track over a swath width of 2200 km. IASI provides a global coverage twice a day with overpass times at 09:30 and 21:30 mean local solar time. The instrument is also characterised by a high spectral resolution which allows the retrieval of numerous gas-phase species (e.g. Clerbaux et al., 2009; Clarisse et al., 2012).

Ozone profiles are retrieved with the Fast Optimal Retrievals on Layers for IASI (FORLI) software developed at ULB/LATMOS. FORLI relies on a fast radiative transfer and on a retrieval methodology based on the optimal estimation method (Rodgers, 2000). In the version used in this study (FORLI-O₃ v20140922), the O₃ profile is retrieved for individual IASI measurement on a uniform 1 km vertical grid on 40 layers from surface up to 40 km. The a priori information (a priori profile and a priori covariance matrix) is built from

the Logan/Labow/McPeters climatology (McPeters et al., 2007) and only one single O₃ a priori profile and variance–covariance matrix are used. The retrieval parameters and performances are detailed in Hurtmans et al. (2012). The FORLI-O₃ profiles and/or total and partial columns have undergone validation using available ground-based, aircraft, O₃ sonde and other satellite observations (Anton et al., 2011; Dufour et al., 2012; Gazeaux et al., 2013; Parrington et al., 2012; Pommier et al., 2012; Scannell et al., 2012; Oetjen et al., 2014). Generally, the results show good agreement between FORLI-O₃ and independent measurements with a low bias (< 10 %) in the total column and in the vertical profile, except in the UTLS where a positive bias of 10–15 % is reported (Dufour et al., 2012; Gazeaux et al., 2013; Oetjen et al., 2014).

In this study, only daytime O₃ IASI observations from good spectral fits (root-mean square of the spectral residual lower than $3.5 \times 10^{-8} \text{ W}/(\text{cm}^2 \text{ sr cm}^{-1})$) have been analysed. Daytime IASI observations (determined with a solar zenith angle to the sun < 80°) are characterised by a better vertical sensitivity to the troposphere associated with a higher surface temperature and a higher thermal contrast (Clerbaux et al., 2009; Boynard et al., 2009). Furthermore, cloud contaminated scenes with cloud cover < 13 % (Hurtmans et al., 2012) were removed using cloud information from the EU-METCast operational processing (August et al., 2012).

An example of typical FORLI-O₃ averaging kernel functions for one mid-latitude observation in July (45° N/66° E) is represented on Fig. 1. The layers have been defined as ground–300 hPa (MLT), 300–150 hPa (UTLS), 150–25 hPa (MLST) and above 25 hPa (UST), so that they are characterised by a DOFS (degree of freedom for signal) close to 1 with a maximum sensitivity approximatively in the middle of the layers, except for the 300–150 hPa layer which has a reduced sensitivity. Taken globally, the DOFS for the entire profile ranges from ~2.5 in cold polar regions to ~4.5 in hot tropical regions, depending mostly on surface temperature, with a maximum sensitivity in the upper troposphere and in the lower stratosphere (Hurtmans et al., 2012). In the MLT, the maximum of sensitivity is around 4–8 km altitude for almost all situations (Wespes et al., 2012). The sharp decrease of sensitivity down to the surface is inherent to nadir thermal IR sounding in cases of low surface temperature or low thermal contrast and indicates that the retrieved information principally comes from the a priori in the lowest layer. Figure 2 presents July 2010 global maps of averaged FORLI-O₃ partial columns for two partial layers (MLT and MLST), and of the associated DOFS and a priori contribution (calculated as $X_a - \mathbf{A}(X_a)$, where X_a is the a priori profile and \mathbf{A} , the averaging kernel matrix, following the formalism of Rodgers, 2000). The two layers exhibit different sensitivity patterns; in the MLT, the DOFS typically ranges from 0.4 in the cold polar regions to 1 in regions characterised by high thermal contrast with medium humidity, such as the mid-latitude continental Northern Hemisphere (NH) (Clerbaux et

al., 2009). Lower DOFS values in the intertropical belt are explained by overlapping water vapour lines. In contrast, the DOFS for the MLST is globally almost constant and close to 1, with only slightly lower values (0.9) over polar regions. The a priori contribution is anti-correlated with the sensitivity, as expected. It ranges between a few to ~30 % and does not exceed 20 % on 20° zonal averages in the troposphere (see Supplement; Fig. S3, dashed lines), while the a priori contribution is smaller than ~12 % in the middle stratosphere. These findings indicate that the IASI MLST time series should accurately represent stratospheric variations, while the time series in the troposphere may reflect variations from the upper layers in addition to the real variability in the troposphere to some extent. In order to quantify this effect, the contribution of the stratosphere into the tropospheric ozone as seen by IASI has been estimated with a global 3-D chemical transport model (MOZART-4). Details of the model–observation comparisons can be found in the Supplement (see Figs. S2 and S3). We interestingly show that the stratospheric contribution to the MLT columns measured by IASI varies between 30 and 60 %, depending on latitude and season (Fig. S5). The limited vertical sensitivity of IASI contributes to this by a smaller part (~10–20 %) than the natural stratospheric influence (~20–45 %) (See Figs. S4 and S5). In addition, we find that the contribution of the natural variability (from both the troposphere and the stratosphere) on the MLT O₃ columns is larger than 50 % everywhere. In the 30–50° N band where the DOFS is the largest (see Fig. 2b), this contribution reaches ~85 % from which ~20–35 % originates from the stratosphere and ~55 % from the troposphere (Fig. S6a and b). Nevertheless, the contamination of IASI MLT O₃ with variations in stratospheric O₃ has to be kept in mind when analysing IASI MLT O₃.

3 Fitting method

3.1 Statistical model

In order to characterise the changes in ozone measured by IASI and to allow a proper separation of trend, we use a multiple linear regression model accounting for a linear trend and for interannual, seasonal and non-seasonal variations related to physical processes that are known to affect the ozone records. More specifically, the time series analysis is based on the fitting of daily (or monthly) median partial columns in different latitude bands following

$$O_3(t) = \text{Cst} + x_1 \cdot \text{trend} + \sum_{n=1,2} \left[a_n \cdot \cos(n\omega t) + b_n \cdot \sin(n\omega t) \right] + \sum_{j=2}^m x_j X_{\text{norm},j}(t) + \varepsilon(t), \quad (1)$$

where t is the number of days (or months), x_1 is the 6-year trend coefficient in the data, $\omega = 2\pi/365.25$ for the daily




Figure 1. Typical IASI FORLI-O₃ averaging kernels, in partial column units, corresponding to one mid-latitude observation in July (45° N/66° E) for each 1 km retrieved layers from ground to 40 km altitude (colour scale) and for four merged layers: ground–300 hPa; 300–150 hPa; 150–25 hPa; 25–3 hPa (grey lines). The total DOFS and the DOFS for each merged layers are also indicated.

model (or $2\pi/12$ for the monthly model) and $X_{\text{norm},j}$ are independent geophysical variables, the so-called “explanatory variables” or “proxies”, which are normalised over the period of IASI observation (2008–2013) in this study, as

$$X_{\text{norm}}(t) = 2[X(t) - X_{\text{median}}]/[X_{\text{max}} - X_{\text{min}}]. \quad (2)$$

$\varepsilon(t)$ in Eq. (1) represents the residual variation which is not described by the model and which is assumed to be autoregressive with time lag of 1 day (or 1 month). The constant term (Cst) and the coefficients a_n , b_n , x_j are estimated by a least-squares method and their standard errors (σ_ε) are calculated from the covariance matrix of the coefficients and corrected to take the uncertainty due to the autocorrelation of the noise residual into account as discussed in Santer et al. (2000) and references therein:

$$\sigma_\varepsilon^2 = (\mathbf{Y}^T \mathbf{Y})^{-1} \cdot \frac{\sum_t [\text{O}_3(t) - y\mathbf{Y}(t)]^2}{n - m} \cdot \frac{1 + \Phi}{1 - \Phi}, \quad (3)$$

where \mathbf{Y} the matrix with the covariates (trend, $\cos(n\omega t)$, $\sin(n\omega t)$, $X_{\text{norm},j}$) sorted by column, y is the vector of the regression coefficients corresponding to the columns of \mathbf{Y} , n is the number of daily (or monthly) data points in the time series, m is the number of the fitted parameters and Φ is the lag -1 autocorrelation of the residuals.

The median is used as a statistical average since it is more robust against the outliers than the normal mean (Kyrölä et

al., 2006, 2010). Note that similarly to Kyrölä et al. (2010), the model has been applied on O₃ mixing ratios rather than on partial columns but without significant improvement on the fitting residuals and R values.

3.2 Geophysical variables

In Eq. (1), harmonic time series with a period of a year and a half year are used to account for the Brewer–Dobson circulation and the solar insolation (a_1 and b_1 coefficients), and for the meridional circulation (a_2 and b_2 coefficients), respectively (Kyrölä et al., 2010). While these effects are of a periodic nature, the geophysical variables (X_j) are used here to parameterise the ozone variations on non-seasonal timescales. The chosen proxies are $F_{10.7}$, QBO^{10} , QBO^{30} , El Niño/Southern Oscillation (ENSO) and the North Atlantic Oscillation/Antarctic Oscillation (NAO/AAO), the first three being the most commonly used (“standard”) proxies to describe the natural ozone variability, i.e. the solar radio flux at 10.7 cm and the quasi-biennial oscillation (QBO) which is represented by two orthogonal zonal components of the equatorial stratospheric wind measured at 10 and 30 hPa, respectively (e.g. Randel and Wu, 2007). The three other proxies, ENSO, NAO and AAO, are used to account for other important fluctuating dynamical features: the El Niño/Southern Oscillation, the North Atlantic Oscillation and the Antarctic Oscillation, respectively. Table 1 lists the selected proxies, their sources and their resolutions. The time series of these proxies normalised over the 2000–2013 period following Eq. (2) are shown in Fig. 3a and b and they are shortly described hereafter.

- Solar flux: the $F_{10.7}$ solar radio flux at 10.7 cm is an excellent indicator of solar activity and is commonly used to represent the 11-year solar cycle. It is available from continuous consistent routine measurements at the Penticton Radio Observatory in British Columbia which are corrected for the variable Sun–Earth distance resulting from the eccentric orbit of the Earth around the Sun. Over the period 2008–2013, the radio solar flux increases from about 65 units in 2008 to 180 units in 2013 and is characterised by a specific daily “fingerprint” (see Fig. 3a). Note that because the period of IASI observations does not cover a full 11-year solar cycle, it could affect the determination of the trend in the regression procedure. The difficulty in discriminating the solar flux and linear trend terms is a known problem for such multivariate regression; it feeds into their uncertainties and it can lead to biases in the coefficients determination (e.g. Soukharev and Hood, 2006).
- QBO terms: the QBO of the equatorial winds is a main component of the dynamics of the tropical stratosphere (Chipperfield et al., 1994; Chipperfield, 2003; Randel and Wu, 1996, 2007; Logan et al., 2003; Tian et al., 2006; Fadnavis and Beig, 2009; Hauchecorne et al.,




Figure 2. Distributions of (a) O_3 columns, (b) DOFS and (c) a priori contribution (given as a %) in the ground–300 hPa (MLT) and 150–25 hPa (MLST) layers for IASI O_3 , averaged over July 2010 daytime data. Note that the scales are different.

Table 1. List of the proxies used in this study and their sources.

Proxy	Description (resolution)	Sources
$F_{10.7}$	The 10.7 cm solar radio flux (daily or monthly)	NOAA National Weather Service Climate Prediction Center: https://www.ngdc.noaa.gov/stp/solar/flux.html
QBO ¹⁰ QBO ³⁰	Quasi-biennial oscillation index at 10 and 30 hPa (monthly)	Free University of Berlin: www.geo.fu-berlin.de/en/met/ag/strat/produkte/qbo/
ENSO	El Niño/Southern Oscillation – Niño 3.4 Index (3-monthly averages)	NOAA National Weather Service Climate Prediction Center: http://www.cpc.noaa.gov/data/indices/
NAO	North Atlantic Oscillation index (daily or monthly)	http://www.cpc.ncep.noaa.gov/products/precip/CWlink/pna/nao.shtml
AAO	Antarctic Oscillation index (daily or monthly)	http://www.cpc.ncep.noaa.gov/products/precip/CWlink/daily_ao_index/ao.shtml

2010). It strongly influences the distributions of stratospheric O_3 propagating alternatively westerly and easterly with a mean period of 28 to 29 months. Positive and negative vertical gradients alternate periodically. At the top of the vertical QBO domain, there is a predominance of easterlies, while, at the bottom, westerly winds are more frequent. In order to account for the out-of-phase relationship between the QBO periodic oscillations in the upper and in the lower stratosphere, orthogonal zonal winds measured at 10 hPa (Fig. 3a; orange) and 30 hPa (Fig. 3a; green) by the ground station in Singapore have been considered here (Randel and Wu, 1996; Hood and Soukharev, 2006).

- NAO, AAO and ENSO: the El Niño/Southern Oscillation is represented by the 3-month running mean of sea surface temperature (SST) anomalies (in degrees Cel-

sus) in the Niño region 3.4 (region bounded by 120–170° W and 5° S–5° N). Raw data are taken from marine ships and buoys observations. The North Atlantic and Antarctic oscillations are described by the daily (or monthly) NAO and AAO indices which are constructed from the daily (or monthly) mean 500 hPa geopotential height anomalies in the 20–90° N region and 700 hPa height anomalies in the SH (Southern Hemisphere), respectively. Detailed information for these proxies can be found at <http://www.cpc.ncep.noaa.gov/>. These proxies describe important dynamical features which affect ozone distributions in both the troposphere and the lower stratosphere (e.g. Weiss et al., 2001; Frossard et al., 2013; Rieder et al., 2013; and references therein). The daily or 3-monthly average indexes used to parameterise these fluctuations are shown in Fig. 3b. The NAO




Figure 3. Normalised proxies as a function of time for the period 2000–2013 for the solar $F_{10.7}$ cm radio flux (blue) and the equatorial winds at 10 (green) and 30 hPa (orange), respectively (top panel), and for the El Niño (red), North Atlantic Oscillation (purple) and Antarctic Oscillation (light blue) indexes (bottom panel).

and AAO indexes are used for the NH and the SH, respectively (both are used for the equatorial band). These proxies have been included in the statistical model for completeness even if they are expected to only have a weak apparent contribution to the IASI ozone time series due to their large spatial variability in a zonal band (e.g. Frossard et al., 2013; Rieder et al., 2013). We have verified that including a typical time lag relation between ozone and the ENSO variable from 0 to 4 months did not improve the regression model in terms of residuals and uncertainty of the fitted parameters. As a consequence, a time lag has not been taken into account in our study.

- Effective equivalent stratospheric chlorine (EESC): the EESC is a common proxy used for describing the influence of the ODSs in O_3 variations. However, because the IASI time series starts several years after the turnaround for the ozone hole recovery in 1996/1997 (WMO, 2010), their influence is not represented by a dedicated proxy but is rather accounted for by the linear trend term.

Even if some of the above proxies are only specific to processes occurring in the stratosphere, we adopt the same approach (geophysical variables, model and regression procedure) for adjusting the IASI O_3 time series in the troposphere. This proves useful in particular to account for the stratospheric contribution to the tropospheric layer (~ 30 – 60 %; see Sect. 2 and Supplement, Fig. S5) due to

stratosphere–troposphere exchanges (STEs) and to the fact that this tropospheric layer is not perfectly decorrelated from the stratosphere. This has to be kept in mind when analysing the time series in the troposphere in Sect. 4. Specific processes in the troposphere such as emissions of ozone precursors, long-range transport and in situ chemical processing are taken into account in the model in the harmonic and the linear trend terms of the Eq. (1) (e.g. Logan et al., 2012). Including harmonic terms with 4- and 3-month periods in the model has been tested to describe O_3 dependency on shorter scales (e.g. Gebhardt et al., 2014), but this did not improve the results in terms of residuals and uncertainty of correlation coefficients.

3.3 Iterative backward variable selection

Similarly to previous studies (e.g. Steinbrecht et al., 2004; Mäder et al., 2007, 2010; Knibbe et al., 2014), we perform an iterative stepwise backward elimination approach, based on p values of the regression coefficients for the rejection, to select the most relevant combination of the above described regression variables (harmonic, linear and explanatory) to fit the observations. The minimum p value for a regression term to be removed (exit tolerance) is set at 0.05, which corresponds to a significance of 95%. The initial model which includes all regression variables is fitted first. Then, at each iteration, the variables characterised by p values larger than 5% are rejected. At the end of the iterative process, the remaining terms are considered to have significant influence on the measured O_3 variability, while the rejected variables are




Figure 4. (a) Daily IASI O_3 profiles (1×10^{12} molecules cm^{-3}) for the period 2008–2013 and over the range of the retrieved profiles as a function of time and altitude, in three latitude bands: 30–50° N (top), 10° S–10° N (middle) and 30–50° S (bottom). Superimposed daily IASI O_3 partial columns (scatters) and the associated fits (solid lines) from the multivariate regressions for the MLT (ground–300 hPa), UTLS (300–150 hPa), MLST (150–25 hPa) and UST (above 25 hPa) layers. The IASI measurements and the fits have been scaled for clarity. (b) Estimated total retrieval errors (%) associated with daily IASI O_3 profiles.

considered to be non-significant. The correction accounting for the autocorrelation in the noise residual is then applied to give more confidence in the coefficients' determination.

4 Ozone variations observed by IASI

In this section, we first examine the ozone variations in IASI time series during 2008–2013 in the four layers defined in the troposphere and the stratosphere to match the IASI sensitivity (Sect. 2). The performance of the multiple linear model is evaluated in Sect. 4.2 in terms of residuals errors, regression coefficients and associated uncertainties determined from the regression procedure (Sect. 3). Based on this, we characterise the principal physical processes that affect the IASI ozone records. Finally, the ability of IASI to derive apparent trends is examined in Sect. 4.3.

4.1 O_3 time series from IASI

Figure 4a shows the time development of daily O_3 number density over the entire altitude range of the retrieved profiles based on daily medians. The time series cover the 6 years of available IASI observations and are separated in three 20° latitude belts: 30–50° N (top panel), 10° N–10° S (middle panel) and 30–50° S (bottom panel). The figure shows the well-known seasonal cycle at mid-latitudes in the troposphere and the stratosphere, with maxima observed

in spring–summer and in winter–spring, respectively, and a strong stability of ozone layers with time in the equatorial belt. At high latitudes of both hemispheres, the high ozone concentrations and the large amplitude of the seasonal cycle observed in MLST and UTLS are mainly the consequence of the large-scale downward poleward Brewer–Dobson circulation which is prominent in late winter below 25 km.

Figure 4b presents the estimated statistical uncertainty on the O_3 profiles retrieved from FORLI. This total error depends on the latitude and the season, reflecting, amongst other things, the influence of signal intensity, of interfering water lines and of thermal contrast under certain conditions (e.g. temperature inversion, high thermal contrast at the surface). It usually ranges between 10 and 30 % in the troposphere and in the UTLS (upper troposphere–lower stratosphere), except in the equatorial belt due to the low O_3 amounts (see Fig. 4a) which leads to larger relative errors. The retrieval errors are usually less than 5 % in the stratosphere.

The relative variability (given as the standard deviation) of the daily median O_3 time series presented in Fig. 4a is shown in Fig. 5, as a function of time and altitude. It is worth noting that except in the UTLS over the equatorial band, the variability is larger than the estimated retrieval errors of the FORLI- O_3 data ($\sim 25\%$ vs. $\sim 15\%$ and $\sim 10\%$ vs. $\sim 5\%$, on average over the troposphere and the stratosphere, respectively), reflecting that the high natural temporal variability of

O₃ in zonal bands is well captured with FORLI (Dufour et al., 2012; Hurtmans et al., 2012). The standard deviation is larger in the troposphere and in the stratosphere below 20 km, where dynamic processes play an important role. The largest values (> 70 %) principally in the northern latitudes during winter) are measured around 9–15 km altitude. They highlight the influence of tropopause height variations and the STE processes. In the stratosphere, the variability is always lower than 20 % and becomes negligible in the equatorial region. Interestingly, the lowest troposphere of the NH (below 700 hPa; < 4 km) is marked by an increase in both O₃ concentrations (Fig. 4a) and standard deviations (between ~ 30 and ~ 45 %) in spring–summer, the latter being larger than the total retrieval error (less than 25%; see Fig. 4b). The lower tropospheric column (e.g. ground–700 hPa) can generally not be well discriminated because of the weak sensitivity of IASI in the lowermost layers (Sect. 2). However, the measurements in northern mid-latitudes in spring–summer are characterised by a larger sensitivity. In the ground–700 hPa columns, we find that the a priori contributions do not exceed 40 % and they range between 10 and 20 % over the continental regions. In addition, the stratosphere–troposphere exchanges are usually the weakest in summer. The stratospheric contributions into the IASI MLT columns are estimated to be the lowest in the summer mid-latitudes NH (e.g. ~ 35 % in the 30–50° N band; see Fig. S5b of the Supplement) and, as mentioned in Sect. 2, the real natural contribution originating from the troposphere reaches ~ 55 % (cf. Fig. S6b in the Supplement). This certainly helps in detecting the real variability of O₃ in the NH troposphere, and, the increase in the observed concentrations and in the variability may likely indicate a photochemical production of O₃ associated with anthropogenic precursor emissions (e.g. Logan, 1985; Fusco and Logan, 2003; Dufour et al., 2010; Cooper et al., 2010; Wilson et al., 2012; Safieddine et al., 2013). Changes in biomass and biogenic emissions of NO_x, CO and non-methane organic volatile compounds (NMVOCs) may also play a role. However, they only represent a small part of the total emissions for NO_x and CO (e.g. ~ 23 % vs. 72 % for the anthropogenic NO_x emissions and ~ 40 % vs. 60 % for the anthropogenic CO emissions from the emissions data set used in the Supplement), while the biogenic emissions of NMVOCs represent the largest contribution to the total (~ 80 %).

The zonal representation of the O₃ variability seen by IASI is given in Fig. 6. It shows the daily number density at altitude levels corresponding to maximum of sensitivity in the four analysed layers in most of the cases (600 hPa–~ 6 km; 240 hPa–~ 10 km; 80 hPa–~ 20 km; 6 hPa–~ 35 km) (Sect. 2). The top panel (~ 35 km) reflects the photochemical O₃ production by sunlight well, with the highest values in the equatorial belt during the summer (~ 3 × 10¹² molecules cm⁻³). The middle panels (~ 20 and ~ 10 km) shows the transport of ozone-rich air to high latitudes in late winter (up to ~ 6 × 10¹² molecules cm⁻³ in the




Figure 5. Daily IASI O₃ variability (%), expressed as $[\sigma(O_3(t))/O_3(t)] \times 100\%$, where σ is the standard deviation, as a function of time and altitude in three latitude bands: 30–50° N (top), 10° S–10° N (middle) and 30–50° S (bottom).

NH), which is induced by the Brewer–Dobson circulation. The fact that the patterns at ~ 10 km are similar to those at ~ 20 km mainly reflects the low sensitivity of IASI to that level compared to the others. Finally, the lower panel (~ 6 km) presents high O₃ levels in spring at high latitudes (~ 1.4 × 10¹² molecules cm⁻³ in the NH), which likely reflects both the STE processes and the contribution from the stratosphere due to the medium IASI sensitivity to that layer (see Sect. 2 and Supplement), and a shift from high to middle latitudes in summer which could be attributed to anthropogenic O₃ production. The MLT panel also reflects the seasonal oscillation of the intertropical convergence zone (ITCZ) around the equator and the large fire activity in spring around 20–40° S.

4.2 Multivariate regression results: seasonal and explanatory variables

Figure 4a shows the time series of the partial columns (dots) for the four layers (colour contours) superimposed on the time series of the IASI ozone concentration profile. The adjusted daily time series to these columns with the regression model defined by Eq. (1) is also overlaid and shown by coloured lines. The model represents the ozone variations in the four layers reasonably well, with, as illustrated for three latitude bands, good correlation coefficients (e.g. $R_{MLT} = 0.94$; $R_{UTLS} = 0.91$; $R_{MLT} = 0.90$ and $R_{US} = 0.91$ for the 30–50° N band) and low residuals (< 8 %) in all cases. The regression model explains a large fraction of the variance in the daily IASI data over the troposphere (~ 85–95 %) and the stratosphere (~ 85–95 % in all cases, except for the UST




Figure 6. Daily IASI O_3 number density (1×10^{12} molecules cm^{-3}) at 35 km (top row), 19 km (second row), 10 km (third row) and 6 km (bottom row) as a function of time and latitude. Note that the colour scales are different.

with ~ 70 – 95 %), as estimated from $\frac{\sigma(\text{O}_3^{\text{Fitted_model}}(t))}{\sigma(\text{O}_3(t))}$, where σ is the standard deviation relative to the fitted regression model and to the IASI O_3 time series.

However, note that the fit fails to reproduce the highest ozone values ($> 5 \times 10^{12}$ molecules cm^{-3}) above the seasonal maxima for the 30 – 50° N latitude band, especially in the MLST during spring 2009 and 2010. This could be associated with occasional downward transport of upper atmospheric NO_x -rich air occurring in winter and spring at high latitudes (Brohede et al., 2008) following the strong subsidence within the intense Arctic vortex in 2009–2010 (Pitts et al., 2011) or with the missing time lags in the regression model between the QBO and ENSO variables and the response of mid-latitude lower stratospheric ozone (Neu et al., 2014).

Figure 7 displays the annual cycle averaged over the 6 years recorded by IASI (dots) for the studied layers and bands, as well as that from the fit of the daily O_3 columns (lines). The regression model follows the O_3 variations perfectly in terms of timing of O_3 maxima and of amplitude of the cycle. The fit is generally characterised by low residuals (< 10 %) and good correlation coefficients (0.70–0.95),

which indicates that the regression model is suitable to describe the zonal variations. An exception is found over the southern latitudes (residual up to 15 % and R down to 0.61), probably because of the variation induced by the ozone hole formation which is not parameterised in the regression model, and because of the low temporal sampling of daytime IASI measurements in this region.

From Fig. 7, the following general patterns in the O_3 seasonal cycle can be isolated from the zonally averaged IASI data sets.

1. In the UST (top left panel), the maximum is in the equatorial belt, around 4.7×10^{18} molecules cm^{-2} throughout the year; and the amplitudes are small compared to the averaged O_3 values. The largest amplitude in the annual cycle is found in the NH between 30 and 50° N, where O_3 peaks in July after the highest solar elevation (in June) following a progressive build-up during spring–summer. In agreement with FTIR observations (e.g. Steinbrecht et al., 2006a; Vigouroux et al., 2008), a shift of the O_3 maximum from spring (March–April) to late summer (August–September) is found as one moves from high to low latitudes in the NH. In the SH, the general shape of the annual cycle, which shows a peak in October–November before the highest solar elevation (in December), results from loss mechanisms depending on the annual cycle of temperatures and other trace gases. Other effects such as changing Brewer–Dobson circulation, light absorption and tropical stratopause oscillations may also considerably impact the cycle in this layer (Brasseur and Solomon, 1984; Schneider et al., 2005).
2. In the lower stratosphere (MLST and UTLS, top right and bottom left panels), the pronounced amplitudes of the annual cycle is dominated by the influence of the Brewer–Dobson circulation with the highest O_3 values observed over polar regions (reaching $\sim 6 \times 10^{18}$ molecules cm^{-2} on average vs. $\sim 2 \times 10^{18}$ molecules cm^{-2} on average in the equatorial belt). The maximum is shifted from late winter at high latitudes to spring at lower latitudes.
3. In MLT (bottom right panel), we clearly see a large hemispheric difference with the highest values over the NH (also in the UTLS). Maxima are observed in spring, reflecting more effective STE processes. A particularly broad maximum from spring to late summer is observed in the 30 – 50° N band. It probably points to anthropogenic production of O_3 . This has been further investigated in the Supplement through a MOZART4–IASI comparison by using constant anthropogenic emissions in the model settings (see Fig. S2). The results show clear differences between the modelled and the observed MLT seasonal cycles, which highlights the need for further investigation of the role of anthropogenically




Figure 7. Monthly medians of measured (scatters) and of fitted (line) IASI O₃ columns averaged over the period 2008–2013, for the UST, MLST, UTLS and MLT layers and for each 20° latitude band (colour scale in the top-right panel). The fit is based on daily medians. Error bars give the 1σ standard deviation relative to the monthly median values. Correlation coefficients (*R*) between the daily median observations and the fit are also indicated. Note that the scales are different.

produced O₃ and the realism of anthropogenic emissions inventories.

Figure 8 presents all the fitted regression parameters included in Eq. (1) (Sect. 3) in the four layers as a function of latitude. The uncertainty in the 95 % confidence limits which accounts for the autocorrelation in the noise residual is given by error bars. The constant term (Fig. 8a) is found to be statistically significant (uncertainty < 10 %) in all cases. It captures the two ozone maxima in the stratosphere: one over the northern polar regions in the MLST and one at equatorial latitudes in the UST ($\sim 4.5 \times 10^{18}$ molecules cm⁻²), the important decrease of O₃ in the lower stratospheric layers (UTLS and MLST) moving from high to equatorial latitudes, and the weak negative and strong positive gradients in the northern MLT and in the UST, respectively. The sum of the constant terms of the four layers varies between 7.50×10^{18} (equatorial region) and 9.50×10^{18} molecules cm⁻² (polar regions) and is similar to the one of the fitted total column (relative differences < 3.5 %) (red line). Note that the constant terms in the UTLS region in the mid-latitudes and in the tropics are certainly affected by the fact that the FORLI-O₃ profiles are biased high by ~ 10 –15 % in this layer and latitude bands (Dufour et al., 2012; Gazeaux et al., 2013). The representativeness of the 20° zonal averages in terms of spatial variability has been examined by fitting the IASI time series

for specific locations in the NH (results shown with stars in Fig. 8a); the constant terms are found to be consistent, within their uncertainties, with those averaged per latitude bands in all cases. Over the polar region where O₃ shows a large natural variability, the regression coefficient is characterised by a large uncertainty.

The regression coefficients for other variables (harmonic and proxy terms) which are retained in the regression model by the stepwise elimination procedure are shown in Fig. 8b. They are scaled by the fitted constant term and the error bars represent the uncertainty in the 95 % confidence limits, accounting for the autocorrelation in the noise residual. A positive (or negative) sign of the coefficients indicates that the associated variables are correlated (or anti-correlated) with the IASI O₃ time series. Note that if the uncertainty is larger than its associated estimate (i.e. larger than 100 %, corresponding to an error bar overlapping the zero line), it means that the estimate becomes statistically non-significant when accounting for the autocorrelation in the noise residuals at the end of the elimination process. This is summarised in Table S1 of the Supplement. The contribution of the fitted variables into the IASI O₃ variations is estimated as $\frac{\sigma([a_n; b_n; x_j] \cdot [\cos(n\omega t); \sin(n\omega t); X_{\text{norm}, j}]}{\sigma(O_3(t))}$, where σ is the standard deviation relative to the fitted signal of harmonic or proxy




Figure 8. (a) Fitted constant factors (Cst; see Eq. 1, Sect. 3) from the 6-year IASI daily O₃ time series for the 20° latitude belts, separately given for the four layers and for the total column. The stars correspond to the constant factors fitted above ground-based measurement stations: Ny-Ålesund (79° N), Kiruna (68° N), Harestua (60° N), Jungfrauoch (47° N) and Izana (28° N). (b) Regression coefficients of the variables retained by the stepwise procedure, given in % as $[\text{regression_coefficient}/\text{fitted_cst}] \times 100\%$. Identification for the variables: annual (top left) and semi-annual variation (top right) terms, QBO at 10 and 30 hPa (bottom left) and solar flux (bottom right). Note that the scales are different. The associated fitting uncertainties (95 % confidence limits) are also represented (error bars).

terms and to the IASI O₃ time series. From Fig. 8, we find the following.

1. The annual harmonic term (upper left) is the main driver of the O₃ variability and largely dominates (scaled $a_1 + b_1$ around $\pm 40\%$) over the semi-annual one (upper right; scaled $a_2 + b_2$ around $\pm 15\%$). In the UTLS and MLST, its amplitude decreases from high to low latitudes, likely following the cycle induced by the Brewer–Dobson circulation (cf. Figs. 6 and 7); and the sign of the coefficient accounts for the winter–spring max-

ima in both hemispheres (negative values in the SH and positive ones in the NH). The annual term contributes importantly around 45–85 % of the observed O₃ variations, except in the 10–30° N and equatorial bands (10–30 %), while the influence of the semi-annual variation on O₃ is smaller (10–25 %) and highly variable between the bands. In the UST, the amplitudes vary only slightly (around -5 to 5%) and account for the weak summer maximum. The contributions of the annual harmonic term are estimated between 5 and 30 %. As expected,

the uncertainties associated with the annual terms are very weak, and most of the harmonic terms (annual and seasonal) are statistically significant.

2. The QBO and solar flux proxies are generally minor (scaled coefficients $< 10\%$ and contributions $< 15\%$) and they are often statistically non-significant contributors to O_3 variations after accounting for the autocorrelation in the noise residual (see Table S1 in the Supplement), except in the equatorial region (scaled coefficients of 10–15% in the UTLS and contributions up to 75 and 21% for QBO and SF, respectively) where they are important drivers of O_3 variations (e.g. Logan et al., 2003; Steinbrecht et al., 2006b; Soukharev and Hood, 2006; Fadnavis and Beig, 2009). Previous studies have indeed supported the solar influence on the lower stratospheric equatorial dynamics (e.g. Soukharev and Hood, 2006; McCormack et al., 2007). Note that the QBO³⁰ proxy (data not shown) has negative coefficients for the mid-latitudes, which is in line with Frossard et al. (2013).

The contributions described by the ENSO and NAO/AAO proxies are generally very weak (< 10 and $< 5\%$, respectively), with scaled coefficients lower than 5%, and, in many cases for the NAO/AAO proxies, they are even not statistically significant when taking into account the correlation in the noise residuals (see Table S1 in Supplement). Despite of this, it is worth pointing out that their effects on the O_3 variations are comparable to the results published in the previous studies. The negative ENSO coefficient in the tropical UTLS is consistent with results from Neu et al. (2014). Rieder et al. (2013) and Frossard et al. (2013) have also shown large regions of negative coefficients for NAO north of 40° N, and large regions of positive and negative coefficient estimates for ENSO, north of 30° N and south of 30° S, respectively.

We note that the non-representation of time lags in the proxy time series may be underestimating the role of some geophysical variables on O_3 variations, in particular that of ENSO and QBO in zonal bands outside the regions where these geophysical quantities are measured (i.e. the Niño region 3.4 for ENSO and Singapore for QBO). Finally, we see in Fig. 8b, large uncertainties associated with the regression coefficients in the UTLS in comparison with other layers, and in polar regions in comparisons with other bands. We interpret this as an effect from the high natural variability of O_3 measured by IASI in the UTLS (see Fig. 5) and from missing parameterisations and low temporal sampling of daytime IASI measurements over the poles, respectively.

As a general feature, the results demonstrate the representativeness of the fitted models in each layer and latitude band. This good performance of the regression procedure allows

examination of the adjusted linear trend term in Sect. 4.3 below.

4.3 Multivariate regression results: trend over 2008–2013

An additional goal of the multivariate regression method applied to the IASI O_3 time series is to determine the linear trend term and its associated uncertainty. Despite the fact that more than 10 years of observations, corresponding to the large scale of solar cycle, is usually required to perform such a trend analysis, we could argue that statistically relevant trends could possibly be derived from the first 6 years of IASI observations, owing to the high spatiotemporal frequency (daily) of IASI global observations, to the daily fingerprint in the solar flux (see Fig. 3a), possibly making it distinguishable from a linear trend, and to its weak contribution to O_3 variations (see Sect. 4.2 and references therein). To verify the specific advantage of IASI in terms of frequency sampling, we compare, in the subsections below, the statistical relevance of the trends when retrieved from the monthly averaged IASI data sets vs. the daily averages as above, in the 20° zonal bands for the four partial and total columns.

4.3.1 Regressions applied on daily vs. monthly averages

Figure 9 (top) provides, as an example, the 6-year time series of IASI O_3 daily averages (left panels) compared to the monthly averages (right panels) for the 30 – 50° S latitude band in the UST (dark blue), along with the results from the regression procedure (light blue). Note that either daily or monthly $F_{10.7}$, NAO and AAO proxies (see Table 1) are used depending on the frequency of the IASI O_3 averages to be adjusted. The second row in Fig. 9 provides the deseasonalised IASI and fitted time series, calculated by subtracting the model seasonal cycle from the time series, as well as the residuals (red curves). The averaged residuals relative to the deseasonalised IASI time series strongly vary with the layers and latitudinal bands and usually range between 30 and 60%. The fitted signal in DU of each proxy is shown in the bottom panels. The O_3 time series and the solar flux signal resulting from the adjustment without the linear term trend in the regression model are also represented (orange lines in second and bottom panels, respectively). When it is not included in the regression model, the linear trend term is only partly compensated by the solar flux term in the daily averages. This leads to an offset between the fitted O_3 time series resulting from the both regression models (with and without the linear term), which corresponds well to a trend over the IASI period, and, consequently, to larger residuals (e.g. 80% without vs. 44% with the linear term for this example and 94% without vs. 58% with the linear trend term for the 30 – 50° S band in the MLST illustrated in Fig. S1 of the Supplement). This offset is observed for a lot of layers and latitudinal bands. On the contrary, the linear term can largely be




Figure 9. Daily (a) and monthly (b) time series of O₃ measurements and of the fitted regression model in the UST for the 30–50° S latitude band (top row), of the deseasonalised O₃ (second row), of the difference of the fitted models with and without the linear term (third row) and of the fitted signal of proxies [regression_coefficient × proxy]: SF (blue), QBO (QBO¹⁰ + QBO³⁰; green), ENSO (red) and AAO (purple) (bottom) (given in DU). The averaged residuals relative to the deseasonalised IASI time series are also indicated (%).

compensated by the solar flux term in the monthly averages; the offset is weak and the relative difference between the both fitted models is smaller (averaged differences relative to the deseasonalised IASI time series of 10 % in monthly data vs. 17 % in daily data for this example). In this example, the linear and solar flux terms are even not simultaneously retained in the iterative stepwise backward procedure when applied on the monthly averages while they are when applied on daily averages. This effective colinearity of the linear and the monthly solar flux terms translates to larger model fit residuals (44 % in daily averages vs. 60 % in monthly averages in the UST, relative to the deseasonalised IASI time series), to smaller relative differences between both regression models (with and without the linear term) (17 % in daily vs. 10 % in monthly data), and to larger uncertainty on the trend coefficients when using the monthly data in comparison with the daily data. This even leads, in this specific example, to a non-statistically significant linear term of $1.21 \pm 1.30 \text{ DU year}^{-1}$ when derived from monthly averages vs. a significant trend of $1.74 \pm 0.77 \text{ DU year}^{-1}$ from daily averages.

The same conclusions can be drawn from the fits in other layers and latitude bands, especially those where the solar cycle variation of ozone is large (MLST and UTLS) or where the ozone recovery occurs (UST). A larger trend uncertainty associated with monthly data vs. daily data is found in all situations (see Table 2, Sect. 4.3.2 below).

This brings us to the important conclusion that thanks to the unprecedented sampling of IASI, apparent trends can be detected in FORLI-O₃ time series, even for a short period of measurements. This supports the need for regular and high-frequency measurements for observing ozone variations underlined in other studies (e.g. Saunois et al., 2012). The O₃ trends from the daily averages of IASI measurements are dis-

cussed and compared with results from the monthly averages in the subsection below.

4.3.2 O₃ trends from daily averages

Table 2 summarises the trends and their uncertainties in the 95 % confidence limit, calculated for each 20° zonal band and for the four partial and total columns. In the northern and southern polar regions, the polar night period is not covered because only IASI observations during sunlight (over February–October and October–April for NH and SH, respectively) are used in this study (see Sect. 2). For the sake of comparison, the trends are reported for both the daily (top values) and the monthly (bottom values) averages, and their uncertainties account for the autocorrelation in the noise residuals, considering a time lag of 1 day or 1 month, respectively. We show that the daily and monthly trends in all layers and all latitude bands fall within each other uncertainties, but that the use of daily median strongly helps in reducing the uncertainty associated with the trends everywhere, for the reasons discussed above (Sect. 4.3.1). This is particularly observed in the UST where the ozone hole recovery would occur, but also in the MLST and the UTLS where the solar cycle variation of ozone is the largest (see Fig. 8). As a consequence, the UST trends in monthly averages are shown to be mostly non-significant in comparison with those from daily averages. Table 3 summarises the trends in the daily averages for two 3-month periods: June–July–August (JJA) and December–January–February (DJF).

From Tables 2 and 3, we observe very different trends according to the latitude and the altitude. From Table 2, we find for the total columns that the trends derived from the daily medians are only significant at high northern latitudes and

Table 2. Ozone trends and associated uncertainties (95 % confidence limits; accounting for the autocorrelation in the noise residuals), given in DU year⁻¹, for 20° latitude bands, based on daily (top values) and monthly (bottom values) medians over 6 years of IASI observations. Bold (underlined) values refer to significant (positive) trends. Values marked with a star (*) refer to trends which are rejected by the iterative backward elimination procedure^a.

DU year ⁻¹	No. days	Ground–300 hPa (MLT)	300–150 hPa (UTLS)	150–25 hPa (MLST)	25–3 hPa (UST)	Total columns
70–90° N (Feb–Oct)	1493	-0.13 ± 0.10 -0.03 ± 0.29*	<u>1.28 ± 0.82</u> 0.70 ± 0.92*	<u>2.81 ± 2.27</u> -0.04 ± 2.60*	-0.16 ± 0.97* -1.81 ± 2.81*	<u>3.90 ± 2.93</u> 1.37 ± 3.62*
50–70° N	2103	-0.08 ± 0.09 0.17 ± 0.35*	<u>0.73 ± 0.51</u> 1.24 ± 1.24	0.97 ± 1.30 2.28 ± 4.24*	<u>0.55 ± 0.36</u> 0.66 ± 0.76	<u>1.93 ± 1.71</u> 4.72 ± 5.58
30–50° N	2105	-0.19 ± 0.05 -0.15 ± 0.13	<u>0.34 ± 0.18</u> 0.75 ± 0.75	-0.34 ± 0.77 -0.37 ± 1.65*	<u>0.89 ± 0.41</u> <u>0.87 ± 0.52</u>	0.91 ± 1.24 0.33 ± 2.25*
10–30° N	2105	0.10 ± 0.11 0.12 ± 0.15*	-0.03 ± 0.10* 0.05 ± 0.12*	-0.73 ± 0.29 -0.55 ± 0.62*	<u>0.95 ± 0.65</u> <u>1.25 ± 0.74</u>	0.21 ± 0.30* 0.82 ± 1.01
10° S–10° N	2104	-0.41 ± 0.12 -0.25 ± 0.14	-0.25 ± 0.07 -0.08 ± 0.10	-0.11 ± 0.26* -0.11 ± 0.64*	<u>0.44 ± 0.19</u> 0.61 ± 0.64	-0.16 ± 0.34 0.13 ± 0.83*
30–10° S	2106	-0.22 ± 0.10 -0.15 ± 0.13	-0.08 ± 0.04 -0.09 ± 0.07	-0.61 ± 0.26 -0.45 ± 0.36	<u>0.89 ± 0.58</u> 0.80 ± 1.23	-0.04 ± 0.31* -0.01 ± 1.26*
50–30° S	2105	-0.19 ± 0.07 -0.18 ± 0.09	-0.22 ± 0.08 -0.27 ± 0.12	-2.17 ± 0.58 -2.36 ± 1.80	<u>1.74 ± 0.77</u> 1.21 ± 1.30	-0.79 ± 0.96 -0.64 ± 1.45*
70–50° S	2105	-0.13 ± 0.05 -0.22 ± 0.12	0.09 ± 0.16 0.05 ± 0.32*	0.56 ± 0.82 0.02 ± 1.15*	<u>0.54 ± 0.29</u> <u>0.57 ± 0.82</u>	1.15 ± 1.28 0.51 ± 1.75*
90–70° S (Oct–Apr)	738	-0.15 ± 0.21* -0.17 ± 0.40*	0.01 ± 0.61* 0.25 ± 0.73*	0.00 ± 2.36* 2.59 ± 3.80*	<u>1.04 ± 0.57</u> 0.91 ± 2.10*	1.50 ± 3.15* 3.28 ± 5.12*

^a The trend values result from the adjustment of the regression model where the linear term is kept whatever its *p* value calculated during the iterative process is.

that they are interestingly of the same order as those obtained from other satellites and assimilated satellite data (Weatherhead and Anderson, 2006; Knibbe et al., 2014) or from ground-based measurements (Vigouroux et al., 2008) calculated over longer time periods. The non-significant trends calculated for the mid-latitudes and low latitudes of the NH are also comparable to the results published in the previous studies (Reinsel et al., 2005; Andersen et al., 2006; Vigouroux et al., 2008). Regarding the individual layers, we find the following.

In the US, significant positive trends are observed in both hemispheres from the daily medians, particularly over the mid-latitudes and high latitudes of both hemispheres (e.g. 1.74 ± 0.77 DU year⁻¹ in the 30–50° S band, i.e. 12 % decade⁻¹), where the changes in ozone trends before and after the turnaround in 1997 have been found to be the highest. K yrola et al. (2013) and Laine et al. (2014) report for instance a change of up to 10 % decade⁻¹ in O₃ trends between 1997 and 2011 vs. between 1984 and 1997. Positive trends in the UST are consistent with many previous observations if one considers the fact that the period covered by IASI is later than those reported in previous studies and that the recovery rate has seemed to have increased since the beginning of the turnaround (Knibbe et al., 2014, reports a factor of 2 increase in the recovery rate between 1997 and 2010 with ~ 0.7 DU year⁻¹ and 2001–2010 with ~ 1.4 DU year⁻¹ in the SH). They could indicate a levelling off of the nega-

tive trends that have been observed since the second half of the 1990s, mostly from satellites and ground-based monthly mean data (e.g. WMO, 2006, 2011; Randel and Wu, 2007; Vigouroux et al., 2008; Steinbrecht et al., 2009; Jones et al., 2009; McLinden et al., 2009; Bourassa et al., 2014; Laine et al., 2014; Nair et al., 2013). The causes of this turnaround remain, however, uncertain. If the compensating impact of decreasing chlorine in recent years and maximum solar cycle (over 2011–2012 in the period studied here) is probably part of the answer (e.g. Steinbrecht et al., 2004), the effects of changing stratospheric temperatures and Brewer–Dobson circulation (Salby et al., 2002; Reinsel et al., 2005; Dhomse et al., 2006; Manney et al., 2006) could also contribute and should be further investigated. The long-lasting cold winter/spring 2011 in the Arctic leading to unprecedented ozone loss (Manney et al., 2011), could explain the non-significant trend in the 70–90° N band. This is supported by the results in winter (Table 3). From Table 3, we generally find significant positive trends in summer NH and weaker positive or even non-significant trends in winter SH. A non-significant trend is also calculated for the 70–90° S band in spring (data not shown). This could indicate the strong influence of changing stratospheric temperatures on ozone depletion from year to year (e.g. Dhomse et al., 2006), leading to larger uncertainties in our trend estimations and larger fitting residuals (see Sect. 4.2) due to the fact that the stratospheric temperature

Table 3. Same as Table 2 but for seasonal O₃ trends and associated uncertainties based on daily medians during JJA (top values) and DJF (bottom values) periods. Values marked with a star (*) refer to trends which are rejected by the iterative backward elimination procedure^a.

DU year ⁻¹	No. days	Ground–300 hPa (MLT)	300–150 hPa (UTLS)	150–25 hPa (MLST)	25–3 hPa (UST)	Total columns
70–90° N	613	-0.18 ± 0.08	1.13 ± 0.65	-0.91 ± 1.52	1.72 ± 0.51	1.36 ± 1.15
(Feb–Oct)	48	–	–	–	–	–
50–70° N	551	-0.23 ± 0.07	1.03 ± 0.37	0.62 ± 1.64	1.67 ± 0.48	3.01 ± 1.64
	527	-0.09 ± 0.12*	1.74 ± 1.30	0.73 ± 1.73*	-0.66 ± 0.79	1.56 ± 2.66*
30–50° N	551	-0.30 ± 0.10	0.42 ± 0.30	-0.30 ± 0.65*	0.84 ± 0.25	1.17 ± 1.35
	529	-0.24 ± 0.09	0.28 ± 0.28	-0.82 ± 0.90	0.62 ± 0.49	-0.81 ± 1.05
10–30° N	551	-0.05 ± 0.16*	0.17 ± 0.05	-0.34 ± 0.30	0.36 ± 0.27	-0.09 ± 0.54*
	529	0.18 ± 0.14	0.01 ± 0.09*	-1.05 ± 0.45	0.49 ± 0.54*	-1.14 ± 0.44
10° S–10° N	551	-0.06 ± 0.10	0.04 ± 0.05*	-0.84 ± 0.86	0.32 ± 0.42	-0.56 ± 0.74
	529	-0.70 ± 0.23	-0.32 ± 0.10	1.64 ± 1.77	0.53 ± 0.59	0.34 ± 0.93*
30–10° S	551	-0.26 ± 0.09	-0.06 ± 0.07	-0.56 ± 0.40	1.06 ± 0.55	0.24 ± 0.43
	530	-0.15 ± 0.11	0.06 ± 0.12*	-0.12 ± 0.31*	1.48 ± 0.53	1.56 ± 0.92
50–30° S	551	-0.21 ± 0.05	-0.16 ± 0.09	-0.52 ± 0.54	0.49 ± 0.59	-0.44 ± 0.83
	529	-0.10 ± 0.06	-0.14 ± 0.06	-2.83 ± 0.64	3.40 ± 0.85	0.47 ± 0.52
70–50° S	551	-0.25 ± 0.06	1.03 ± 0.60	2.63 ± 1.65	0.98 ± 0.62	3.44 ± 2.47
	529	-0.10 ± 0.04	0.19 ± 0.24*	0.52 ± 0.48	1.66 ± 0.70	1.72 ± 0.74
90–70° S	–	–	–	–	–	–
(Oct–Apr)	523	-0.21 ± 0.20	-0.46 ± 0.80*	0.16 ± 2.53*	1.18 ± 0.67	0.98 ± 3.27*

^a The trend values result from the adjustment of the regression model where the linear term is kept whatever its *p* value calculated during the iterative process is.

is not taken into account as an explanatory variable in the model.

In the MLST, one can see that, except in the high latitude bands, the trends are either non-significant or significantly negative. This is in agreement with the trend analysis of Jones et al. (2009) for the 20–25 km altitude range over the 1997–2008 period, as well as with other studies at NH latitudes, which investigated O₃ changes in the 18–25 km range between 1996 and 2005 (Miller et al., 2006; Yang et al., 2006; Kivi et al., 2007). The results derived separately for summer and winter in Table 3 are also in line with those of Kivi et al. (2007) which reported contrasted trends in the Arctic MLST depending on season.

In the UTLS, negative trends are calculated in the tropics and significant positive trends are found in the mid-latitudes and high latitudes of NH, the latter falling within the uncertainties of those reported by Kivi et al. (2007) for the tropopause–150 hPa layer between 1996 and 2003. The large positive trends calculated at northern latitudes (e.g. 1.28 ± 0.82 DU year⁻¹ in the 70–90° N band) contribute ~ 30 % to the positive trend in the total column. This result is consistent with Yang et al. (2006) which reported that the UTLS contributes 50 % to positive trends for the total columns measured in the mid-latitudes of the NH from ozonesondes. In that study, these positive trends were linked to changes in atmospheric dynamics either related to natural variability induced by potential vorticity and tropopause height variations or related to anthropogenic climate change. Hence, the apparent increase in total ozone in the mid-

latitudes of the NH seen by IASI would reflect the combined contribution of dynamical variability and declining ozone-depleting substances (e.g. Weatherhead and Andersen, 2006; WMO, 2006; Harris et al., 2008; Nair et al., 2013). It is worth keeping in mind that these effects are not independently accounted for in the regression model. Previous studies have reported, however, that dynamical and chemical processes are physically coupled in the atmosphere, making it difficult to define such drivers in a statistical model unambiguously (e.g. Mäder et al., 2007; Harris et al., 2008). On a seasonal basis (see Table 3), the trends seen by IASI at northern latitudes in summer are all significantly positive and increase towards the pole. Note that the trends in upper layers may contribute to the ones calculated in the UTLS due to the medium IASI sensitivity to that layer (cf. Sect. 2).

In the MLT, most of the trends are significantly negative (Tables 2 and 3). The non-significant trends in polar regions could be partly related to the lack of IASI sensitivity to tropospheric O₃ (see Sect. 2, Fig. 2). On a seasonal basis, we see that the negative trends are more pronounced during the JJA period (around -0.25 ± 0.10 DU year⁻¹) for all bands except between 30° N and 10° S. In the NH, these results tend to confirm the levelling off of tropospheric ozone observed in recent years during the summer months (Logan et al., 2012). This trend, however, remains difficult to interpret because it could be linked to a variety of processes including most importantly, the decline of anthropogenic emissions of ozone precursors, the increase of ultraviolet (UV)-induced O₃ destruction in the troposphere and STE processes (Isaksen et

al., 2005; Logan et al., 2012; Parrish et al., 2012; Hess and Zbinden, 2013). As a consequence, it is hard to reconcile the trends in tropospheric ozone with changes in emissions of ozone precursors. However, trends in emissions have already been able to qualitatively explain measured ozone trends over some regions but with inconsistent magnitude between observations and model simulations (e.g. Cooper et al., 2010; Logan et al., 2012; Wilson et al., 2012). It is also worth keeping in mind that due to medium sensitivity of IASI to the troposphere, the a priori contribution and ozone variations in stratospheric layers may largely influence the trends seen by IASI in the MLT layer (cf. Sect. 2 and Supplement).

4.3.3 O₃ trends from IASI vs. FTIR data

In order to validate the trends inferred from IASI in the UST and in the total columns, we compare them with those obtained from ground-based FTIR measurements at several NDACC stations (Network for the Detection of Atmospheric Composition Change, available at http://www.ndsc.ncep.noaa.gov/data/data_tbl/) by using the same fitting procedure and taking into account the autocorrelation in the noise residuals. A box of $1^\circ \times 1^\circ$ centred on the stations has been used for the collocation criterion. The regression model is applied on the daily FTIR data for a series of time periods starting after the turnaround point (from 1998 for mid-latitude stations and from 2000 for polar stations), as well as for the same periods as recently studied in Vigouroux et al. (2015) for the sake of comparison. Note that because we are not interested here in validating the IASI columns, which was achieved in previous papers (e.g. Dufour et al., 2012; Oetjen et al., 2014), but in validating the trends obtained from IASI, we did not correct biases between IASI and FTIR due to different vertical sensitivity and a priori information. The results are given in DU year⁻¹ in Table 4. We see large significant positive total column trends from IASI at middle and polar stations (e.g. 5.26 ± 4.72 DU year⁻¹ at Ny-Ålesund), especially during spring. These values are consistent with those reported in Knibbe et al. (2014) for the 2001–2010 period in spring in the Antarctic (around 3–5 DU year⁻¹). This trend is not obtained from the FTIR data for which trends are found to be mostly non-significant (even not retained in the stepwise elimination procedure in some cases) as reported in Vigouroux et al. (2015), except at Jungfrauoch which shows a trend of 5.28 ± 4.82 DU year⁻¹ over the 2008–2012 period. For the periods starting before 2000, we calculated from FTIR, in agreement with Vigouroux et al. (2015), a significantly negative trend at Ny-Ålesund for the total column and significantly positive trends at polar stations for the US. In addition, we see from Table 4 a levelling off of O₃ at polar stations in the UST after 2003, as previously reported in Vigouroux et al. (2015), which is explained by a compensation effect between the decrease of solar cycle after its maximum in 2001–2002 and a positive trend. These trends are,

however, non-significant and inferred only from few FTIR measurements (see number of days column, Table 4).

From IASI, it is worth pointing out that, in all cases, positive trends are calculated in the UST (even if some are not significant) and that these trends are consistent with those calculated from FTIR data covering a ~ 11 -year period and starting after the turnaround (e.g. at Thule; 1.24 ± 1.09 DU year⁻¹ from IASI for the period 2008–2013 vs. 1.42 ± 0.78 DU year⁻¹ from the FTIR over 2001–2012). This is illustrated for three stations (Ny-Ålesund, Thule and Kiruna) in Fig. 10 which compares the time series from IASI (2008–2013, in red) with those from FTIR covering periods starting after the turnaround (in blue). Their associated trends as well as the trend calculated from FTIR covering the IASI period (in green) are also indicated.

In order to better characterise the effect of the temporal frequency on determining statistical trends, the IASI time series have been subsampled to match the temporal resolution of FTIR. The associated trend values are also indicated in Table 4 (second row). In all cases, we observe that the fitted trends inferred from both IASI and FTIR with the same temporal samplings are within the uncertainties of each other and that those associated with the subsampled IASI data sets are significantly larger than those obtained with the daily ones, leading to statistically non-significant trends.

Even if validating the IASI fitted trends with independent data sets is challenging due to the short-time period of available IASI measurements and the insufficient number of usable correlative measurements over such a short period, the results obtained for IASI vs. FTIR tend to confirm the conclusion drawn in Sects. 4.3.1 and 4.3.2, that the high temporal sampling of IASI provides good confidence in the determination of the trends even for periods shorter than those usually required from other observational means.

5 Summary and conclusions

In this study, we have analysed 6 years of IASI O₃ profile measurements as well as the total O₃ columns based on the profile. Four layers have been defined following the ability of IASI to provide reasonably independent information on the ozone partial columns: the mid–lower troposphere (MLT), the upper troposphere–lower stratosphere (UTLS), the mid–lower stratosphere (MLST) and the upper stratosphere (UST). Based on daily values of these four partial or total columns in 20° zonal averages, we have demonstrated the capability of IASI of capturing large-scale ozone variability (seasonal cycles and trends) in these different layers. We have presented daytime vertical and latitudinal distributions for O₃ as well as their evolution with time and we have examined the underlying dynamical or chemical processes. The distributions were found to be controlled by photochemical production, leading to a maximum in summer in the equatorial region in the UST, while they reflect the impact of

Table 4. Ozone trends and associated uncertainties (95 % confidence limits), given in DU year⁻¹ over NDACC (Network for the Detection of Atmospheric Composition Change) stations in the NH based on daily medians of IASI (within a grid box of 1° × 1° centred on stations, two first rows) and FTIR observations (successive rows for different time intervals). Italic values (second row) refer to trends inferred from subsampled IASI data and bold values refer to statistically significant trends. Values marked with a star (*) refer to trends which are rejected by the iterative backward elimination procedure^a.

DU year ⁻¹	Data periods	No. days	25–3 hPa (US)	Total columns
Ny-Ålesund (79° N) Mar–Sep	2008–2013	1239	0.56 ± 0.73	5.26 ± 4.72
	<i>Subsamp.</i>	82	<i>-0.29 ± 4.58</i>	<i>6.26 ± 18.11</i>
	2008–2012	84	-3.58 ± 4.58	2.24 ± 20.78*
	2008–2012	168	-0.17 ± 0.70*	-4.84 ± 3.01
	2003–2012	288	0.64 ± 0.60	-1.02 ± 2.40*
	2000–2012	320	0.62 ± 0.55	-2.35 ± 1.40
	1999–2012	383	1.03 ± 0.66	1.31 ± 2.39*
	1995–2012	167	1.25 ± 1.05	3.33 ± 3.41
Thule (77° N) Mar–Sep	2008–2013	1094	1.24 ± 1.09	4.97 ± 4.72
	<i>Subsamp.</i>	231	<i>1.31 ± 2.69</i>	<i>0.10 ± 7.36</i>
	2008–2012	340	-2.10 ± 2.89	0.39 ± 11.59*
	2008–2012	697	0.86 ± 0.89	-2.77 ± 2.99
	2003–2012	776	1.33 ± 0.86	-1.29 ± 1.73
	2000–2012	779	1.69 ± 0.88	-1.25 ± 1.74
	1999–2012	138	3.73 ± 2.90	4.86 ± 10.13*
	1999–2003			
Kiruna (68° N) Mar–Sep	2008–2013	1236	0.21 ± 1.42	4.41 ± 4.00
	<i>Subsamp.</i>	226	<i>0.97 ± 4.05</i>	<i>3.78 ± 6.03</i>
	2008–2012	254	-1.97 ± 6.04*	-3.75 ± 6.64*
	2008–2012	678	0.15 ± 0.67*	2.26 ± 3.68
	2003–2012	913	1.60 ± 1.29	3.69 ± 4.20
	2000–2012	984	1.10 ± 0.98	-0.43 ± 1.64*
	1999–2012	1183	1.11 ± 0.54	1.82 ± 1.77
	1996–2012	596	1.26 ± 1.21	1.12 ± 3.77*
Jungfraujoch (47° N)	2008–2013	1580	2.95 ± 0.61	5.64 ± 3.15
	<i>Subsamp.</i>	524	<i>3.72 ± 1.14</i>	<i>5.61 ± 5.11</i>
	2008–2012	565	1.60 ± 1.80	5.28 ± 4.82
	2008–2012	1582	0.10 ± 0.35	-0.28 ± 0.86*
	1998–2012	1771	0.02 ± 0.33*	0.85 ± 0.79
	1995–2012			
Zugspitze (47° N)	2008–2013	1729	3.17 ± 0.56	5.53 ± 2.92
	<i>Subsamp.</i>	538	<i>3.56 ± 1.63</i>	<i>5.99 ± 4.49</i>
	2008–2012	597	0.71 ± 1.22	3.46 ± 3.79
	2008–2012	1472	0.08 ± 0.32*	0.81 ± 0.98
	1998–2012	1525	0.23 ± 0.32	1.36 ± 1.01
	1995–2012			
Izana (28° N)	2008–2013	1803	0.56 ± 0.65	1.28 ± 0.77
	<i>Subsamp.</i>	380	<i>0.32 ± 1.28</i>	<i>0.11 ± 1.95</i>
	2008–2012	443	0.24 ± 0.80*	0.91 ± 2.44*
	2008–2012	1257	0.46 ± 0.25	0.20 ± 0.33*
	1999–2012			

^a The trend values result from the adjustment of the regression model where the linear term is kept whatever its *p* value calculated during the iterative process is.




Figure 10. Daily time series of O₃ FTIR (blue symbols) and IASI (red symbols) measurements in the UST at Ny-Ålesund (top), Thule (middle) and Izana (bottom), covering the 1995–2012 and the 1999–2012 periods, respectively (given in DU). The fitted regression models (dark blue and dark red lines, for FTIR and IASI, respectively) and the linear trends calculated for periods starting after the turnaround over 1999/2000–2012 and over 2008–2012 for FTIR (light blue and green lines), and the 2008–2013 period for IASI (orange line) are also represented (DU year⁻¹). The trend values given in DU year⁻¹ are indicated.

the Brewer–Dobson circulation with a maximum in winter–spring at mid-latitudes and high latitudes in the MLST and in the troposphere. The effect of the photochemical production of O₃ from anthropogenic precursor emissions was also observed in the troposphere with a shift in the timing of the maximum from spring to summer in the mid-latitudes of the NH.

The dynamical and chemical contributions contained in the daily time development of IASI O₃ have been analysed by fitting the time series in each layer and for the total column with a set of parameterised geophysical variables, a constant factor and a linear trend term. The model was shown to perform well in term of residuals (< 10 %), correlation coefficients (between 0.70 and 0.99) and statistical uncertainties (< 7 %) for each fitted proxies. The annual harmonic terms (seasonal behaviour) were found to be largely dominant in all layers but the UST, with fitted amplitudes decreasing from

high to low latitudes in agreement with the Brewer–Dobson circulation. The QBO and solar flux terms were calculated to be important only in the equatorial region, while other dynamical proxies accounted for in the regression (ENSO, NAO, AAO) were found to be negligible.

Despite the short time period of the available IASI data set used in this study (2008–2013) and the potential ambiguity between the solar and the linear trend terms, statistically significant trends were derived from the 6 first years of daily O₃ partial columns measurements (as opposed to monthly averages which lead to mostly non-significant trends). This result, which was strengthened from comparisons with the regression applied on local FTIR measurements, is remarkable as it demonstrates the added value of IASI exceptional frequency sampling for monitoring medium- to long-term changes in global ozone concentrations. We found two important apparent trends.

1. There are significant positive trends in the upper stratosphere, especially at high latitudes in both hemispheres (e.g. $1.74 \pm 0.77 \text{ DU year}^{-1}$ in the $30\text{--}50^\circ \text{ S}$ band), which are consistent with a probable turnaround for upper stratospheric O_3 recovery (even if the causes of such a turnaround are still under investigations). In addition, the trends calculated for some local stations are in line with those calculated from FTIR measurements after the turnaround.
2. There are negative trends in the troposphere at mid-latitudes and high northern latitudes, especially during summer (e.g. $-0.26 \pm 0.11 \text{ DU year}^{-1}$ in the $30\text{--}50^\circ \text{ N}$ band), which are in line with the decline of ozone precursor emissions.

To confirm the above findings beyond the 6 first years of IASI measurements and to better disentangle the effects of dynamical changes of the 11-year solar cycle and of the equivalent effective stratospheric chlorine (EESC) decline on the O_3 time series, further years of IASI observations will be required, and more complete fitting procedures (including, among others, proxies to account for the decadal trend in the EESC, for the ozone hole formation and for changes in the Brewer–Dobson circulation, as well as including time lags in ENSO and QBO proxies) will have to be explored. Further investigation on the regressors' uncertainties and on the total error on ozone measurements should be performed as well to understand the unexplained variations in IASI O_3 records.

This will be achievable with the long-term homogeneous records obtained by merging measurements from the three successive IASI instruments on MetOp-A (2006), -B (2012) and -C (2018), and by IASI successor on EPS-SG after 2021 (Clerbaux and Crevoisier, 2013; Crevoisier et al., 2014).

Data availability

The ground-based FTIR measurements are accessible from http://www.ndsc.ncep.noaa.gov/data/data_tbl/.

The Supplement related to this article is available online at doi:10.5194/acp-16-5721-2016-supplement.

Acknowledgements. IASI has been developed and built under the responsibility of the Centre National d'Etudes Spatiales (CNES, France). It is flown onboard the MetOp satellites as part of the EUMETSAT Polar System. The IASI L1 data are received through the EUMETCast near-real-time data distribution service. Ozone data used in this paper are freely available upon request to the corresponding author. We acknowledge support from the O_3 -CCI project funded by ESA and by the O3M-SAF project funded by EUMETSAT. P.-F. Coheur and C. Wespes are, respectively, Senior Research Associate and Postdoctoral Researcher with

F.R.S.-FNRS. The research in Belgium was also funded by the Belgian State Federal Office for Scientific, Technical and Cultural Affairs and the European Space Agency (ESA Prodex IASI Flow and BO_3MSAF). The National Center for Atmospheric Research is funded by the National Science Foundation.

Edited by: M. Van Roozendael

References

- Andersen, S. B., Weatherhead, E. C., Stevermer, A., Austin, J., Brühl, C., Fleming, E. L., de Grandpré, J., Grewe, V., Isaksen, I., Pitari, G., Portmann, R. W., Rognerud, B., Rosenfield, J. E., Smyshlyaev, S., Nagashima, T., Velders, G. J. M., Weisenstein, D. K., and Xia, K.: Comparison of recent modeled and observed trends in total column ozone, *J. Geophys. Res.*, 111, D02303, doi:10.1029/2005JD006091, 2006.
- Anton, M., Loyola, D., Clerbaux, C., Lopez, M., Vilaplana, J., Banon, M., Hadji-Lazaro, J., Valks, P., Hao, N., Zimmer, W., Coheur, P., Hurtmans, D., and Alados-Arboledas, L.: Validation of the Metop-A total ozone data from GOME-2 and IASI using reference ground-based measurements at the Iberian peninsula, *Remote Sens. Environ.*, 115, 1380–1386, 2011.
- August, T., Klaes, D., Schlüssel, P., Hultberg, T., Crapeau, M., Ariaga, A., O'Carroll, A., Coppens, D., Munro, R., and Calbet, X.: IASI on Metop-A: Operational Level 2 retrievals after five years in orbit, *J. Quant. Spectrosc. Ra.*, 114, 1340–1371, 2012.
- Barret, B., Le Flochmoen, E., Sauvage, B., Pavelin, E., Matricardi, M., and Cammas, J. P.: The detection of post-monsoon tropospheric ozone variability over south Asia using IASI data, *Atmos. Chem. Phys.*, 11, 9533–9548, doi:10.5194/acp-11-9533-2011, 2011.
- Bourassa, A. E., Degenstein, D. A., Randel, W. J., Zawodny, J. M., Kyrölä, E., McLinden, C. A., Sioris, C. E., and Roth, C. Z.: Trends in stratospheric ozone derived from merged SAGE II and Odin-OSIRIS satellite observations, *Atmos. Chem. Phys.*, 14, 6983–6994, doi:10.5194/acp-14-6983-2014, 2014.
- Boynard, A., Clerbaux, C., Coheur, P.-F., Hurtmans, D., Turquety, S., George, M., Hadji-Lazaro, J., Keim, C., and Meyer-Arnek, J.: Measurements of total and tropospheric ozone from IASI: comparison with correlative satellite, ground-based and ozonesonde observations, *Atmos. Chem. Phys.*, 9, 6255–6271, doi:10.5194/acp-9-6255-2009, 2009.
- Brasseur, G. and Solomon, S.: *Aeronomy of the Middle Atmosphere*, 441 pp., D. Reidel Publishing Company, Dordrecht, the Netherlands, 1984.
- Brohede, S., McLinden, C. A., Urban, J., Haley, C. S., Jonsson, A. I., and Murtagh, D.: Odin stratospheric proxy NO_y measurements and climatology, *Atmos. Chem. Phys.*, 8, 5731–5754, doi:10.5194/acp-8-5731-2008, 2008.
- Chipperfield, M. P.: A three-dimensional model study of long-term mid-high latitude lower stratosphere ozone changes, *Atmos. Chem. Phys.*, 3, 1253–1265, doi:10.5194/acp-3-1253-2003, 2003.
- Chipperfield, M. P., Kinnersley, J. S., and Zawodny, J.: A two-dimensional model study of the QBO signal in SAGE II NO_2 and O_3 , *Geophys. Res. Lett.*, 21, 589–592, 1994.

- Clarisse, L., Hurtmans, D., Clerbaux, C., Hadji-Lazaro, J., Ngadi, Y., and Coheur, P.-F.: Retrieval of sulphur dioxide from the infrared atmospheric sounding interferometer (IASI), *Atmos. Meas. Tech.*, 5, 581–594, doi:10.5194/amt-5-581-2012, 2012.
- Clerbaux, C. and Crevoisier, C.: New Directions: Infrared remote sensing of the troposphere from satellite: Less, but better, *Atmos. Environ.*, 72, 24–26, 2013.
- Clerbaux, C., Boynard, A., Clarisse, L., George, M., Hadji-Lazaro, J., Herbin, H., Hurtmans, D., Pommier, M., Razavi, A., Turquety, S., Wespes, C., and Coheur, P.-F.: Monitoring of atmospheric composition using the thermal infrared IASI/MetOp sounder, *Atmos. Chem. Phys.*, 9, 6041–6054, doi:10.5194/acp-9-6041-2009, 2009.
- Cooper, O., Parrish, D., Stohl, A., Trainer, M., Nédélec, P., Thouret, V., Cammas, J., Oltmans, S., Johnson, B., and Tarasick, D.: Increasing springtime ozone mixing ratios in the free troposphere over western North America, *Nature*, 463, 344–348, doi:10.1038/nature08708, 2010.
- Crevoisier, C., Clerbaux, C., Guidard, V., Phulpin, T., Armante, R., Barret, B., Camy-Peyret, C., Chaboureaud, J.-P., Coheur, P.-F., Crépeau, L., Dufour, G., Labonne, L., Lavanant, L., Hadji-Lazaro, J., Herbin, H., Jacquinet-Husson, N., Payan, S., Péquignot, E., Pierangelo, C., Sellitto, P., and Stubenrauch, C.: Towards IASI-New Generation (IASI-NG): impact of improved spectral resolution and radiometric noise on the retrieval of thermodynamic, chemistry and climate variables, *Atmos. Meas. Tech.*, 7, 4367–4385, doi:10.5194/amt-7-4367-2014, 2014.
- de Laat, A. T. J., van der A, R. J., and van Weele, M.: Tracing the second stage of ozone recovery in the Antarctic ozone-hole with a “big data” approach to multivariate regressions, *Atmos. Chem. Phys.*, 15, 79–97, doi:10.5194/acp-15-79-2015, 2015.
- Dhomse, S., Weber, M., Wohltmann, I., Rex, M., and Burrows, J. P.: On the possible causes of recent increases in northern hemispheric total ozone from a statistical analysis of satellite data from 1979 to 2003, *Atmos. Chem. Phys.*, 6, 1165–1180, doi:10.5194/acp-6-1165-2006, 2006.
- Dufour, G., Eremenko, M., Orphal, J., and Flaud, J.-M.: IASI observations of seasonal and day-to-day variations of tropospheric ozone over three highly populated areas of China: Beijing, Shanghai, and Hong Kong, *Atmos. Chem. Phys.*, 10, 3787–3801, doi:10.5194/acp-10-3787-2010, 2010.
- Dufour, G., Eremenko, M., Griesfeller, A., Barret, B., LeFlochmoën, E., Clerbaux, C., Hadji-Lazaro, J., Coheur, P.-F., and Hurtmans, D.: Validation of three different scientific ozone products retrieved from IASI spectra using ozonesondes, *Atmos. Meas. Tech.*, 5, 611–630, doi:10.5194/amt-5-611-2012, 2012.
- Fadnavis, S. and Beig, J.: Quasi-biennial Oscillation in Ozone and Temperature over Tropics, *J. Atmos. Sol.-Terr. Phys.*, 71, 1450–1455, doi:10.1016/j.jastp.2008.11.012, 2009.
- Frossard, L., Rieder, H. E., Ribatet, M., Staehelin, J., Maeder, J. A., Di Rocco, S., Davison, A. C., and Peter, T.: On the relationship between total ozone and atmospheric dynamics and chemistry at mid-latitudes – Part 1: Statistical models and spatial fingerprints of atmospheric dynamics and chemistry, *Atmos. Chem. Phys.*, 13, 147–164, doi:10.5194/acp-13-147-2013, 2013.
- Fusco, A. C. and Logan, J. A.: Analysis of 1970–1995 trends in tropospheric ozone at Northern Hemisphere midlatitudes with the GEOS-CHEM model, *J. Geophys. Res.*, 108, 4449, doi:10.1029/2002JD002742, 2003.
- Gardiner, T., Forbes, A., de Mazière, M., Vigouroux, C., Mahieu, E., Demoulin, P., Velasco, V., Notholt, J., Blumenstock, T., Hase, F., Kramer, I., Sussmann, R., Stremme, W., Mellqvist, J., Strandberg, A., Ellingsen, K., and Gauss, M.: Trend analysis of greenhouse gases over Europe measured by a network of ground-based remote FTIR instruments, *Atmos. Chem. Phys.*, 8, 6719–6727, doi:10.5194/acp-8-6719-2008, 2008.
- Gazeaux, J., Clerbaux, C., George, M., Hadji-Lazaro, J., Kuttippurath, J., Coheur, P.-F., Hurtmans, D., Deshler, T., Kovilakam, M., Campbell, P., Guidard, V., Rabier, F., and Thépaut, J.-N.: Inter-comparison of polar ozone profiles by IASI/MetOp sounder with 2010 Concordiasi ozonesonde observations, *Atmos. Meas. Tech.*, 6, 613–620, doi:10.5194/amt-6-613-2013, 2013.
- Gebhardt, C., Rozanov, A., Hommel, R., Weber, M., Bovensmann, H., Burrows, J. P., Degenstein, D., Froidevaux, L., and Thompson, A. M.: Stratospheric ozone trends and variability as seen by SCIAMACHY from 2002 to 2012, *Atmos. Chem. Phys.*, 14, 831–846, doi:10.5194/acp-14-831-2014, 2014.
- Hadjinicolaou, P., Pyle, J. A., and Harris, N. R. P.: The recent turnaround in stratospheric ozone over northern middle latitudes: A dynamical modeling perspective, *Geophys. Res. Lett.*, 32, L12821, doi:10.1029/2005GL022476, 2005.
- Harris, N. R. P., Kyrö, E., Staehelin, J., Brunner, D., Andersen, S.-B., Godin-Beekmann, S., Dhomse, S., Hadjinicolaou, P., Hansen, G., Isaksen, I., Jrrar, A., Karpetchko, A., Kivi, R., Knudsen, B., Krizan, P., Lastovicka, J., Maeder, J., Orsolini, Y., Pyle, J. A., Rex, M., Vanicek, K., Weber, M., Wohltmann, I., Zanis, P., and Zerefos, C.: Ozone trends at northern mid- and high latitudes – a European perspective, *Ann. Geophys.*, 26, 1207–1220, doi:10.5194/angeo-26-1207-2008, 2008.
- Hauchecorne, A., Bertaux, J. L., Dalaudier, F., Keckhut, P., Lemenais, P., Bekki, S., Marchand, M., Lebrun, J. C., Kyrölä, E., Tamminen, J., Sofieva, V., Fussen, D., Vanhellemont, F., Fanton d’Andon, O., Barrot, G., Blanot, L., Fehr, T., and Saavedra de Miguel, L.: Response of tropical stratospheric O₃, NO₂ and NO₃ to the equatorial Quasi-Biennial Oscillation and to temperature as seen from GOMOS/ENVISAT, *Atmos. Chem. Phys.*, 10, 8873–8879, doi:10.5194/acp-10-8873-2010, 2010.
- Hess, P. G. and Zbinden, R.: Stratospheric impact on tropospheric ozone variability and trends: 1990–2009, *Atmos. Chem. Phys.*, 13, 649–674, doi:10.5194/acp-13-649-2013, 2013.
- Hofmann, D. J.: Recovery of Antarctic ozone hole, *Nature*, 384, 222–223, doi:10.1038/384222a0, 1996.
- Hood, L. L. and Soukharev, B. E.: Solar induced variations of odd nitrogen: Multiple regression analysis of UARS HALOE data, *Geophys. Res. Lett.*, 33, L22805, doi:10.1029/2006GL028122, 2006.
- Hurtmans, D., Coheur, P., Wespes, C., Clarisse, L., Scharf, O., Clerbaux, C., Hadji-Lazaro, J., George, M., and Turquety, S.: FORLI radiative transfer and retrieval code for IASI, *J. Quant. Spectrosc. Ra.*, 113, 1391–1408, 2012.
- Isaksen, I. S. A., Zerefos, C. S., Kourtidis, K., Meleti, C., Dalsøren, S. B., Sundet, J. K., Grini, A., Zanis, P., and Balis, D.: Tropospheric ozone changes at unpolluted and semipolluted regions induced by stratospheric ozone changes, *J. Geophys. Res.*, 110, D02302, doi:10.1029/2004JD004618, 2005.

- Jiang, X., Pawson, S., Camp, C. D., Nielsen, E., Shia, R., Liao, T., Jeev, K., Limpasuvan, V., and Yung, Y. L.: Interannual variability and trends in extratropical ozone. Part II: Southern Hemisphere, *J. Atmos. Sci.*, 65, 3030–3041, 2008.
- Jones, A., Urban, J., Murtagh, D. P., Eriksson, P., Brohede, S., Haley, C., Degenstein, D., Bourassa, A., von Savigny, C., Sonkaew, T., Rozanov, A., Bovensmann, H., and Burrows, J.: Evolution of stratospheric ozone and water vapour time series studied with satellite measurements, *Atmos. Chem. Phys.*, 9, 6055–6075, doi:10.5194/acp-9-6055-2009, 2009.
- Kivi, R., Kyrö, E., Turunen, T., Harris, N. R. P., von der Gathen, P., Rex, M., Andersen, S. B., and Wohltmann, I.: Ozone sondes observations in the Arctic during 1989–2003: Ozone variability and trends in the lower stratosphere and free troposphere, *J. Geophys. Res.*, 112, D08306, doi:10.1029/2006JD007271, 2007.
- Knibbe, J. S., van der A, R. J., and de Laat, A. T. J.: Spatial regression analysis on 32 years of total column ozone data, *Atmos. Chem. Phys.*, 14, 8461–8482, doi:10.5194/acp-14-8461-2014, 2014.
- Kramarova, N. A., Nash, E. R., Newman, P. A., Bhartia, P. K., McPeters, R. D., Rault, D. F., Seftor, C. J., Xu, P. Q., and Labow, G. J.: Measuring the Antarctic ozone hole with the new Ozone Mapping and Profiler Suite (OMPS), *Atmos. Chem. Phys.*, 14, 2353–2361, doi:10.5194/acp-14-2353-2014, 2014.
- Kuttippurath, J., Lefèvre, F., Pommereau, J.-P., Roscoe, H. K., Goutail, F., Pazmiño, A., and Shanklin, J. D.: Antarctic ozone loss in 1979–2010: first sign of ozone recovery, *Atmos. Chem. Phys.*, 13, 1625–1635, doi:10.5194/acp-13-1625-2013, 2013.
- Kuttippurath, J., Bodeker, G. E., Roscoe, H. K., and Nair, P. J.: A cautionary note on the use of EESC-based regression analysis for ozone trend studies, *Geophys. Res. Lett.*, 42, 162–168, doi:10.1002/2014GL02142, 2015.
- Kyrölä, E., Tamminen, J., Leppelmeier, G. W., Sofieva, V., Hassinen, S., Seppälä, A., Verronen, P. T., Bertaux, J.-L., Hauchecorne, A., Dalaudier, F., Fussen, D., Vanhellemont, F., d'Andon, O. F., Barrot, G., Mangin, A., Theodore, B., Guirlet, M., Koopman, R., Saavedra, L., Snoeij, P., and Fehr, T.: Night-time ozone profiles in the stratosphere and mesosphere by the Global Ozone Monitoring by Occultation of Stars on Envisat, *J. Geophys. Res.*, 111, D24306, doi:10.1029/2006JD007193, 2006.
- Kyrölä, E., Tamminen, J., Sofieva, V., Bertaux, J. L., Hauchecorne, A., Dalaudier, F., Fussen, D., Vanhellemont, F., Fanton d'Andon, O., Barrot, G., Guirlet, M., Fehr, T., and Saavedra de Miguel, L.: GOMOS O₃, NO₂, and NO₃ observations in 2002–2008, *Atmos. Chem. Phys.*, 10, 7723–7738, doi:10.5194/acp-10-7723-2010, 2010.
- Kyrölä, E., Laine, M., Sofieva, V., Tamminen, J., Päivärinta, S.-M., Tukiainen, S., Zawodny, J., and Thomason, L.: Combined SAGE II–GOMOS ozone profile data set for 1984–2011 and trend analysis of the vertical distribution of ozone, *Atmos. Chem. Phys.*, 13, 10645–10658, doi:10.5194/acp-13-10645-2013, 2013.
- Laine, M., Latva-Pukkila, N., and Kyrölä, E.: Analysing time-varying trends in stratospheric ozone time series using the state space approach, *Atmos. Chem. Phys.*, 14, 9707–9725, doi:10.5194/acp-14-9707-2014, 2014.
- Logan, J. A.: Tropospheric Ozone: Seasonal behaviour, Trends, and Anthropogenic Influence, *J. Geophys. Res.*, 90, 10463–10482, 1985.
- Logan, J. A., Jones, D. B. A., Megretskaia, I. A., Oltmans, S. J., Johnson, B. J., Vömel, H., Randel, W. J., Kimani, W., and Schmidlin, F. J.: Quasi-biennial oscillation in tropical ozone as revealed by ozonesonde and satellite data, *J. Geophys. Res.*, 108, 4244, doi:10.1029/2002JD002170, 2003.
- Logan, J. A., Staehelin, J., Megretskaia, I. A., Cammas, J.-P., Thouret, V., Claude, H., De Backer, H., Steinbacher, M., Scheel, H.-E., Stübi, R., Fröhlich, M., and Derwent R.: Changes in ozone over Europe: Analysis of ozone measurements from sondes, regular aircraft (MOZAIC) and alpine surface sites, *J. Geophys. Res.*, 117, D09301, doi:10.1029/2011JD016952, 2012.
- Mäder, J. A., Staehelin, J., Brunner, D., Stahel, W. A., Wohltmann, I., and Peter, T.: Statistical modelling of total ozone: Selection of appropriate explanatory variables, *J. Geophys. Res.*, 112, D11108, doi:10.1029/2006JD007694, 2007.
- Mäder, J. A., Staehelin, J., Peter, T., Brunner, D., Rieder, H. E., and Stahel, W. A.: Evidence for the effectiveness of the Montreal Protocol to protect the ozone layer, *Atmos. Chem. Phys.*, 10, 12161–12171, doi:10.5194/acp-10-12161-2010, 2010.
- Manney, G. L., Santee, M. L., Froidevaux, L., Hoppel, K., Livesey, N. J., and Waters, J. W.: EOS MLS observations of ozone loss in the 2004–2005 Arctic winter, *Geophys. Res. Lett.*, 33, L04802, doi:10.1029/2005GL024494, 2006.
- Manney, G., Santee, M. L., Rex, M., Livesey, N. J., Pitts, M. C., Veefkind, P., Nash, E. N., Wohltmann, I., Lehmann, R., Froidevaux, L., Poole, L. R., Schoeberl, M. R., Haffner, D. P., Davies, J., Dorokhov, V., Gernandt, H., Johnson, B., Kivi, R., Kyrö, E., Larsen, N., Levelt, P. F., Makshtas, A., McElroy, C. T., Nakajima, H., Parrondo, M. C., Tarasick, D. W., von der Gathen, P., Walker, K. A., and Zinoviev, N. S.: Unprecedented Arctic ozone loss in 2011, *Nature*, 478, 469–475, doi:10.1038/nature10556, 2011.
- McCormack, J. P., Siskind, D. E., and Hood, L. L.: Solar-QBO interaction and its impact on stratospheric ozone in a zonally averaged photochemical transport model of the middle atmosphere, *J. Geophys. Res.*, 112, D16109, doi:10.1029/2006JD008369, 2007.
- McLinden, C. A., Tegtmeier, S., and Fioletov, V.: Technical Note: A SAGE-corrected SBUV zonal-mean ozone data set, *Atmos. Chem. Phys.*, 9, 7963–7972, doi:10.5194/acp-9-7963-2009, 2009.
- McPeters, R. D., Labow, G. J., and Logan, J. A.: Ozone climatological profiles for satellite retrieval algorithms, *J. Geophys. Res.-Atmos.*, 112, D05308, doi:10.1029/2005JD006823, 2007.
- Miller, A. J., Cai, A., Tiao, G., Wuebbles, D. J., Flynn, L. E., Yang, S.-K., Weatherhead, E. C., Fioletov, V., Petropavlovskikh, I., Meng, X.-L., Guillas, S., Nagatani, R. M., and Reinsel, G. C.: Examination of ozonesonde data for trends and trend changes incorporating solar and Arctic oscillation signals, *J. Geophys. Res.*, 111, D13305, doi:10.1029/2005JD006684, 2006.
- Nair, P. J., Godin-Beekmann, S., Kuttippurath, J., Ancellet, G., Goutail, F., Pazmiño, A., Froidevaux, L., Zawodny, J. M., Evans, R. D., Wang, H. J., Anderson, J., and Pastel, M.: Ozone trends derived from the total column and vertical profiles at a northern mid-latitude station, *Atmos. Chem. Phys.*, 13, 10373–10384, doi:10.5194/acp-13-10373-2013, 2013.
- Neu, J. L., Flury, T., Manney, G. L., Santee, M. L., Livesey, N. J., and Worden, J.: Tropospheric ozone variations governed by changes in stratospheric circulation, *Nat. Geosci.*, 7, 340–344, doi:10.1038/ngeo2138, 2014.

- Newchurch, M. J., Yang, E.-S., Cunnold, D. M., Reinsel, G. C., and Zawodny, J. M.: Evidence for slowdown in stratospheric ozone loss: first stage of ozone recovery, *J. Geophys. Res.-Atmos.*, 108, 4507, doi:10.1029/2003JD003471, 2003.
- Oetjen, H., Payne, V. H., Kulawik, S. S., Eldering, A., Worden, J., Edwards, D. P., Francis, G. L., Worden, H. M., Clerbaux, C., Hadji-Lazaro, J., and Hurtmans, D.: Extending the satellite data record of tropospheric ozone profiles from Aura-TES to MetOp-IASI: characterisation of optimal estimation retrievals, *Atmos. Meas. Tech.*, 7, 4223–4236, doi:10.5194/amt-7-4223-2014, 2014.
- Oltmans, S. J., Lefohn, A. S., Scheel, H. E., Harris, J. M., Levy II, H., Galbally, I. E., Brunke, E., Meyer, C. P., Lathrop, J. A., Johnson, B. J., Shadwick, D. S., Cuevas, E., Schmidlin, F. J., Tarasick, D. W., Claude, H., Kerr, J. B., Uchino, O., and Mohner, V.: Trends of ozone in the troposphere, *Geophys. Res. Lett.*, 25, 139–142, doi:10.1029/97GL03505, 1998.
- Oltmans, S. J., Lefohn, A. S., Harris, J. M., Galbally, I., Scheel, H. E., Bodeker, G., Brunke, E., Claude, H., Tarasick, D., Johnson, B. J., Simmonds, P., Shadwick, D., Anlauf, K., Hayden, K., Schmidlin, F., Fujimoto, T., Akagi, K., Meyer, C., Nichol, S., Davies, J., Redondas, A., and Cuevas, E.: Long-term changes in tropospheric ozone, *Atmos. Environ.*, 40, 3156–3173, 2006.
- Parrington, M., Palmer, P. I., Henze, D. K., Tarasick, D. W., Hyer, E. J., Owen, R. C., Helmig, D., Clerbaux, C., Bowman, K. W., Deeter, M. N., Barratt, E. M., Coheur, P.-F., Hurtmans, D., Jiang, Z., George, M., and Worden, J. R.: The influence of boreal biomass burning emissions on the distribution of tropospheric ozone over North America and the North Atlantic during 2010, *Atmos. Chem. Phys.*, 12, 2077–2098, doi:10.5194/acp-12-2077-2012, 2012.
- Parrish, D. D., Law, K. S., Staehelin, J., Derwent, R., Cooper, O. R., Tanimoto, H., Volz-Thomas, A., Gilge, S., Scheel, H.-E., Steinbacher, M., and Chan, E.: Long-term changes in lower tropospheric baseline ozone concentrations at northern mid-latitudes, *Atmos. Chem. Phys.*, 12, 11485–11504, doi:10.5194/acp-12-11485-2012, 2012.
- Pitts, M. C., Poole, L. R., Dörnbrack, A., and Thomason, L. W.: The 2009–2010 Arctic polar stratospheric cloud season: a CALIPSO perspective, *Atmos. Chem. Phys.*, 11, 2161–2177, doi:10.5194/acp-11-2161-2011, 2011.
- Pommier, M., Clerbaux, C., Law, K. S., Ancellet, G., Bernath, P., Coheur, P.-F., Hadji-Lazaro, J., Hurtmans, D., Nédélec, P., Paris, J.-D., Ravetta, F., Ryerson, T. B., Schlager, H., and Weinheimer, A. J.: Analysis of IASI tropospheric O₃ data over the Arctic during POLARCAT campaigns in 2008, *Atmos. Chem. Phys.*, 12, 7371–7389, doi:10.5194/acp-12-7371-2012, 2012.
- Randel, W. J. and Wu, F.: Isolation of the ozone QBO in SAGE II data by singular-value decomposition, *J. Atmos. Sci.*, 53, 2546–2559, 1996.
- Randel, W. J. and Wu, F.: A stratospheric ozone profile data set for 1979–2005: Variability, trends, and comparisons with column ozone data, *J. Geophys. Res.-Atmos.*, 112, D06313, doi:10.1029/2006JD007339, 2007.
- Reinsel, G. C., Weatherhead, E. C., Tiao, G. C., Miller, A. J., Nagatani, R. M., Wuebbles, D. J., and Flynn, L. E.: On detection of turnaround and recovery in trend for ozone, *J. Geophys. Res.*, 107, ACH 1-1–ACH 1-12, doi:10.1029/2001JD000500, 2002.
- Reinsel, G. C., Miller, A. J., Weatherhead, E. C., Flynn, L. E., Nagatani, R. M., Tiao, G. C., and Wuebbles, D. J.: Trend analysis of total ozone data for turnaround and dynamical contributions, *J. Geophys. Res.*, 110, D16306, doi:10.1029/2004JD004662, 2005.
- Rieder, H. E., Frossard, L., Ribatet, M., Staehelin, J., Maeder, J. A., Di Rocco, S., Davison, A. C., Peter, T., Weihs, P., and Holawe, F.: On the relationship between total ozone and atmospheric dynamics and chemistry at mid-latitudes – Part 2: The effects of the El Niño/Southern Oscillation, volcanic eruptions and contributions of atmospheric dynamics and chemistry to long-term total ozone changes, *Atmos. Chem. Phys.*, 13, 165–179, doi:10.5194/acp-13-165-2013, 2013.
- Rodgers, C. D.: Inverse methods for atmospheric sounding: Theory and Practice, Series on Atmospheric, Oceanic and Planetary Physics, Vol. 2, World Scientific Publishing Co., Singapore, 2000.
- Safieddine, S., Clerbaux, C., George, M., Hadji-Lazaro, J., Hurtmans, D., Coheur, P.-F., Wespes, C., Loyola, D., Valks, P., and Hao, N.: Tropospheric ozone and nitrogen dioxide measurements in urban and rural regions as seen by IASI and GOME-2, *J. Geophys. Res.*, 118, 10555–10566, 2013.
- Salby, M., Callaghan, P., Keckhut, P., Godin, S., and Guirlet, M.: Interannual changes of temperature and ozone: Relationship between the lower and upper stratosphere, *J. Geophys. Res.*, 107, 4342, doi:10.1029/2001JD000421, 2002.
- Salby, M., Titova, E., and Deschamps, L.: Rebound of Antarctic ozone, *Geophys. Res. Lett.*, 38, L09702, doi:10.1029/2011GL047266, 2011.
- Santer, B. D., Wigley, T. M. L., Boyle, J. S., Gaffen, D. J., Hnilo, J. J., Nychka, D., Parker, D. E., Parker, D. E., and Taylor, K. E.: Statistical significance of trends and trend differences in layer-average atmospheric temperature time series, *J. Geophys. Res.*, 105, 7337–7356, 2000.
- Saunois, M., Emmons, L., Lamarque, J.-F., Tilmes, S., Wespes, C., Thouret, V., and Schultz, M.: Impact of sampling frequency in the analysis of tropospheric ozone observations, *Atmos. Chem. Phys.*, 12, 6757–6773, doi:10.5194/acp-12-6757-2012, 2012.
- Scannell, C., Hurtmans, D., Boynard, A., Hadji-Lazaro, J., George, M., Delcloo, A., Tuinder, O., Coheur, P.-F., and Clerbaux, C.: Antarctic ozone hole as observed by IASI/MetOp for 2008–2010, *Atmos. Meas. Tech.*, 5, 123–139, doi:10.5194/amt-5-123-2012, 2012.
- Schneider, M., Blumenstock, T., Hase, F., Höpfner, M., Cuevas, E., Redondas, A., and Sancho, J. M.: Ozone profiles and total column amounts derived at Izana Tenerife Island, from FTIR solar absorption spectra, and its validation by an intercomparison to ECC-sonde and Brewer spectrometer measurements, *J. Quant. Spectrosc. Ra.*, 91, 245–274, doi:10.1016/j.jqsrt.2004.05.067, 2005.
- Shepherd, T. G., Plummer, D. A., Scinocca, J. F., Hegglin, M. I., Fioletov, V. E., Reader, M. C., Remsberg, E., von Clarmann, T., Wang, H. J.: Reconciliation of halogen-induced ozone loss with the total-column ozone record, *Nat. Geosci.*, 7, 443–449, doi:10.1038/ngeo2155, 2014.
- Soukharev, B. E. and Hood, L. L.: Solar cycle variation of stratospheric ozone: Multiple regression analysis of long-term satellite data sets and comparisons with models, *J. Geophys. Res.-Atmos.*, 111, D20314, doi:10.1029/2006JD007107, 2006.

- Steinbrecht, W., Claude, H., and Winkler, P.: Enhanced upper stratospheric ozone: Sign of recovery or solar cycle effect?, *J. Geophys. Res.*, 109, D02308, doi:10.1029/2003JD004284, 2004.
- Steinbrecht, W., Claude, H., Schonenborn, F., McDermid, I. S., Leblanc, T., Godin, S., Song, T., Swart, D. P. J., Meijer, Y. J., Bodeker, G. E., Connor, B. J., Kampf, N., Hocke, K., Calisesi, Y., Schneider, N., de la Noe, J., Parrish, A. D., Boyd, I. S., Brühl, C., Steil, B., Giorgetta, M. A., Manzini, E., Thomason, L. W., Zawodny, J. M., McCormick, M. P., Russell III, J. M., Bhartia, P. K., Stolarski, R. S., and Hollandsworth-Frith, S. M.: Long-term evolution of upper stratospheric ozone at selected stations of the Network for the Detection of Stratospheric Change (NDSC), *J. Geophys. Res.*, 111, D10308, doi:10.1029/2005JD006454, 2006a.
- Steinbrecht, W., Haßler, B., Brühl, C., Dameris, M., Giorgetta, M. A., Grewe, V., Manzini, E., Matthes, S., Schnadt, C., Steil, B., and Winkler, P.: Interannual variation patterns of total ozone and lower stratospheric temperature in observations and model simulations, *Atmos. Chem. Phys.*, 6, 349–374, doi:10.5194/acp-6-349-2006, 2006b.
- Steinbrecht, W., Claude, H., Schönenborn, F., McDermid, I. S., Leblanc, T., Godin-Beekmann, S., Keckhut, P., Hauchecorne, A., Van Gijsel, J. A. E., Swart, D. P. J., Bodeker, G. E., Parrish, A., Boyd, I. S., Kämpfer, N., Hocke, K., Stolarski, R. S., Frith, S. M., Thomason, L. W., Remsberg, E. E., Von Savigny, C., Rozanov, A., and Burrows, J. P.: Ozone and temperature trends in the upper stratosphere at five stations of the Network for the Detection of Atmospheric Composition Change, *Int. J. Remote Sens.*, 30, 3875–3886, doi:10.1080/01431160902821841, 2009.
- Stolarski, R. S. and Frith, S. M.: Search for evidence of trend slowdown in the long-term TOMS/SBUV total ozone data record: the importance of instrument drift uncertainty, *Atmos. Chem. Phys.*, 6, 4057–4065, doi:10.5194/acp-6-4057-2006, 2006.
- Tian, W., Chipperfield, M. P., Gray, L. J., and Zawodny, J. M.: Quasi-biennial oscillation and tracer distributions in a coupled chemistry-climate model, *J. Geophys. Res.*, 111, D20301, doi:10.1029/2005JD006871, 2006.
- Varai, A., Homonnai, V., Jánosi, I. M., and Müller, R.: Early signatures of ozone trend reversal over the Antarctic, *Earth's Future*, 3, 95–109, doi:10.1002/2014EF000270, 2015.
- Vigouroux, C., De Mazière, M., Demoulin, P., Servais, C., Hase, F., Blumenstock, T., Kramer, I., Schneider, M., Mellqvist, J., Strandberg, A., Velasco, V., Notholt, J., Sussmann, R., Stremme, W., Rockmann, A., Gardiner, T., Coleman, M., and Woods, P.: Evaluation of tropospheric and stratospheric ozone trends over Western Europe from ground-based FTIR network observations, *Atmos. Chem. Phys.*, 8, 6865–6886, doi:10.5194/acp-8-6865-2008, 2008.
- Vigouroux, C., Blumenstock, T., Coffey, M., Errera, Q., García, O., Jones, N. B., Hannigan, J. W., Hase, F., Liley, B., Mahieu, E., Mellqvist, J., Notholt, J., Palm, M., Persson, G., Schneider, M., Servais, C., Smale, D., Thölix, L., and De Mazière, M.: Trends of ozone total columns and vertical distribution from FTIR observations at eight NDACC stations around the globe, *Atmos. Chem. Phys.*, 15, 2915–2933, doi:10.5194/acp-15-2915-2015, 2015.
- Weatherhead, E. C. and Andersen, S. B.: The search for signs of recovery of the ozone layer, *Nature*, 441, 39–45, doi:10.1038/nature04746, 2006.
- Weiss, A. K., Staehelin, J., Appenzeller, C., and Harris, N. R. P.: Chemical and dynamical contributions to ozone profile trends of the Payerne (Switzerland) balloon soundings, *J. Geophys. Res.*, 106, 22685–22694, 2001.
- Wespes, C., Hurtmans, D., Clerbaux, C., Santee, M. L., Martin, R. V., and Coheur, P. F.: Global distributions of nitric acid from IASI/MetOP measurements, *Atmos. Chem. Phys.*, 9, 7949–7962, doi:10.5194/acp-9-7949-2009, 2009.
- Wespes, C., Emmons, L., Edwards, D. P., Hannigan, J., Hurtmans, D., Saunio, M., Coheur, P.-F., Clerbaux, C., Coffey, M. T., Batchelor, R. L., Lindenmaier, R., Strong, K., Weinheimer, A. J., Nowak, J. B., Ryerson, T. B., Crouse, J. D., and Wennberg, P. O.: Analysis of ozone and nitric acid in spring and summer Arctic pollution using aircraft, ground-based, satellite observations and MOZART-4 model: source attribution and partitioning, *Atmos. Chem. Phys.*, 12, 237–259, doi:10.5194/acp-12-237-2012, 2012.
- Wilson, R. C., Fleming, Z. L., Monks, P. S., Clain, G., Henne, S., Konovalov, I. B., Szopa, S., and Menut, L.: Have primary emission reduction measures reduced ozone across Europe? An analysis of European rural background ozone trends 1996–2005, *Atmos. Chem. Phys.*, 12, 437–454, doi:10.5194/acp-12-437-2012, 2012.
- WMO: Scientific Assessment of Ozone Depletion: 2002, Global Ozone Research and Monitoring Project – Report No. 47, World Meteorological Organization, Geneva, Switzerland, 2003.
- WMO: Scientific Assessment of Ozone Depletion: 2006, Global Ozone Research and Monitoring Project – Report 50, World Meteorological Organization, Geneva, Switzerland, 2007.
- WMO: Scientific Assessment of Ozone Depletion: 2010, Global Ozone Research and Monitoring Project – Report 52, World Meteorological Organization, Geneva, Switzerland, 2011.
- WMO: Scientific Assessment of Ozone Depletion: 2014, Global Ozone Research and Monitoring Project – Report 56, World Meteorological Organization, Geneva, Switzerland, 2014.
- Yang, E.-S., Cunnold, D. M., Salawitch, R. J., McCormick, M. P., Russell, J., Zawodny, J. M., Oltmans, S., and Newchurch, M. J.: Attribution of recovery in lower-stratospheric ozone, *J. Geophys. Res.*, 111, D17309, doi:10.1029/2005JD006371, 2006.