

# Mapping of a buried basement combining aeromagnetic, gravity and petrophysical data: The substratum of southwest Paris Basin, France

Julien Baptiste, Guillaume Martelet, Michel Faure, Laurent Beccaletto,

Pierre-Alexandre Reninger, José Perrin, Yan Chen

## ▶ To cite this version:

Julien Baptiste, Guillaume Martelet, Michel Faure, Laurent Beccaletto, Pierre-Alexandre Reninger, et al.. Mapping of a buried basement combining aeromagnetic, gravity and petrophysical data: The substratum of southwest Paris Basin, France. Tectonophysics, 2016, 683, pp.333-348. 10.1016/j.tecto.2016.05.049. insu-01339986

## HAL Id: insu-01339986 https://insu.hal.science/insu-01339986

Submitted on 1 Jul 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

1	Mapping of a buried basement combining aeromagnetic, gravity
2	and petrophysical data: the substratum of southwest Paris
3	Basin, France
4	
5	*Julien Baptiste <sup>1,2</sup> , Guillaume Martelet <sup>2</sup> , Michel Faure <sup>1</sup> , Laurent Beccaletto <sup>2</sup> ,
6	Pierre-Alexandre Reninger <sup>2</sup> , José Perrin <sup>2</sup> , Yan Chen <sup>1</sup> ,
7	
8	<sup>1</sup> Institut des Sciences de la Terre d'Orléans, UMR 7327, Université d'Orléans, Campus
9	Géosciences, 1A rue de la Férollerie, 45071 Orléans Cedex 2
10	<sup>2</sup> Bureau de Recherche Géologique et Minière (BRGM), UMR 7327, 3 avenue Claude
11	Guillemin, BP 36009, 45060 Orléans Cedex 2.
12	* Corresponding author: j.baptiste@brgm.fr
13	

## 15 **Abstract**

Aeromagnetic and gravity data have proven to be among the most effective methods for 16 17 mapping deeply buried basin/basement interfaces. However, the data interpretation 18 generally suffers from ambiguities, due to the non-uniqueness of the gravity and magnetic 19 signatures. Here, we tie the gravity and magnetic signatures with a petrophysical 20 characterization of the lithologies outcropping around the French Paris Basin. Our 21 methodology investigates the lithology and structure of its hidden Variscan substratum at 22 the junction between the Armorican Massif and Massif Central. Our approach is based on 23 the combination of potential field data, magnetic susceptibilities measured in the field, 24 density values of sample rocks and information documented in boreholes, in order to 25 propose a new interpretative geological map of the buried substratum of the Paris Basin. The petrophysical description is combined with geophysical patterns of the substratum, 26 27 mapped through statistical unsupervised classification of suitably selected magnetic and 28 gravity maps. The first step of interpretation consists in extending the outcropping major 29 structures below the Meso-Cenozoic sedimentary cover of the Paris Basin. The litho-30 structural units, in between these major structures, are then interpreted separately. The 31 second step consists in assigning lithologies within each unit, with respect to its 32 magnetization and density (as derived from the petrophysical compilation), and mapping its 33 extension under cover, integrating punctual borehole information. Overall, with a special 34 emphasis on relating geophysical signatures and petrophysical characteristics of litho-35 structural units, this methodology permits a precise structural and lithological cartography of 36 a segment of the buried Variscan substratum. In the southwestern part of the Paris Basin, 37 this approach reveals: i) the limited eastward extension of Central Brittany, ii) the eastward 38 extension of the major Cholet fault, iii) the emphasis on N150E-N160E striking fault and their N30E conjugates, controlling the opening of Permo-Carboniferous basins, and iv) the
eastward extension of the Eo-Variscan suture.

41 Keywords: aeromagnetic, gravity, Paris Basin, Variscan substratum, petrophysical data,
42 undercover mapping

## 43 1. Introduction

The European Variscan belt belongs to a several thousand-km-long Late Paleozoic orogen 44 45 extending from the Appalachians in the eastern North America, to Polish Sudetes, through the Mauritanides in West Africa. This belt, formed by Proterozoic to Carboniferous rocks, 46 47 constitutes the Pre-Mesozoic basement of a large part of Western and Central Europe. In 48 France, the Variscan belt presently crops out in several massifs, namely: Massif Central, 49 Armorican, Ardennes, and Vosges Massifs and in the basement of the Cenozoic Alpine and 50 Pyrenean belts. The continuity between these massifs is hidden by several Mesozoic to 51 Cenozoic sedimentary basins, such as the Paris or Aguitaine Basin (Fig. 1)

52 The Paris Basin is an intraplate sedimentary basin, set up on the Variscan substratum that 53 crops out in the above massifs (Pomerol, 1978; Mégnien; 1980, Perrodon and Zabek, 1990; 54 Guillocheau et al., 2000; Chantraine et al., 2003) (Fig. 1). It is well known that lithologies 55 and structures of the southern part of the Armorican Massif and the Massif Central are 56 closely related (Autran and Lameyre, 1980; Matte and Hirn, 1988; Virlogeux et al., 1999; Faure et al., 2005; Cartannaz et al., 2006; Gébelin et al., 2007; Ballèvre et al., 2009; Rolin 57 58 et al., 2009); however, their connection is still poorly known, as it hidden by the Meso-59 Cenozoic Paris Basin sedimentary cover. Both massifs are composed of several lithotectonic units separated by crustal-scale shear zones, such as the North Armorican Shear 60 Zone (NASZ) and the South Armorican Shear Zone (SASZ), in the Armorican Massif or the 61 62 Marche Fault in the Massif Central (Fig. 1). In addition, the Nort-sur-Erdre fault of the

63 Armorican Massif (NSE F.; Fig. 1) is acknowledged as the Eo-Variscan suture (Fig. 1) 64 (Matte, 1986; Le Corre et al., 1991; Ballèvre et al., 1992; Lardeux and Cavet, 1994; Faure 65 et al., 1997; Cartier et al., 2001; Bitri et al., 2003). The eastward extension of the Armorican 66 litho-structural units has already been integrated in large-scale geodynamic reconstructions of the Variscan belt, either based on geological evidence (e.g. Matte, 1986; Faure et al., 67 68 2005; Ballèvre et al., 2009; Martínez-Catalán, 2012, Edel et al., 2015) or on low to medium-69 resolution regional geophysics (e.g. Edel, 2008). There is no agreement on the limits and 70 nature of the units, and geodynamic significance is still under debate. A way to ascertain 71 these models would be to fill the geological observation gap caused by the Meso-Cenozoic 72 sedimentary cover of the Paris Basin by new high resolution data.

In the second half of the 20<sup>th</sup> century, the Paris Basin substratum started being investigated 73 using gravity data (Goguel and Francia, 1954), deep boreholes (Lienhardt, 1961) or deep 74 75 seismic profiles (Matte and Hirn, 1988). A combination of gravity and intermediate to low 76 resolution aeromagnetic data was used in the southwestern part of the Paris Basin 77 (Weber, 1973). The extension of this last study to the entire Paris Basin led to the first 78 version of a pseudo-lithological and structural sketch of the pre-Mesozoic substratum of the 79 Paris Basin (Debeglia and Weber, 1980). More recently, on the basis of the combination of 80 new high resolution magnetic data and updated gravity data, Martelet et al., (2013) 81 proposed a method to characterize the substratum geometry of the central-south part of the 82 Paris Basin, but no detailed lithological interpreted map was produced yet.

In addition, a map of the buried substratum around the Poitou High was proposed on the base of detailed structural analysis and drilling information (Rolin and Colchen, 2001). Because we now benefit from recent high resolution aeromagnetic data on the entire southwestern part of the Paris Basin, we are now able to address the litho-structural pattern of its buried substratum with an unprecedented resolution. Complemented by a

characterization of the field petrophysical properties of the various litho-structural units, our work ascertain cartographic interpretations of the geophysical signatures. Our methodology emphasizes the processing of interpretation of potential field data to derive a highresolution structural map. Combining the latter with field petrophysical properties, lithologies were then interpreted in each litho-structural unit, leading to a new geological map of the Paris Basin substratum.

94 The extensions of the litho-structural units as well as the lithologies of the Variscan and 95 their regional geological implications are further discussed.

96

## 97 2. Geological setting

The Paris Basin is a low subsidence Meso-Cenozoic sedimentary basin. It is composed of silico-clastic and calcareous rocks (Pomerol, 1978; Mégnien, 1980; Perrodon and Zabek, 100 1990; Guillocheau et *al.*, 2000; Beccaletto et *al.*, 2011). It is set up on a Variscan substratum including Permo-Carboniferous basins. In the study area, the formations 102 constituting the Variscan substratum of the Paris Basin laterally outcrop in the Armorican 103 Massif to the west and the Massif Central, to the south, respectively (Fig. 2a).

104 The Armorican Massif is composed of several litho-structural units separated by crustal-105 scale faults and characterized by distinct lithologies and tectonic evolutions (Fig. 2a). The 106 North Armorican Shear Zone (NASZ; Chauris, 1969; Watts and Williams, 1979), the 107 northern and the southern branches of the South Armorican Shear Zone (NBSASZ and 108 SBSASZ, respectively) (Jégouzo, 1980) and the Nort-sur-Erdre fault (NSE) are the main 109 Variscan structures of the central and southeastern part of the Armorican Massif (Fig. 2a). 110 The NASZ and the SBSASZ delimit the Central Brittany to the north and south, respectively 111 (Fig. 2a). Central Brittany consists of folded and weakly metamorphosed Neoproterozoic sediments, unconformably overlain by weakly deformed Paleozoic sediments, intruded by Carboniferous granites (Fig. 2b) (Vernhet et *al.*, 2009). Located along the NASZ, the southerly early Carboniferous Laval basin is superimposed on the Neoproterozoic to Paleozoic series (Fig. 2b). It is composed of early Carboniferous sedimentary rocks interbedded with acidic and basic volcanic rocks (Le Hérissé and Plaine, 1982). The Laval basin was folded during the Late Carboniferous (Houlgatte *et al.*, 1988).

118 Between the NBSASZ and the NSE fault, the Paleozoic series of the St-Georges-sur-Loire 119 unit overthrusts to the NW the Lanvaux unit (Fig. 2a). The NW-SE striking Lanvaux unit is composed of Neoproterozoic and early Cambrian metasediments overlain by Paleozoic 120 121 weakly metamorphosed sediments (Lardeux and Cavet, 1994), and intruded by an early 122 Ordovician granite deformed into an orthogneiss (Fig. 2b), called the St-Clément-de-la-123 Place (Vidal, 1980). This orthogneiss is intruded by two granitic plutons: i) the Bécon 124 granite (Chauris and Lucas, 1964; Cavet et al., 1970, 1976) and ii) the St-Lambert 125 granodiorite showing S/C structures, emplaced during to the Carboniferous dextral shearing 126 of the NBSASZ (Faure and Cartier, 1998). The Lanvaux unit experienced a polyphase 127 deformation (Faure and Cartier, 1998). The foliation attitude documents an antiform, the 128 core of which is constituted by the orthogneiss. Outcropping at the junction between the 129 Lanvaux and St-Georges-sur Loire unit, the Questembert leucogranite, emplaced during the 130 late Carboniferous (Tartèse et al., 2011a, 2011b) is a syn-tectonic granite related to the 131 SASZ shearing (Berthé et al., 1979; Bernard-Griffiths et al., 1985).

To the south, the St-George-sur-Loire unit is divided into two sub-units (Fig. 2a): the blocky sub-unit in the south overthrusts to the NW the northerly Sandstone-Pelite sub-unit (Cartier et *al.*, 2001; Cartier and Faure, 2004).

The south of Nort-sur-Erdre fault is a metamorphic nappe stack: described in the southern
part of the Armorican Massif (Burg, 1981; Matte, 1991; Bosse et *al.*, 2000; Le Hébel et *al.*,

137 2002) and in the northern part of the Massif Central (Quenardel and Rolin, 1984; Faure et 138 al., 1990). The structure and the lithology of the metamorphic Champtoceaux Complex in 139 the Armorican Massif and of the Aigurande Plateau in the Massif Central are guite similar 140 (Fig. 2b) (e.g. Faure et al., 2005; Ballèvre et al., 2009). However the Armorican Massif 141 exposes peculiar units absent in the Massif Central. The uppermost unit of the nappe stack, 142 called the Mauges nappe (Fig. 2a), consists of Neoproterozoic metagrauwackes 143 interbedded with meta-volcanics (Wyns and Le Métour, 1983; Cabanis and Wyns, 1986; 144 Wyns et al., 1998), and unconformably overlain by Cambrian sedimentary rocks and felsic 145 volcanites, and Ordovician sandstones (Fig. 2b). This Paleozoic sedimentary and volcanic 146 series is widely exposed in the Choletais area (Cavet et al., 1966). The Cambrian Thouars 147 microgranitic massif associated with basic rocks (gabbros, quartz diorites) intrudes within 148 the Neoproterozoic micaschists, the volcanic series and the dyke complex (Fig. 2b; 149 Thiéblemont et al., 1987, 2001). To the north of the Mauges nappe, the Ancenis basin is 150 located along the NSE fault (Fig. 2a). It is made of Devonian to early Carboniferous 151 deposits (Fig. 2b) (Cavet et al., 1971; Ballèvre and Lardeux., 2005), superimposing the 152 Neoproterozoic micaschists (Fig. 2b).

153 Another unit, exposed uniquely in the Armorican Massif is the "Drain Unit" consisting of 154 serpentinite, gabbro, basalt and siliceous sedimentary rocks, interpreted as dismembered 155 ophiolites along the Eo-variscan suture (Marchand, 1981; Ballèvre et al., 1994; Faure et al., 156 2008). The underlying unit, called the Champtoceaux complex, is an imbrication of crustal-157 scale thrust sheets characterized by highly deformed and metamorphosed gneiss and 158 eclogites (Marchand, 1981; Ballèvre et al., 1989, 1994; Bosse et al., 2000). Lastly, the 159 lowermost unit exposed in this area consists of a micaschist and paragneiss suite, named 160 the Mauves-sur-Loire series (Fig. 2b). The entire stack of nappes from the Mauges nappe 161 to the Mauves-sur-Loire series is folded in a km-scale antiform with a steeply eastward

plunging axis probably related to the dextral shearing of the SASZ (Martelet et *al.*, 2004).
Carboniferous plutons occupy the core of this antiform (e.g. Wyns et *al.*, 1998).

164 The Aigurande Plateau also consists of a stack of metamorphic nappes, refolded as 165 an ENE-SSW striking antiform, and intruded by several two-mica granitic plutons 166 (Quenardel and Rolin, 1984; Faure et al., 1990). From top to bottom, the litho-tectonic units 167 are characterized by: i) the Upper Gneiss Unit (UGU) composed of a bimodal magmatic 168 series, named leptynite-amphibolite complex with eclogites, and migmatites; ii) the Lower 169 Gneiss Unit (LGU) made of metagrauwackes, micaschists, metarhyolites and amphibolites 170 that never experienced a HP metamorphism; iii) the Para-autochthonous Unit composed of 171 low-grade micaschists (Fig. 2b). The left-lateral Marche fault, main Variscan structure of the 172 northern part of the Massif Central (e.g. Quenardel and Rolin, 1984), is the southern 173 boundary of the Aigurande Plateau (Fig. 2a). A reasonable correlation between the 174 Champtoceaux Complex and the Mauves-sur-Loire Unit in the Armorican Massif and the 175 UGU and LGU in the Massif Central, respectively has been proposed (e.g. Faure et al., 176 2005).

177 Previous structural and lithological analyses within the Poitou High (Rolin and 178 Colchen, 2001) documented the close connection between the Haut Bocage unit and the 179 Confolentais area. In the Poitou High, the substratum that crops out in rare valleys is made 180 of Carboniferous granite. Below the sedimentary cover of the Poitou High, the SBSASZ 181 eastward extension splits into several branches which separate structures and lithologies of 182 the Haut Bocage and Confolentais units (Fig. 2a) (Rolin and Colchen, 2001). The Haut 183 Bocage unit is composed of anatectic gneiss and Neoproterozoic micaschists (Fig. 2b). The 184 Confolentais unit is made of granites, metavolcanites and metasediments belonging to the 185 UGU (Fig. 2b). Both units are intruded by late Devonian gabbro, granodiorites and diorites 186 that also crop out along the left-lateral Marche fault (Peiffer, 1986; Cuney et al., 1993; Pin

and Paquette, 2002). In the Confolentais area can be found the westernmost expression of
the Limousin tonalite belt (Bernard-Griffiths et *al.*, 1985; Peiffer, 1986).

189

## 190 3. Aeromagnetic and gravity data

## 191 3.1. Processing of aeromagnetic data

192 Magnetic measurements monitor the spatial variations of magnetic properties of the 193 underground, from the surface of the Earth down to several kilometers. The French 194 Geological Survey (BRGM) conducted a fixed-wing magnetic survey in 1998 over Brittany 195 (Bonijoly et al., 1999; Truffert et al., 2001) and from August 2008 to October 2010 over the 196 Pays de la Loire (PaL) and Région Centre (Martelet et al., 2013) covering the Armorican 197 Massif and the southwestern part of the Paris Basin (Fig. 3a). These surveys were flown at 198 an average 85 m and 120 m ground clearance for Région Centre, PaL and Brittany, 199 respectively. For the three surveys, the flight path was oriented N-S, with a line-spacing of 200 500 m reduced to 250 m over key areas for the Brittany survey and 1 km reduced to 500 m 201 for PaL and Région Centre surveys; perpendicular control tie lines were also flown every 202 10 km. The surveys overlap with each other on a 3 to 5 km band at their periphery. In order 203 to get rid of punctual artifacts related to human activities, the data were upward continued 204 to an elevation of 600 m. This removed the short wavelength cultural noise without 205 significantly smoothing the data, with regards to the aims and regional extent of the study. 206 Magnetic anomaly 250 m regular grids were produced using the minimum curvature 207 gridding method (Taylor and Mason, 1972), separately, for each survey.

In addition three grid transforms were applied to emphasize various properties of the buriedsubstratum:

The reduction to the pole (RTP; Fig. 3b): it contributes to simplify the magnetic signal
 interpretation (Blakely, 1996). Taking into account the Earth field direction, this

operator relocates the magnetic anomaly on top of its causative body. When induced
 magnetization predominates, anomaly bipolarities are removed, so that RTP positive
 anomalies indicate a local increase of magnetic susceptibility at depth.

The vertical derivative (Fig. 3c): it enhances local signal gradients while regional trends are removed. Magnetization in the Meso-Cenozoic Paris Basin is weak and sedimentary cover thickness vary smoothly over large distances; therefore, they generate long wavelength components in the magnetic map, which are removed by the vertical gradient operator. Consequently the map of the magnetic vertical gradient highlights the magnetic contrasts within the underlying substratum (e.g. Weber, 1973; .Martelet et *al.*, 2013).

The Tilt derivative (TILT) or Tilt angle processing (e.g. Miller and Singh, 1994;
 Verduzco et *al.*, 2004) is powerful for structural interpretation as it depicts equally the
 edge of deep and shallow magnetic sources (Miller and Singh, 1994). It is an
 effective method for mapping contacts or faults, weakly contrasted magnetized
 bodies, located under a sedimentary cover (Fairhead et *al.*, 2011). The TILT map
 was used to highlight the structures of the underlying substratum and the magnetic
 body contours which are defined by the zero value in the TILT map (Fig. 3d).

The vertical derivative and the tilt derivative operators were applied on the anomaly reduced to the pole (RTP). In order to avoid artifacts at survey junctions, the magnetic transforms were applied separately, survey by survey, before merging. The merge of the three magnetic surveys was carefully achieved, using a standard grid stitching algorithm.

233

## 234 3.2. Processing of gravity data

235 Gravity data derive from the compilation of ground gravity surveys conducted in France 236 since the middle of the 20th century and compiled in the Banque Gravimétrique de la 237 France (BGF) (Martelet, et al., 2009). In the study area, the average station coverage is 238 about 1 station/km<sup>2</sup>. Data are tied to the CGF65 base station network. In order to derive the 239 Bouguer anomaly, all standard corrections are included, with a reference density of 240 2600 kg/m3, and terrain corrections computed to a distance of 167 km (Martelet et al., 241 2002). Taking into account the accuracy of 1) the network, 2) the gravity measurements 242 and their positioning and 3) the terrain corrections, the RMS error on the Bouquer anomaly 243 is 0.32 mGal in the study area. The Bouguer anomaly map presented in Fig. 4a locates the 244 main regional density contrasts, from the surface down to several kilometers at depth.

245 A map of the vertical derivative of the Bouguer anomaly is presented in Fig. 4b. The first 246 order of the vertical gradient of the Bouguer anomaly has long been used for separating 247 close structures (Elkins, 1951; Gérard and Griveau, 1972; Goguel, 1972). Here, this 248 operator is used to highlight density contrasts within the Paris Basin substratum. As for the 249 magnetic map, the regional effect of the Meso-Cenozoic sedimentary pile results in smooth 250 and long wavelength signals and is therefore strongly attenuated by this operator (see 251 Martelet et al., 2013 for more details). Positive and negative signals of the vertical gradient 252 feature relatively high and low density rocks at depth, respectively.

253

## **4. Measurements of field rock properties**

Gravity and magnetic map usually display "averaged" geophysical signatures of geological bodies as compared to the lithological variations at the outcrop scale; the discrimination of the magnetic and gravity causative lithologies suffering from ambiguities. An extensive campaign of petrophysical sampling and measurements was conducted in order to take into account this scale effect in our geophysical maps interpretation. It was designed to derive
 reliable petrophysical "statistic signatures" for the main lithologies encountered in the study
 area.

262

#### 263 4.1 Magnetic susceptibility

264 The magnetic susceptibility of a rock refers to its ability to become magnetized by an 265 external magnetic field such as the Earth's field (e.g. Dearing, 1999). Rocks have various 266 magnetic responses due to their magnetic properties, which to the first order, depend on the volume content of magnetite (Clark and Emerson, 1991). Magnetic susceptibility field 267 268 measurements were carried out using a hand-held kappameter (KT-9, Exploranium, 269 Canada). The measuring range of KT-9 susceptibility meters is from -999 to 999 x 10–3 SI 270 units with a sensitivity of 1 x 10-5 SI. Because reproducibility of measures is influenced by 271 the irregularity of the rock surface (Lecoanet et al., 1999), measures were achieved in the 272 "Pin-mode" of the kappameter which takes into account a geometric factor to reduce 273 roughness effects.

274 About 4050 magnetic susceptibility measurements were taken directly on 130 outcrops, all 275 along the southeastern border of the Armorican Massif and the northwestern border of the 276 Massif Central and Poitou High (Fig. 3a). Within each litho-structural unit, most of the 277 outcropping lithologies were sampled. The compilation of these measurements is presented 278 in Fig. 5. For each litho-structural unit, we numerated the magnetic susceptibility 279 measurements within constant intervals of variation. In Fig. 5, the colored bars highlight the 280 most represented magnetic susceptibility ranges, whereas the grey intervals indicate 281 ranges with few randomly distributed data.

282 As an aid for the interpretation, the magnetic susceptibility was subdivided into three 283 representative ranges, based on the classification of Clark and Emerson (1991) and the 284 separation range of Théveniaut and Clarke, (2013): i) from negative values to 4 x 10-4 SI, ii) 285 from 4 x 10-4 to 5 x 10-3 SI and iii) above 5 x 10-3 SI, designated as low, intermediate and 286 high magnetic susceptibility ranges, respectively. Fe-rich sandstones in Central Brittany 287 exhibit the highest magnetic susceptibility of the study area (Fig. 5). Basic rocks (gabbro-288 diorite, granodiorite and amphibolite), and Cambrian felsic volcanites are also within the 289 high magnetic susceptibility range, due to their high amount of ferrimagnetic minerals, such 290 as magnetite, in their mineralogical composition (Thiéblemont et al., 2011). Granite, 291 leptynite, orthogneiss, migmatite, Neoproterozoic metasedimentary and Paleozoic 292 sedimentary rocks, micaschists and metasediments are within the low magnetic 293 susceptibility range, given that they are mainly composed of diamagnetic (quartz, 294 plagioclase) and paramagnetic minerals. Some Paleozoic sedimentary rocks, basalts, 295 metabasites, meta-gabbro, diorites and amphibolites are within the intermediate range, 296 since they contain paramagnetic minerals, and a small amount of iron-bearing minerals 297 (amphibole, biotite or clay minerals).

298 Induced magnetization is predominant in the area, but several bimodal anomalies in the 299 RTP map (Fig. 3b) indicate some remnantly magnetized rocks. To the north, the Fe-rich 300 Ordovician sandstones of Central Brittany, mainly composed of magnetite, have been studied in detail (Corpel and Weber, 1970): their Koenigsberger ratio is around 6 but the 301 302 remnant component of magnetization is almost collinear to the ambient magnetic field and 303 therefore does not strongly affect the mapping of this unit. Also, the granodiorite within the 304 Choletais area and the diorite plutons within the Confolentais area are partly remnantly 305 magnetized, but their magnetic anomalies almost perfectly match their field cartographic 306 limits (Fig. 3b). This suggests that i) the effect of the remnant magnetization is weak, or ii) their direction of remnant magnetization is close to the induced magnetization. In these three cases, the location of the magnetic causative bodies is only slightly affected by the remnant magnetization and these lithologies also display a high magnetic susceptibility in the field (Fig. 5). Consequently, we made the assumption that the interpretation of the magnetic maps could be achieved considering the magnetic susceptibility only.

312 Globally, there is a significant overlap between the magnetic susceptibility ranges of the 313 various lithologies; however, this overlap is rather limited between lithologies within each 314 litho-structural unit. This observation is crucial for the geophysical maps interpretation, as 315 described in the following paragraphs.

## 316 4.2 Density

Density is the petrophysical property influencing the gravity data. For this study, 54 unweathered rock samples from most of the lithologies were collected all along the Armorican and Massif Central borders in 48 outcrops (Fig. 3a). The densities of these samples were measured using the double weighting method, with a ca. 0.01 g/cm<sup>3</sup> uncertainty. The density determination of some lithologies was not possible due to bad outcropping conditions. In this case, a density value was affected with respect to the density average of the same lithology in the other litho-structural units.

As for the magnetic susceptibilities, the densities were subdivided into three representative groups: i) from 2.55 to 2.65 g/cm<sup>3</sup>, ii) from 2.65 to 2.8 g/cm<sup>3</sup>, and iii) above 2.8 g/cm<sup>3</sup> for low, intermediate and high densities, respectively (e.g. Edel, 2008) (Fig. 5).

327

## **5. Geophysical signatures of the Paris Basin substratum**

329 The previous studies investigating the substratum of the Paris Basin using magnetic and 330 gravity data manually outlined the main geophysical anomalies. They were interpreted with 331 simplified lithological attributions, based on the substratum nature documented in some 332 boreholes as well as some rock property data (Weber, 1973; Debeglia and Weber, 1980). 333 More recently, Martelet et al., (2013) proposed a map of petrophysical signatures of the 334 substratum of the south-central part of the Paris Basin, based on a numerical classification 335 combining gravity and magnetic data. We used the same approach to achieve a simplified 336 magnetic-gravity signature of the substratum of the study area.

337 We agree with previous studies that considered the magnetic effect of the Meso-Cenozoic 338 sedimentary cover of the Paris Basin, almost "transparent" for the magnetic field 339 (Weber, 1973); therefore the magnetic map mostly features the buried substratum. 340 Nevertheless the Meso-Cenozoic sedimentary pile at least attenuates the intensity of the 341 magnetic response of the substratum and increases the wavelength of the substratum 342 anomalies, as the sedimentary cover gets thicker; from 0 to about 2000m in the study area. 343 The gravity field contains both the effects of the substratum and of the sedimentary basin. 344 The vertical gradient of the Bouguer anomaly used for our classification attenuates the long 345 wavelengths of the Meso-Cenozoic sedimentary pile (Debeglia and Weber 1985; Martelet 346 et al., 2013). Aiming at the same goals as Martelet et al., (2013), the map of the magnetic 347 anomaly reduced to the pole that was introduced in our classification, followed two 348 considerations: 1) displaying information of the structure and magnetization of the 349 substratum as detailed as possible; 2) being physically as homogeneous as possible with 350 the gravity first vertical derivative. We added a third consideration: 3) reducing as much as 351 possible the variable smoothing and attenuating effect of the Meso-Cenozoic sedimentary 352 pile. Combining the characteristics of RTP and the TILT fulfils the three conditions. These 353 two magnetic maps were combined with the vertical gradient of the Bouguer anomaly to 354 obtain synthetized signatures of magnetic and gravity data. The three layers were 355 combined into a ternary image, using a standard image fusion procedure. We then 356 performed a numerical classification of this ternary image using unsupervised isodata 357 clustering (e.g. Venkateswarlu and Raju, 1992). Based on their gravity and magnetic 358 signatures, all pixels of the map were statistically distributed among 6 classes (Fig. 6a) 359 which well figure the geophysical signatures of the outcropping geology. This map displays 360 self-consistent cartographic bodies, which are compatible with known geological patterns. It 361 features 2 levels of magnetic intensity (from light to dark green) and an intermediate 362 average magnetic/gravity signature (in white) as well as 3 levels of gravity intensity (from 363 light to dark blue). These synthetized geophysical signatures can be related to the 364 simplified 3-levels categorization of the petrophysical magnetization/density parameters 365 (Fig. 5). This map combines magnetic and gravity signatures and it is used as a support for 366 the following structural and lithological interpretations.

367

## 368 6. Geological map of the pre-Mesozoic substratum

369 The first step of the interpretation consists in extending below the Paris Basin sedimentary cover the major structures recognized in the field (Fig. 2a) in order to delineate the Variscan 370 371 litho-structural units under cover. The structural interpretation (Fig. 6b) uses all geophysical 372 enhanced maps presented in Section 3 supported by the synthetized geophysical 373 signatures map (Fig. 6a). The second step consists in interpreting the lithological nature of 374 the hidden substratum, using the combined petrophysical characteristics of rocks (Fig. 5), 375 the synthetized geophysical signatures of the substratum (Fig. 6a) and the structural sketch 376 map including available boreholes (Fig. 6b).

#### 377 6.1 Undercover delineation of structural features

Manually interpreted geophysical trends deriving from the magnetic and gravity maps are outlined in red in Fig. 6a. They underline the N110E-N120E and N90E striking structural directions of the Armorican Massif and the Massif Central, respectively, known in the field. In addition to these trends, main geophysical structures and discontinuities were interpreted. The interpretative structural map (Fig. 6b) showing the extension of the structural units below the Paris Basin cover is discussed from north to south.

384 Central Brittany is limited to the north by the NASZ. Its extension constitutes the northern 385 limit of the study area; it is defined regionally by a major N110°E-oriented disharmony 386 between the strong magnetic signals of Central Brittany to the south, and the northern weak 387 magnetic signals (Fig. 3b). It also outlined by a successive E-W striking chaplet of magnetic 388 anomalies bounded the NASZ to the north, which can be outlined from the field to the 389 easternmost part of the studied area (Baptiste et al., 2015) (Fig. 3c, Fig. 3d). The Lanvaux 390 unit is marked by a well-defined NW-SE striking elongated low density anomaly, known as 391 Lanvaux orthogneiss (Fig. 4b); it is bounded to the north by the NBSASZ. Following the 392 magnetic and gravity trends under cover (Fig. 6a), the strike of the NBSASZ changes 393 eastwards from NW-SE to NE-SW and joins the NASZ, limiting Central Brittany to the east 394 (Fig. 6b). The southern border of the noticeable Lanvaux low gravity anomaly defines the 395 limit between the Lanvaux unit and the northern part of St-Georges-sur-Loire unit (Fig. 6b). 396 In the field, the St-Georges-sur-Loire unit is characterized by low magnetic (Fig. 3b) and 397 gravity signal (Fig. 4b), and by scattered moderate intensity magnetic anomalies (Fig. 3c) 398 and a high intensity gravity anomaly, to the north and south, respectively. Predominant NW-399 SE/E-W striking geophysical trends are also observed (Fig. 6b). Under the Paris Basin

401 data. The southern border of the high intensity gravity anomaly defines the southern limit of

sedimentary cover, the northern and southern part cannot be separated by geophysical

400

402 the St-Georges-sur-Loire unit (Fig. 4b); it defines the cartographic trace of the NSE
403 Eo-Variscan suture (Fig. 6b).

404 The Choletais area, marked by an E-W striking high gravity elongated anomaly, extending 405 150km eastwards below the Paris Basin sedimentary cover (Fig. 4b), is bounded to the 406 south, by the northern branch of the Cholet fault (Fig. 6b). This noticeable anomaly defines 407 the southward boundary of the Mauges nappe. Along this fault, the magnetic and gravity 408 trends strike E-W in continuity along more than 150 km (Fig. 6a). Northwards, the 409 geophysical trends become less and less continuous (Fig. 6a), suggesting the decreasing 410 gradient of deformation away from the fault, as observed in the Armorican Massif 411 (Thiéblemont et al., 2011). To the east (around 2°E), the E-W striking Cholet fault marks the 412 northern limit of NW-SE geophysical trends of the Aigurande Plateau. Altogether, these 413 features suggest that the Cholet fault can be considered as a major shear zone.

In the east of the study area (around 2°E), the NSE fault and the northern branch of the Cholet fault almost meet, closing the Mauges nappe and the Choletais area, to the east (Fig. 6a). To the south of the Cholet fault, low gravity anomalies (Fig. 4b) and signatures (Fig. 6a) and NW-SE striking geophysical trends (Fig. 6a) are predominant; this refers to the Haut Bocage unit and Confolentais area connection. To the east of the Haut Bocage unit, the direction of geophysical trends progressively changes from NW-SE to E-W, featuring the connection between the Haut Bocage unit and Aigurande Plateau.

In the easternmost part of the studied area, all these units are interrupted by NE-SW
striking structures bounding the Permo-Carboniferous Contres basin, well characterized on
seismic profiles and in deep boreholes (Fig. 6b; Beccaletto et *al.*, 2015).

424 At the regional scale, all structures are offset by N150E-N160E-striking faults and, to a 425 lesser extent, by their conjugate N20E-N30E striking faults (Fig. 6b). In agreement with 426 Martelet et *al.*, (2013), two kinematics are interpreted along the N150E-N160E faults: i) a

427 dextral motion highlighted by the offset of preexisting structures and lithological markers, 428 and ii) a vertical movement documented by the attenuation and the spreading of the 429 magnetic signal from west to east. In the southeastern part of the Armorican Massif, the 430 N150E striking Partenay fault is described as a middle to upper Visean dextral shear zone 431 (Rolin et al., 2009). The seismic information confirms the role of these faults during the 432 opening of Carboniferous or Permian Arpheuilles and Contres basins (Fig. 6b; Beccaletto et 433 al., 2015). Moreover, they were interpreted as Permian or Triassic fracture zones 434 reactivated during the opening of the North Atlantic Ocean and Gulf of Biscave (Vigneresse. 435 1988). They are also known throughout the Armorican Massif, where they bound small 436 Tertiary basins. Based on these information, the N150E-N160E striking faults can be 437 interpreted as Variscan faults, reactivated during the tectonic evolution of the Paris Basin, 438 strongly affecting present-day geometry in the southwestern part of the basin.

The N30E striking normal faults, mainly located in the Mauges nappe (Fig. 6b), are also interpreted as Variscan structures reactivated during the Permo-Carboniferous, controlling the geometry of the Arpheuilles basin (Fig. 6b; Beccaletto et *al.*, 2015).

## 442 **6.2** Interpretation of the undercover lithologies

Based on the geophysical signatures and the structural information, we propose a geological map displaying the interpreted lithologies assigned to the dominant geophysical signatures (Fig.7a) and an interpretative cross section based on geological information observed in the field (Fig. 7b). This geological map as well as its tectonic implications are discussed in each litho-structural unit.

#### 448 6.2.1 Central Brittany

449 Central Brittany is mainly composed of Neoproterozoic metasedimentary rocks and 450 Paleozoic sedimentary rocks intruded by Carboniferous granites (Fig. 2). In this litho-structural unit, high magnetization and density signatures (Fig. 6a) unambiguously
refer to the Fe-rich Ordovician sandstone (up to 0.8 SI and 3.11 g/cm<sup>3</sup>; Fig. 5). This marker
extends eastwards, bounding the NBSASZ (Fig. 8a); its presence is confirmed at 300m
depth under cover by a borehole (Fig. 6b).

455 Contrary to previous models (Weber, 1971, 1973; Debeglia and Weber, 1980), the Paleozoic rocks can be discriminated from the Neoproterozoic ones using the petrophysical 456 457 information. Paleozoic rocks including Fe-rich sandstones displaying low magnetization and density (from 1.5 to 5 x  $10^{-4}$  SI and 2.71 g/cm<sup>3</sup>; Fig. 5) can be traced eastwards (Fig. 7a). In 458 459 the field, geophysical signatures of the Neoproterozoic metasedimentary rocks are heterogeneous, with intermediate magnetic susceptibility and low density (from 10<sup>-4</sup> to 460  $3 \times 10^{-4}$  SI and 2.55 g/cm<sup>3</sup>; Fig. 5). They are also intruded by moderately magnetic and 461 462 dense (3.11 g/cm<sup>3</sup>) gabbroic dykes (Verhnet et al., 2009). Thus the map exhibits a 463 succession of thin E-W trends with intermediate to high magnetic and intermediate gravity 464 signatures that can be identified under the sedimentary cover and grouped with the 465 Neoproterozoic metasedimentary rocks (Fig. 6a). The Laval basin has intermediate to low 466 magnetic and gravity signatures as well as peculiar E-W trending texture (Fig. 6a) well 467 visible in the magnetic vertical gradient (Fig. 3c). This E-W trending signature is likely related to interbedded basalts with intermediate magnetization and density (up to 10<sup>-2</sup> SI 468 and 2.85 g/cm<sup>3</sup>; Fig. 5) as observed in the field (Fig. 8b). This feature allows delimiting the 469 470 southern extension of the Laval basin under the Paris Basin cover (Fig. 8b). Furthermore, 471 the low magnetic and gravity signatures (Fig. 6a) are related to Carboniferous granite intrusions (less than  $5 \times 10^{-4}$  SI and 2.65 g/cm<sup>3</sup>; Fig. 5). Witnesses of these granitic plutons 472 473 are mapped in the field, bounded and affected by the dextral shearing of the NASZ: the Pertre granite emplaced at 343 ± 3Ma (Verhnet et al., 2009) and the Craon granite (e.g. Le 474

Gall et *al.*, 2011; Trautmann et *al.*, 2011). The location and the shape of the undercover granites suggest that they probably are affected by the NASZ shearing (Fig. 8c).

477 Overall, the Central Brittany litho-structural unit, delimited to the south by clear markers of 478 Fe-rich Ordovician sandstones, appears also limited to the east, at the junction between the 479 NASZ and the NBSASZ (Fig. 6b). In the southwestern part of the Paris Basin, these 480 interpretations are consistent with the tectonic sketch of major Armorican shear zones 481 proposed by Martinez Catalàn et al., (2012). The southern part of Central Brittany is 482 structured by patterns of Neoproterozoic and Paleozoic sedimentary rocks. In its 483 northwestern part, the early-Carboniferous Laval basin is developed along the NASZ; it is 484 delimited by low magnetic and gravity signatures (Fig. 6b) related to granitic bodies (Fig. 485 7a), to the south.

486

## 487 6.2.2 The Lanvaux unit

488 In agreement with previous works, the eastern extension of the Lanvaux negative gravity 489 anomaly (Fig. 4b) is well documented along more than 200 km (Weber, 1973; Debeglia and Weber, 1980; Autran et al., 1994; Martelet et al., 2013). This anomaly is interpreted as 490 491 deriving from the low magnetization and low density signatures of the Lanvaux orthogneiss, confirmed by the petrophysical measurements (from  $10^{-5}$  to 0.7 x  $10^{-4}$  SI and 2.55 g/cm<sup>3</sup>; 492 493 Fig. 5). In the field, the Carboniferous Bécon and the St-Lambert granites display low 494 magnetic susceptibilities and densities comparable to those of the orthogneiss (from 2 x 10<sup>-5</sup> to 10<sup>-4</sup> SI; Fig. 5); it is therefore not possible to discriminate the older Lanvaux 495 496 orthogneiss (477 ± 18 Ma; Guerrot et al., in Janjou et al., 1998) from the Carboniferous 497 granites. The Carboniferous granites being mainly located along the SBSASZ (Fig. 8c), we 498 consider that the low magnetic and gravity signatures are related to the Lanvaux 499 orthogneiss (Fig. 8c). Eastwards, under cover, the strike of the NW-SE Lanvaux 500 orthogneiss evolves to an E-W and progressively NE-SW direction, parallel to the NBSASZ 501 (Fig. 8c). Lanvaux unit is an antiform (Faure and Cartier, 1998), with Neoproterozoic 502 metasedimentary and Paleozoic sedimentary rocks similar to those of the Central Brittany 503 and with comparable petrophysical characteristics (Fig. 5). The succession of geophysical 504 signatures of metasedimentary and sedimentary rocks as well as the orthogneiss observed 505 in the field extends eastwards under the Paris Basin sedimentary cover. This suggests that 506 the unit has the same antiformal structure, throughout its eastward extension (Fig. 7b). At 507 the southern border of the Lanvaux unit, the magnetic signature of the Paleozoic rocks 508 highlights the tectonic limit with the St-Georges-sur-Loire unit (Fig. 6b).

509

510 6.2.3 The St-Georges-sur-Loire unit

511 Whereas cartographically well marked under cover, the eastern extension of the St-512 Georges-sur-Loire unit has various geophysical signatures (Fig. 6a), rather magnetic and 513 Cartographically, these signatures cannot be formally related to the dense. northern/southern parts of the unit, as observed in the field (Fig. 2a). The northern part of 514 515 the unit is marked by low magnetic and gravity signatures (Fig. 6a) related to a granitic pluton (up to 10<sup>-4</sup> SI and 2.65 g/cm<sup>3</sup>; Fig. 5), located at the junction between the Lanvaux 516 517 and the St-Georges-sur-Loire units, mostly hidden below the Paleozoic series (up to 5 x 10<sup>-</sup> 518 <sup>4</sup> SI and 2.68 g/cm<sup>3</sup>). Granites located along the SBSASZ (Fig. 8c) belong to a leucogranite 519 belt displaying dextral shearing, associated with the emplacement of the leucogranite 520 plutons (Berthé et al., 1979; Jégouzo, 1980; Vigneresse and Brun, 1983; Gapais et al., 521 1993; Turillot et al., 2009). Among them, Questembert and Lizio leucogranites (Fig. 8c) 522 emplaced at 316 ± 3 Ma and 316 ± 6 Ma (Tartèse et al., 2011b, 2011a), in agreement with

523 the St-Lambert granite located to the Lanvaux unit, which displays S/C structures 524 suggesting an emplacement during the dextral shearing of the NBSASZ at 312 ± 3 Ma 525 (Faure and Cartier, 1998). Under cover, these low magnetic and gravity signatures are not 526 observed (Fig. 6a), thus these granites cannot be extended eastward, suggesting a limited 527 eastward extension of the leucogranitic belt.

528 The southern part of the unit, the blocky sub-unit is characterized by scattered high 529 magnetic and various low to high gravity signatures (Fig. 6a). Below the Paris Basin 530 sedimentary cover, the high magnetic and gravity signatures elongated eastwards 531 throughout the map, likely correspond to basic rocks, as interpreted by Martelet et al., 532 (2013). These basic rocks can be variously interpreted in terms of lithology: i) basaltic or 533 gabbroic olistostoliths as described in this unit in the Armorican Massif (Cartier and Faure, 534 2004), ii) interbedded basalts related to the opening of a back-arc basin (Ducassou et al., 535 2011) or iii) mafic rocks such as the ophiolites which sporadically crop out along the NSE 536 fault (Marchand, 1981; Faure et al., 2008; Ballèvre et al., 2009). In the eastern part of the 537 St-Georges-sur-Loire unit, the depth of the contact between the substratum and the 538 sedimentary cover is defined at about 500 m, and the bodies responsible for the magnetic 539 signal are located at more than 1500 m depth (Martelet et al., 2013). Consequently the 540 magnetic susceptibility of the magnetic source has to be strong enough to produce such a 541 magnetic signature. According to the magnetic susceptibility measured in the field in St-542 Georges-sur-Loire unit, we cannot interpret the source as basalts since they do not yield a high magnetic susceptibility (from 6 to 8 x  $10^{-4}$  SI; Fig. 5). The third hypothesis better 543 544 complies with the petrophysical signatures and with other evidence reported in the 545 literature. Indeed, the interpretation of the Armor2 seismic profile, located in the southern 546 part of the Armorican Massif (Bitri et al., 2003), suggested the presence of the northern part 547 of the Champtoceaux complex beneath St-Georges-sur-Loire unit. The northern part of this

548 complex crops out as ophiolitic series along the NSE fault; it is made of high magnetic 549 susceptibility and high density rocks, such as amphibolite, and micaschists (from 1.1 to 1.5 550  $\times 10^{-2}$  SI, 3.02 g/cm<sup>3</sup> and up to 0.3 SI, 2.79 g/cm<sup>3</sup>, respectively; Fig. 5). Therefore, contrary 551 to previous sketch (Weber, 1973; Debeglia and Weber, 1980), our results reveal the 552 eastward extension of ophiolitic series marking the NSE Eo-Variscan suture (Fig. 8d); it is 553 supported by high magnetic and gravity signatures which extend up to the eastern part of 554 the study area limiting St-Georges-sur-Loire unit to the south.

555

#### 556 **6.2.4 The Mauges nappe and the Champtoceaux Complex**

557 The Mauges nappe, mainly composed of Neoproterozoic micaschists (Fig. 2), exhibits both 558 low magnetic and gravity signatures (Fig. 6a) that may account for hidden granitic plutons. 559 Granitic rocks crop out locally, as for instance the Chemillé pluton intruding the Mauges micaschists, displaying low magnetic susceptibility and density (up to 8 x 10<sup>-4</sup> SI and 560 2.55 g/cm<sup>3</sup>; Fig. 5). Similar low geophysical signatures depicted in the undercover 561 562 extension of the Mauges nappes (Fig. 6a) suggest the presence of punctual granitic 563 plutons, probably similar to the Chemillé granite (Fig. 8c). Also, metabasites interbedded in the micaschists yield higher magnetic susceptibility and density ranges (from 5 to 7 x  $10^{-4}$ 564 SI and 2.82 g/cm<sup>3</sup>; Fig. 5) than the surrounding micaschists. These signatures trace the 565 566 eastward extension of the Mauges nappe under the sedimentary cover of the southern part 567 of the Paris Basin (Fig. 7a).

The Choletais area is composed of Cambrian acidic volcanites, intruded by microgranite, granodiorite and gabbro-diorite, unconformably covering the Neoproterozoic rocks (Fig. 2). Among them, the Vézins granodiorite (Fig. 8c) emplaced at  $345 \pm 5$  Ma (Thiéblemont et *al.*, 2011) is supposed to be a syn-kinematics intrusion related to the dextral shearing of the

572 Cholet fault (Rolin et al., 2009). In the Choletais area, the magnetic susceptibilities and 573 density measurements do not discriminate the gabbro-diorite suites from the granodioritic plutons (from 4 to 7 x  $10^{-2}$  SI, 2.83 g/cm<sup>3</sup> and from 2 to 5 x  $10^{-2}$  SI, 2.73 g/cm<sup>3</sup>, respectively; 574 Fig. 5). For this reason, these lithologies are grouped together as "basic rocks" in the 575 576 interpretative geological map (Fig. 7a). The Cambrian volcanics are discrimated by a lower density compared to basic rocks (2.65 g/cm<sup>3</sup> and from 2.73 to 2.83 g/cm<sup>3</sup>, respectively; Fig. 577 578 5). As previously described (Weber, 1971), in the southern part of the Mauges nappe, the 579 northern branch of the Cholet fault is bounded to the north by an unexpected E-W striking 580 high density elongated anomaly, which extends 200 km eastwards (Fig. 4b). It is 581 superimposed, in the Armorican Massif, to the low magnetization and low density Thouars microgranite (from 1 to 7 x  $10^{-4}$  SI and 2.65 g/cm<sup>3</sup>; Fig. 5). The high intensity anomaly can 582 583 however be explained by the close association of acidic and basic magmatism composing 584 the Thouars massif (Mathieu, 1943, 1958; Weber, 1971) emplaced at 519 ± 10 Ma 585 (Thiéblemont, et al., 2011). The high resolution geophysical data enhance this dual 586 anomaly: it highlights punctual high magnetization signatures associated with high density anomaly (Fig. 6a) related to basics rocks (from 3 to 6 x  $10^{-2}$  SI and up to 2.83 g/cm<sup>3</sup>; Fig. 5) 587 and the presence of high density intrusive dolerite (2.97 g/cm<sup>3</sup>; Fig. 5) located along the 588 589 Cholet fault, at the junction between the Mauges nappe and the Haut Bocage unit.

The early Carboniferous Ancenis basin is marked by a low gravity anomaly (Fig. 4b) likely related to a hidden granitic pluton (Fig. 8c) as previously described by Martelet et *al.*, (2013) as well as E-W striking high magnetic trends well defined in the vertical gradient of the magnetic anomaly. This contrast can reasonably be related to the northern extension of the Champtoceaux complex, corresponding to the ophiolitic nappe, buried below the Ancenis basin, along the NSE fault, also interpreted in Armor2 seismic profile (Bitri et *al.*, 2003).

596 Following the NSE fault to the east, the Permo-Carboniferous Arpheuilles basin, recognized 597 by seismic profiles under cover (Beccaletto et al., 2015) and two boreholes (Fig. 6b), 598 exhibits a peculiar texture in the magnetic first vertical derivative (Fig. 3c) and tilt (Fig. 3d) 599 maps. This texture allows delimiting the extension of the Arpheuilles basin (Fig. 8b). 600 Superimposed on the magnetic texture, high intensity magnetic anomalies are interpreted 601 as the presence of Cambrian volcanic rocks (Fig. 7a) underlying Arpheuilles basin, 602 accompanied by granitic plutons highlighted by their low gravity signatures (Fig. 6a). Our 603 results show that, to the north, Arpheuilles basin is bounded by the NSE fault and controlled 604 by N150E striking faults as well as their N30E-N40E conjugates (Fig. 8b). This suggests 605 that the Arpheuilles basin may be the lateral equivalent of the early Carboniferous Ancenis 606 basin.

607

## 608 6.2.5 The Aigurande Plateau and its connection with the Haut Bocage unit

This area consists in a stack of metamorphic nappes intruded by granitic plutons (Fig. 2).
Under cover, the Cholet fault delimits the northern contact of the high grade metamorphic
nappes with the Mauges nappe, composed of low grade micaschists (Fig. 8d).

612 In the northern part of the Aigurande Plateau, the Lower Gneiss Unit (LGU), is mainly 613 composed of low magnetic susceptibility and density micaschists and metagrauwackes (up to  $4 \times 10^{-4}$  SI and 2.8 g/cm<sup>3</sup>; Fig. 5), and intermediate magnetic and high density 614 615 amphibolite (from 5 to 9 x 10<sup>-4</sup> SI, 3.01 g/cm<sup>3</sup>). The Upper Gneiss Unit (UGU) was 616 discriminated from the LGU by both the high magnetic and density signatures (Fig. 6a) deriving from the amphibolites within the leptynite-amphibolite complex (from  $1.5 \times 10^{-2}$  to 5 617 x 10<sup>-2</sup> SI and 2.98 g/cm<sup>3;</sup> Fig. 5). Consequently, the LGU/UGU contact can be mapped 618 619 under the Paris Basin sedimentary cover, bounded to the north by the northern branch of the Cholet fault (Fig. 7a). The LGU micaschists are associated with the Haut Bocage unit
described in the field (Fig. 6b; Fig. 7a), which consist of a stack of metamorphic nappes,
refolded in an ENE-SSW striking antiform/synform succession, intruded by Carboniferous
granites (Fig. 7b).

The micaschists of the Para-autochtonous unit displaying low to intermediate magnetic susceptibility and density (up to  $4 \times 10^{-4}$  SI and 2.68 g/cm<sup>3</sup>), related to low to intermediate magnetic and gravity signatures (Fig. 6a), has no characteristic signature that can be mapped under the Paris Basin sedimentary cover.

In this area, granitic plutons belong to the Hercynian Mortagne – Marche leucogranites belt
(Vigneresse, 1988; Gapais et *al.*, 1993; Vigneresse, 1999; Rolin and Colchen, 2001; Rolin
et *al.*, 2009; Edel et *al.*, 2015; Gapais et *al.*, 2015). These granites are largely represented
in the Haut Bocage unit and Massif connection (Fig. 7a).

632

## 633 6.2.6 The Poitou High: the Haut Bocage unit and the Confolentais area junction

634 Largely represented in the field, in the Haut Bocage unit and the Confolentais area 635 (Fig. 6b), the low magnetic and gravity signatures (Fig. 6a) are related to granitic plutons (from 0.8 x 10<sup>-5</sup> to 10<sup>-4</sup> SI and 2.65 g/cm<sup>3</sup>; Fig. 5). Under cover, equivalent magnetic and 636 637 gravity signatures related to granitic plutons described in the Poitou High (from 2.5 to  $4 \times 10^{-5}$  SI and 2.62 g/cm<sup>3</sup>; Fig. 5), mark the connection between the southern part of the 638 639 Haut Bocage unit and Confolentais area (Fig. 7a). As largely recovered in boreholes (Fig. 640 6b), granitic plutons represent the main rocks of the substratum in this area (Fig. 7a). These 641 granitic plutons consist in leucomonzogranites, leucogranites, granodiorites and calk-642 alkaline diorites. They belong to a granitic belt emplaced from late Devonian to early-643 Carboniferous and are associated with the shearing of the various branches of the SASZ 644 (see Rolin et al., 2009 for more information). Using the geophysical signatures of the
645 substratum (Fig. 6a), it is not possible to discriminate the various leucomonzogranites,
646 leucogranites and granodiorites.

647 Under the sedimentary cover, the southern part of the Poitou High is marked by punctual 648 high magnetic and gravity signatures (Fig. 6a) related to the calk-alkaline diorite plutons 649 exposed in the field, both in Confolentais area and Haut Bocage unit (Fig. 6b). These diorite 650 plutons emplaced at 373  $\pm$  10 Ma (Cuney et al., 1993) and from 360  $\pm$  3 Ma to 349  $\pm$  5 Ma (Bertrand et al., 2001; Alexandre et al., 2002) for the Montcoutant and various Poitou High 651 652 diorite plutons, respectively, are associated with the dextral shearing of the SBSASZ (Fig. 653 8c) (see Rolin et al., 2009 for more information). In the southern Aigurande Plateau, Huriel 654 diorite intrusion (Fig. 8c) emplaced at 361 ± 1 Ma (Pin and Paquette, 2002) and post-date 655 the dextral shearing of the Marche fault (Rolin et al., 2009). Diorite plutons have consistent high magnetic susceptibility and density (up to  $1.5 \times 10^{-2}$  SI and 2.80 g/cm<sup>3</sup> and from 3 to 6 656 x  $10^{-3}$  SI, and 2.80 g/cm<sup>3</sup>, respectively; Fig. 5). These basic rocks are discriminated i) from 657 658 the intermediate magnetic and low density signatures corresponding to the LGU host 659 micaschists, in the Haut Bocage unit, and ii) from the high magnetic and intermediate 660 signatures related to the metavolcanites belonging to the UGU (up to 0.3 SI; Fig. 5), in the 661 Confolentais area (Fig. 7a).

662

## 663 7. Summary and Conclusion

664 Our paper outlines the benefit of a joint interpretation of potential fields (high-resolution 665 aeromagnetic and gravity data) and petrophysical characterization (magnetic susceptibility 666 and density measurements on rock samples), in order to derive reliable lithological and 667 structural mapping of a buried substratum. 668 In order to propose a geological map of the hidden substratum of the southwestern part of 669 the Paris Basin, our methodology, is divided into five successive stages: i) the potential field 670 data were processed, with the aim to get specific information (geophysical contrasts, 671 structural features...), ii) magnetic susceptibilities and densities were measured on field 672 rock samples along the eastern border of the Armorican Massif and the northern border of 673 the Massif Central, leading to a petrophysical library of lithologies, iii) using selected 674 magnetic and gravity maps, a map of geophysical signatures was synthetized using an 675 unsupervised classification, featuring 6-levels of magnetic/gravity intensities, iv) the 676 combined analysis and interpretation of magnetic and gravity trends with the synthetized 677 geophysical signatures, allowed extending the Variscan litho-structural units below the 678 Paris Basin sedimentary pile, v) relating the geophysical signatures to the petrophysical 679 characteristics (density and magnetization) within each litho-structural unit, allowed 680 interpreting a geological map of the substratum. This updated study reveals new geological 681 information: i) the limited eastward extension of Central Brittany, bordered to the east by the 682 NE-SW striking NBSASZ; ii) the eastward extension, along ca. 150 km, of the Cholet fault, 683 interpreted as a major fault, delineating the northern limit of the Aigurande Plateau; iii) the 684 emphasis on a series of N150E-N160E and N30E striking normal Variscan fault, reactivated 685 during the tectonic history of the Paris Basin, especially controlling the opening of Permo-Carboniferous basins; iv) the extension of the Nort-sur-Erdre fault considered as an 686 687 ophiolitic suture, documented by the presence of high magnetic and density rocks along the 688 southern part of the St-Georges-sur-Loire unit.

Overall, our methodology provides keys for extensive mapping of buried basement using magnetic, gravity and petrophysical data. In the near future, this study will be extended to the entire Paris Basin in order to propose a complete geological map of the pre-Mesozoic substratum of the Paris Basin.

#### 693 Acknowledgments

This work is part of a PhD Thesis co-funded by Région Centre and BRGM. We thank an anonymous reviewer and J.B. Edel for useful and constructive comments, which contributed to improving the manuscript. Geophysical maps were drawn using Geosoft© software.

698

## 699 References

- Alexandre, P., Le Carlier de Veslud, C., Cuney, M., Ruffet, G., Virlogeux, D., Cheilletz, A.,
- 701 2002. Datation 40Ar/39Ar des leucogranites sous couverture du complexe plutonique
- de Charroux–Civray (Vienne). Comptes Rendus Geoscience 334, 1141–1148.
  doi:10.1016/S1631-0713(02)01859-X
- Autran, A., Lameyre, J., 1980. Conclusions générales. Les granitoïdes de France.
  Mémoires BRGM 107, 87–99.
- Autran, A., Lefort, J.P., Debeglia, N., Edel, J.B., Vigneresse, J.L., 1994. Gravity and
  Magnetic Expression of Terranes in France and Their Correlation Beneath Overstep
  Sequences, in: Chantraine, J., Rolet, J., Santallier, D.S., Piqué, A., Keppie, J.D.
  (Eds.), Pre-Mesozoic Geology in France and Related Areas, IGCP-Project 233.
  Springer Berlin Heidelberg, pp. 49–72.
- Ballèvre, M., Pinardon, J.-L., Kienast, J.-R., Vuichard J.-P., 1989. Reversal of Fe-Mg
  partitioning between garnet and staurolite in eclogite-facies metapelites from the
  Champtoceaux nappe (Brittany, France). Journal of Petrology 30, 1321–1349.
- 714 Ballèvre, M., Paris, F., Robardet, M., 1992. Corrélations ibéro-armoricaines au
  715 Paléozoïque : une confrontation des données paléobiogéographiques et
  716 tectonométamorphiques. Comptes Rendus de l'Académie des sciences. Série 2,

717 Mécanique, Physique, Chimie, Sciences de l'univers, Sciences de la Terre 315, 1783–
718 1789.

- Ballèvre, M., Marchand, J., Godard, G., Goujou, J.-C., Christian, J., Wyns, R., 1994.
  Eo-Hercynian Events in the Armorican Massif, in: Chantraine, J., Rolet, J., Santallier,
  D.S., Piqué, A., Keppie, J.D. (Eds.), Pre-Mesozoic Geology in France and Related
  Areas, IGCP-Project 233. Springer Berlin Heidelberg, pp. 183–194.
- Ballèvre, M., Lardeux, H., 2005. Signification paléoécologique et paléogéographique des
  bivalves du Carbonifère inférieur du bassin d'Ancenis (Massif armoricain). Comptes
  Rendus Palevol 4, 109–121. doi:10.1016/j.crpv.2004.11.011.
- Ballèvre, M., Bosse, V., Ducassou, C., Pitra, P., 2009. Palaeozoic history of the Armorican
  Massif: Models for the tectonic evolution of the suture zones. Comptes Rendus
  Géoscience 341, 174–201. doi:10.1016/j.crte.2008.11.009.
- Baptiste, J., Martelet, G., Faure., Beccaletto, L., Perrin, J., 2015. Up-to-date regional gravity
  and aeromagnetic data to unravel the geological patterns of the pre-Mesozoic
  substratum of the Paris Basin. Variscan 2015 : The Variscan belt : correlations and
  plate dynamics, Jun 2015, Rennes, France.
- 733 Beccaletto, L., Hanot, F., Serrano, O., Marc, S., 2011. Overview of the subsurface structural 734 pattern of the Paris Basin (France): Insights from the reprocessing and interpretation of 735 regional seismic lines. Marine and Petroleum Geology, Thematic Set on the 736 Implications of basin dynamics on petroleum systems 28, 861-879. 737 doi:10.1016/j.marpetgeo.2010.11.006
- Beccaletto, L., Capar, L., Serrano, O., Marc, S., 2015. Structural evolution and sedimentary
  record of the Stephano-Permian basins occurring beneath the Mesozoic sedimentary
  cover in the southwestern Paris basin (France). Bulletin de la Société Géologique de
  France. 6, 429-450.

Bernard-Griffiths, J., Peucat, J.J., Sheppard, S., Vidal, P., 1985. Petrogenesis of Hercynian
leucogranites from the southern Armorican Massif: contribution of REE and isotopic (Sr,
Nd, Pb and O) geochemical data to the study of source rock characteristics and ages.

745Earth and Planetary Science Letters 74, 235–250. doi:10.1016/0012-821X(85)90024-X

- Berthé, D., Choukroune, P., Jegouzo, P., 1979. Orthogneiss, mylonite and non coaxial
  deformation of granites: the example of the South Armorican Shear Zone. Journal of
  Structural Geology 1, 31–42. doi:10.1016/0191-8141(79)90019-1
- Bertrand, J.M., Leterrier, J., Cuney, M., Brouand, M., Stussi, J.M., Delaperrière, E.,
  Virlogeux, D., 2001. Géochronologie U-Pb sur zircons de granitoïdes du Confolentais,
- du massif de Charroux-Civray (seuil du Poitou) et de Vendée. Géologie de la France 1,
  167–189.
- Bitri, A., Ballèvre, M., Brun, J.-P., Chantraine, J., Gapais, D., Guennoc, P., Gumiaux, C.,
  Truffert, C., 2003. Imagerie sismique de la zone de collision hercynienne dans le
  Sud-Est du Massif armoricain (projet Armor 2/programme Géofrance 3D). Comptes
  Rendus Géoscience 335, 969–979. doi:10.1016/j.crte.2003.09.002.
- 757 Blakely, R.J. 1996. Potential Theory in Gravity and Magnetic Applications (Cambridge758 University Press).
- Bonijoly, D., Perrin, J., Truffert, C., Asfirane, F., 1999. Couverture géophysique aéroportée
  du Massif armoricain, magnétisme et radiométrie spectrale. Rapport BRGM R 40471.
- Bosse, V., Feraud, G., Ruffet, G., Ballèvre, M., Peucat, J.-J., De Jong, K., 2000. Late
  Devonian subduction and early-orogenic exhumation of eclogite-facies rocks from the
  Champtoceaux Complex (Variscan belt, France). Geological Journal. 35, 297–325.
  doi:10.1002/gj.864.

- Burg, J.-P., 1981. "Tectonique tangentielle hercynienne en Vendée littorale : Signification
  des linéations d'étirement E-W dans les porphyroïdes à foliation horizontale." Comptes
  Rendus de l'Académie des Sciences de Paris, 293, 849-854.
- Cabanis, B., Wyns, R., 1986. Le volcanisme précambrien des Mauges (sud-est du Massif
  armoricain) et ses caractères géochimiques. Hercynica, Rennes, (II), 1, 71-78.
- 770 Cartannaz, C., Rolin, P., Le Métour, J., Fabbri, O., 2006. Fammenian–Tournaisian dextral
- ductile shear in the French Variscan belt. Comptes Rendus Géoscience 338, 214–221.
  doi:10.1016/j.crte.2005.12.003
- 773 Cartier, C., Faure, M., Lardeux, H., 2001. The Hercynian Orogeny in the South Armorican
- Massif (Saint-Georges-sur-Loire Unit, Ligerian Domain, France): rifting and welding of
  continental stripes. Terra Nova. 13, 143–149.
- Cartier, C., Faure, M., 2004. The Saint-Georges-sur-Loire olistostrome, a key zone to
  understand the Gondwana-Armorica boundary in the Variscan belt (Southern Brittany,
  France). Int. J. Earth Sci. 93, 945–958. doi:10.1007/s00531-004-0398-3.
- 779 Cavet, P., Gruet, M., Pillet, J., 1966. Sur la présence du Cambrien à Paroxides à Cléré-sur-
- 780 Layon (Maine et Loire) dans le Nord-Est du Bocage vendéen (Massif Armoricain).
- 781 Comptes Rendus Hebdomadaires des séances de l'Académie des Sciences. 263,
  782 1685.
- 783 Cavet, P., Arnaud, A., Barbaroux, L., Blaise, J., Brossé, R., Chauris, L., Gruet, M., Lardeux,
- H., Rivière, J.-M., 1970. Carte géologique de la France (1/50000), *feuille Chalonnes- sur-Loire* (453), BRGM, Orléans.
- Cavet, P., Lardeux, H., Philippot, A., 1971. Ordovicien et Silurien aux environs de Montjean
  et Chalonnes (Maine-et-Loire, Sud-Est du Massif armoricain). Mémoires BRGM 73,
  199-212.

- Cavet, P., Arnaud, A., Blaise, J., Brossé, R., Chauris, L., Gruet, M., Lardeux, H., 1976.
  Carte géologique de la France (1/50000), *feuille Angers* (454), BRGM, Orléans.
- Chantraine, J., Autran, A., Cavelier, C. and collab., 2003. Carte Géologique de la France au
   1/100000<sup>e</sup>. BRGM (6<sup>ème</sup> édition), Orléans. France.
- Chauris, L., and Lucas, G., 1964. Les environs de Bécon-les-Granits (Maine-et-Loire)
  (Feuille d'Ancenis au 80.000e). Bulletin de la Carte Géologique de France, LX-277
  25–33.
- Chauris, L., 1969. Sur un important accident structural dans le Nord-Ouest de l'Armorique.
  Comptes Rendus de l'Académie des Sciences. 268, 2859–2861.
- Clark, D., Emerson, D., 1991. Notes on rock magnetization characteristics in applied
   geophysical studies. Exploration Geophysics 22, 547–555. doi:10.1071/EG991547.
- 800 Corpel, J., Weber, C., 1970. Contribution de l'étude de l'aimantation rémanente à la 801 gîtologie du synclinal ferrifère de Segré. Rapport BRGM/70-SGN-029-GPH, 23 pp.
- Cuney, M., Stussi, J.-M., Brouand, M., Dautel, D., Michard, A., Gros, Y., Poncet, D.,
  Bouton, P., Colchen, M., Vervialle, J.-P., 1993. Géochimie et géochronologie U/Pb
  des diorites quartziques du Tallud et de Moncoutant : nouveaux arguments pour une
  extension de la Ligne Tonalitique Limousine en Vendée. Comptes Rendus de
  l'Académie des Sciences. Série 2, Mécanique, Physique, Chimie, Sciences de
  l'univers, Sciences de la Terre 316, 1383–1390.
- B08 Dearing, J., 1999. Magnetic susceptibility, in: Environmental Magnetism: A Practical Guide.
  809 Quaternary Research Association London, pp. 35–62.
- Broglia, N., Weber, C., 1980. Synthèse géologique du bassin de Paris. Mémoires du
  BRGM n°102.

Debeglia, N., Weber, C., 1985. Geologic mapping of the basement of the Paris basin
(France) by gravity and magnetic data interpretation. In: The Utility of regional gravity
and magnetic anomaly maps, (ed. W.J. Hinze). Society of Exploration Geophysicists.

815 Ducassou, C., Ballèvre, M., Lardeux, H., Robin, C., 2011. Evidence for pre-orogenic, Early

816 Devonian rifting in the Variscan belt: stratigraphy and structure of the Palaeozoic cover

of the Mauges Unit (Upper Allochthon, Armorican massif, France). Int. J. Earth Sci.

818 (Geol Rundsch) 100, 1451–1475. doi:10.1007/s00531-010-0605-3.

Edel, J.B., 2008. Structure et nature du socle anté-permien du Bassin de Paris d'après les
données gravimétriques et magnétiques – Le problème de l'Anomalie Magnétique du
Bassin de Paris (AMBP). Géochronique, 105, pp. 31-37.

Edel, J.-B., Schulmann, K., Lexa, O., Diraison, M., Géraud, Y., 2015. Permian clockwise
rotations of the Ebro and Corso-Sardinian blocks during Iberian–Armorican oroclinal
bending: Preliminary paleomagnetic data from the Catalan Coastal Range (NE Spain).
Tectonophysics 657, 172–186. doi:10.1016/j.tecto.2015.07.002

Elkins, T., 1951. The second derivative method of gravity interpretation. Geophysics. 16,
29–50. doi:10.1190/1.1437648.

Fairhead, J.D., Salem, A., Cascone, L., Hammill, M., Masterton, S., Samson, E., 2011. New
developments of the magnetic tilt-depth method to improve structural mapping of
sedimentary basins. Geophysical Prospecting 59, 1072–1086. doi:10.1111/j.13652478.2011.01001.x.

Faure, M., Prost, A.E., Lasne, E., 1990. Deformation ductile extensive d'age namurowestphalien dans le plateau d'Aigurande, Massif central francais. Bulletin de la Société
Géologique de France VI, 189–197. doi:10.2113/gssgfbull.VI.1.189.

835 Faure M., Leloix C., Roig J.Y., 1997. L'évolution polycyclique de la chaîne hercynienne.

Bulletin de la Société géologique de France 168, 695–705.

Faure, M., and Cartier, C., 1998. Déformations ductiles polyphasées dans l'antiforme
orthogneissique de St-Clément-de-la-Place (unité de Lanvaux, Massif armoricain).
Comptes Rendus de l'Académie des Sciences - Series IIA - Earth and Planetary
Science 326, 795–802. doi:10.1016/S1251-8050(98)80245-0.

- Faure, M., Mezeme, E.B., Duguet, M., Cartier, C., Talbot, J.-Y., 2005. Paleozoic tectonic
  evolution of medio-Europa from the example of the French Massif Central and Massif
  Armoricain. Journal of the virtual Explorer 19, 1–25.
- Faure, M., Bé Mézème, E., Cocherie, A., Rossi, P., Chemenda, A., Boutelier, D., 2008.
  Devonian geodynamic evolution of the Variscan Belt, insights from the French Massif
  Central and Massif Armoricain. Tectonics 27, TC2005. doi:10.1029/2007TC002115.
- Faure M., 2014. Le substratum de la France métropolitaine : de la formation du Gondwana
  à la constitution de la Pangée, une histoire de 600 Ma. Géologues, 180, pp. 13-21.
- Gapais, D., Lagarde, J.L., Le Corre, C., Audren, C., Jegouzo, P., Casas Sainz, A., Van Den
  Driessche, J., 1993. La zone de cisaillement de Quiberon: témoin d'extension de la
  chaine varisque en Bretagne méridionale au Carbonifère. Comptes Rendus de
  l'Académie des Sciences, Paris, série II 316, 1123–1129.
- Gapais, D., Brun, J.-P., Gumiaux, C., Cagnard, F., Ruffet, G., Veslud, C.L.C.D., 2015.
  Extensional tectonics in the Hercynian Armorican belt (France). An overview. Bulletin
- de la Société Géologique de France 186, 117–129. doi:10.2113/gssgfbull.186.2-3.117
- 856 Gébelin, A., Brunel, M., Monié, P., Faure, M., Arnaud, N., 2007. Transpressional tectonics
- and Carboniferous magmatism in the Limousin, Massif Central, France: Structural and
  40Ar/39Ar investigations. Tectonics 26.
- Gérard, A., and Griveau, P., 1972. Interpretation quantitative en gravimétrie ou magnétisme
  à partir de cartes transformées en gradient vertical. Geophysical Prospecting 20, 459–
  481. doi:10.1111/j.1365-2478.1972.tb00648.x.

- Goguel, J., and Francia, 1954. Levé gravimétrique détaillé du Bassin parisien. Ministère de
  l'Industrie et de l'Energie. Direction des Mines. Imprimerie Nationale.
- Goguel, J., 1972. Tendances modernes dans l'interprétation géologique des données
  gravimétriques. Bull. BRGM 43–50.
- 866 Guillocheau, F., Robin, C., Allemand, P., Bourquin, S., Brault, N., Dromart, G., Friedenberg,
- R., Garcia, J.-P., Gaulier, J.-M., Gaumet, F., others, 2000. Meso-Cenozoic geodynamic
  evolution of the Paris Basin: 3D stratigraphic constraints. Geodinamica Acta 13, 189–
  245.
- Houlgatte, E., Le Hérissé, A., Pelhate, A., Rolet, J., 1988. Evolution géodynamique du
  Bassin carbonifère de Laval. Géologie de la France 1, 27–46.
- Jégouzo, P., 1980. The South Armorican Shear Zone. Journal of Structural Geology, Shear
  zones in rocks 2, 39–47. doi:10.1016/0191-8141(80)90032-2.
- 874 Lardeux, H., Cavet, P., 1994. Paleozoic of the Ligerian Domain, in: Chantraine, J., Rolet, J.,
- 875 Santallier, D.S., Piqué, A., Keppie, J.D. (Eds.), Pre-Mesozoic Geology in France and
  876 Related Areas, IGCP-Project 233. Springer Berlin Heidelberg, pp. 152–156.
- Lecoanet, H., Lévêque, F., Segura, S., 1999. Magnetic susceptibility in environmental
  applications: comparison of field probes. Physics of the Earth and Planetary Interiors
  115, 191–204. doi:10.1016/S0031-9201(99)00066-7.
- Le Corre, C., Auvray, B., Ballèvre, M., & Robardet, M., 1991. Le massif
  armoricain. Scientifical Geological Bulletin. 44, 31-103.
- Le Gall, J., Vernhet, Y., Lacquement, F., Gauquelin, J.-L., Robert, A., Cocherie, A.,
  Naveau, J, 2011 Notice explicative, Carte géologique de France (1/50 000), feuille
  Laval (319). Orléans : BRGM, 261 p. Carte géologique par Le Gall J., Gigot P.,
  Savaton P., Lacquement F., Poprawsky Y., Vernhet Y. (2011).

- Le Hébel, F., Vidal, O., Kienast, J.-R., Gapais, D., 2002. Les « Porphyroïdes » de Bretagne
  méridionale : une unité de HP–BT dans la chaîne hercynienne. Comptes Rendus
  Geoscience 334, 205–211. doi:10.1016/S1631-0713(02)01746-7.
- Le Hérissé, A., and Plaine, J., 1982. Volcanisme basique dans le Carbonifère inférieur du
  Synclinorium de Laval (Massif armoricain, France). Comptes Rendus de l'Académie
  des Scences de Paris, 294, 1199-1202.
- Lienhardt, M.J., 1961. Étude stratigraphique, pétrographique et structurale du socle anté permien du bassin de Paris. Annales de la Société Géologique du Nord 81, 233-241.
- Marchand, J., 1981. Ecaillage d'un "mélange tectonique" profond: le complexe
  cristallophyllien de Champtoceaux (Bretagne méridionale). Comptes Rendus de
  l'Académie des Sciences de Paris 293, 223–228.
- Martelet, G., Debeglia, N., Truffert, C., 2002. Updating and validating the French gravity
  terrain corrections out to a distance of 167 km. Comptes Rendus Géoscience. 334,
  449–454.
- 900 Martelet, G., Calcagno, P., Gumiaux, C., Truffert, C., Bitri, A., Gapais, D., Brun, J.P., 2004.
- 901 Integrated 3D geophysical and geological modelling of the Hercynian Suture Zone in
  902 the Champtoceaux area (south Brittany, France). Tectonophysics 382, 117–128.
  903 doi:10.1016/j.tecto.2003.12.009.
- Martelet, G., Pajot, G., Debeglia, N., 2009. Nouvelle carte gravimétrique de la France;
  RCGF09 Réseau et Carte Gravimétrique de la France, 2009. Rapport BRGM/RP57908-FR, 77 p.
- Martelet, G., Perrin, J., Truffert, C., Deparis, J., 2013. Fast mapping of magnetic basement
   depth, structure and nature using aeromagnetic and gravity data: combined methods
   and their application in the Paris Basin: Fast mapping of a buried basement using

- 910 magnetic and gravity data. Geophysical Prospecting 61, 857–873. doi:10.1111/1365911 2478.12024.
- 912 Martínez Catalán J.R., 2012. The Central Iberian arc, an orocline centered in the Iberian
- 913 Massif and some implications for the Variscan belt. Int. J. Earth Sci. 101, 1299-1314.
- 914 doi:10.1007/s00531-011-0715-6.
- 915 Mathieu, G., 1943. Révision de la feuille de Saumur au 1/80000ème. Bulletin du Service de
  916 la Carte géologique de France, 212, p119-129.
- 917 Mathieu, G., 1958. Les grandes lignes de la Vendée. Ibid., 253, 46p.
- Matte, P., 1986. Tectonics and plate tectonics model for the Variscan belt of Europe.
  Tectonophysics 126, 329–374. doi:10.1016/0040-1951(86)90237-4.
- Matte, P., Hirn, A., 1988. Seismic signature and tectonic cross section of the Variscan crust
  in western France. Tectonics 7, 141–155.
- Matte, P., 1991. Accretionary history and crustal evolution of the Variscan belt in Western
  Europe. Tectonophysics 196, 309–337. doi:10.1016/0040-1951(91)90328-P.
- 924 Mégnien, C., 1980. Synthèse géologique du Bassin de Paris, I, Stratigraphie et 925 paléogéographie, Mémoires du BRGM, 101, 466pp.
- Miller, H. G. and Singh, V., 1994. Potential field tilt; a new concept for location of potential
  field sources. Journal of Applied Geophysics. 32, 213–217.
- Peiffer, M.-T., 1986. La signification de la ligne tonalitique du Limousin. Son implication
  dans la structuration varisque du Massif Central français. Comptes Rendus de
  l'Académie des Sciences. Série 2, Mécanique, Physique, Chimie, Sciences de
  l'univers, Sciences de la Terre 303, 305–310.
- 932 Perrodon, A. and Zabek, J., 1990. Paris Basin; Interior cratonic basins. AAPG Memoir, 51,933 633-679.

- Pin, C., and Paquette, J.-L., 2002. Le magmatisme basique calcoalcalin d'âge dévonodinantien du nord du Massif Central, témoin d'une marge active hercynienne :
  arguments géochimiques et isotopiques Sr/Nd. Geodinamica Acta 15, 63–77.
  doi:10.1016/S0985-3111(01)01079-8.
- Pomerol, C., 1978. Paleogeographic and structural evolution of the Paris Basin, from the
  Precambrian to the present day, in relation to neighboring regions. Geologie En
  Mijnbouw Journal of Geosciences, 57, 533-543.
- Quenardel, J.-M., Rolin, P., 1984. Palaeozoic evolution of the Plateau d'Aigurande (NW
  Massif Central, France). Geological Society, London, Special Publications 14, 63–70.
  doi:10.1144/GSL.SP.1984.014.01.06.
- 944 Rolin, P., Colchen, M., 2001. Les cisaillements hercyniens de la Vendée au Limousin.
  945 Géologie de la France, n° 1-2. 87-116.
- Rolin, P., Marquer, D., Colchen, M., Cartannaz, C., Cocherie, A., Thiery, V., Quenardel, J.M., Rossi, P., 2009. Famenno-Carboniferous (370-320 Ma) strike slip tectonics
  monitored by syn-kinematic plutons in the French Variscan belt (Massif Armoricain and
  French Massif Central). Bulletin de la Société Géologique de France 180, 231–246.
  doi:10.2113/gssgfbull.180.3.231.
- 951 Tartèse, R., Boulvais, P., Poujol, M., Vigneresse, J.-L., 2011a. Granite petrogenesis
  952 revealed by combined gravimetric and radiometric imaging. Tectonophysics 501, 98–
  953 103. doi:10.1016/j.tecto.2011.02.003
- Tartèse, R., Poujol, M., Ruffet, G., Boulvais, P., Yamato, P., Košler, J., 2011b. New U-Pb
  zircon and 40Ar/39Ar muscovite age constraints on the emplacement of the Lizio syntectonic granite (Armorican Massif, France). Comptes Rendus Geoscience 343, 443–
  453. doi:10.1016/j.crte.2011.07.005

- Taylor, H.L., Mason, M.C., 1972. A Systematic Approach to Well Surveying Calculations.
  Society of Petroleum Engineers Journal 12, 474–488. doi:10.2118/3362-PA.
- 960 Théveniaut, H., Clarke, B., 2013. Large scale magnetic susceptibility soil mapping: a proxy
  961 for geological mapping and exploration from Bogoso (Ghana). Exploration Geophysics
  962 44, 48. doi:10.1071/EG12019.
- 963 Thiéblemont, D., Cabanis, B., Le Métour, J., 1987. Étude géochimique d'un magmatisme
  964 de distension intracontinentale: la série bimodale ordovico-silurienne du Choletais
  965 (Massif vendéen), Géologie de la France 1, 65–76.
- Thiéblemont, D., Guerrot, C., Le Métour, J., Jézéquel, P., 2001. Le complexe de CholetThouars: un ensemble volcano-plutonique cambrien moyen au sein du bloc
  précambrien des Mauges. Géologie de la France, n°1-2. 7-17.
- Thiéblemont, D., Augier, R., Ferry, J.-N., Laurent-Charvet, S., Maleyx, C., Ravoux, A.,
  Lebret, P., Guerrot, C., Chretien, P., 2011 Notice explicative, Carte géologique de.
  France (1/50 000), feuille Vihiers (511). Orléans : BRGM, 122 p.
- 972 Trautmann, F., Lacquement, F., Vernhet, Y., Pivette, B., avec la collaboration de Cocherie
  973 A., Guerrot C., Thiéblemont D., Tegyey M., Denis E., 2011. Notice explicative, Carte
  974 géologique de France (1/50 000), feuille Vitré (318). Orléans : BRGM, 131 p. Carte
  975 géologique par Lacquement F., Trautmann F., Vernhet Y. (2011).
- Truffert, C., Gumiaux, C., Chantraine, J., Perrin, J., Galdeano, A., Gapais, D., Ballèvre, M.,
  Asfirane, F., Guennoc, P., Brun, J.-P., 2001. Levé géophysique aéroporté dans le SudEst du Massif armoricain (programme GéoFrance3D Armor2). Magnétisme et
  radiométrie spectrale. Comptes Rendus de l'Académie des Sciences-Series IIA-Earth
  and Planetary Science 333, 263–270.

981 Turrillot, P., Augier, R., Faure, M., 2009. The top-to-the-southeast Sarzeau shear zone and
982 its place in the late-orogenic extensional tectonics of southern Armorica. Bulletin de la
983 Société Géologique de France 180, 247–261. doi:10.2113/gssgfbull.180.3.247

Venkateswarlu, N.B., Raju, P.S.V.S.K., 1992. Fast isodata clustering algorithms. Pattern
Recognition 25, 335–342. doi:10.1016/0031-3203(92)90114-X

- 986 Verduzco, B., Fairhead, J.D., Green, C.M., MacKenzie, C., 2004. New insights into
  987 magnetic derivatives for structural mapping. The Leading Edge 23, 116–119.
- 988 Vernhet, Y., Plaine, J., Trautmann, F., Pivette, B., avec la collaboration de Chevremont, P.,

Bourdillon, C., Cocherie, A., 2009 – Notice explicative, Carte géologique de France
(1/50 000), feuille Cossé-le-Vivien (355). Orléans : BRGM, 222 p. Carte géologique
par Vernhet, Y., Plaine, J., Trautmann, F., Clément J.P., 2009.

992 Vidal, P., 1980. L'évolution polyorogénique du Massif Armoricain : apport de la géochronologie et de la géochimie isotopique du strontium. Mémoire de la Société
994 géologique Minérale de Bretagne, Rennes, 21,162-958

- Vigneresse, J.-L., Brun, J.-P., 1983. Les leucogranites armoricains marqueurs de la deformation regionale; apport de la gravimetrie. Bulletin de la Société Géologique de
   France S7–XXV, 357–366. doi:10.2113/gssgfbull.S7-XXV.3.357
- Vigneresse, J.-L., 1988. La fracturation post-hercynienne du Massif armoricain d'après les
  données géophysiques. Géologie de la France 4, 3–10.
- Vigneresse, J.L., 1999. Intrusion level of granitic massifs along the Hercynian belt:
  balancing the eroded crust. Tectonophysics 307, 277–295. doi:10.1016/S00401951(99)00104-3
- 1003 Virlogeux, D., Roux, J. ets Guillemot, D., 1999. Apport de la géophysique à la connaissance
   1004 géologique du Massif de Charroux-Civray et du socle Poitevin. Dans « Etude du Massif

- 1005 de Charroux-Civray ». Actes Journées Sciences CNRS-ANDRA, Poitiers, 13-14.
  1006 10,33-61.
- Watts, M.S., and Williams, G.D., 1979. Faults rocks as indicators of progressive shear
  deformation in the Guingamp region, Brittany. Journal of Structural Geology, 1, 323332.
- Weber, C., 1971. Le socle antépermien sous la bordure sud du bassin de Paris d'après les
  données géophysiques. Bulletin BRGM, section I n°3, 177–189.
- 1012 Weber, C., 1973. Le socle antétriasique sous la partie sud du Bassin de Paris d'après les
  1013 données géophysiques. Bulletin BRGM, sect. II, n°3 et 4, 219-343.
- 1014 Wyns, R., Le Métour, J., 1983. Le Précambrien du massif vendéen. Etude détaillée de deux
  1015 coupes de référence (coupes de l'Evre et de la Divatte) et synthèse des données
  1016 récentes. Document BRGM, 68.
- 1017 Wyns, R., Lardeux, H., Moguedet, G., Duermael, G., Gruet, M., Biagi, R. avec la 1018 collaboration de Ballèvre, M., Chevremont, P., 1998. Notice explicative, carte 1019 géologique de la France (1/50 000), feuille Chemillé (483). BRGM, Orléans, 72 p.

1021



**Fig. 1**: Map of the Pre-Permian basement in France with Variscan sutures (modified from Faure, 2014). In dark grey: outcrops of the Variscan basement. NASZ: North Armorican Shear Zone, NBSASZ: Northern Branch of the South Armorican Shear Zone, SBSASZ: Southern Branch of the South Armorican Shear Zone, NSE F: Nort sur-Erdre Fault corresponding to the Eo-Variscan suture, PBMA: Paris Basin Magnetic Anomaly. In light grey: outcrop of the Meso-Cenozoic Paris Basin.



**Fig. 2**: Geological map of the southeastern part of the Armorican Massif and the northwestern part of the Massif Central: a) Major structures and litho-structural unit delimitations, b) Main lithologies within the litho-structural units (modified from Chantraine et al., 2003).



**Fig. 3**: **a)** Location map of the three aeromagnetic surveys used in this study. From west to east: Brittany, Pays de Loire (PaL), Région Centre. Crosses: location of the petrophysical field samples: in red: magnetic susceptibility only, in purple: magnetic susceptibility and density. **b)** Map of the magnetic anomaly reduced to the pole. **c)** Map of the magnetic vertical gradient. **d)** Map of the magnetic tilt derivative. Structural and lithological contours (thin black lines) and massif boundaries (thick black lines) are superimposed (modified from Chantraine et al., 2003).



**Fig. 4**: **a)** Map of the Bouguer anomaly. **b)** Map of the vertical gradient of the Bouguer anomaly. Structural and lithological contours (thin black lines) and massif boundaries (thick black lines) are superimposed (modified from Chantraine et al., 2003).



**Fig. 5**: Magnetic susceptibilities and average density values of the main lithologies sampled in the field, within each litho-structural unit. L: Low, I: Intermediate; H: High densities: refer to the text for explanations. In brackets: number of magnetic susceptibility measurements and number of density samples. Location of the sampling sites is documented in Fig. 3a.



Fig. 6: a) Map of synthetized geophysical signatures resulting from the classification of the magnetic (magnetic anomaly reduced to the pole and magnetic tilt derivative) and gravity (vertical gradient of the Bouguer anomaly) signatures. Magnetic and gravity trends are superimposed (red lines).
b) Interpretative structural sketch map of the Pre-Mesozoic substratum, of the southwestern part of the Paris Basin, with the location of the Permo-Carboniferous basins, the lithology of the studied outcrops (colored stars) and the lithology of the substratum in boreholes (colored circles).



**Fig. 7**: **a)** Interpretative geological map of the Pre-Mesozoic substratum of the southwestern part of the Paris Basin. This map synthesizes the information coming from the potential field data (aeromagnetic and gravity enhanced maps), the classified geophysical signatures, the structural sketch map and the petrophysical characteristics of rock samples. For clarity reasons, Permo-Carboniferous basins are not represented in this map. The thick black dotted line locates the cross section presented in Fig.7b. **b)** Interpretative geological cross-section of the Pre-Mesozoic substratum of the southwestern part of the Paris Basin inferred from the geophysical information processed in this study, and the extension of the structures known in the field.



**Fig. 8**: Decomposition of the geological map into its main lithological ensemble: **a**) Sedimentary rocks and volcanites; **b**) Permo-Carboniferous basins, using geophysical and seismic data (modified from Beccaletto et al., 2015); **c**) magmatic intrusions; **d**) metamorphic nappes. Blue lines : massif boundaries (modified from Chantraine et *al.*, 2003)