



**HAL**  
open science

## Laboratory measurements for the wavelength dependence of the linear polarization with the PROGRA2 instruments

Jean-Baptiste Renard, Edith Hadamcik, Anny Chantal Levasseur-Regourd, Nathalie Carrasco, Benoit Couté, Vincent Duverger, Vincent Guerrini

► **To cite this version:**

Jean-Baptiste Renard, Edith Hadamcik, Anny Chantal Levasseur-Regourd, Nathalie Carrasco, Benoit Couté, et al.. Laboratory measurements for the wavelength dependence of the linear polarization with the PROGRA2 instruments. EGU General Assembly 2017, Apr 2017, Vienna, Austria. pp. EGU2017-2909. insu-01516152

**HAL Id: insu-01516152**

**<https://insu.hal.science/insu-01516152>**

Submitted on 12 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

## Laboratory measurements for the wavelength dependence of the linear polarization with the PROGRA2 instruments

Jean-Baptiste Renard (1), Edith Hadamcik (2), Anny-Chantal Levasseur-Regourd (3), Nathalie Carrasco (4), Benoit Couté (1), Vincent Duverger (1), and Vincent Guerrini (1)

(1) LPC2E-CNRS, Orléans cedex 2, France (jbrenard@cnrs-orleans.fr), (2) LATMOS-IPSL/CNRS, Guyancourt, France, (3) LATMOS-IPSL / UMPC (Université Pierre et Marie Curie), Paris, France, (4) LATMOS-IPSL, UVSQ, Guyancourt, France

Clouds of solid particles are present in many regions of the Solar System (comets, interplanetary dust cloud, planetary atmospheres). These clouds can be remotely studied by the light they scatter. There is a need for a data base of the light scattered by a large variety of samples at different wavelengths to interpret such measurements. The PROGRA2 instruments are used for this purpose.

The PROGRA2 instruments, PROGRA2-VIS and PROGRA2-IR, are imaging polarimeters with a rotating arm to change the phase angle (angle between directions of illumination and observation). They allow to retrieve the complete polarization phase function between  $10^\circ$  and  $165^\circ$ . The light sources are at around 550 and 650 nm for PROGRA2-VIS, and 1000 and 1500 nm for PROGRA2-IR. The detectors are cameras, with a spatial resolution between 20 and 40 micrometers per pixel (PROGRA2-IR uses now new high sensitivity cameras). Measurements are conducted in the laboratory by an air draught technique for grains smaller than about 20 micrometers (which can be included in porous aggregates), and during parabolic flights on-board the A300 ZeroG and now the A310 ZeroG for larger grains; these flights campaigns are funded by the French and European Space Agencies. Hundreds of scattering phase functions have been obtained since 20 years, for a large number of samples (sands, silicon carbide, basalt, volcanic ashes, lunar and Martian simulants, tholins, meteoritic material, black carbon, carbonaceous compounds, . . .); the main results are available at [www.icare.univ-lille1.fr/progra2/](http://www.icare.univ-lille1.fr/progra2/).

Several samples have been already studied by the two instruments, showing a large diversity of wavelength dependences, from close-to-zero dependence for yellow and ocher sand grains to high dependence for silicon carbide and anthracite grains. These variations should be related to the wavelength dependence of the real and imaginary parts of the refractive index of the particles.

We present a summary of the main results of the wavelength dependence of polarization already obtained with PROGRA2. We present also the interest of conducting new measurements for meteoritic material, tholins, and carbonaceous particles that can be analogues of the cometary and interplanetary grains. Since PROGRA2 is well adapted to study the large (porous) particles, as those detected by Rosetta, these new measurements will be useful to tentatively relate the wavelength dependence of linear polarization detected within several cometary comae with the main composition and nature of the particles.

References : Renard et al., JQSRT, 146, 424-430, 2014 ; Hadamcik et al., Earth Planets Space, 65, 1175-1184, 2013.