



HAL
open science

The northwest-directed "Bretonian phase" in the French Variscan Belt (Massif Central and Massif Armoricain): A consequence of the Early Carboniferous Gondwana-Laurussia collision

Michel Faure, Xian-Hua Li, Wei Lin

► **To cite this version:**

Michel Faure, Xian-Hua Li, Wei Lin. The northwest-directed "Bretonian phase" in the French Variscan Belt (Massif Central and Massif Armoricain): A consequence of the Early Carboniferous Gondwana-Laurussia collision. *Comptes Rendus Géoscience*, 2017, 349, pp.126-136. 10.1016/j.crte.2017.05.002 . insu-01540215

HAL Id: insu-01540215

<https://insu.hal.science/insu-01540215>

Submitted on 4 Jul 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

1 The northwest-directed "Bretonian phase" in the French Variscan Belt (Massif
2 Central and Massif Armoricain): a consequence of the Early Carboniferous
3 Gondwana-Laurussia collision

4

5 Michel Faure¹, Xian-Hua Li^{2,3}, Wei Lin^{2,3}

6 1: Institut des Sciences de la Terre d'Orléans, CNRS-Université UMR 7327, Campus

7 Géosciences, 1A Rue de la Férollerie, 45071 Orléans Cedex 2, France

8 2: State Key Laboratory of Lithospheric Evolution, Institute of Geology and Geophysics,

9 Chinese Academy of Sciences, Beijing, China

10 3: College of Earth Sciences, University of Chinese Academy of Sciences, Beijing 100049,

11 China

12

13

14 **ABSTRACT**

15 In the Variscan French Massif Central and Massif Armoricain, the tectonic
16 significance of a widespread NW-SE trending stretching lineation, coeval with medium
17 pressure-medium temperature metamorphism, is an open question. Based on a structural
18 analysis in the Southern part of the Massif Central, we show that this top-to-the-NW shearing
19 is a deformation event, referred to as D2, that followed a D1 top-to-the-South shearing
20 Devonian phase, and was itself re-deformed by a Late D3 Viséan-Serpukhovian southward
21 thrusting event. We date the D2 phase at 360 Ma (Famennian-Tournaisian boundary). In the
22 Massif Armoricain, D2 is the "Bretonian phase" recorded in the metamorphic series and
23 sedimentary basins. Geodynamically, D2 is related to a general northwestward shearing
24 during the Laurussia-Gondwana collision, which occurred after the closure of the Rheic
25 Ocean, as indicated by the emplacement of the Lizard ophiolitic nappe in Britain. The left-

26 lateral Nort-sur-Erdre fault accommodated the absence of ductile shearing in Central
27 Armorica.

28

29 **Keywords:** Variscan belt, Bretonian phase, French Massif Central, Massif Armoricain

30

31 **1. Introduction**

32

33 It is widely acknowledged that the Variscan Belt of Western Europe formed as a result
34 of the multiple collisions that occurred between Gondwana to the South and Laurussia to the
35 North, from **Early Devonian to Carboniferous** times. These collisions stacked together
36 continents and micro-blocks such as Armorica or Mid-German Crystalline Rise that had been
37 previously drifted from Gondwana (e.g. Holder and Leveridge, 1986; Pin, 1990; Matte, 1986,
38 2001; Franke, 2000; Faure et al., 2005, 2008; Lardeaux et al., 2014). In the French Massif
39 Central (FMC, Fig. 1A), a polyphase syn-metamorphic deformation is well documented (e.g.
40 Faure et al., 2009 and enclosed references). By analogy with the Himalayas, it was initially
41 proposed that a ductile south-directed shearing, active from Late Devonian to Middle
42 Carboniferous accommodated the nappe stacking with younging from North to South (e.g.
43 Mattauer 1974; Mattauer and Etchecopar, 1977; Matte, 1986; Ledru et al., 1989). Top-to-the-
44 South shearing is indeed well established in the southern part of the French Massif Central
45 (Montagne Noire and Cévennes, Fig. 1) –where it is referred to as a “D3 event”, but more to
46 the North and West, in Lot, Rouergue, Lyonnais, Sioule and Limousin, the dominant structure
47 is instead a NW-SE stretching lineation with a top-to-the-NW shearing. The tectonic
48 significance of this NW-SE deformation and its relations with the widespread southward D3
49 shearing are not clear. According to Brun and Burg, (1982) and Burg et al., (1987), the top-to-
50 the-NW shearing would be due to a combination of southward thrusting and sinistral
51 wrenching during the ibero-armoricain oroclinal bending. Mattauer et al., (1988) considered

52 this deformation as a consequence of extensional tectonics, whereas Bouchez and Jover
53 (1986), Friedrich et al., (1988), Faure et al. (2005), and Bellot and Roig (2007) argued that it
54 corresponds to a nappe stacking. This event is referred to as the "D2 event" (Faure et al.,
55 2009).

56 Here we present structural and dating data in the Rouergue-Albigeois (Fig. 1B) area,
57 which allow us to discriminate and date the S- and NW-ward displacement events (D3 and
58 D2). We report similar observations in the Massif Armoricain. We propose a geodynamic
59 interpretation of the D2 event in the two massifs in the general framework of the Laurussia-
60 Gondwana continental collision.

61

62

63 **2. The architecture of the Southern Massif Central**

64 The French Massif Central stack of metamorphic nappes belongs to the northern
65 Gondwana margin. From top-to-bottom, the following units are recognized: 1) the Upper
66 Gneiss Unit, 2) the Lower Gneiss Unit, 3) the Para-Autochthonous domain, 4) the Fold-and-
67 Thrust Belt, and 5) the Southern Foreland. South of the Margeride pluton, the nappe
68 architecture exhibits several particularities, which we describe below (Figs. 1, 2).

69 *2.1 The Upper Gneiss Unit* (UGU) crops out as several klippes (Marvejols, Vibal,
70 Lévézou, Najac, and Decazeville). It is made of an association of felsic-mafic-ultramafic
71 rocks and sedimentary rocks, called the leptynite-amphibolite complex. This complex
72 experienced a high pressure-medium temperature (HP/MT) metamorphism (Nicollet and
73 Leyreloup, 1978; Burg et al., 1986, 1989; Bodinier and Burg, 1980-1981) at 415 ± 5 Ma
74 (U/Pb) or 408 ± 7 Ma (Sm/Nd), thus in Late Silurian-Early Devonian (Pin and Lancelot, 1982;
75 Paquette et al., 1995). During their exhumation, most of the eclogites were retrogressed in the
76 amphibolite facies and the melting of the Al-rich part produced migmatite. Due to its circular
77 shape in map view and the abundance of anatexites in its center, the southernmost Lévézou

78 klippe has long been considered as a diapiric dome (Collomb, 1970; Burg, 1987; Collomb and
79 Meyzindi, 1991). However, structural and gravimetric data show that the amphibolites and
80 migmatites are not rooted in the Lower Gneiss Unit but rather overlie it (Matte, 1986; Bayer
81 and Hirn, 1987). The UGU klippe have thus been displaced to the SW. As a matter of fact,
82 northwestward overturned folds deform the UGU foliations. As we explain below, they are
83 attributed to the D2 deformation phase.

84 2.2. *The Lower Gneiss Unit (LGU)* consists of metagreywackes and metapelites that
85 never experienced any HP/MT metamorphism. Several plutons that intrude the LGU, such as
86 the Caplongue diorite (557 ± 10 Ma, U/Pb on zircon, Lafon, 1984), the Rodez alkaline granite,
87 and the Pinet and Comps porphyritic monzogranites, have been transformed into augen
88 orthogneiss.

89 2.3. *The Para-autochthonous Unit (PAU)*, well exposed in Cévennes, consists of a
90 thrust sheet-imbrication of greenschist facies metapelites, quartzites, and metagrauwackes
91 with subordinate layers of conglomerate, felsic and mafic lava, and rare intrusions (Pin and
92 Marini, 1993; Caron, 1994). In the Albigeois area, the PAU has been subdivided into the S^t-
93 Sernin-sur-Rance and the S^t-Salvi-de-Carcavès nappes (Guéranzé-Lozes and Guéranzé, 1984;
94 Guéranzé-Lozes, 1987; Guéranzé-Lozes and Burg, 1990). The lithostratigraphy of these two
95 nappes is similar, with, from base to top, Cambrian greywacke, rhyolite and ignimbrite,
96 Ordovician white quartzite, and black metapelite. However, a biotite-garnet \pm staurolite
97 assemblage is fairly common in the former but is absent in the latter.

98 2.4. *The Fold-and Thrust Belt*, developed in the Montagne Noire and the Viganais, is
99 composed of unmetamorphosed fossiliferous Paleozoic sedimentary rocks terminated by a
100 Late Viséan to Serpukhovian turbiditic basin. The series have been deformed, by South-
101 directed thrust sheets and km-scale recumbent folds (Gèze, 1949; Arthaud, 1970; Engel et al.,
102 1980-81). The Montagne Noire axial zone is a migmatitic dome that cross-cuts the folds and
103 thrusts. This area is beyond the scope of this paper.

104

105

106 **3. Polyphase shearing in the Southern Massif Central**

107 The pervasive foliation developed in the UGU, LGU, and PAU units results from
108 three synmetamorphic ductile deformations, D1, D2, D3, each associated with specific
109 kinematics and P-T conditions. The three deformation phases are described below from the
110 youngest to the oldest.

111 *3.1. The D3 Carboniferous southward thrusting phase.* Spectacular km-scale,
112 southward overturned recumbent folds characterize the Paleozoic Fold-and-Thrust Belt (e.g.
113 Arthaud, 1970, Arthaud and Matte, 1977). In the Southern Montagne Noire, folding is coeval
114 with horizontal cleavage and rare N-S to N30E trending mineral and stretching lineation (Fig.
115 1). A more dominant lineation exists, that strikes NE-SW (Lee et al 1988; Echtler and
116 Malavielle, 1990). This NE microstructure is a late feature since it overprints the already
117 deformed series. The NE lineation is coeval with a HT/LP metamorphic event that has been
118 related to the doming of the axial zone (e.g. Courtillot et al., 1986; Echtler and Malavieille,
119 1990; Faure et al. 2014). In the Northern Montagne Noire and the S^t-Salvi-de-Carcavès
120 nappe, the submeridian stretching lineation is coeval with a top-to-the-S sense of shear
121 (Brunel et al., 1974; Guérangé-Lozes, 1987). In map view, the basal thrust of the D3 S^t-Salvi-
122 de-Carcavès nappe truncates the bedding surface (Figs. 1, 2). The D3 event recognized in the
123 PAU series in Cévennes and at the base of the UGU series in Marvejols is dated at ca 335-
124 325 Ma and 340-335 Ma, respectively (Pin and Lancelot, 1982; Caron, 1994; Faure et al.,
125 2001). Therefore, the D3 event is clearly younging southward. Moreover, the D3 structures
126 are locally overprinted by a top-to-the-north shearing related to the Late Carboniferous
127 extensional tectonics (Arnaud and Burg, 1993) that are not considered here (cf. Faure et al.,
128 2009 for details).

129 3.2. *The D2 Early Carboniferous top-to-the-NW shearing phase.* The main
130 deformation of the S^t-Sernin-sur-Rance nappe is characterized by a flat-lying foliation and a
131 NW-SE trending mineral and stretching lineation ascribed to the D2 phase. Near Naucelle
132 (Fig. 1), top-to-the NW shear criteria coeval with a biotite-garnet-staurolite metamorphism is
133 observed (Fig. 3C, E, F; Burg et al., 1986; 1989). A flat-lying mylonitic fabric with NW-SE
134 trending folds with axes parallel to the stretching lineation developed during the ductile
135 shearing that accommodated the tectonic superposition of the S^t-Sernin-sur-Rance nappe
136 upon the LGU (Fig. 3A, B). Further east, the dip of the foliation increases progressively to
137 become SW east of Naucelle, and vertical south of the Lévézou klippe. The kinematic
138 consistency from a top-to-the-NW shearing developed in the flat-lying foliation to dextral one
139 observed in vertical surfaces with subhorizontal stretching lineation suggests a wrench-thrust
140 structure. The D2 deformation phase is not recognized in the S^t-Salvi-de-Carcavès nappe and
141 farther south, but in contrast, it is clearly observed in the north in the LGU, and to a lesser
142 extent, in the UGU units.

143 3.3. *The D1 early southwestward shearing phase.* The D2 top-to-the-NW shearing
144 event was preceded by an earlier deformation phase that is well observed in the UGU and
145 LGU series, in the S^t-Sernin-sur-Rance and S^t-Salvi-de-Carcavès nappes in the Rouergue
146 area. Where the units are not deformed by the later D2 and D3 events, a N-S to N30°E
147 trending stretching lineation is observed, associated with intrafolial folds and top-to-the-SW
148 shearing (Figs. 1, 3D). Yet, at many places, the D2 NW-SE trending folds deformed this early
149 mineral lineation (Fig. 3A,B D). In the S^t-Sernin-sur-Rance nappe, km-scale S-verging
150 recumbent folds appear re-deformed by the D2 deformation, and hence are attributed to the
151 D1 phase (Guérangé-Lozes, 1987). The D1 top-to-the-S stretching lineation is observed in the
152 Rodez orthogneiss of the LGU series, and in the UGU gneiss, amphibolite and migmatite of
153 the Lévézou and Vibal klippen.

154 The foliated Pinet orthogneiss has recorded both the D1 and the D2 deformation
155 events. Although the D2 NW-SE lineation is dominant in many places (Fig. 1, Burg and
156 Teyssier, 1983), an older N-S trending D1 mineral lineation is observed where the D2
157 overprint is lacking. Petro-structural analysis documents a top-to-the-south shearing coeval
158 with a high-temperature deformation, whereas low-temperature top-to-the-NW shear bands
159 are ascribed to the D2 phase (Fig. 3D, Duguet and Faure, 2004).

160

161

162 **4. Chronological constraints**

163 *4.1 Timing of the polyphase shearing in the Southeast Massif Central.* Due to their
164 similar submeridian trend and top-to-the-South sense of shear, the D1 and D3 deformation
165 events have been so far considered as the result of a single, long duration intracontinental
166 shearing period, operating during the "Himalayan thrusting" (Mattauer, 1974, Mattauer and
167 Etchecopar, 1977). However, we have shown that the D1 and D3 events are diachronous
168 since they are separated in time by nearly 40 Ma. The age of the D3 event is well constrained
169 as this event deforms the Late Viséan-Early Serpukhovian (330-325 Ma) syntectonic flysch
170 basin of the Southern Montagne Noire. Muscovite populations from the Mendic orthogneiss
171 and the S^t-Salvi-de-Carcavès nappe metasandstone yield ⁴⁰Ar/³⁹Ar ages of 330±3 Ma and
172 333±4 Ma, respectively (Costa, 1990). These Viséan ages are similar to those available in the
173 Cévennes area (Caron, 1994; Najoui et al., 2000).

174 The biotites, muscovites and amphiboles aligned along the NW-SE lineation reveal
175 ⁴⁰Ar/³⁹Ar ages in the range 354-346 Ma. Since the closure temperature of micas is lower than
176 that of the metamorphic climax, the above ages provide the minimum age of the D2 event. In
177 the S^t-Sernin-sur-Rance nappe, muscovites from quartzites yield a 345±3.5 Ma age. In the
178 UGU and LGU series, biotites, muscovites and amphiboles yield consistent ⁴⁰Ar/³⁹Ar ages in

179 the range 354-346 Ma (Costa, 1990). The age of the D1 event is still poorly constrained with
180 only one $^{40}\text{Ar}/^{39}\text{Ar}$ muscovite age of 380 ± 10 Ma (Guérangé-Lozes, 1987).

181 Whatever the precise absolute age of the three D1, D2 and D3 events, their relative
182 timing allows us to draw an interpretative general cross-section (Fig. 2). The top-to-the-SW
183 D1 structures were folded in the Late Devonian-Early Carboniferous by the northwestward
184 D2 deformation. Later, during the Middle Carboniferous, the D3 southward shearing event re-
185 deformed the whole stack. This new interpretation makes the Pinet orthogneiss a pre-D1
186 pluton.

187 *4.2. New zircon U-Pb age of the Pinet orthogneiss.* The Pinet orthogneiss is a
188 monzogranite or syenogranite foliated in a variable way, that includes both nearly
189 undeformed facies at local spots, and mylonitic to ultramylonitic shear zones at other spots. A
190 detailed petro-structural study of the Pinet orthogneiss has revealed that the pluton
191 experienced two deformation phases, ascribed to the D1 and D2 events (Duguet and Faure,
192 2004). The early deformation phase is coeval with a pervasive post-solidus planar fabric and a
193 N-S striking lineation. The second deformation phase is characterized by a local foliation
194 cross-cutting the previous one, and a NW-SE striking lineation with top-to-the-NW shearing.
195 Zircon populations in the granite yielded a TIMS U-Pb age of $ca\ 360\pm 20$ Ma (Pin, 1981),
196 whereas a biotite single grain was dated at 346 ± 7 Ma using the $^{40}\text{Ar}/^{39}\text{Ar}$ method (Maluski
197 and Monié, 1988).

198 We separated zircon concentrates from two samples (FR 41 and FR 42) of the
199 porphyritic orthogneiss, located along the Tarn river, at $44^\circ\ 03'\ 16.27''/002^\circ\ 45'\ 56.35''$ and
200 $44^\circ\ 02'\ 43.57''/002^\circ\ 46'\ 47.70''$, respectively. We applied analytical procedures and data
201 reduction as in Li et al. (2009) and Do Couto et al. (2016). Cathodoluminescence images of
202 the zircon grains show a well developed oscillatory zoning representative of a magmatic
203 origin of the grains (Fig. 5).

204 Zircons from the two rock samples yield consistent Ordovician ages of 477 ± 5 Ma and
205 475 ± 4 Ma (Fig. 5); these ages are interpreted as dating the emplacement of the porphyritic
206 granite. This new result shows that the Pinet orthogneiss is not a syn-kinematic pluton
207 emplaced during the D1 or D2 events, but instead formed in the Early Ordovician pre-
208 orogenic magmatism phase that is widespread in the Variscan belt. The ca 350 Ma $^{40}\text{Ar}/^{39}\text{Ar}$
209 age of post-folial biotite corresponds to the D2 event.

210

211

212 **5. Discussion-conclusion**

213 *5.1. Significance of the three D1, D2 and D3 tectono-metamorphic events.* The D1
214 event is recognized in the Lyonnais, Sioule, Limousin, and Plateau d'Aigurande areas of the
215 Massif Central. It is coeval with the Middle Devonian (385-375 Ma) migmatization, that
216 was interpreted as the result of the exhumation of the HP rocks of the UGU series (cf Faure et
217 al., 2008 for details).

218 The Late Devonian-Early Carboniferous (360-350 Ma) D2 event with a top-to-the-
219 NW shearing is also recognized in many places of the Massif Central (Fig. 6). In the LGU
220 series of Marvejols (Fig. 1), biotites from metadiorite and micaschist, aligned along the NW-
221 SE lineation, yield consistent $^{40}\text{Ar}/^{39}\text{Ar}$ ages of 352 ± 2 and 351 ± 3 Ma (Costa, 1990). The
222 Limousin, Sioule and Plateau d'Aigurande series are affected by a consistent medium
223 temperature-medium pressure metamorphism coeval with a top-to-the-NW shearing (e.g.
224 Floc'h 1983; Bouchez and Jover, 1986; Friedrich et al., 1988; Faure et al., 1990, 1993, 2005;
225 Roig and Faure, 2000; Bellot and Roig, 2007). Since the $^{40}\text{Ar}/^{39}\text{Ar}$ dates are very sensitive to
226 the temperature, the monazite U-Th-Pb chemical chronometer that records high temperature
227 events has also been carried out in the UGU, LGU, and PAU metapelites of Limousin and
228 Sioule areas (Melleton et al., 2009; Do Couto et al., 2016). The 360-350 Ma ages yielded by
229 the syn-D2 metamorphic monazites confirm the Famennian-Tournaisian age of the D2 event.

230 Furthermore, in eastern Massif Central, the Brévenne ophiolitic nappe was emplaced from SE
231 to NW under amphibolite facies conditions, before the unconformable deposition of the Early
232 Visean (i.e. 345 Ma) Le Goujet, terrigenous formation (Leloix et al., 1999). Thus the
233 Brévenne ophiolitic nappe formed also during the D2 event.

234 The D3 event is identified only in the South Massif Central. The Visean to
235 Serpukhovian southward migration from middle to shallow crustal levels is acknowledged
236 since a long time as the consequence of the post-collisional nappe stacking (Matte, 1986;
237 Ledru et al., 1989, Faure et al., 2009).

238

239 *5.2. Geodynamic significance of the D2 event in the French Variscan segment.* In the
240 southern Massif Armoricaïn that belongs to the metamorphic zone of the Variscan chain, the
241 D2 synmetamorphic shearing event is recognized in Vendée and south Brittany coast (Burg,
242 1981; Cannat and Bouchez, 1986; Fig. 6). There, an E-W to NW-SE striking mineral lineation
243 develops on a flat-lying foliation, coeval with a MP/MT metamorphism characterised by a
244 biotite-garnet-staurolite assemblage. Top-to-the-NW shearing shown by asymmetric pressure
245 shadows and shear bands is conspicuous.

246 More to the North, in Central Armorica, top-to-the-NW shearing is described only in
247 the S^t-Georges-sur-Loire and Lanvaux units (Bouchez and Blaise, 1976; Diot et al., 1983;
248 Cogné et al., 1983; Faure and Cartier, 1998; Cartier et al., 2001; Cartier and Faure, 2004).
249 The pre-Visean age of the deformation is attested by the age of the terrigenous deposits of the
250 Ancenis basin that did not record the D2 event. Therefore, the top-to-the-NW shearing, that is
251 widespread in the Massif Armoricaïn and South Massif Central, is a major feature of the
252 Variscan orogeny.

253 The MP/MT D2 metamorphism dated at ca 360 Ma, is coeval with the Devonian-
254 Carboniferous "Bretonian phase" long recognized in the central part of the Massif
255 Armoricaïn, through the Tournaisian unconformity and the widespread erosion of the Late

256 Devonian formations (e.g. Stille, 1929 in Rolet, 1982). In the 70's and 80's, on the basis of the
257 gentle folding of the pre-Tournaisian formations, and the small unconformity angle (lower
258 than 20°), the Bretonian phase was considered as a km-scale upright folding coeval with
259 vertical movements, responsible for erosion and clastic sedimentation in the Laval and
260 Châteaulin basins (e.g. Cogné 1965, 1974; Pelhâte, 1971; Paris et al., 1982; Houlgatte et al.,
261 1988, Le Gall et al. 1992; Fig. 6). However, recumbent folds, ductile shear zones, and low
262 angle thrust faults identified in the Brest area (Rolet et al., 1986) allow us to reassess the
263 importance of the Bretonian phase.

264 The Variscan orogeny is the consequence of the closure of the Rheic ocean, and
265 subsequent collision of Laurussia that was driven by a southward oceanic and then
266 continental subduction below Gondwana. The Rheic suture is today hidden below the English
267 Channel, but the Lizard ophiolitic nappe of SW Britain that overthrusts to the NNW the
268 Laurussia foreland at ca 360 Ma is a remnant of the Rheic Ocean (e.g. Holder and Leveridge,
269 1986; Le Gall and Darboux, 1986; Sandeman et al., 1994). Therefore we argue here that the
270 ductile and synmetamorphic, top-to-the-NW, shearing observed in the French Massif Central
271 and South Armorica, and more broadly the Famennian-Tournaisian deformation ascribed to
272 the Bretonian phase, are the result of the collision between Laurussia and Gondwana. **Top-to-**
273 **the NW shearing observed at the microtectonic scale results of the Variscan collision.**

274 Although close to the Rheic suture, (Fig. 6), the syn-metamorphic ductile deformation
275 does not exist in Central Armorica. A possible explanation would be to consider the
276 contrasted mechanical behaviour between the Massif Central - South Armorican domain with
277 a widespread D2 event, and the Central Armorican domain where D2 is absent. Due to its
278 crustal rigidity, possibly related to the Neoproterozoic Cadomian orogeny, and the thick sub-
279 continental mantle lithosphere, the Central Armorican domain behaved as a rigid plate
280 characterized by sinistral wrenching, responsible for the opening of the Châteaulin and Laval
281 basins (Fig. 7). In its southern margin, along the Nort-sur-Erdre strike-slip fault, and the S^t-

282 Georges-sur-Loire and Lanvaux units, the sinistral shearing accommodated the decoupling
283 between the ductile and brittle domains (Cogné et al., 1983; Diot et al., 1983; Lardeux and
284 Cavet, 1994; Faure et al., 1997; 2005; Cartier et al., 2001; Cartier and Faure, 2004). This
285 interpretation, which is at variance to the previous geodynamic models (e.g. Ballèvre et al.,
286 2009), provides a satisfactory solution for the significance of the D2 event in the French
287 Variscan orogen.

288

289

290 **Acknowledgements**

291 This article synthesizes numerous works conducted since 20 years, in the framework
292 of PhD and Master theses supported by ISTO and BRGM. The dating of the Pinet orthogneiss
293 has been funded by the National Natural Science Foundation of China (grants 41273070,
294 41472193, 41225009), and the Ministry of Science and technology (MOST) of China
295 (projects 2016YFC0600401, and 2016YFC0600102). Scientific and Editorial comments by I.
296 Manighetti to improve the early version of this paper are greatly acknowledged.

297

298 **References**

299 Arnaud, F., Burg J-P, 1993. Microstructures des mylonites schisteuses: cartographie des
300 chevauchements varisques dans les Cévennes et détermination de leur cinématique. C. R.
301 Acad. Sci., Paris 317, 1441-1447.

302

303 Arthaud, F. 1970. Etude tectonique et microtectonique comparée de deux domaines
304 hercyniens: les nappes de la Montagne Noire (France) et l'anticlinorium de l'Iglesiente
305 (Sardaigne). Thèse d'Etat, Univ. de Montpellier, 175 pp.

306

- 307 Arthaud, F., Matte, P., 1977. Synthèse provisoire sur l'évolution tectonique et les raccords
308 entre les segments hercyniens situés autour du bassin nord-Baléare (sud de la France,
309 Espagne, bloc corso-sarde). In La Chaîne Varisque d'Europe moyenne et occidentale, coll.
310 intern. CNRS 243, 497-513.
311
- 312 Ballèvre, M., Bosse, V., Ducassou, C., Pitra P., 2009. Palaeozoic history of the Armorican
313 Massif: Models for the tectonic evolution of the suture zones. C. R. Geoscience 341, 174-201.
314
- 315 Bayer, R., Hirn, A., 1987. Données géophysiques sur la structure profonde de la croûte
316 hercynienne dans l'arc ibéro-armoricain et le Massif Central français. Bull. Soc. Géol. France
317 8, 561-574.
318
- 319 Bellot, J-P, Roig, J-Y. 2007. Episodic exhumation of HP rocks inferred from structural data
320 and P-T paths from the southwestern part Massif Central (Variscan belt, France). J. Struct.
321 Geol. 29,1538–1557.
322
- 323 Bodinier, J.L., Burg, J.P., 1980-1981. Evolution métamorphique et tectonique des séries
324 cristallophyliennes du Rouergue occidental : mise en évidence d'un chevauchement dans la
325 région de Najac (Aveyron). Bull. BRGM 2, 315-339.
326
- 327 Bouchez, J-L., Blaise, J., 1976. Une structure hercynienne liée à un accident ductile:
328 l'anticlinal de Lanvaux-Les-Ponts-de-Cé, aux environs d'Angers (Massif Central français).
329 Bull. soc. Géol. France 7, 145-157.
330
- 331 Bouchez, J-L., Jover, O., 1986. Le Massif central: un chevauchement de type himalayen vers
332 l'Ouest-Nord-ouest. C. R. Acad. Sci. Paris 302, 675-680.

333

334 Brun, J-P., Burg, J-P., 1982. Combined thrusting and wrenching in the Ibero-Armorican arc: a
335 corner effect during continental collision. *Earth Planet Sci. Letters* 61, 319-332.

336

337 Brunel, M., 1974. La phase hercynienne majeure dans le versant nord de la Montagne Noire
338 (Est de Lacaune, Massif Central français). *C. R. Acad. Sciences, Paris* 278, 991-994.

339

340 Burg, J-P., 1981. Tectonique tangentielle en Vendée littorale: signification des linéations
341 d'étirement E-W dans les porphyroïdes à foliation horizontale. *C. R. Acad. Sciences, Paris*,
342 293 849-854.

343

344 Burg, J.P., 1987. Regional shear variation in relation to diapirism and folding. *J. Struct. Geol.*
345 9, 925-934.

346

347 Burg, J.P., Teyssier, C., 1983. Contribution à l'étude tectonique et microtectonique des séries
348 crystallophylliennes du Rouergue oriental: La déformation des laccolites syntectoniques, type
349 Pinet. *Géologie de la France* 1, 3-30.

350

351 Burg, J.P., Delor, C., Leyreloup, A., 1986. Le massif du Lévézou et les séries adjacentes du
352 Rouergue oriental. Nouvelles données pétrographiques et structurales. *Géologie de la France*,
353 3 229-272.

354

355 Burg, J-P., Balé, P., Brun, J.-P., Girardeau, J., 1987. Stretching lineation and transport
356 direction in the Ibero-Armorican arc during the Siluro-Devonian collision. *Geodin. Acta* 1,
357 71-87.

358

- 359 Burg, J.P., Delor C., Leyreloup, A., Rommey, F., 1989. Inverted metamorphic zonation and
360 Variscan thrust tectonics in the Rouergue area (Massif Central, France): P-T-t record from
361 mineral to regional scale. In: Evolution of metamorphic belts Daly, J.S., Cliff, P., and B.W.D.
362 Yardley, B.W.D. (Eds). Geol. Soc. London Spec. Publ. 43, 423-439.
363
- 364 Cannat, M., Bouchez, J-L., 1986. Linéations N-S et E-W en Vendée littorale (Massif
365 Armoricaïn). Episodes tangentiels succesifs éo-hercyniens en France occidentale. Bull. Soc.
366 Géol. France 8, 299-310.
367
- 368 Caron, C., 1994. Les minéralisations Pb-Zn associées au Paléozoïque inférieur d'Europe
369 méridionale. Traçage isotopique Pb-Pb des gîtes de l'Iglesiente (SW Sardaigne) et des
370 Cévennes et évolution du socle encaissant par la géochronologie U-Pb, 40Ar-39Ar et K-Ar.,
371 Thèse, Univ. Montpellier II, 226 pp.
372
- 373 Cartier, C., Faure, M., 2004. Structure and Geodynamic evolution of the Gondwana-Armorica
374 boundary in the Ligerian domain (Armorican massif, France), Int. J. Earth Sci. 93, 945-958.
375
- 376 Cartier, C., Faure, M., Lardeux, H., 2001. The Hercynian orogeny in the South Armorican
377 Massif (France): Rifting and welding of continental stripes, Terra Nova 13, 143 – 149.
378
- 379 Cogné J. 1965. Observations sur l'âge et la signification de la Phase bretonne. C. R. session
380 extraordinaire Bull. Soc. Belge Géol. Pal. Hydro. 73, 239-243.
381
- 382 Cogné, J. 1974. Le Massif Armoricaïn. In: Debelmas, J. (Ed.), Géologie de la France, Doin,
383 Paris, France, pp. 105-161.
384

- 385 Cogné, J-P., Choukroune, P., Cogné, J., 1983. Cisaillements varisques superposés dans le
386 massif de Lanvaux (Bretagne centrale). C. R. Acad. Sci. Paris 296, 773-777.
387
- 388 Collomb, P., 1970. Etude géologique du Rouergue cristallin. Mem. Serv. Carte Géol. France,
389 Paris, 419 pp.
390
- 391 Collomb, P., Meyzindi, C., 1991. Données lithostratigraphiques et structurales nouvelles sur
392 les confins du Rouergue et de l'Albigeois cristallins (Sud-Ouest du Massif Central français).
393 C. R. Acad. Sci. Paris 312, 509-515.
394
- 395 Costa, S., 1990. De la collision continentale à l'extension tardi-orogénique: 100 Ma d'histoire
396 varisque dans le Massif Central français. Une étude chronologique par la méthode $^{40}\text{Ar}/^{39}\text{Ar}$.
397 Thèse Univ. Montpellier 442 pp.
398
- 399 Courtillot, V., Chambon, P., Brun, J-P., Rochette, P., 1986. A magnetotectonic study of the
400 Hercynian Montagne Noire (France). Tectonics 5, 733-751.
401
- 402 Diot, H., Bouchez, J.L., Blaise, J. 1983. La bande ductile Lanvaux-Les-Ponts-de-Cé, entre
403 Redon et Angers. Une composante du cisaillement hercynien du Massif Armoricaïn (France).
404 Bull. Soc. Géol. France 25, 155-167.
405
- 406 Do Couto, D., Faure, M., Augier, R., Cocherie, A., Rossi, P., Li, X.H., Lin, W., 2016.
407 Monazite U-Th-Pb EPMA and zircon U-Pb SIMS chronological constraints on the tectonic,
408 metamorphic, and thermal events in the inner part of the Variscan orogen, example from the
409 Sioule series, French Massif Central. Inter. J. Earth Sci. 105 557-579.
410

- 411 Duguet, M., Faure, M. 2004. Granitoid emplacement during a thrusting event: structural
412 analysis, microstructural and quartz-c-axis patterns. An example from Hercynian plutons in
413 the French Massif Central. *J. Struct. Geol.* 26, 927-945.
- 414
- 415 Echtler, H., Malavieille, J., 1990. Extensional tectonics, basement uplift and Stephano-
416 Permian collapse basin in a late Variscan metamorphic core complex (Montagne
417 Noire, southern Massif Central). *Tectonophysics* 17, 125-138.
- 418
- 419 Engel, W., Feist, R., Franke, W., 1980-81. Le Carbonifère anté-stéphanien de la Montagne
420 Noire : rapports entre mise en place des nappes et sédimentation. *Bull. BRGM* 2, 341-389.
- 421
- 422 Faure, M., Prost, A., Lasne, E., 1990. Déformation ductile extensive d'âge namuro-
423 westphalienne dans le plateau d'Aigurande, Massif Central français. *Bull. Soc. Géol. France*
424 161, 189-197.
- 425
- 426 Faure, M., Grolier, J., Pons, J., 1993. Extensional ductile tectonics in the Sioule metamorphic
427 series (Variscan French Massif Central), *Geol. Rundschau* 82, 461-474.
- 428
- 429 Faure, M., Leloix, C., Roig, J-Y., 1997. L'évolution polycyclique de la chaîne hercynienne.
430 *Bull. Soc Géol. France* 168, 3-14.
- 431
- 432 Faure, M., Cartier, C., 1998. Déformations ductiles polyphasées dans l'antiforme
433 orthogneissique de St-Clément-de-la-Place (unité de Lanvaux, Massif Armoricaïn). *C. R.*
434 *Acad. Sci, Paris* 326, 795-802.
- 435

- 436 Faure, M., X. Charonnat, A. Chauvet, Y. Chen, J-Y. Talbot, G. Martelet, G. Courrioux, P.
437 Monié, J-P. Milesi. 2001. Tectonic evolution and ore bearing fluids circulation in the
438 Cévennes para-autochthonous domain of the Hercynian Belt (French Massif Central). Bull.
439 Soc. Géol. France 172, 687-696.
440
- 441 Faure, M., Bé Mézème, E., Duguet, M., Cartier, C., Talbot, J-Y., 2005. Paleozoic tectonic
442 evolution of medio-Europa from the example of the French Massif Central and Massif
443 Armoricain, In: Carosi, R. Dias, R., Iacopini, D., Rosenbaum, G. (Eds.), The southern
444 Variscan belt, J. Virtual Explor. 19, paper 5.
445
- 446 Faure, M., Bé Mézème, E., Cocherie, A., Rossi, P., Chemenda, A. Boutelier, D., 2008.
447 Devonian geodynamic evolution of the Variscan Belt, insights from the French Massif
448 Central and Massif Armoricain, Tectonics 27.
449
- 450 Faure, M., Lardeaux, J.M., Ledru, P., 2009. A review of the pre-Permian geology of the
451 French Massif Central. C. R. Géoscience 341, 202–213.
452
- 453 Faure, M., Cocherie, A., Gaché, J., Esnault, C., Guerrot, C., Rossi, P., Lin, W., Li, Q., 2014.
454 Middle Carboniferous intracontinental subduction in the Outer Zone of the Variscan belt
455 (Montagne Noire Axial Zone, French Massif Central): multimethod geochronological
456 approach of polyphase metamorphism. In: Schulmann, K., Martínez Catalán, J.R., Lardeaux,
457 J.M., Janoušek, V., Oggiano, G. (Eds.), The Variscan orogeny: extent, timescale and the
458 formation of the European crust. Geol. Soc. London, Spec. Publ. 405, pp. 289–311.
459
- 460 Floc'h, J-P., 1983. La série métamorphique du Limousin central. Thèse d'Etat, Univ. Limoges,
461 455 pp.

462

463 Franke, W., 2000. The Mid-European segment of the Variscides: tectonostratigraphic units,
464 terrane boundaries and plate tectonic evolution. *Geol. Soc. London Spec. Publ.* 179, 35–61.

465

466 Friedrich, M., Marignac, C. and Floc'h, J-P. 1988. Sur l'existence de trois chevauchements
467 ductiles "himalayens" successifs à vergence NW en Limousin. *C. R. Acad. Sci. Paris* 306,
468 663-669.

469

470 Gèze, B., 1949. Etude géologique de la Montagne Noire et des Cévennes méridionales.

471 *Mém. Soc. Géol. Fr.* 62, 1–215.

472

473 Guérangé-Lozes, J., 1987. Les nappes varisques de l'Albigeois cristallin. Lithostratigraphie,
474 volcanisme et déformations. *Doc. Bureau Recherche Géol. Min.*, 135, BRGM ed., Orléans,
475 259 pp.

476

477 Guérangé-Lozes, J., Guérangé, B., 1984. Les nappes de charriage de l'Albigeois cristallin
478 (Tarn et Aveyron, France). *C. R. Acad. Sci.*, Paris 299, 471-474.

479

480 Guérangé-Lozes, J., Burg, J-P. 1990. Les nappes varisques du sud-ouest du Massif Central.
481 *Géologie de la France* 3-4, 71-106.

482

483 Holder, M. and Leveridge, B., 1986. A model for the tectonic evolution of South Cornwall. *J.*
484 *Geol. Soc.*, London 143, 125-134.

485

486 Houlgatte, E., Le Hérisse, A., Pelhâte, A., J. Rolet, J. 1988. Evolution géodynamique du bassin
487 carbonifère de Laval. *Géol. de la France* 1, 27-46.

488

489 Lafon, J.M., 1984. La granodiorite de Caplongue nouveau témoin d'un magmatisme cambrien
490 dans le Rouergue oriental. C. R. Acad. Sci. Paris, 298, 595-600.

491

492 Lardeaux, J.M., Schulmann, K., Faure, M., Janousek, V., Lexa, O., Skrzypek, E., Edel, J.-B.,
493 Stipska, P., 2014. The Moldanubian Zone in French Massif Central, Vosges/Schwarzwald and
494 Bohemian Massif revisited: differences and similarities. In: Schulmann, K., Martínez Catalán,
495 J.R., Lardeaux, J.M., Janoušek, V., Oggiano, G. (Eds.), The Variscan Orogeny: Extent,
496 Timescale and the Formation of the European Crust, Geol. Soc. London Spec. Publ., 405, pp.
497 7-44.

498

499 Lardeux, H., Cavet, P., 1994. Paleozoic of the Ligérien domain. In: J. D. Keppie, J.D. (Ed.),
500 Pre-Mesozoic Geology in France and related areas, Springer-Verlang, Berlin, pp. 152-156.

501

502 Lee, B-J., Faure, M., Cluzel, D., Cadet, J-P., 1988. Mise en evidence d'un cisaillement ductile
503 d'Ouest en Est dans les nappes du versant sud de la Montagne Noire (sud du Massif central).
504 C. R. Acad. Sciences 306, II, 455-462.

505

506 Le Gall, B., Darboux, J-R., 1986. Variscan strain pattern in the Paleozoic series at the Lizard
507 front, SW England, Tectonics 5, 589-606.

508

509 Le Gall, B., Loboziak, S., Le Hérisse, A., 1992. Le flanc sud du synclinorium carbonifère de
510 Châteaulin (Massif armoricain, France): une bordure de bassin reactive en context décro-
511 chevauchant. Bull. Soc. Géol. France 163, 13-26.

512

- 513 Ledru, P., Lardeaux, J-M., Santallier, D., Autran, A., Quenardel, J-M., Floc'h J-P., Lerouge,
514 G., Maillet, N., Marchand, J., Ploquin, A., 1989. Où sont les nappes dans le Massif Central
515 français? Bull. Soc. Géol. France 8, 605-618.
516
- 517 Leloix, C., Faure, M., Feybesse, J-L., 1999. Hercynian polyphase tectonics in the northeast
518 French Massif Central: the closure of the Brévenne Devonian-Dinantian rift. Int. J. Earth
519 Sciences 88, 409-421.
520
- 521 Li, X.H., Liu, Y., Li, Q.L., Guo, C.H., Chamberlain, K.R., 2009. Precise determination of
522 Phanerozoic zircon Pb/Pb age by multi-collector SIMS without external standardization.
523 Geochem. Geophys. Geosyst. 10, Q04010, doi.org/10.1029/cGC002400.
524
- 525 Maluski, H., Monié, P., 1988. $^{39}\text{Ar}/^{40}\text{Ar}$ laser probe multi-dating inside single biotites of a
526 Variscan orthogneiss (Pinet, Massif Central, France). Chem. Geol. 73, 245-263.
527
- 528 Mattauer, M., 1974. Existe t-il des chevauchements de type himalayen dans la chaîne
529 hercynienne du Sud de la France? 2^o R.A.S.T., Pont à Mousson, Abstract vol. 279.
530
- 531 Mattauer, M., and Etchecopar, A., 1977. Argument en faveur de chevauchements de type
532 himalayen dans la chaîne hercynienne du Massif Central français, Coll. Int. C.N.R.S. 268,
533 261-267.
534
- 535 Mattauer, M., Brunel, P. Matte. 1988. Failles normales ductiles et grands
536 chevauchements: Une nouvelle analogie entre l'Himalaya et la chaîne hercynienne du Massif
537 Central français, Comptes R. Acad. Sci. Paris, Ser. 2, 306, 671-676.
538

- 539 Matte, P., 1986. La chaîne varisque parmi les chaînes paléozoïques péri-atlantiques, modèle
540 d'évolution et position des grands blocs continentaux au Permo-Carbonifère. Bull. Soc. Géol.
541 France 8, 9-24.
542
- 543 Matte, P., 2001. The Variscan collage and orogeny (480–290 Ma) and the tectonic definition
544 of the Armorica microplate. Terra Nova 13, 122–128.
545
- 546 Melleton, J., Faure, M., Cocherie A. 2009. Monazite U-Th/Pb chemical dating of the Early
547 Carboniferous syn-kinematic MP/MT metamorphism in the Variscan French Massif Central.
548 Bull. Soc. Géol. France 180, 283-292.
549
- 550 Najoui, K., Leyreloup, A. and Monié, P., 2000. Conditions et âges $^{40}\text{Ar}/^{39}\text{Ar}$ de mise en place
551 des granitoïdes de la zone externe sud du massif central français : exemple des granodiorites
552 de St-Guiral et du Liron (Cévennes, France). Bull. Soc. Géol. France 171, 495-510.
553
- 554 Nicollet, C., Leyreloup, A., 1978. Pétrologie des niveaux trondjémittiques de haute pression
555 associés aux éclogites et amphibolites des complexes leptyno-amphiboliques du Massif
556 Central. Can. J. Earth Sci. 15, 696-707.
557
- 558 Paquette, J-L, Monchoux, P., Couturier, M. 1995. Geochemical and isotopic study of a norite-
559 eclogite transition in the European variscan Belt: Implications for U-Pb zircon systematics in
560 metabasic rocks. Geochimica et Cosmochimica Acta 59, 1611-1622.
561
- 562 Paris, F., Le Hérisse, A., Pelhâte, A., Weyant, M., 1982. Les formations carbonifères et la
563 phase bretonne dans le synclinorium du Menez-Belair: essai de synthèse. Bull. Soc. géol.
564 minéral., Bretagne 14, 19-33.

565

566 Pelhâte; Y., 1971. Le Carbonifère inférieur du bassin de Laval, Massif Armoricaïn. Mem.
567 Soc. géol. minéral. Bretagne 15, 315 pp.

568

569 Pin, C., 1981. Old inherited zircons in two synkinematic variscan granitoids: the "granite du
570 Pinet" and the "orthogneiss de Marvejols" (southern French Massif Central). N. Jb. Miner.
571 Abh. 142, 27-48.

572

573 Pin, C., 1990. Variscan oceans: ages, origins and geodynamic implications inferred from
574 geochemical and radiometric data. Tectonophysics 177, 215-227.

575

576 Pin, C., Lancelot, J., 1982. U-Pb dating of an early Paleozoic bimodal magmatism in the
577 French Massif Central and its further metamorphic evolution. Contrib. Mineral. Petrol. 79, 1-
578 12.

579

580 Pin C., Marini F.1993. Early Ordovician continental break-up in Variscan Europe: Nd
581 Sr isotope and trace element evidence for bimodal igneous associations of the southern
582 Massif Central, France. Lithos 29, 177-196.

583

584 Roig, J.Y., Faure, M., 2000. La tectonique polyphasée du Sud Limousin (Massif Central
585 français) et son interprétation dans un modèle d'évolution polycyclique de la chaîne
586 hercynienne. Bull. Soc. Géol. France 171, 295-307.

587

588 Rolet, J., La phase bretonne en Bretagne. 1982. Bull. Soc. géol. minéral. Bretagne 14, 63 -71.

589

590 Rolet J., Le Gall, B., Darboux, J.R., Thonon, P. Gravelle, M., 1986. L'évolution
591 géodynamique dévono-carbonifère de l'extrémité occidentale de la chaîne hercynienne
592 d'Europe sur le transect Armorique-Cornwall. Bull. Soc. Géol. France 2, 43-54.
593
594 Sandeman, H., Chen, Y., Clark, A. and Farrar, E., 1994. Constraints on the P-T conditions
595 and age of emplacement of the Lizard ophiolite, Cornwall: amphibole-plagioclase thermo-
596 barometry and $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology of basal amphibolites. Can. J. Earth Sci. 32, 261-272.

597

598

599 Figure Captions

600 Fig. 1: A Insert showing the Variscan belt in France. NASZ: North Armorican Shear Zone,
601 NBSASZ: North Branch of the South Armorican Shear Zone, SBSASZ: South Branch of the
602 South Armorican Shear Zone, NPF: North Pyrenean Fault, MGCR: Mid-German Crystalline
603 Rise. B: simplified structural sketch of the French Massif Central (purple areas: Upper Gneiss
604 Unit; the Lower Gneiss Unit and Para-Autochthonous Unit are not distinguished). C: Structural
605 map of the S part of the French Massif Central with emphasis on the compressional
606 structures. Syn- to late orogenic extensional structures are not considered here. Captions are
607 the same as for Fig. 2.

608

609 Fig. 2: Crustal scale cross-section through the Montagne Noire-Rouergue area (location in
610 Fig. 1). The bulk architecture results of three shearing events, D1, D2, and D3, with different
611 ages and displacement directions.

612

613 Fig. 3: Field and thin section photographs showing D1 and D2 deformation in the S^t-Sernin-
614 sur-Rance nappe and Lower Gneiss Unit. A: NW-SE trending D2 fold (L2) reworking an
615 early NE-SW trending L1 lineation; B: NW-SE trending "a-type" fold deforming the L1

616 lineation; C: Low-temperature D2 top-to-the-NW shear zone reworking the foliated Pinet
617 granodiorite; D: N-S striking intrafolial D1 fold; E: Thin section of staurolite porphyroblast
618 with top-to-the-NW, D2, asymmetric biotite pressure shadows; F: Garnet with top-to-the NW,
619 D2, pressure shadows.

620

621 Fig. 4: Cathodoluminescence images of representative zircons for SIMS U–Pb dating. White
622 ellipses indicate the in-situ analytical spots of $^{206}\text{Pb}/^{238}\text{U}$ age (shown nearby the spot). SIMS
623 spots are 30 μm in length for scale.

624

625 Fig. 5: Zircon ion-probe U-Pb age of the Pinet orthogneiss. Samples FR 41 and FR 42 are
626 located at $44^\circ 03' 16.27''/002^\circ 45' 56.35''$, and $44^\circ 02' 43.57''/002^\circ 46' 47.70''$, respectively.



627

628 Fig. 6: Structural map of the French-Britain segment of the Variscan belt with emphasis on
629 the D2 NW-SE shearing developed from Albigeois up to the Lizard nappe. The Late
630 Carboniferous, south-directed, D3 event develops in southern Massif Central (Cévennes and
631 Montagne Noire). At the scale of the entire Variscan belt, a top-to-the-North-directed
632 deformation, characterized by thrust and faults, deforms the Ardenne Massif and the northern
633 Variscan front.

634

635 Fig. 7: 3D Schematic geodynamic interpretative model to account for the tectonic difference
636 between the Massif Central-Massif Armorica areas dominated by the top-to-the D2
637 synmetamorphic ductile shearing, and the Central Armorican Domain, where D2 deformation
638 is represented by brittle structures only. This difference in rheological behaviour can be
639 explained by a tectonic decoupling between the two domains. The Neoproterozoic Cadomian
640 orogeny, and the thick subcontinental lithosphere mantle may explain the rigid behaviour of
641 the Central Armorican Domain in which the brittle deformation is associated with the opening

642 of the Châteaulin and Laval basins. The left-lateral Nort-sur-Erdre strike-slip fault can be
643 seen as a transfer fault between the S. Gondwana and Central Armorican domains.



- Serpukhovian-Bashkirian granites (326-315 Ma)
- Migmatite and granites in the Montagne Noire dome
- Orthogneiss derived from Cambrian or Ordovician granite (e.g. Mendic, Pinet)
- Upper Gneiss Unit (UGU)
- Lower Gneiss Unit (LGU)
- Para-autochthonous (PAU)
- St-Salvi-de-Carcavès Nappe
- Paleozoic Fold and Thrust Belt
- Southern Foreland basin
- St-Sernin-sur-Rance Nappe
- South-directed D1 event (ca 380 Ma)
- NW-directed D2 event (ca 360Ma)
- South-directed D3 event (340-320 Ma)




Fig. 1




Fig. 2










Fig. 3








Fig. 5



-  Post-orogenic formations:
Late Visean "tufs anthracifères"
Red granites and microgranites
-  360 Ma lineation related to the D2 event
-  Upper Gneiss Unit with
420 Ma high pressure rocks
-  Late Visean-Namurian thrust with
nappe displacement sense (D3 event)

Fig. 6




Fig. 7