Geophysical Research Abstracts Vol. 16, EGU2014-16046, 2014 EGU General Assembly 2014 © Author(s) 2014. CC Attribution 3.0 License.



## Remote sensing of a NTC radio source from a Cluster tilted spacecraft pair

Pierrette Décréau (1), Séna Kougblénou (1), Guillaume Lointier (1), Jean Louis Rauch (1), Jean Gabriel Trotignon (1), Xavier Vallières (1), Patrick Canu (2), Sandrine Rochel Grimald (3), Farida El-Lemdani Mazouz (4), and Fabien Darrouzet (5)

 (1) Laboratoire de Physique et Chimie de l'Environnement et de l'Espace (LPC2E), UMR7328, Orléans, France (pierrette.decreau@cnrs-orleans.fr), (2) Laboratoire de Physique des Plasmas (LPP), Ecole Polytechnique, Palaiseau, France,
(3) Office National d'Études et de Recherches Aérospatiales (ONERA), The French Aerospace Laboratory, Toulouse, France,
(4) Laboratoire ATmosphère, Milieux, Observations Spatiales (LATMOS), Guyancourt, France, (5) Belgian Institute for Space Aeronomy (IASB-BIRA), Brussels, Belgium

The non-thermal continuum (NTC) radiation is a radio wave produced within the magnetosphere of a planet. It has been observed in space around Earth since the '70s, and within the magnetospheres of other planets since the late '80s. A new study using ESA's Cluster mission has shown improved precision in determining the source of various radio emissions produced by the Earth. The experiment involved tilting one of the four identical Cluster spacecraft to measure the electric field of this emission in three dimensions for the first time. Our analysis of a NTC case event pinpointed a small deviation from the generally assumed (circular) polarization of this emission. We show that classical triangulation, in this case using three of the spacecraft located thousands of kilometres apart, can lead to an erroneous source location. A second method, using the new 3D electric field measurements, indicated a source located along the plasmapause at medium geomagnetic latitude, far away from the source location estimated by triangulation. Cluster observations reveal that this NTC source emits from the flank of the plasmapause towards the polar cap. Understanding the source of NTC waves will help with the broader understanding of their generation, amplification, and propagation.