

Effects of early diagenesis on the isotopic signature of wood (δ 13C and δ 15N): incubation in aquatic microcosm

Romain Tramoy, Thanh Thuy T Nguyen Tu, Veronique Vaury, Mathieu Sebilo, Laurence Millot-Cornette, Céline Roose-Amsaleg, Johann Schnyder

▶ To cite this version:

Romain Tramoy, Thanh Thuy T Nguyen Tu, Veronique Vaury, Mathieu Sebilo, Laurence Millot-Cornette, et al.. Effects of early diagenesis on the isotopic signature of wood (δ 13C and δ 15N): incubation in aquatic microcosm. Goldschmidt-2017, Aug 2017, Paris, France. 2017. insu-01624744

HAL Id: insu-01624744 https://insu.hal.science/insu-01624744

Submitted on 30 Oct 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Effects of early diagenesis on the isotopic signature of wood (δ^{13} C and δ^{15} N): incubation in aquatic microcosm



B



metis Milieux Environnementaux Transferts et Interactions dans les hydrosystèmes et les sols







Romain Tramoy¹, Thanh Thuy Nguyen Tu², Véronique Vaury³, Mathieu Sebilo³, Laurence Millot Cornette², Celine Roose-Amsaleg², Johann Schnyder¹.

¹Sorbonne universités, CNRS, UPMC, UMR 7193, ISTEP, France ; ² Sorbonne universités, CNRS, UPMC, EPHE, UMR 7619, METIS, France ; ³ Sorbonne universités, CNRS, UPMC, INRA, IRD-Paris Diderot-UPEC, UMR7618, IEES, France ;


*Work published as Tramoy et al. (2017) in *Environmental Chemistry*




		Average	2σ	Range	Average	2σ	Range
Carbon	11	47.5	0.9	46.7;48.3	-24.1	0.2	-24.3 ; -23.9
Nitrogen	10	0.08	0.02	0.07 ; 0.1	-0.2	1	-0.8 ; 1.1









Loss of ¹³C-depleted compounds (*cf.* flotting particles; tannins, lignin, other non-polar compounds; Melillo et al., 1989)

• **Biotic degradation** by (micro-)organisms like fungi

Bacteria are **NOT** the **main decomposers** Fungi are the main decomposers

• Powders: complex dynamic

► Key-role of respiration leading to ¹³C-enrichment ?


water ?) thanks to fungi activity. Higher NO3- δ^{15} N supports this hypothesis. N accumulation according to Melillo et al. (1989)

• Nitrogen loss in pieces in DW + ¹⁵N-depletion

► Amino-acids and proteins (¹⁵N-enriched) consumption by fungi, which exports nitrogen from the wood through mycellium.

In both type of water, mycellium of fungi likely constitute a nitrogen transport network and their activity may proceed toward isotopic uniformisation of a system (e.g. wood-fungi-water) as long as nitrogen is maintained in the bulk system

Conclusions



 $\blacktriangleright \delta^{13}$ C values of organic matter has lower variability than δ^{15} N values, which confirms its interest as a source and environment indicator

• Without invalidating the use of $\delta^{15}N_{org}$ as a paleoenvironmental marker, this study shows that early diagenesis leads to the integration of isotopic compositions from multiple environmental origins that should be addressed when interpreting $\delta^{15}N_{org}$ in soils and sediments

Références

Garland, J.L., Mills, A.L., 1991. Classification and characterization of heterotrophic microbial communities on the basis of patterns of community-level-sole-carbon-source-utilization. Appl. Environ. Microbiol. 57, 2351-2359. Melillo, J.M., Aber, J.D., Linkins, A.E., Ricca, A., Fry, B., Nadelhoffer, K.J., 1989. Carbon and nitrogen dynamics along the decay continuum: Plant litter to soil organic matter, in: Clarholm, M., Bergström, L. (Eds.), Ecology of Arable Land — Perspectives and Challenges, Developments in Plant and Soil Sciences. Springer Netherlands, pp. 53–62. Tramoy, R., Sebilo, M., Nguyen Tu, T.T., Schnyder, J., 2017. Carbon and nitrogen dynamics in decaying wood: paleoenvironmental implications. Environ. Chem. 14, 9–18. doi:10.1071/EN16049 Zak, J.C., Willig, M.R., Moorhead, D.L., Wildman, H.G., 1994. Functional diversity of microbial communities: A quantitative approach. Soil Biol. Biochem. 26, 1101–1108. Zhao, D., Li, F., Yang, Q., Wang, R., Song, Y., Tao, Y., 2013. The influence of different types of urban land use on soil microbial biomass and functional diversity in Beijing, China. Soil Use Manag. 29, 230–239. doi:10.1111/sum.12034 **Acknowledgement**

We thank Julien Legrand, who collected the wood used. We are grateful to Véronique Vaury (Institute of Ecology and Environmental Sciences of Paris; IEES-UPMC) for analyses. This study was supported by the EMERGENCE project from UPMC. We also thank Sylvie Derenne for access to experimental facilities and financial support for congress.