



Plant diversity affects GHG fluxes in an ecological engineering experiment in a disturbed Sphagnum peatland (La Guette, France)

Sébastien Gogo, Fatima Laggoun-Défarge, Fabien Leroy, Christophe Guimbaud, Léonard Bernard-Jannin

► To cite this version:

Sébastien Gogo, Fatima Laggoun-Défarge, Fabien Leroy, Christophe Guimbaud, Léonard Bernard-Jannin. Plant diversity affects GHG fluxes in an ecological engineering experiment in a disturbed Sphagnum peatland (La Guette, France). EGU General Assembly 2017, Apr 2017, Vienne, Austria. EGU, p.12194, 2017. insu-01664606

HAL Id: insu-01664606

<https://insu.hal.science/insu-01664606>

Submitted on 15 Dec 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Plant diversity affects GHG fluxes in an ecological engineering experiment in a disturbed *Sphagnum* peatland (La Guette, France).

Sébastien Gogo^{1,2,3}, Fatima Laggoun-Défarge^{1,2,3}, Fabien Leroy^{1,2,3}, Christophe Guimbaud^{4,5}, Léonard Bernard-Jannin^{1,2,3}


¹Université d'Orléans, ISTO, UMR 7327, 45071, Orléans, France - ²CNRS, ISTO, UMR 7327, 45071 Orléans, France - ³BRGM, ISTO, UMR 7327, BP 36009, 45060 Orléans, France - ⁴Université d'Orléans, LPC2E, UMR 7328, 45071, Orléans, France - ⁵CNRS, LPC2E, UMR 7328, 45071, Orléans, France

A: Aim of the study

***Sphagnum* peatlands** are huge carbon (C) store, participating in the regulation of the global climate. Their **C sink function is under threat** because of global changes. In addition to their direct effects on ecosystems functioning, these perturbations induce **vegetation change toward increasing vascular plant abundance**. To reduce vegetation modification, hydrological restoration can be done to reduce water table fluctuations. Furthermore, ecological engineering actions can be undertaken to promote species efficient in storing C.

The aims of this study were to set up an **in situ experiment to test the effect of different vegetation treatment on the C fluxes** in a disturbed peatland.

B: Study site



SNO Tourbières
Fonctionnement des tourbières tempérées et impact des changements globaux




It is one of the four sites of the **French Peatland Observatory Service (SNO Tourbières-CNRS)**

La Guette peatland is in the Centre region, 100 km south of Paris.

It is an acidic fen invaded by ***Molinia caerulea***, ***Betula* spp** and ***Pinus sylvestris***. Typical peatland vegetation consists of ***Sphagnum*** (e.g. *S. rubellum*, *S. cuspidatum* and *S.*), *Eriophorum angustifolium*, *Rhynchospora alba*, *Gentiana pneumonanthe*, *Erica tetralix*.

C: Hydrological disturbance / restoration

Water table depth monitoring in the piezometer close to the outlet ★



Hydrological restoration




Work done in January 2014

D: Vegetation restoration experiment

Vegetation treatment that could promote biodiversity and C storing

- ⇒ 8 intact plots serve as **CONTROL** (4 DOWN, 4 UP)
- ⇒ 8 **BARE PEAT** plots after vegetation and peat stripping
- ⇒ 8 bare peat plots where 1 kg of fresh **SPHAGNUM** spp was added

Work done in January 2014




Response variables

- ⇒ Ecosystem respiration
- ⇒ Net Ecosystem Exchange
- ⇒ CH₄ flux
- ⇒ Vegetation % cover
- 6 campaigns per year

E: Vegetation and C fluxes dynamics

Monitoring of vegetation % cover in C fluxes measurement plots



Control
Summer total %: 100 - 150


Bare peat
Summer total %: 20 - 50

Bare peat + Sphagnum
Summer total %: 30 - 60

Vegetation in **CONTROL** is stable with time. *Eriophorum angustifolium* and *Trichophorum caespitosum* colonised the **BARE PEAT**, whereas only the former was present with the moss in **SPHAGNUM** plots.


With time, ER and NEE increased in **BARE PEAT**, and **SPHAGNUM** plots. In winter (red arrows), **SPHAGNUM** plots tend to function as a C sink, unlike the other plots. CH₄ fluxes increased in **BARE PEAT**, and **SPHAGNUM** treatments.

Monitoring of CO₂ and CH₄ fluxes




CH₄ fluxes (μmol.m⁻².s⁻¹)

CH₄ fluxes




F: Sensitivity of ER to temperature and GPP to vegetation % cover

Variation of Q₁₀ of ecosystem respiration with time



Q₁₀ coefficient increases faster with time in **BARE PEAT** and **SPHAGNUM** plots than in the control plots.

Gross Primary Production versus vegetation % cover



GPP increases faster with vegetation % cover in **CONTROL** and **BARE PEAT** plots than in **SPHAGNUM** plots.

G: Discussion

The results suggested that:

- When no *Sphagnum* were added, the colonising vegetation may be more sensitive to temperature than control vegetation. Increased CO₂ emissions in early colonisation stages can be a negative point of the vegetation restoration.
- Although *Sphagnum* GPP is less sensitive than other plots to vegetation % cover, this plots were the most efficient to store C in winter.
- Also, the **BARE PEAT**, and **SPHAGNUM** treatments tended to increased CH₄ emissions compared to **CONTROL**. This is due to the fact that as peat was removed, the water table was higher, promoting CH₄ production and limiting its consumption.

As a conclusion, **BARE PEAT**, and **SPHAGNUM** treatments promotes plant diversity, but their positive effect on C fluxes are still not observable after 2 (CH₄) and 3 (CO₂) years of monitoring. However, the addition of *Sphagnum* is advised as it may restore C storing functioning faster than when colonisation is let free to proceed such as in the **BARE PEAT** plots.