



**HAL**  
open science

## Thermal dependency of CO<sub>2</sub> VUV absorption cross section and application to warm exoplanetary atmospheres

O. Venot, Yves Bénilan, Nicolas Fray, Marie-Claire Gazeau, Franck Montmessin, Franck Lefèvre, C. Bahrini, Et-Touhami Es-Sebbar, Martin Schwell, Eric Hébrard, et al.

► **To cite this version:**

O. Venot, Yves Bénilan, Nicolas Fray, Marie-Claire Gazeau, Franck Montmessin, et al.. Thermal dependency of CO<sub>2</sub> VUV absorption cross section and application to warm exoplanetary atmospheres. European Planetary Science Congress 2017, Sep 2017, Riga, Latvia. pp.EPSC2017-42. insu-01672369

**HAL Id: insu-01672369**

**<https://insu.hal.science/insu-01672369v1>**

Submitted on 24 Dec 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Thermal dependency of CO<sub>2</sub> VUV absorption cross section and application to warm exoplanetary atmospheres

O. Venot (1), Y. Bénéilan (1), N. Fray (1), M.-C. Gazeau (1), F. Montmessin (2), F. Lefèvre (3), C. Bahrini (1), Et. Es-sebbar (3), M. Schwell (1), E. Hébrard (4), M. Lefevre (1), I. P. Waldmann (5)

(1) Laboratoire Interuniversitaire des Systèmes Atmosphériques, UMR CNRS 7583, Universités Paris Est Créteil (UPEC) et Paris Diderot (UPD), Créteil, France, (2) Laboratoire Atmosphères, Milieux, Observations Spatiales (LATMOS), CNRS/IPSL/UVSQ, Guyancourt, France, (3) Laboratoire Atmosphères, Milieux, Observations Spatiales (LATMOS), CNRS/IPSL/UPMC, Paris, France, (4) Paul Scherrer Institute, Laboratory of Thermal Processes and Combustion, CH-5232 Villigen PSI, Switzerland, (5) School of Physics and Astronomy, University of Exeter, EX4 4QL, Exeter, UK, (6) University College London, Department of Physics and Astronomy, Gower Street, London WC1E 6BT, UK (olivia.venot@lisa.u-pec.fr)

### Abstract

Most of exoplanets detected so far have atmospheric temperatures significantly higher than 300 K. These exoplanets are often close to their star and thus receive an intense UV photons flux, triggering important photodissociation processes. However, the temperature dependency of VUV absorption cross sections, which are essential data to model photolyses in atmospheric models, are barely known. Thus, by lack of appropriate data, absorption cross sections at room temperature are used in photochemical models of extrasolar planets, leading to a non-measurable uncertainty. With the future space- or ground-based telescopes that will be developed in the coming years (JWST, E-ELT...) investigating these research fields becomes urgent [1]. In this context, we quantified the temperature dependency of the VUV absorption cross section of carbon dioxide (CO<sub>2</sub>). We performed experimental measurements on the range (115-230 nm) between 150 and 800 K. The absorption cross section of CO<sub>2</sub> increases with the temperature (Fig. 1). At 200 nm, there are more than four orders of magnitude between the one at 300 K and the one at 800 K. We also determined a parametrisation to calculate the continuum of the absorption cross section on this wavelength range. We used these new data in our photo-thermochemical model for exoplanets [2] and studied the impact on the results (photolyses rates and chemical composition) as well as on the observables (synthetic transmission spectra) [3, 4]. We will present these experimental results and their consequences on the modelling of exoplanet atmospheres.




Figure 1: Absorption cross section of CO<sub>2</sub> (cm<sup>2</sup>) at temperature between 150 and 800 K.

### References

- [1] Fortney, J. J., Robinson, T. D., Domagal-Goldman, S., et al.: The Need for Laboratory Work to Aid in The Understanding of Exoplanetary Atmospheres, arXiv e-prints 1602.06305, 2016
- [2] Venot, O., Hébrard, E., Agúndez, M., et al.: A chemical model for the atmosphere of hot Jupiters, *A&A*, 546, A43, 2012
- [3] Waldmann, I. P., Tinetti, G., Rocchetto, M., et al.: TauREx I: A Next Generation Retrieval Code for Exoplanetary Atmospheres, *The Astrophysical Journal*, 802, 107, 2015
- [4] Waldmann, I. P., Rocchetto, M., Tinetti, G., et al.: TauREx II: Retrieval of Emission Spectra, *The Astrophysical Journal*, 813, 13, 2015