



HAL
open science

Mapping Venus Mineralogy and Chemistry In Situ from Orbit with Six-Window VNIR Spectroscopy

M. D. Dyar, J. Helbert, T. Boucher, D. Wendler, I. Walter, Thomas Widemann, Emmanuel Marcq, A. Maturilli, S. Ferrari, M. d'Amore, et al.

► **To cite this version:**

M. D. Dyar, J. Helbert, T. Boucher, D. Wendler, I. Walter, et al.. Mapping Venus Mineralogy and Chemistry In Situ from Orbit with Six-Window VNIR Spectroscopy. 15th Meeting of the Venus Exploration Analysis Group (VEXAG), Nov 2017, Laurel, Maryland, United States. LPI Contribution No. 2061, p.8004. insu-01672492

HAL Id: insu-01672492

<https://insu.hal.science/insu-01672492>

Submitted on 20 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MAPPING VENUS MINERALOGY AND CHEMISTRY *IN SITU* FROM ORBIT WITH SIX-WINDOW VNIR SPECTROSCOPY. M. D. Dyar¹, J. Helbert², T. Boucher³, D. Wendler², I. Walter⁴, T. Widemann⁵, E. Marcq⁶, A. Maturilli², S. Ferrari^{7,1}, M. D'Amore², N. Müller⁸, and S. Smrekar⁸, ¹Planet. Sci. Inst., 1700 East Fort Lowell, Tucson, AZ 85719 USA (mdyar@mtholyoke.edu); ²Inst. Planet. Res., DLR, Rutherfordstrasse 2, 12489 Berlin, Germany; ³Col. of Inform. and Computer Sci., Univ. of Massachusetts Amherst, Amherst, MA, 01003, USA; ⁴Inst. Optical Sensorsystems, DLR, Rutherfordstrasse 2, 12489 Berlin, Germany, ⁵LESIA; ⁶LATMOS; ⁷Center of Studies and Activities for Space G. Colombo, University of Padova, Via Venezia 15, 35131 Padova, Italy ; ⁸Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Dr., Pasadena CA, 91109.

Introduction: How can Venus' surface mineralogy and geochemistry best be studied? Technological advances in surface techniques since prior Venus landed missions such as LIBS, Raman [1], and XRF [2] make in-situ instrumentation attractive. However they are limited to short analysis times, autonomous sample collection, and provide local versus global information.

Although not designed to image the Venusian surface, both the Galileo spacecraft and VIRTIS on Venus Express were able to image the surface through transparent windows in the CO₂ spectrum near 1 μm [3-7]. To leverage this capability, the Venus Emissivity Mapper (VEM) was developed specifically to study the surface of Venus through six different windows at 0.86, 0.91, 0.99, 1.02, 1.11, and 1.18 μm. This project explores what can be learned about Venus surface geochemistry and mineralogy using those six windows from an orbiter *in situ* at Venus, with two focuses:

- 1) the ability of VEM-window data to distinguish among key rock types on Venus, and
- 2) their capability to evaluate redox state and transition metal contents of Venus surface rocks.

Data: VNIR data were collected at Venus temperatures in the Planetary Spectroscopy Laboratory at DLR [8-10]. Fe³⁺/Fe²⁺ ratios were measured using Mössbauer spectroscopy [11] and standard methods. Samples studied were reported in [12].




Figure 1. Laboratory emissivity data with %Fe³⁺ (left) and wt.% Fe²⁺ (right, indicated as determined by combining Mössbauer, x-ray fluorescence, and EMPA).

Rock Type Distinctions: As seen in **Figure 1**, samples containing Fe oxides (rhyolite, granites, and the oxides themselves) have negative slopes between the two lowest wavelength bands. The wavelength region ~0.99 and 1.02 μm allows felsic rocks to be distinguished from mafic ones. In this region, Fe (and other transition metals) in silicate minerals causes elevated emissivity, causing felsic rocks to group separately, and oxides to have low emissivities. Binary classifiers [1] demonstrate that at current best estimate errors, basalt spectra can be confidently discriminated from basaltic andesites, andesites, and rhyolite/granite.




Figure 2. Assessment of weathering based on emissivity at 1.18 μm; other bands can also be used to support this trend.

Redox state: Surface weathering based on oxidation state can be inferred from the 1.18 μm and other bands (**Figure 2**). This distinction can also be made using the 0.99 and 1.02 μm bands, where hematite is easily distinguished from magnetite.

References: [1] Esposito L. W. et al. (2017) *LPS XLVIII*, Abstract #2444. [2] Glaze L.S. et al. (2017) *LPS XLVIII*, Abstract #2288. [3] Mueller N. et al. (2008) *JGR*, 113, 1–21. [4] Helbert J. et al. (2008) *GRL*, 35, L11201. [5] Hashimoto G. L. et al. (2008) *JGR*, 113, E00B24. [6] Piccioni G. et al. (2007) *ESA Special Pub.* 1295. [7] **Mueller et al VEXAG**. [8] Helbert J. et al. (2013) *EPSL*, 369-370, 233-238. [9] Maturilli A. and Helbert J. (2014) *J. Appl. Remote Sens.*, 8, 084985. [10] Maturilli A. et al. (2014) *EPSL*, 398, 58-65. [11] Dyar M. D. et al. (2017) *LPS XLVIII*, Abstract #3014. [12] Dyar M. D. et al. (2016) *LPS XLVII*, Abstract #2205.