

First Search for Nontensorial Gravitational Waves from Known Pulsars

B P Abbott, R. Abbott, T. D Abbott, F. Acernese, K. Ackley, C. Adams, T. Adams, P. Addesso, R. X Adhikari, V B Adya, et al.

To cite this version:

B P Abbott, R. Abbott, T. D Abbott, F. Acernese, K. Ackley, et al.. First Search for Nontensorial Gravitational Waves from Known Pulsars. Physical Review Letters, 2018, 120 (3), 13 p. 10.1103/physrevlett.120.031104 . insu-01887103

HAL Id: insu-01887103 <https://insu.hal.science/insu-01887103v1>

Submitted on 3 Oct 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

First Search for Nontensorial Gravitational Waves from Known Pulsars

B. P. Abbott et al.^{*}

(LIGO Scientific Collaboration and Virgo Collaboration)

(Received 12 October 2017; revised manuscript received 16 November 2017; published 19 January 2018)

We present results from the first directed search for nontensorial gravitational waves. While general relativity allows for tensorial (plus and cross) modes only, a generic metric theory may, in principle, predict waves with up to six different polarizations. This analysis is sensitive to continuous signals of scalar, vector, or tensor polarizations, and does not rely on any specific theory of gravity. After searching data from the first observation run of the advanced LIGO detectors for signals at twice the rotational frequency of 200 known pulsars, we find no evidence of gravitational waves of any polarization. We report the first upper limits for scalar and vector strains, finding values comparable in magnitude to previously published limits for tensor strain. Our results may be translated into constraints on specific alternative theories of gravity.

DOI: [10.1103/PhysRevLett.120.031104](https://doi.org/10.1103/PhysRevLett.120.031104)

Introduction.—The first gravitational waves detected by the Advanced Laser Interferometer Gravitational-Wave Observatory (aLIGO) and Virgo have already been used to place some of the most stringent constraints on deviations from the general theory of relativity (GR) in the highly dynamical and strong-field regimes of gravity [1–4]. However, even though some partial progress has been made with the observation of GW170814 [5,6] and in spite of the wealth of new information provided by GW170817 [7,8], it has not yet been possible to unambiguously confirm GR's prediction that the associated metric perturbations are of a tensor nature (helicity ± 2), rather than vector (helicity ± 1), or scalar (helicity 0) [9]. This is unfortunate, since the presence of nontensorial modes is a key prediction of many extensions to GR [10–14]. Most importantly, the detection of a scalar or vector component, no matter how small, would automatically point to physics beyond Einstein's theory [12,13].

In order to experimentally study gravitational-wave polarizations directly, one needs a local measurement of their geometric effect (i.e., which directions are stretched and squeezed) that breaks degeneracies between the five distinguishable (to differential-arm instruments) modes supported by a generic metric theory of gravity [10,11]. For transient waves like those detected so far, this cannot be fully achieved with the LIGO-Virgo network, as at least five nonco-oriented differential-arm antennas are required to break all such degeneracies[13,15]. Constraints on the magnitude of non-GR polarizations inferred from indirect measurements, like the rate of orbital decay of binary pulsars, are only meaningful in the context of specific theories (see, e.g., Refs. [16,17] or Refs. [18,19] for reviews).

Theory-independent polarization measurements could instead be carried out with current detectors in the presence of signals sufficiently long to probe the detector antenna patterns, which are themselves polarization sensitive [20–23]. Such is the case, for instance, for the continuous, almost-monochromatic waves expected from spinning neutron stars with an asymmetric moment of inertia [24]. Known galactic pulsars are one of the main candidates for searches for such signals in data from ground-based detectors, and analyses targeting them have already achieved sensitivities that are comparable to, or even surpass, their canonical spin-down limit (i.e., the strain that would be produced if the observed slowdown in the pulsar's rotation was completely due to gravitational radiation) [25].

However, all previous targeted searches have been, by design, restricted to tensorial gravitational polarizations only. This leaves open the possibility that, due to a departure from GR, the neutron stars targeted in previous searches may indeed be emitting strong continuous waves with nontensorial content, in spite of the null results of standard searches.

In this Letter, we present results from a search for continuous gravitational waves in aLIGO data that makes no assumptions about how the gravitational field transforms under local spatial rotations, and is thus sensitive to any of the five measurable polarizations allowed by a generic metric theory of gravity. We targeted 200 known pulsars using data from aLIGO's first observation run (O1), and assuming emission at twice the rotational frequency of the source.

Our data provide no evidence for the emission of gravitational signals of tensorial or nontensorial polarization from any of the pulsars targeted. For sources in the most sensitive band of our detectors, we constrain the strain of the scalar and vector modes to be below 1.5×10^{-26} at 95% credibility. These are the first direct upper limits for *

Full author list given at the end of this article.

scalar and vector strain, and may in principle be used to constrain beyond-GR theories of gravity.

Analysis.—We search aLIGO O1 data from the Hanford (H1) and Livingston (L1) detectors for continuous waves of any polarization (tensor, scalar, or vector) by applying the Bayesian time-domain method of Ref. [26], generalized to non-GR modes as described in Ref. [21] and summarized below. Our analysis follows closely that of Ref. [25], and uses the exact same interferometric data.

Calibrated detector data are heterodyned and filtered using the timing solutions obtained from electromagnetic observations for each pulsar. The maximum calibration uncertainties estimated over the whole run give a limit on the combined H1 and L1 amplitude uncertainties of 14% this is the conservative level of uncertainty on the strain upper limits [25,27].

The data streams start on 2015 Sep 11 at 01∶25:03 UTC for H1 and 18∶29:03 UTC for L1, and finish on 2016 Jan 19 at 17∶07:59 UTC at both sites. The pulsar timing solutions used are also the same as in Ref. [25] and were obtained from the 42-ft telescope and Lovell telescope at Jodrell Bank (UK), the 26-m telescope at Hartebeesthoek (South Africa), the Parkes radio telescope (Australia), the Nancay Decimetric Radio Telescope (France), the Arecibo Observatory (Puerto Rico), and the Fermi Large Area Telescope (LAT).

As described in detail in Ref. [21], we construct a Bayesian hypothesis that captures signals of any polarization content (our *any-signal* hypothesis, \mathcal{H}_S) by combining the sub-hypotheses corresponding to the signal being composed of tensor, vector, scalar modes, or any combination thereof. Each of these sub-hypotheses corresponds to a different signal model; in particular, the least restrictive template includes contributions from all polarizations and can be written as

$$
h(t) = \sum_{p} F_p(t; \alpha, \delta, \psi) h_p(t), \qquad (1)
$$

where the sum is over the five independent polarizations: plus $(+)$, cross (x) , vector x (x) , vector y (y) , and scalar (s) [11]. The two scalar modes in the most common basis, breathing and longitudinal, are degenerate for networks of quadrupolar antennas [13], so we do not make a distinction between them.

Each term in Eq. (1) is the product of an antenna pattern function F_p and an intrinsic strain function h_p . We define the different polarizations in a wave frame such that the ζ axis points in the direction of propagation, χ lies in the plane of the sky along the line of nodes (here defined to be the intersection of the equatorial plane of the source with the plane of the sky), and y completes the right-handed system, such that the polarization angle ψ is the angle between the y axis and the projection of the celestial North onto the plane of the sky (see, e.g., Ref. [28]). We can thus write the F_p 's as implicit functions of the source's right ascension α , declination δ , and polarization ψ . (For the sources targeted here, α and δ are always known to high accuracy, while ψ is usually unknown.) The antenna patterns acquire their time dependence from the sidereal rotation of Earth; explicit expressions for the F_p 's are given in Refs. [20–22,29,30].

For a continuous wave, the polarizations take the simple form

$$
h_p(t) = a_p \cos[\phi(t) + \phi_p],
$$
 (2)

where a_p is a time-independent strain amplitude, $\phi(t)$ is the intrinsic phase evolution, and ϕ_p a phase offset for each polarization. The nature of these three quantities depends on the specifics of the underlying theory of gravity and the associated emission mechanism (for different emission mechanisms within GR, see, e.g., Refs. [31–33]). While we treat a_p and ϕ_p as free parameters, we take $\phi(t)$ to be the same as in the traditional GR analysis [25]:

$$
\phi(t) = 2\pi \sum_{j=0}^{N} \frac{\partial_t^{(j)} f_{\text{GW},0}}{(j+1)!} [t - T_0 + \delta t(t)]^{(j+1)}, \quad (3)
$$

where $\partial_t^{(j)} f_{\text{GW},0}$ is the *j*th time derivative of $f_{\text{GW},0}$, the emission frequency measured at the fiducial time T_0 ; $\delta t(t)$ is the time delay from the observatory to the solar system barycenter (including the known Rømer, Shapiro and Einstein delays), and can also include binary system corrections to transform the time coordinate to a frame approximately inertial with respect to the source; N is the order of the series expansion (1 or 2 for most sources).

The gravitational-wave frequency f_{GW} is related to the rotational frequency of the source f_{rot} , which is in turn known from electromagnetic observations. Although arbitrary theories of gravity and emission mechanisms may predict gravitational emission at any multiple of the rotational frequency, here we assume $f_{GW} = 2f_{rot}$, in accordance with the most favored emission model in GR [24]. This restriction arises from practical considerations affecting our specific implementation, and will be relaxed in future studies.

For convenience, we define effective strain amplitudes for tensor, vector, and scalar modes, respectively, by

$$
h_t \equiv \sqrt{a_+^2 + a_{\times}^2},\tag{4}
$$

$$
h_v \equiv \sqrt{a_x^2 + a_y^2},\tag{5}
$$

$$
h_s \equiv a_s,\tag{6}
$$

in terms of the intrinsic a_p amplitudes of Eq. (2). These quantities may serve as proxy for the total power in each polarization group.

One may recover the GR hypothesis considered in previous analyses by setting

$$
a_{+} = h_0(1 + \cos^2 t)/2, \qquad \phi_{+} = \phi_0, \tag{7}
$$

$$
a_{\times} = h_0 \cos \iota, \qquad \phi_{\times} = \phi_0 - \pi/2, \tag{8}
$$

$$
a_x = a_y = a_s = 0,\t\t(9)
$$

where ι is the inclination (angle between the line of sight and the spin axis of the source), and h_0 , ϕ_0 are free parameters. (As with ψ , *i* is unknown for most pulsars.) This corresponds to the standard triaxial-star emission mechanism (see, e.g., Ref. [34]). We use this parameterization only when we wish to incorporate known orientation information as explained below; otherwise, we parametrize the tensor polarizations directly in terms of a_+ , a_{\times} , ϕ_+ and ϕ_{\times} .

Templates of the form of Eq. (1), together with appropriate priors, allow us to compute Bayes factors (marginalized-likelihood ratios) for the presence of signals in the data vs Gaussian noise. We do this using an extension of the nested sampling implementation presented in Ref. [35] (see Ref. [21] for details specific to non-GR polarizations). The Bayes factors corresponding to each signal model may be combined into the odds \mathcal{O}_N^S that the data contain a continuous signal of any polarization vs Gaussian noise:

$$
\mathcal{O}_N^S = P(\mathcal{H}_S | \mathbf{B}) / P(\mathcal{H}_N | \mathbf{B}),\tag{10}
$$

i.e., the ratio of the posterior probabilities that the data B contain a signal of any polarizations (\mathcal{H}_S) vs just Gaussian noise (\mathcal{H}_N) . We compute these odds by setting model priors such that $P(\mathcal{H}_S) = P(\mathcal{H}_N)$; then, by Bayes' theorem, $\mathcal{O}_N^S = \mathcal{B}_N^S$, with the Bayes factor

$$
\mathcal{B}_N^S \equiv P(\mathbf{B}|\mathcal{H}_S)/P(\mathbf{B}|\mathcal{H}_N). \tag{11}
$$

Built into the astrophysical signal hypothesis, \mathcal{H}_s , is the requirement of coherence across detectors, which must be satisfied by a real gravitational wave. In order to make the analysis more robust against non-Gaussian instrumental features in the data, we also define an instrumental feature hypothesis, \mathcal{H}_I , that identifies non-Gaussian noise artifacts by their lack of coherence across detectors [25,36]. In particular, we define \mathcal{H}_I to capture Gaussian noise or a detector-incoherent signal (i.e., a feature that mimics an astrophysical signal in a single instrument, but is not recovered consistently across the network) in each detector [21]. We may then compare this to $\mathcal{H}_\mathcal{S}$ by means of the odds \mathcal{O}_I^S . For D detectors, this is given by

$$
\log \mathcal{O}_I^S = \log \mathcal{B}_N^S - \sum_{d=1}^D \log \left(\mathcal{B}_{N_d}^{S_d} + 1 \right),\tag{12}
$$

where $\mathcal{B}_{N_d}^{S_d}$ is the signal vs noise Bayes factor computed only from data from the dth detector. This choice implicitly assigns prior weight to the models such that $P(\mathcal{H}_S) =$ $P(\mathcal{H}_I) \times 0.5^D$ [21]. For an in depth analysis of the behavior of the different Bayesian hypotheses considered here, in

TABLE I. Existing orientation information for pulsars in our band, obtained from observations of the pulsar wind nebulae (see Table 3 in Ref. [42], and Refs. [43,44] for measurement details).

		W
$J0534 + 2200$	62° : 2 + 1.99	$35\degree 2 + 1\degree 5$
$J0537 - 6910$	$92^{\circ}8 + 0^{\circ}9$	$41^\circ 0 + 2^\circ 2$
$J0835 - 4510$	$63\% + 0\%$	$40\% + 0\%$
J1833-1034	$85^\circ\!\!.4 \pm 0^\circ\!\!.3$	$45^{\circ} + 1^{\circ}$
$J1952 + 3252$		$-11^{\circ}5 + 8^{\circ}6$

the presence and absence of simulated signals of all polarizations, we again refer the reader to the methods paper [21].

We compute likelihoods by taking source location, frequency, and frequency derivatives as known quantities. In computing Bayes factors, we employ priors uniform in the logarithm of amplitude parameters $(h_0 \text{ or } a_p \text{'s})$, since these are the least informative priors for scaling coefficients [37]; we bound these amplitudes to the $10^{-28} - 10^{-24}$ range [38]. On the other hand, flat amplitude priors are used to compute upper limits, to facilitate comparison with published GR results in Ref. [25]. In all cases, flat priors are placed over all phase offsets (ϕ_0 and all the ϕ_p 's).

For those few cases in which some orientation information exists (see Table I), we analyze the data a second time using the triaxial parametrization of tensor modes, Eqs. (7) and (8), taking that information into account by marginalizing over ranges of $\cos i$ and ψ in agreement with measurement uncertainties. Following previous work [25], we only consider orientation constraints obtained from pulsar wind nebulae. However, pulsar orientations can also be inferred from other measurements, especially if the object is in a binary (e.g., Refs. [39–41]). We will consider incorporating such constraints in future searches.

Results.—We find no evidence of continuous-wave signals of any polarization, tensorial or otherwise, from any of the 200 pulsars analyzed. The main quantity of interest is $\log_{10} \tilde{O}_I^S$, defined in Eq. (12), since it encodes the probability that the data contain a signal vs just instrumental noise (Gaussian or otherwise). This quantity, together with the log(odds) for signal vs Gaussian noise, is presented as a function of assumed emission frequency for each pulsar in Fig. 1, and histogrammed in Fig. 2. Importantly, the outliers in Fig. 1 lose significance once $\log_{10} \mathcal{O}_I^S$ is taken into account; indeed, Figs. 1 and 2 reveal the usefulness of $\log_{10} \mathcal{O}_I^S$ in increasing the robustness of the search against non-Gaussian instrumental artifacts.

Based on the intrinsic probabilistic meaning of $\log_{10} \mathcal{O}_I^S$ in terms of betting odds, it is standard to demand at least $\log_{10} \mathcal{O}_I^S > 1$ to conclude that the signal model is favored (see, e.g., the table in Sec. 3.2 of Ref. [46], or Jeffrey's original criteria in Ref. [47] or [48]). Since none of the odds obtained meet this criterion, we conclude that there is no evidence for signals from any of the pulsars targeted.

FIG. 1. Log(odds) vs emission frequency. Log(odds) comparing the any-signal hypothesis to the instrumental (top) and Gaussian noise (bottom) hypotheses, as a function of assumed gravitational-wave frequency, $f_{GW} = 2f_{rot}$, for each pulsar. Looking at the top plot for $\log_{10} \mathcal{O}_I^S$, notice that the instrumental noise hypothesis is clearly favored for all pulsars except one, for which the analysis is inconclusive. (This is $J1932 + 17$, the same nonsignificant outlier identified in Ref. [25].) These results were obtained without incorporating any information on the source orientation, and are tabulated in Table II in the Supplemental Material [45]. Expressions for both odds are given in Eq. (10) and Eq. (12).

In most cases, $\log_{10} \mathcal{O}_I^S < 0$ and the noise model is clearly favored; the single exception is $J1932 + 17$, for which $\log_{10} \mathcal{O}_I^S \sim 0$, so that we can make no conclusive statement about which hypothesis is preferred. (The presence of this nonsignificant outlier is to be expected, as it was already identified in Ref. [25].)

The distribution of the odds corresponding to the subhypotheses making up \mathcal{H}_s are summarized in the box plots of Fig. 3. These correspond to tensor-only (t) , scalaronly (s) , vector-only (v) , scalar-vector (sv) , scalar-tensor (st) , vector-tensor (vt) , and scalar-vector-tensor (stv) models. The mean of these distributions decreases with the number of degrees of freedom in the model, which is to be expected from the associated Occam penalties [21]. The rightmost panel in Fig. 3 shows the distribution of $\log_{10} \mathcal{O}_N^S$, which results from the combination of all the other odds; this is the same quantity histogrammed on the abscissa of Fig. 2.

In the absence of any discernible signals, we produce upper limits for the magnitude of scalar, vector, and tensor polarizations, with a 95% credibility. As usual in Bayesian analyses, upper limits are obtained by integrating posterior probability distributions for the relevant parameters up to the desired credibility (see, e.g., Ref. [21]). Using the effective amplitude definitions of Eqs. (4) – (6) , these quantities are presented in Fig. 4 as a function of assumed emission frequency, and the Supplemental Material [45].

FIG. 2. Log(odds) distributions. Distributions of log(odds) comparing the any-signal hypothesis to the instrumental (ordinate axis, right) and Gaussian noise (abscissa axis, top) hypotheses for all pulsars. This plot contains the same information as Fig. 1 and displays the same nonsignificant outlier. These results were obtained without incorporating any information on the source orientation, and are tabulated in Table II in the Supplemental Material [45]. Expressions for both odds in this plot are given in Eq. (10) and Eq. (12). We underscore that, although this plot looks similar to Fig. 2 in Ref. [25], the signal hypothesis here incorporates scalar, vector, and tensor modes, in all possible combinations.

The plotted upper limits are computed under the assumption of a signal model that includes all five independent polarizations (\mathcal{H}_{svt}) ; limits obtained assuming other signal models may be found online in Ref. [45]. Previous work has demonstrated that the presence or absence of a GR component does not affect the non-GR upper limits (Fig. 13 in Ref. [21]).

As expected, the upper limits presented here are comparable in magnitude to the upper limits on the GR strain obtained by the traditional searches [25]. However, constraints on the scalar amplitude are, on average, around 20% less stringent than those on the vector or tensor amplitudes. This is a consequence of the fact that, for most source locations in the sky, the LIGO detectors are intrinsically less sensitive to continuous waves of scalar (breathing or longitudinal) polarization [21].

Technically, traditional all-sky searches for continuous gravitational waves are also sensitive to nontensorial modes, because they are generally designed to look for any signal of sidereal and half-sidereal periodicities in the data, without assuming knowledge of phase evolution or source sky location [49–53]. However, as can be seen by comparing the magnitude of all-sky upper limits

FIG. 3. Sub-hypothesis odds. Box plots for the distribution of the signal vs noise log(odds) for each of the sub-hypotheses considered, for all of the pulsars analyzed. The sub-hypotheses are (st) , vector-tensor (tv) , scalar-vector-tensor tensor-only (t) , scalar-only (s) , vectoronly (v) , scalar-vector (sv) , scalar-tensor (st) , vector-tensor (vt) , and scalar-vector-tensor (stv) ; these are all combined into the signal hypothesis (S). The quantity represented is $\log_{10} B_N^m$, which is the same as $\log_{10} O_N^m$ if neither \mathcal{H}_m nor \mathcal{H}_N are favored a priori (hence, the label on the ordinate axis). The horizontal red line marks the median of the distribution, while each gray box extends from the lower to upper quartile, and the whiskers mark the full range of the distribution of $\log_{10} \mathcal{O}_N^m$ for the 200 pulsars analyzed. These results were produced without incorporating any information on the source orientation, and are tabulated in Table II in the Supplemental Material [45].

(e.g., Fig. 9 in Ref. [49]) to those shown here in Fig. 4, the sensitivity of these searches would be substantially poorer than that of a targeted search like this one—if only because they are not targeted to a specific source. This is especially true if the search is optimized for a given signal polarization (e.g., circular combination of plus and cross).

Odds and 95%-credible upper limits are summarized in the Supplemental Material: Table I, for pulsars with measured orientations (using the triaxial parameterization of tensor modes), and Table II, for all pulsars without incorporating any orientation information (using the unconstrained parameterization of tensor modes) [45]. Odds values are reported with an error of 5% at 90% confidence; errors on the upper limits due to the use of finite samples in estimating posterior probability distributions are at most 10% at 90% confidence, which is slightly less than the 15% error expected from calibration uncertainties.

Conclusion.—We have presented the results of the first direct search for nontensorial gravitational waves. This is also the first search targeted at known pulsars that is sensitive to any of the five measurable polarizations of the gravitational perturbation allowed by a generic metric theory of gravity. From the analysis of O1 data from both aLIGO observatories, we have found no evidence of signals from any of the 200 pulsars targeted.

In the absence of a clear signal, we have produced the first direct upper limits for scalar and vector strains (Fig. 4, and tables in the Supplemental Material [45]). The values of the 95%-credible upper limits are comparable in magnitude to previously published GR constraints, reaching $h \sim 1.5 \times$ 10[−]²⁶ for pulsars whose frequency is in the most sensitive band of our instruments. This means that, to 95% credibility, none of the pulsars in our set is emitting gravitational waves (tensorial or otherwise) at the frequencies analyzed with enough power for them to reach Earth with amplitudes larger than our upper limits.

Our results have been obtained in a theory-independent fashion. However, our upper limits on nontensorial strain can be translated into model-dependent constraints on beyond-GR theories by picking a specific alternative theory and emission mechanism. To do so, one should use the upper limits produced under the assumption of a signal model that incorporates the same polarizations allowed by the theory one wishes to constrain; these may not necessarily be those in Fig. 4 (e.g., for limits on a scalar-tensor theory, one needs upper limits from \mathcal{H}_{st}). However, this also requires nontrivial knowledge of the dynamics of spinning neutron stars under the theory of interest.

While it is conventional to compare the sensitivity of continuous wave searches to the canonical spin-down limit for each pulsar, it is not possible to do so here without committing to a specific theory of gravity. This is because doing so would require specific knowledge of how each polarization contributes to the effective gravitational-wave stress energy, how matter couples to the gravitational field, how the waves propagate (dispersion and dissipation), and what the angular dependence of the emission pattern is. However, analogs of the canonical spin-down limit for specific theories may be obtained from the results presented here by using the strain upper limits obtained assuming the sub-hypotheses with polarizations corresponding to that theory, as mentioned above.

We have demonstrated the robustness of searches for generalized polarization states (tensor, vector, or scalar) in gravitational waves from spinning neutron stars. Furthermore, even in the absence of a detection, we were able to obtain novel constraints on the strain amplitude of nontensorial polarizations. In the future, once a signal is detected, similar methods will allow us to characterize the gravitational polarization content and, in so doing, perform novel tests of general relativity. Although this search assumed an emission frequency of twice the rotational

FIG. 4. Non-GR upper limits vs emission frequency. Circles mark the 95%-credible upper limit on the scalar, $h_s^{95\%}$ (top), and the effective vector, $h_v^{95\%}$ (middle), and tensor $h_t^{95\%}$ (bottom) strain amplitudes as a function of assumed gravitational-wave frequency for each of the 200 pulsars in our set. The upper limits are obtained assuming a signal model including all five independent polarizations (\mathcal{H}_{str}) , and incorporating no information on the orientation of the source (Table II in the Supplemental Material [45]). The effective amplitude spectral density (ASD) of the detector noise is also displayed for reference; this is the harmonic mean of the H1 and L1 spectra; the scaling is obtained from linear regression to the upper limits.

frequency of the source, this restriction will be relaxed in future analyses.

The authors gratefully acknowledge the support of the United States National Science Foundation (NSF) for the construction and operation of the LIGO Laboratory and Advanced LIGO as well as the Science and Technology Facilities Council (STFC) of the United Kingdom, the Max-Planck-Society (MPS), and the State of Niedersachsen, Germany for support of the construction of Advanced LIGO and construction and operation of the GEO600 detector. Additional support for Advanced LIGO was provided by the Australian Research Council. The authors gratefully acknowledge the Italian Istituto Nazionale di Fisica Nucleare (INFN), the French Centre National de la Recherche Scientifique (CNRS), and the Foundation for Fundamental Research on Matter supported by the Netherlands Organisation for Scientific Research, for the construction and operation of the Virgo detector and the creation and support of the EGO consortium. The authors also gratefully acknowledge research support from these agencies as well as by the Council of Scientific and Industrial Research of India, Department of Science and Technology, India, Science & Engineering Research Board (SERB), India, Ministry of Human Resource Development, India, the Spanish Ministerio de Economía y Competitividad, the Conselleria d'Economia i Competitivitat and Conselleria d'Educació, Cultura i Universitats of the Govern de les Illes Balears, the National Science Centre of Poland, the European Commission, the Royal Society, the Scottish Funding Council, the Scottish Universities Physics Alliance, the Hungarian Scientific Research Fund (OTKA), the Lyon Institute of Origins (LIO), the National Research Foundation of Korea, Industry Canada and the Province of Ontario through the Ministry of Economic Development and Innovation, the Natural Science and Engineering Research Council Canada, Canadian Institute for Advanced Research, the Brazilian Ministry of Science, Technology, and Innovation, Fundaçao de Amparo à Pesquisa do Estado de São Paulo (FAPESP), Russian Foundation for Basic Research, the Leverhulme Trust, the Research Corporation, Ministry of Science and Technology (MOST), Taiwan, and the Kavli Foundation. The authors gratefully acknowledge the support of the NSF, STFC, MPS, INFN, CNRS, and the State of Niedersachsen, Germany for provision of computational resources.

This paper has been assigned LIGO Document Number LIGO-P1700009.

- [1] B. P. Abbott et al. (LIGO Scientific Collaboration and Virgo Collaboration), Phys. Rev. Lett. 116[, 061102 \(2016\)](https://doi.org/10.1103/PhysRevLett.116.061102).
- [2] B. P. Abbott et al. (LIGO Scientific Collaboration and Virgo Collaboration), Phys. Rev. Lett. 116[, 241103 \(2016\)](https://doi.org/10.1103/PhysRevLett.116.241103).
- [3] B. P. Abbott et al. (LIGO Scientific Collaboration and Virgo Collaboration), Phys. Rev. X 6[, 041015 \(2016\)](https://doi.org/10.1103/PhysRevX.6.041015).
- [4] B. P. Abbott et al. (LIGO Scientific Collaboration and Virgo Collaboration), Phys. Rev. Lett. 118[, 221101 \(2017\)](https://doi.org/10.1103/PhysRevLett.118.221101).
- [5] B. P. Abbott et al. (LIGO Scientific Collaboration and Virgo Collaboration), Phys. Rev. Lett. 119[, 141101 \(2017\)](https://doi.org/10.1103/PhysRevLett.119.141101).
- [6] M. Isi, Technical Note No. LIGO-P1700276, 2017, [https://](https://arxiv.org/abs/1710.03794) arxiv.org/abs/1710.03794.
- [7] B. P. Abbott et al. (LIGO Scientific Collaboration and Virgo Collaboration), Phys. Rev. Lett. 119[, 161101 \(2017\)](https://doi.org/10.1103/PhysRevLett.119.161101).
- [8] B. P. Abbott et al. (LIGO Scientific Collaboration, Virgo Collaboration, Fermi Gamma-ray Burst Monitor, and INTEGRAL), [Astrophys. J.](https://doi.org/10.3847/2041-8213/aa920c) 848, L13 (2017).
- [9] B. P. Abbott et al. (LIGO Scientific Collaboration and Virgo Collaboration), Phys. Rev. Lett. 116[, 221101 \(2016\)](https://doi.org/10.1103/PhysRevLett.116.221101).
- [10] D. M. Eardley, D. L. Lee, A. P. Lightman, R. V. Wagoner, and C. M. Will, [Phys. Rev. Lett.](https://doi.org/10.1103/PhysRevLett.30.884) 30, 884 (1973).
- [11] D. Eardley, D. Lee, and A. Lightman, [Phys. Rev. D](https://doi.org/10.1103/PhysRevD.8.3308) 8, 3308 [\(1973\).](https://doi.org/10.1103/PhysRevD.8.3308)
- [12] C. M. Will, Theory and Experiment in Gravitational Physics (Cambridge University Press, Cambridge, England, 1993), revised ed.
- [13] C. M. Will, [Living Rev. Relativity](https://doi.org/10.12942/lrr-2014-4) 17, 4 (2014).
- [14] E. Berti, E. Barausse, V. Cardoso, L. Gualtieri, P. Pani, U. Sperhake, L. C. Stein, N.Wex, K. Yagi, T. Baker, C. P. Burgess, F. S. Coelho, D. Doneva, A. D. Felice, P. G. Ferreira, P. C. C. Freire, J. Healy, C. Herdeiro, M. Horbatsch, B. Kleihaus, et al., [Classical Quantum Gravity](https://doi.org/10.1088/0264-9381/32/24/243001) 32, 243001 (2015).
- [15] K. Chatziioannou, N. Yunes, and N. Cornish, [Phys. Rev. D](https://doi.org/10.1103/PhysRevD.86.022004) 86[, 022004 \(2012\).](https://doi.org/10.1103/PhysRevD.86.022004)
- [16] J. M. Weisberg, D. J. Nice, and J. H. Taylor, [Astrophys. J.](https://doi.org/10.1088/0004-637X/722/2/1030) 722[, 1030 \(2010\)](https://doi.org/10.1088/0004-637X/722/2/1030).
- [17] P. C. C. Freire, N. Wex, G. Esposito-Farèse, J. P. W. Verbiest, M. Bailes, B. A. Jacoby, M. Kramer, I. H. Stairs, J. Antoniadis, and G. H. Janssen, [Mon. Not. R. Astron. Soc.](https://doi.org/10.1111/j.1365-2966.2012.21253.x) 423[, 3328 \(2012\)](https://doi.org/10.1111/j.1365-2966.2012.21253.x).
- [18] I. H. Stairs, [Living Rev. Relativity](https://doi.org/10.12942/lrr-2003-5) 6, 5 (2003).
- [19] N. Wex, [arXiv:1402.5594.](http://arXiv.org/abs/1402.5594)
- [20] M. Isi, A. J. Weinstein, C. Mead, and M. Pitkin, [Phys. Rev.](https://doi.org/10.1103/PhysRevD.91.082002) D **91**[, 082002 \(2015\)](https://doi.org/10.1103/PhysRevD.91.082002).
- [21] M. Isi, M. Pitkin, and A. J. Weinstein, [Phys. Rev. D](https://doi.org/10.1103/PhysRevD.96.042001) 96, [042001 \(2017\).](https://doi.org/10.1103/PhysRevD.96.042001)
- [22] A. Nishizawa, A. Taruya, K. Hayama, S. Kawamura, and M.-a. Sakagami, Phys. Rev. D 79[, 082002 \(2009\)](https://doi.org/10.1103/PhysRevD.79.082002).
- [23] T. Callister, A. S. Biscoveanu, N. Christensen, M. Isi, A. Matas, O. Minazzoli, T. Regimbau, M. Sakellariadou, J. Tasson, and E. Thrane, Phys. Rev. X 7[, 041058 \(2017\)](https://doi.org/10.1103/PhysRevX.7.041058).
- [24] K. S. Thorne, in Three Hundred Years of Gravitation, edited by S. W. Hawking and W. Israel (Cambridge University Press, Cambridge, England, 1987), Chap. 9, pp. 330–458.
- [25] B. P. Abbott et al. (LIGO Scientific Collaboration and Virgo Collaboration), [Astrophys. J.](https://doi.org/10.3847/1538-4357/aa677f) 839, 12 (2017).
- [26] R. J. Dupuis and G. Woan, Phys. Rev. D **72**[, 102002 \(2005\).](https://doi.org/10.1103/PhysRevD.72.102002)
- [27] B. P. Abbott et al. (LIGO Scientific Collaboration), [Phys.](https://doi.org/10.1103/PhysRevD.95.062003) Rev. D 95[, 062003 \(2017\)](https://doi.org/10.1103/PhysRevD.95.062003).
- [28] W. G. Anderson, P. R. Brady, J. D. E. Creighton, and É. E. Flanagan, Phys. Rev. D 63[, 042003 \(2001\).](https://doi.org/10.1103/PhysRevD.63.042003)
- [29] A. Błaut, Phys. Rev. D **85**[, 043005 \(2012\)](https://doi.org/10.1103/PhysRevD.85.043005).
- [30] E. Poisson and C. M. Will, Gravity: Newtonian, Post-Newtonian, Relativistic (Cambridge University Press, Cambridge, England, 2014).
- [31] M. Zimmermann and E. Szedenits, [Phys. Rev. D](https://doi.org/10.1103/PhysRevD.20.351) 20, 351 [\(1979\).](https://doi.org/10.1103/PhysRevD.20.351)
- [32] B. J. Owen, L. Lindblom, C. Cutler, B. F. Schutz, A. Vecchio, and N. Andersson, Phys. Rev. D 58[, 084020 \(1998\)](https://doi.org/10.1103/PhysRevD.58.084020).
- [33] R. Bondarescu, S. A. Teukolsky, and I. Wasserman, [Phys.](https://doi.org/10.1103/PhysRevD.79.104003) Rev. D 79[, 104003 \(2009\)](https://doi.org/10.1103/PhysRevD.79.104003).
- [34] D. I. Jones and N. Andersson, [Mon. Not. R. Astron. Soc.](https://doi.org/10.1046/j.1365-8711.2002.05180.x) 331[, 203 \(2002\)](https://doi.org/10.1046/j.1365-8711.2002.05180.x).
- [35] M. Pitkin, C. Gill, J. Veitch, E. Macdonald, and G. Woan, [J. Phys. Conf. Ser.](https://doi.org/10.1088/1742-6596/363/1/012041) 363, 012041 (2012).
- [36] D. Keitel, R. Prix, M. A. Papa, P. Leaci, and M. Siddiqi, Phys. Rev. D 89[, 064023 \(2014\)](https://doi.org/10.1103/PhysRevD.89.064023).
- [37] E. Jaynes, [IEEE Trans. Syst. Sci. Cybern.](https://doi.org/10.1109/TSSC.1968.300117) 4, 227 (1968).
- [38] The specific range chosen for the amplitude priors has little effect on our results, as explained in Appendix B of Ref. [21].
- [39] R. D. Ferdman, I. H. Stairs, M. Kramer, R. P. Breton, M. A. McLaughlin, P. C. C. Freire, A. Possenti, B. W. Stappers, V. M. Kaspi, R. N. Manchester, and A. G. Lyne, [Astro](https://doi.org/10.1088/0004-637X/767/1/85)phys. J. 767[, 85 \(2013\).](https://doi.org/10.1088/0004-637X/767/1/85)
- [40] B. J. Rickett, W. A. Coles, C. F. Nava, M. A. McLaughlin, S. M. Ransom, F. Camilo, R. D. Ferdman, P. C. C. Freire, M. Kramer, A. G. Lyne, and I. H. Stairs, [Astrophys. J.](https://doi.org/10.1088/0004-637X/787/2/161) 787, 161 [\(2014\).](https://doi.org/10.1088/0004-637X/787/2/161)
- [41] W. W. Zhu, I. H. Stairs, P. B. Demorest, D. J. Nice, J. A. Ellis, S. M. Ransom, Z. Arzoumanian, K. Crowter, T. Dolch, R. D. Ferdman, E. Fonseca, M. E. Gonzalez, G. Jones, M. L. Jones, M. T. Lam, L. Levin, M. A. McLaughlin, T. Pennucci, K. Stovall, and J. Swiggum, [Astrophys. J.](https://doi.org/10.1088/0004-637X/809/1/41) 809, 41 (2015).
- [42] J. Aasi et al. (LIGO Scientific Collaboration and Virgo Collaboration), [Astrophys. J.](https://doi.org/10.1088/0004-637X/785/2/119) 785, 119 (2014).
- [43] C. Ng and R. W. Romani, [Astrophys. J.](https://doi.org/10.1086/380486) 601, 479 (2004).
- [44] C. Ng and R. W. Romani, [Astrophys. J.](https://doi.org/10.1086/523935) 673, 411 (2008).
- [45] See Supplemental Material at [http://link.aps.org/](http://link.aps.org/supplemental/10.1103/PhysRevLett.120.031104) [supplemental/10.1103/PhysRevLett.120.031104](http://link.aps.org/supplemental/10.1103/PhysRevLett.120.031104) for

expanded tables with odds and upper-limits for all non-GR hypotheses.

- [46] R. E. Kass and A. E. Raftery, [J. Am. Stat. Assoc.](https://doi.org/10.1080/01621459.1995.10476572) 90, 773 [\(1995\).](https://doi.org/10.1080/01621459.1995.10476572)
- [47] H. Jeffreys, *Theory of Probability*, 3rd ed. (Clarendon Press, Oxford, 1998).
- [48] C. P. Robert, N. Chopin, and J. Rousseau, [Stat. Sci.](https://doi.org/10.1214/09-STS284) 24, 141 [\(2009\).](https://doi.org/10.1214/09-STS284)
- [49] B. P. Abbott et al. (LIGO Scientific Collaboration and Virgo Collaboration), Phys. Rev. D 94[, 102002 \(2016\)](https://doi.org/10.1103/PhysRevD.94.102002).
- [50] B. P. Abbott, M. A. Papa, H.-B. Eggenstein, and S. Walsh (LIGO Scientific Collaboration and Virgo Collaboration), Phys. Rev. D 96[, 082003 \(2017\)](https://doi.org/10.1103/PhysRevD.96.082003).
- [51] B. P. Abbott et al. (LIGO Scientific Collaboration and Virgo Collaboration), Phys. Rev. D 94[, 042002 \(2016\)](https://doi.org/10.1103/PhysRevD.94.042002).
- [52] B. P. Abbott et al. (LIGO Scientific Collaboration and Virgo Collaboration), Phys. Rev. D 96[, 062002 \(2017\)](https://doi.org/10.1103/PhysRevD.96.062002).
- [53] J. Aasi et al. (LIGO Scientific Collaboration and Virgo Collaboration), Phys. Rev. D 90[, 062010 \(2014\)](https://doi.org/10.1103/PhysRevD.90.062010).

B. P. Abbott,¹ R. Abbott,¹ T. D. Abbott,² F. Acernese,^{3,4} K. Ackley,⁵ C. Adams,⁶ T. Adams,⁷ P. Addesso,⁸ R. X. Adhikari,¹ V. B. Adya, ⁹ C. Affeldt, ⁹ M. Afrough, ¹⁰ B. Agarwal, ¹¹ M. Agathos, ¹² K. Agatsuma, ¹³ N. Aggarwal, ¹⁴ O. D. Aguiar, ¹⁵ L. Aiello,^{16,17} A. Ain,¹⁸ P. Ajith,¹⁹ G. Allen,¹¹ A. Allocca,^{20,21} P. A. Altin,²² A. Amato,²³ A. Ananyeva,¹ S. B. Anderson,¹ W. G. Anderson,²⁴ S. Antier,²⁵ S. Appert,¹ K. Arai,¹ M. C. Araya,¹ J. S. Areeda,²⁶ N. Arnaud,^{25,27} K. G. Arun,²⁸ S. Ascenzi, $29,17$ G. Ashton, 9 M. Ast, 30 S. M. Aston, 6 P. Astone, 31 P. Aufmuth, 32 C. Aulbert, 9 K. AultONeal, 33 A. Avila-Alvarez,²⁶ S. Babak,³⁴ P. Bacon,³⁵ M. K. M. Bader,¹³ S. Bae,³⁶ P. T. Baker,^{37,38} F. Baldaccini,^{39,40} G. Ballardin,²⁷ S. W. Ballmer,⁴¹ S. Banagiri,⁴² J. C. Barayoga,¹ S. E. Barclay,⁴³ B. C. Barish,¹ D. Barker,⁴⁴ F. Barone,^{3,4} B. Barr,⁴³ L. Barsotti,¹⁴ M. Barsuglia,³⁵ D. Barta,⁴⁵ J. Bartlett,⁴⁴ I. Bartos,⁴⁶ R. Bassiri,⁴⁷ A. Basti,^{20,21} J. C. Batch,⁴⁴ C. Baune,⁹ M. Bawaj, 48,40 M. Bazzan, 49,50 B. Bécsy, 51 C. Beer, 9 M. Bejger, 52 I. Belahcene, 25 A. S. Bell, 43 B. K. Berger, 1 G. Bergmann, 9 C. P. L. Berry,⁵³ D. Bersanetti,^{54,55} A. Bertolini,¹³ J. Betzwieser, S. Bhagwat,⁴¹ R. Bhandare,⁵⁶ I. A. Bilenko,⁵⁷ G. Billingsley,¹ C. R. Billman,⁵ J. Birch,⁶ R. Birney,⁵⁸ O. Birnholtz,⁹ S. Biscans,¹⁴ A. Bisht,³² M. Bitossi,^{27,21} C. Biwer,⁴¹ M. A. Bizouard,²⁵ J. K. Blackburn,¹ J. Blackman,⁵⁹ C. D. Blair,⁶⁰ D. G. Blair,⁶⁰ R. M. Blair,⁴⁴ S. Bloemen,⁶¹ O. Bock,⁹ N. Bode,⁹ M. Boer,⁶² G. Bogaert,⁶² A. Bohe,³⁴ F. Bondu,⁶³ R. Bonnand,⁷ B. A. Boom,¹³ R. Bork,¹ V. Boschi,^{20,21} S. Bose,^{64,18} Y. Bouffanais,³⁵ A. Bozzi,²⁷ C. Bradaschia,²¹ P. R. Brady,²⁴ V. B. Braginsky,^{57,†} M. Branchesi,^{65,66} J. E. Brau,⁶⁷ T. Briant,⁶⁸ A. Brillet,⁶² M. Brinkmann,⁹ V. Brisson,²⁵ P. Brockill,²⁴ J. E. Broida,⁶⁹ A. F. Brooks,¹ D. A. Brown,⁴¹ D. D. Brown,⁵³ N. M. Brown,¹⁴ S. Brunett,¹ C. C. Buchanan,² A. Buikema,¹⁴ T. Bulik,⁷⁰ H. J. Bulten,^{71,13} A. Buonanno,^{34,72} D. Buskulic,⁷ C. Buy,³⁵ R. L. Byer,⁴⁷ M. Cabero,⁹ L. Cadonati,⁷³ G. Cagnoli,^{23,74} C. Cahillane,¹ J. Calderón Bustillo,⁷³ T. A. Callister,¹ E. Calloni,^{75,4} J. B. Camp,⁷⁶ M. Canepa,^{54,55} P. Canizares,⁶¹ K. C. Cannon,⁷⁷ H. Cao,⁷⁸ J. Cao,⁷⁹ C. D. Capano, ⁹ E. Capocasa, ³⁵ F. Carbognani, ²⁷ S. Caride, ⁸⁰ M. F. Carney, ⁸¹ J. Casanueva Diaz, ²⁵ C. Casentini, ^{29,17} S. Caudill,²⁴ M. Cavaglià,¹⁰ F. Cavalier,²⁵ R. Cavalieri,²⁷ G. Cella,²¹ C. B. Cepeda,¹ L. Cerboni Baiardi,^{65,66} G. Cerretani,^{20,21} E. Cesarini,^{29,17} S. J. Chamberlin,⁸² M. Chan,⁴³ S. Chao,⁸³ P. Charlton,⁸⁴ E. Chassande-Mottin,³⁵ D. Chatterjee, 24 B. D. Cheeseboro, 37,38 H. Y. Chen, 85 Y. Chen, 59 H.-P. Cheng, 5 A. Chincarini, 55 A. Chiummo, 27 T. Chmiel, 81 H. S. Cho,⁸⁶ M. Cho,⁷² J. H. Chow,²² N. Christensen,^{69,62} Q. Chu,⁶⁰ A. J. K. Chua,¹² S. Chua,⁶⁸ A. K. W. Chung,⁸⁷ S. Chung, 60 G. Ciani, ⁵ R. Ciolfi, 88,89 C. E. Cirelli, 47 A. Cirone, 54,55 F. Clara, 44 J. A. Clark, 73 F. Cleva, 62 C. Cocchieri, 10 E. Coccia, ^{16,17} P.-F. Cohadon,⁶⁸ A. Colla, ^{90,31} C. G. Collette, ⁹¹ L. R. Cominsky, ⁹² M. Constancio Jr., ¹⁵ L. Conti, ⁵⁰ S. J. Cooper,⁵³ P. Corban,⁶ T. R. Corbitt,² K. R. Corley,⁴⁶ N. Cornish,⁹³ A. Corsi,⁸⁰ S. Cortese,²⁷ C. A. Costa,¹⁵ M. W. Coughlin,⁶⁹ S. B. Coughlin,^{94,95} J.-P. Coulon,⁶² S. T. Countryman,⁴⁶ P. Couvares,¹ P. B. Covas,⁹⁶ E. E. Cowan,⁷³ D. M. Coward,⁶⁰ M. J. Cowart,⁶ D. C. Coyne,¹ R. Coyne,⁸⁰ J. D. E. Creighton,²⁴ T. D. Creighton,⁹⁷ J. Cripe,² S. G. Crowder,⁹⁸ T. J. Cullen,²⁶ A. Cumming,⁴³ L. Cunningham,⁴³ E. Cuoco,²⁷ T. Dal Canton,⁷⁶ S. L. Danilishin,^{32,9} S. D'Antonio,¹⁷ K. Danzmann,^{32,9} A. Dasgupta,⁹⁹ C. F. Da Silva Costa,⁵ V. Dattilo,²⁷ I. Dave,⁵⁶ M. Davier,²⁵ D. Davis,⁴¹

E. J. Daw,¹⁰⁰ B. Day,⁷³ S. De,⁴¹ D. DeBra,⁴⁷ J. Degallaix,²³ M. De Laurentis,^{75,4} S. Deléglise,⁶⁸ W. Del Pozzo,^{53,20,21} T. Denker, 9 T. Dent, 9 V. Dergachev, 34 R. De Rosa, $75,4$ R. T. DeRosa, 6 R. DeSalvo, 101 J. Devenson, 58 R. C. Devine, $37,38$ S. Dhurandhar,¹⁸ M. C. Díaz, $\overline{97}$ L. Di Fiore,⁴ M. Di Giovanni,^{102,89} T. Di Girolamo,^{75,4,46} A. Di Lieto,^{20,21} S. Di Pace,^{90,31} I. Di Palma, $90,31$ F. Di Renzo, $20,21$ Z. Doctor, 85 V. Dolique, 23 F. Donovan, 14 K. L. Dooley, 10 S. Doravari, 9 I. Dorrington, 95 R. Douglas,⁴³ M. Dovale Álvarez,⁵³ T. P. Downes,²⁴ M. Drago,⁹ R. W. P. Drever,^{1,†} J. C. Driggers,⁴⁴ Z. Du,⁷⁹ M. Ducrot,⁷ J. Duncan, 94 S. E. Dwyer, 44 T. B. Edo, 100 M. C. Edwards, 69 A. Effler, 6 H.-B. Eggenstein, 9 P. Ehrens, 1 J. Eichholz, 1 S. S. Eikenberry,⁵ R. A. Eisenstein,¹⁴ R. C. Essick,¹⁴ Z. B. Etienne,^{37,38} T. Etzel,¹ M. Evans,¹⁴ T. M. Evans,⁶ M. Factourovich,⁴⁶ V. Fafone,^{29,17,16} H. Fair,⁴¹ S. Fairhurst,⁹⁵ X. Fan,⁷⁹ S. Farinon,⁵⁵ B. Farr,⁸⁵ W. M. Farr,⁵³ E. J. Fauchon-Jones,⁹⁵ M. Favata,¹⁰³ M. Fays,⁹⁵ H. Fehrmann,⁹ J. Feicht,¹ M. M. Fejer,⁴⁷ A. Fernandez-Galiana,¹⁴ I. Ferrante,^{20,21} E.C. Ferreira,¹⁵ F. Ferrini,²⁷ F. Fidecaro,^{20,21} I. Fiori,²⁷ D. Fiorucci,³⁵ R. P. Fisher,⁴¹ R. Flaminio,^{23,104} M. Fletcher,⁴³ H. Fong,¹⁰⁵ P. W. F. Forsyth,²² S. S. Forsyth,⁷³ J.-D. Fournier,⁶² S. Frasca,^{90,31} F. Frasconi,²¹ Z. Frei,⁵¹ A. Freise,⁵³ R. Frey,⁶⁷ V. Frey,²⁵ E. M. Fries,¹ P. Fritschel,¹⁴ V. V. Frolov,⁶ P. Fulda,^{5,76} M. Fyffe,⁶ H. Gabbard,⁹ M. Gabel,¹⁰⁶ B. U. Gadre,¹⁸ S. M. Gaebel,⁵³ J. R. Gair,¹⁰⁷ L. Gammaitoni,³⁹ M. R. Ganija,⁷⁸ S. G. Gaonkar,¹⁸ F. Garufi,^{75,4} S. Gaudio,³³ G. Gaur, 108 V. Gayathri, 109 N. Gehrels, $^{76, \dagger}$ G. Gemme, 55 E. Genin, 27 A. Gennai, 21 D. George, 11 J. George, 56 L. Gergely, 110 V. Germain,⁷ S. Ghonge,⁷³ Abhirup Ghosh,¹⁹ Archisman Ghosh,^{19,13} S. Ghosh,^{61,13} J. A. Giaime,^{2,6} K. D. Giardina,⁶ A. Giazotto,²¹ K. Gill,³³ L. Glover,¹⁰¹ E. Goetz,⁹ R. Goetz,⁵ S. Gomes,⁹⁵ G. González,² J. M. Gonzalez Castro,^{20,21} A. Gopakumar,¹¹¹ M. L. Gorodetsky,⁵⁷ S. E. Gossan,¹ M. Gosselin,²⁷ R. Gouaty,⁷ A. Grado,^{112,4} C. Graef,⁴³ M. Granata,²³ A. Grant,⁴³ S. Gras,¹⁴ C. Gray,⁴⁴ G. Greco,^{65,66} A. C. Green,⁵³ P. Groot,⁶¹ H. Grote,⁹ S. Grunewald,³⁴ P. Gruning,²⁵ G. M. Guidi,^{65,66} X. Guo,⁷⁹ A. Gupta,⁸² M. K. Gupta,⁹⁹ K. E. Gushwa,¹ E. K. Gustafson,¹ R. Gustafson,¹¹³ B. R. Hall,⁶⁴ E. D. Hall,¹ G. Hammond,⁴³ M. Haney,¹¹¹ M. M. Hanke,⁹ J. Hanks,⁴⁴ C. Hanna,⁸² O. A. Hannuksela,⁸⁷ J. Hanson,⁶ T. Hardwick,² J. Harms,^{65,66} G. M. Harry,¹¹⁴ I. W. Harry,³⁴ M. J. Hart,⁴³ C.-J. Haster,¹⁰⁵ K. Haughian,⁴³ J. Healy,¹¹⁵ A. Heidmann,⁶⁸ M. C. Heintze,⁶ H. Heitmann,⁶² P. Hello,²⁵ G. Hemming,²⁷ M. Hendry,⁴³ I. S. Heng,⁴³ J. Hennig,⁴³ J. Henry, 115 A. W. Heptonstall, ¹ M. Heurs, 9,32 S. Hild, 43 D. Hoak, 27 D. Hofman, 23 K. Holt, 6 D. E. Holz, 85 P. Hopkins, 95 C. Horst,²⁴ J. Hough,⁴³ E. A. Houston,⁴³ E. J. Howell,⁶⁰ Y. M. Hu,⁹ E. A. Huerta,¹¹ D. Huet,²⁵ B. Hughey,³³ S. Husa,⁹⁶ S. H. Huttner,⁴³ T. Huynh-Dinh,⁶ N. Indik,⁹ D. R. Ingram,⁴⁴ R. Inta,⁸⁰ G. Intini,^{90,31} H. N. Isa,⁴³ J.-M. Isac,⁶⁸ M. Isi,¹ B. R. Iyer,¹⁹ K. Izumi,⁴⁴ T. Jacqmin,⁶⁸ K. Jani,⁷³ P. Jaranowski,¹¹⁶ S. Jawahar,¹¹⁷ F. Jiménez-Forteza,⁹⁶ W. W. Johnson,² D. I. Jones, ¹¹⁸ R. Jones, ⁴³ R. J. G. Jonker, ¹³ L. Ju, ⁶⁰ J. Junker, ⁹ C. V. Kalaghatgi, ⁹⁵ V. Kalogera, ⁹⁴ S. Kandhasamy, ⁶ G. Kang, 36 J. B. Kanner, 1 S. Karki, 67 K. S. Karvinen, 9 M. Kasprzack, 2 M. Katolik, 11 E. Katsavounidis, 14 W. Katzman, 6 S. Kaufer,³² K. Kawabe,⁴⁴ F. Kéfélian,⁶² D. Keitel,⁴³ A. J. Kemball,¹¹ R. Kennedy,¹⁰⁰ C. Kent,⁹⁵ J. S. Key,¹¹⁹ F. Y. Khalili,⁵⁷ I. Khan,^{16,17} S. Khan,⁹ Z. Khan,⁹⁹ E. A. Khazanov,¹²⁰ N. Kijbunchoo,⁴⁴ Chunglee Kim,¹²¹ J. C. Kim,¹²² W. Kim,⁷⁸ W. S. Kim,¹²³ Y.-M. Kim,^{86,121} S. J. Kimbrell,⁷³ E. J. King,⁷⁸ P. J. King,⁴⁴ R. Kirchhoff,⁹ J. S. Kissel,⁴⁴ L. Kleybolte,³⁰ S. Klimenko,⁵ P. Koch,⁹ S. M. Koehlenbeck,⁹ S. Koley,¹³ V. Kondrashov,¹ A. Kontos,¹⁴ M. Korobko,³⁰ W. Z. Korth,¹ I. Kowalska,⁷⁰ D. B. Kozak,¹ C. Krämer,⁹ V. Kringel,⁹ B. Krishnan,⁹ A. Królak,^{124,125} G. Kuehn,⁹ P. Kumar,¹⁰⁵ R. Kumar,⁹⁹ S. Kumar,¹⁹ L. Kuo,⁸³ A. Kutynia,¹²⁴ S. Kwang,²⁴ B. D. Lackey,³⁴ K. H. Lai,⁸⁷ M. Landry,⁴⁴ R. N. Lang,²⁴ J. Lange,¹¹⁵ B. Lantz,⁴⁷ R. K. Lanza,¹⁴ A. Lartaux-Vollard,²⁵ P. D. Lasky,¹²⁶ M. Laxen,⁶ A. Lazzarini,¹ C. Lazzaro,⁵⁰ P. Leaci,^{90,31} S. Leavey,⁴³ C. H. Lee,⁸⁶ H. K. Lee,¹²⁷ H. M. Lee,¹²¹ H. W. Lee,¹²² K. Lee,⁴³ J. Lehmann,⁹ A. Lenon,^{37,38} M. Leonardi,^{102,89} N. Leroy,²⁵ N. Letendre,⁷ Y. Levin,¹²⁶ T. G. F. Li,⁸⁷ A. Libson,¹⁴ T. B. Littenberg,¹²⁸ J. Liu,⁶⁰ R. K. L. Lo,⁸⁷ N. A. Lockerbie,¹¹⁷ L. T. London,⁹⁵ J. E. Lord,⁴¹ M. Lorenzini,^{16,17} V. Loriette,¹²⁹ M. Lormand,⁶ G. Losurdo,²¹ J. D. Lough,^{9,32} C. O. Lousto,¹¹⁵ G. Lovelace,²⁶ H. Lück,^{32,9} D. Lumaca,^{29,17} A. P. Lundgren,⁹ R. Lynch,¹⁴ Y. Ma,⁵⁹ S. Macfoy,⁵⁸ B. Machenschalk,⁹ M. MacInnis,¹⁴ D. M. Macleod,² I. Magaña Hernandez,⁸⁷ F. Magaña-Sandoval,⁴¹ L. Magaña Zertuche,⁴¹ R. M. Magee,⁸² E. Majorana,³¹ I. Maksimovic,¹²⁹ N. Man,⁶² V. Mandic,⁴² V. Mangano,⁴³ G. L. Mansell,²² M. Manske,²⁴ M. Mantovani,²⁷ F. Marchesoni,^{48,40} F. Marion,⁷ S. Márka,⁴⁶ Z. Márka,⁴⁶ C. Markakis,¹¹ A. S. Markosyan,⁴⁷ E. Maros,¹ F. Martelli,^{65,66} L. Martellini,⁶² I. W. Martin,⁴³ D. V. Martynov,¹⁴ K. Mason,¹⁴ A. Masserot,⁷ T. J. Massinger,¹ M. Masso-Reid,⁴³ S. Mastrogiovanni,^{90,31} A. Matas,⁴² F. Matichard,¹⁴ L. Matone,⁴⁶ N. Mavalvala,¹⁴ N. Mazumder,⁶⁴ R. McCarthy,⁴⁴ D. E. McClelland,²² S. McCormick,⁶ L. McCuller,¹⁴ S. C. McGuire,¹³⁰ G. McIntyre,¹ J. McIver,¹ D. J. McManus,²² T. McRae,²² S. T. McWilliams,^{37,38} D. Meacher,⁸² G. D. Meadors,^{34,9} J. Meidam,¹³ E. Mejuto-Villa, ⁸ A. Melatos,¹³¹ G. Mendell,⁴⁴ R. A. Mercer,²⁴ E. L. Merilh,⁴⁴ M. Merzougui,⁶² S. Meshkov,¹ C. Messenger,⁴³ C. Messick,⁸² R. Metzdorff,⁶⁸ P. M. Meyers,⁴² F. Mezzani,^{31,90} H. Miao,⁵³ C. Michel,²³ H. Middleton,⁵³ E. E. Mikhailov, 132 L. Milano, 75,4 A. L. Miller, 5 A. Miller, 90,31 B. B. Miller, 94 J. Miller, 14 M. Millhouse, 93 O. Minazzoli, 62

Y. Minenkov,¹⁷ J. Ming,³⁴ C. Mishra,¹³³ S. Mitra,¹⁸ V. P. Mitrofanov,⁵⁷ G. Mitselmakher,⁵ R. Mittleman,¹⁴ A. Moggi,²¹ M. Mohan,²⁷ S. R. P. Mohapatra,¹⁴ M. Montani,^{65,66} B. C. Moore,¹⁰³ C. J. Moore,¹² D. Moraru,⁴⁴ G. Moreno,⁴⁴ S. R. Morriss, 97 B. Mours, 7 C. M. Mow-Lowry, 53 G. Mueller, 5 A. W. Muir, 95 Arunava Mukherjee, 9 D. Mukherjee, 24 S. Mukherjee,⁹⁷ N. Mukund,¹⁸ A. Mullavey, ⁶ J. Munch,⁷⁸ E. A. M. Muniz, ⁴¹ P. G. Murray, ⁴³ K. Napier, ⁷³ I. Nardecchia, ^{29,17} L. Naticchioni,^{90,31} R. K. Nayak,¹³⁴ G. Nelemans,^{61,13} T. J. N. Nelson,⁶ M. Neri,^{54,55} M. Nery,⁹ A. Neunzert,¹¹³ J. M. Newport, 114 G. Newton, $^{43, \dagger}$ K. K. Y. Ng, 87 T. T. Nguyen, 22 D. Nichols, 61 A. B. Nielsen, 9 S. Nissanke, 61,13 A. Nitz, 9 A. Noack,⁹ F. Nocera,²⁷ D. Nolting,⁶ M. E. N. Normandin,⁹⁷ L. K. Nuttall,⁴¹ J. Oberling,⁴⁴ E. Ochsner,²⁴ E. Oelker,¹⁴ G. H. Ogin,¹⁰⁶ J. J. Oh,¹²³ S. H. Oh,¹²³ F. Ohme,⁹ M. Oliver,⁹⁶ P. Oppermann,⁹ Richard J. Oram,⁶ B. O'Reilly,⁶ R. Ormiston,⁴² L. F. Ortega,⁵ R. O'Shaughnessy,¹¹⁵ D. J. Ottaway,⁷⁸ H. Overmier,⁶ B. J. Owen,⁸⁰ A. E. Pace,⁸² J. Page,¹²⁸ M. A. Page,⁶⁰ A. Pai,¹⁰⁹ S. A. Pai,⁵⁶ J. R. Palamos,⁶⁷ O. Palashov,¹²⁰ C. Palomba,³¹ A. Pal-Singh,³⁰ H. Pan,⁸³ B. Pang,⁵⁹ P. T. H. Pang, ⁸⁷ C. Pankow, ⁹⁴ F. Pannarale, ⁹⁵ B. C. Pant, ⁵⁶ F. Paoletti, ²¹ A. Paoli, ²⁷ M. A. Papa, ^{34, 24, 9} H. R. Paris, ⁴⁷ W. Parker,⁶ D. Pascucci,⁴³ A. Pasqualetti,²⁷ R. Passaquieti,^{20,21} D. Passuello,²¹ B. Patricelli,^{135,21} B. L. Pearlstone,⁴³ M. Pedraza,¹ R. Pedurand,^{23,136} L. Pekowsky,⁴¹ A. Pele,⁶ S. Penn,¹³⁷ C. J. Perez,⁴⁴ A. Perreca,^{1,102,89} L. M. Perri,⁹⁴ H. P. Pfeiffer,¹⁰⁵ M. Phelps,⁴³ O. J. Piccinni,^{90,31} M. Pichot,⁶² F. Piergiovanni,^{65,66} V. Pierro,⁸ G. Pillant,²⁷ L. Pinard,²³ I. M. Pinto, 8 M. Pitkin, 43 R. Poggiani, $20,21$ P. Popolizio, 27 E. K. Porter, 35 A. Post, 9 J. Powell, 43 J. Prasad, 18 J. W. W. Pratt, 33 V. Predoi, ⁹⁵ T. Prestegard, ²⁴ M. Prijatelj, ⁹ M. Principe, ⁸ S. Privitera, ³⁴ R. Prix, ⁹ G. A. Prodi, ^{102,89} L. G. Prokhorov, ⁵⁷ O. Puncken,⁹ M. Punturo,⁴⁰ P. Puppo,³¹ M. Pürrer,³⁴ H. Qi,²⁴ J. Qin,⁶⁰ S. Qiu,¹²⁶ V. Quetschke,⁹⁷ E. A. Quintero,¹ R. Quitzow-James, ⁶⁷ F. J. Raab, ⁴⁴ D. S. Rabeling, ²² H. Radkins, ⁴⁴ P. Raffai, ⁵¹ S. Raja, ⁵⁶ C. Rajan, ⁵⁶ M. Rakhmanov, ⁹⁷ K. E. Ramirez, ⁹⁷ P. Rapagnani, ^{90,31} V. Raymond, ³⁴ M. Razzano, ^{20,21} J. Read, ²⁶ T. Regimbau, ⁶² L. Rei, ⁵⁵ S. Reid, ⁵⁸ D. H. Reitze,^{1,5} H. Rew,¹³² S. D. Reyes,⁴¹ F. Ricci,^{90,31} P. M. Ricker,¹¹ S. Rieger,⁹ K. Riles,¹¹³ M. Rizzo,¹¹⁵ N. A. Robertson, 1,43 R. Robie, 43 F. Robinet, 25 A. Rocchi, 17 L. Rolland, 7 J. G. Rollins, 1 V. J. Roma, 67 R. Romano, 3,4 C. L. Romel,⁴⁴ J. H. Romie,⁶ D. Rosińska,^{138,52} M. P. Ross,¹³⁹ S. Rowan,⁴³ A. Rüdiger,⁹ P. Ruggi,²⁷ K. Ryan,⁴⁴ S. Sachdev,¹ T. Sadecki,⁴⁴ L. Sadeghian,²⁴ M. Sakellariadou,¹⁴⁰ L. Salconi,²⁷ M. Saleem,¹⁰⁹ F. Salemi,⁹ A. Samajdar,¹³⁴ L. Sammut,¹²⁶ L. M. Sampson, 94 E. J. Sanchez, V. Sandberg, 44 B. Sandeen, 94 J. R. Sanders, 41 B. Sassolas, 23 B. S. Sathyaprakash, $82,95$ P. R. Saulson,⁴¹ O. Sauter,¹¹³ R. L. Savage,⁴⁴ A. Sawadsky,³² P. Schale,⁶⁷ J. Scheuer,⁹⁴ E. Schmidt,³³ J. Schmidt,⁹ P. Schmidt,^{1,61} R. Schnabel,³⁰ R. M. S. Schofield,⁶⁷ A. Schönbeck,³⁰ E. Schreiber,⁹ D. Schuette,^{9,32} B. W. Schulte,⁹ B. F. Schutz,^{95,9} S. G. Schwalbe,³³ J. Scott,⁴³ S. M. Scott,²² E. Seidel,¹¹ D. Sellers,⁶ A. S. Sengupta,¹⁴¹ D. Sentenac,²⁷ V. Sequino,^{29,17} A. Sergeev,¹²⁰ D. A. Shaddock,²² T. J. Shaffer,⁴⁴ A. A. Shah,¹²⁸ M. S. Shahriar,⁹⁴ L. Shao,³⁴ B. Shapiro,⁴⁷ P. Shawhan,⁷² A. Sheperd,²⁴ D. H. Shoemaker,¹⁴ D. M. Shoemaker,⁷³ K. Siellez,⁷³ X. Siemens,²⁴ M. Sieniawska,⁵² D. Sigg,⁴⁴ A. D. Silva,¹⁵ A. Singer,¹ L. P. Singer,⁷⁶ A. Singh,^{34,9,32} R. Singh,² A. Singhal,^{16,31} A. M. Sintes,⁹⁶ B. J. J. Slagmolen,²² B. Smith,⁶ J. R. Smith,²⁶ R. J. E. Smith,¹ E. J. Son,¹²³ J. A. Sonnenberg,²⁴ B. Sorazu,⁴³ F. Sorrentino,⁵⁵ T. Souradeep,¹⁸ A. P. Spencer,⁴³ A. K. Srivastava,⁹⁹ A. Staley,⁴⁶ M. Steinke,⁹ J. Steinlechner,^{43,30} S. Steinlechner,³⁰ D. Steinmeyer,^{9,32} B. C. Stephens,²⁴ R. Stone,⁹⁷ K. A. Strain,⁴³ G. Stratta,^{65,66} S. E. Strigin,⁵⁷ R. Sturani,¹⁴² A. L. Stuver,⁶ T. Z. Summerscales,¹⁴³ L. Sun,¹³¹ S. Sunil,⁹⁹ P. J. Sutton,⁹⁵ B. L. Swinkels,²⁷ M. J. Szczepańczyk,³³ M. Tacca,³⁵ D. Talukder,⁶⁷ D. B. Tanner,⁵ M. Tápai,¹¹⁰ A. Taracchini,³⁴ J. A. Taylor,¹²⁸ R. Taylor,¹ T. Theeg,⁹ E. G. Thomas,⁵³ M. Thomas, 6 P. Thomas, 44 K. A. Thorne, 6 K. S. Thorne, 59 E. Thrane, 126 S. Tiwari, $16,89$ V. Tiwari, 95 K. V. Tokmakov, 117 K. Toland,⁴³ M. Tonelli,^{20,21} Z. Tornasi,⁴³ C. I. Torrie,¹ D. Töyrä,⁵³ F. Travasso,^{27,40} G. Traylor,⁶ D. Trifirò,¹⁰ J. Trinastic,⁵ M. C. Tringali,^{102,89} L. Trozzo,^{144,21} K. W. Tsang,¹³ M. Tse,¹⁴ R. Tso,¹ D. Tuyenbayev,⁹⁷ K. Ueno,²⁴ D. Ugolini,¹⁴⁵ C. S. Unnikrishnan,¹¹¹ A. L. Urban,¹ S. A. Usman,⁹⁵ H. Vahlbruch,³² G. Vajente,¹ G. Valdes,⁹⁷ M. Vallisneri,⁵⁹ N. van Bakel,¹³ M. van Beuzekom,¹³ J. F. J. van den Brand,^{71,13} C. Van Den Broeck,¹³ D. C. Vander-Hyde,⁴¹ L. van der Schaaf,¹³ J. V. van Heijningen,¹³ A. A. van Veggel,⁴³ M. Vardaro,^{49,50} V. Varma,⁵⁹ S. Vass,¹ M. Vasúth,⁴⁵ A. Vecchio,⁵³ G. Vedovato,⁵⁰ J. Veitch,⁵³ P. J. Veitch,⁷⁸ K. Venkateswara,¹³⁹ G. Venugopalan,¹ D. Verkindt,⁷ F. Vetrano,^{65,66} A. Viceré,^{65,66} A. D. Viets,²⁴ S. Vinciguerra,⁵³ D. J. Vine,⁵⁸ J.-Y. Vinet,⁶² S. Vitale,¹⁴ T. Vo,⁴¹ H. Vocca,^{39,40} C. Vorvick,⁴⁴ D. V. Voss,⁵ W. D. Vousden,⁵³ S. P. Vyatchanin,⁵⁷ A. R. Wade,¹ L. E. Wade,⁸¹ M. Wade,⁸¹ R. Walet,¹³ M. Walker,² L. Wallace,¹ S. Walsh,²⁴ G. Wang,^{16,66} H. Wang,⁵³ J. Z. Wang,⁸² M. Wang,⁵³ Y.-F. Wang,⁸⁷ Y. Wang,⁶⁰ R. L. Ward,²² J. Warner,⁴⁴ M. Was,⁷ J. Watchi,⁹¹ B. Weaver,⁴⁴ L.-W. Wei,^{9,32} M. Weinert,⁹ A. J. Weinstein,¹ R. Weiss,¹⁴ L. Wen,⁶⁰ E. K. Wessel,¹¹ P. Weßels,⁹ T. Westphal,⁹ K. Wette,⁹ J. T. Whelan,¹¹⁵ B. F. Whiting,⁵ C. Whittle,¹²⁶ D. Williams,⁴³ R. D. Williams,¹ A. R. Williamson,¹¹⁵ J. L. Willis,¹⁴⁶ B. Willke,^{32,9} M. H. Wimmer,^{9,32} W. Winkler,⁹ C. C. Wipf,¹ H. Wittel, 9.32 G. Woan, 43 J. Woehler, 9 J. Wofford, 115 K. W. K. Wong, 87 J. Worden, 44 J. L. Wright, 43 D. S. Wu, 9 G. Wu, 6 W. Yam,¹⁴ H. Yamamoto,¹ C. C. Yancey,⁷² M. J. Yap,²² Hang Yu,¹⁴ Haocun Yu,¹⁴ M. Yvert,⁷ A. Zadrożny,¹²⁴ M. Zanolin,³³ T. Zelenova,²⁷ J.-P. Zendri,⁵⁰ M. Zevin,⁹⁴ L. Zhang,¹ M. Zhang,¹³² T. Zhang,⁴³ Y.-H. Zhang,¹¹⁵ C. Zhao,⁶⁰ M. Zhou,⁹⁴ Z. Zhou,⁹⁴ S. J. Zhu,^{34,9} X. J. Zhu,⁶⁰ M. E. Zucker,^{1,14} and J. Zweizig¹

(LIGO Scientific Collaboration and Virgo Collaboration)

S. Buchner, 147,148 I. Cognard, 149,150 A. Corongiu, 151 P. C. C. Freire, 152 L. Guillemot, 149,150 G. B. Hobbs, 153 M. Kerr, 154 A. G. Lyne,¹⁵⁵ A. Possenti,¹⁵¹ A. Ridolfi,¹⁵² R. M. Shannon,^{153,156} B. W. Stappers,¹⁵⁵ and P. Weltevrede¹⁵⁵

¹LIGO, California Institute of Technology, Pasadena, California 91125, USA
 $\frac{2}{5}I$ quisique State University, Pator Pouce, Louisique 70803, USA

 2 Louisiana State University, Baton Rouge, Louisiana 70803, USA 3 Università di Salerno, Fisciano, I-84084 Salerno, Italy

Università di Salerno, Fisciano, I-84084 Salerno, Italy $\frac{4}{NEN}$ Satisma di Napoli Complexes Universitatio di Monte Schnocle J.

INFN, Sezione di Napoli, Complesso Universitario di Monte S.Angelo, I-80126 Napoli, Italy
⁵Llaivarsity of Florida, Cainewilla, Florida 32611, USA

 5 University of Florida, Gainesville, Florida 32611, USA

 ${}^{6}LIGO$ Livingston Observatory, Livingston, Louisiana 70754, USA

 U^{\dagger} Laboratoire d'Annecy-le-Vieux de Physique des Particules (LAPP), Université Savoie Mont Blanc,

CNRS/IN2P3, F-74941 Annecy, France
⁸University of Sannie at Penevante, L82100 Benevante, Italy and INEN. See

University of Sannio at Benevento, I-82100 Benevento, Italy and INFN, Sezione di Napoli, I-80100 Napoli, Italy
⁹ Albert Finatein Institut, Max Blanck Institut für Cravitationsplassik, D. 20167 Hannover, Cermany

²Albert-Einstein-Institut, Max-Planck-Institut für Gravitationsphysik, D-30167 Hannover, Germany

¹¹NCSA, University of Mississippi, University, Mississippi 38677, USA

¹¹NCSA, University of Illinois at Urbana-Champ

²⁹Università di Roma Tor Vergata, I-00133 Roma, Italy
³⁰Universität Hamburg, D-22761 Hamburg, Germany
³¹INFN, Sezione di Roma, I-00185 Roma, Italy
³¹INFN, Sezione di Roma, I-00185 Roma, Italy
³²Leibniz Universitä

³⁶ Korea Institute of Science and Technology Information, Daejeon 34141, Korea
³⁷ West Virginia University, Morgantown, West Virginia 26506, USA
³⁸ Center for Gravitational Waves and Cosmology, West Virginia Univers

⁴⁵Wigner RCP, RMKI, H-1121 Budapest, Konkoly Thege Miklós út 29-33, Hungary ⁴⁶Columbia University, New York, New York 10027, USA ⁴⁷Stanford University, Stanford, California 94305, USA

⁴⁸Università di Camerino, Dipartimento di Fisica, I-62032 Camerino, Italy

PHYSICAL REVIEW LETTERS 120, 031104 (2018)

⁴⁹Università di Padova, Dipartimento di Fisica e Astronomia, I-35131 Padova, Italy
⁵⁰INFN, Sezione di Padova, I-35131 Padova, Italy
⁵¹MTA Eötvös University, "Lendulet" Astrophysics Research Group, Budapest 1117, Hun

⁵⁷Faculty of Physics, Lomonosov Moscow State University, Moscow 119991, Russia

⁵⁸SUPA, University of the West of Scotland, Paisley PA1 2BE, United Kingdom
⁵⁹Caltech CaRT, Pasadena, California 91125, USA

⁶⁰OzGrav, University of Western Australia, Crawley, Western Australia 6009, Australia
⁶¹Department of Astrophysics/IMAPP, Radboud University Nijmegen, P.O. Box 9010, 6500 GL Nijmegen, Netherlands
⁶²Artemis, Universi

⁶³Institut de Physique de Rennes, CNRS, Université de Rennes 1, F-35042 Rennes, France
⁶⁴Washington State University, Pullman, Washington 99164, USA
⁶⁵Università degli Studi di Urbino "Carlo Bo," I-61029 Urbino, Ita

Collège de France, F-75005 Paris, France
⁶⁹Carleton College, Northfield, Minnesota 55057, USA

⁷⁰Astronomical Observatory Warsaw University, 00-478 Warsaw, Poland ⁷¹VU University Amsterdam, 1081 HV Amsterdam, Netherlands ⁷²University of Maryland, College Park, Maryland 20742, USA

⁷²Center for Relativistic Astrophysics and School of Physics, Georgia Institute of Technology, Atlanta, Georgia 30332, USA
⁷³Center for Relativistic Astrophysics and School of Physics, Georgia Institute of Technology,

⁸³National Tsing Hua University, Hsinchu City, 30013 Taiwan, Republic of China
⁸⁴Charles Sturt University, Wagga Wagga, New South Wales 2678, Australia
⁸⁵University of Chicago, Chicago, Illinois 60637, USA
⁸⁶Pusan

⁸⁷The Chinese University of Hong Kong, Shatin, New Territories, Hong Kong
⁸⁸INAF, Osservatorio Astronomico di Padova, Vicolo dell'Osservatorio 5, 1-35122 Padova, Italy
⁸⁹INFN, Trento Institute for Fundamental Physic

 94 Center for Interdisciplinary Exploration & Research in Astrophysics (CIERA), Northwestern University,

Evanston, Illinois 60208, USA

⁹⁵Cardiff University, Cardiff CF24 3AA, United Kingdom

⁹⁶Universitat de les Illes Balears, IAC3—IEEC, E-07122 Palma de Mallorca, Spain
⁹⁷The University of Texas Rio Grande Valley, Brownsville, Texas 78520, USA
⁹⁸Bellevue College, Bellevue, Washington 98007, USA
⁹⁹Inst

¹⁰¹California State University, Los Angeles, 5151 State University Drive, Los Angeles, California 90032, USA
¹⁰²Università di Trento, Dipartimento di Fisica, I-38123 Povo, Trento, Italy
¹⁰³Montclair State University

¹⁰⁷School of Mathematics, University of Edinburgh, Edinburgh EH9 3FD, United Kingdom

¹⁰⁸University and Institute of Advanced Research, Gandhinagar Gujarat 382007, India
¹⁰⁹IISER-TVM, CET Campus, Trivandrum Kerala 695016, India
¹¹⁰University of Szeged, Dóm tér 9, Szeged 6720, Hungary

¹¹¹Tata Institute of Fundamental Research, Mumbai 400005, India
¹¹²INAF, Osservatorio Astronomico di Capodimonte, I-80131, Napoli, Italy
¹¹³University of Michigan, Ann Arbor, Michigan 48109, USA
¹¹⁴American Univer

¹¹⁵Rochester Institute of Technology, Rochester, New York 14623, USA
¹¹⁶University of Biał ystok, 15-424 Biał ystok, Poland
¹¹⁶University of Strathclyde, Glasgow G1 IXQ. United Kingdom
¹¹⁹University of Swathampton

 $^{131}OzGr$ av, University of Melbourne, Parkville, Victoria 3010, Australia

 132 College of William and Mary, Williamsburg, Virginia 23187, USA
 133 Indian Institute of Technology Madras, Chennai 600036, India

¹³⁴IISER-Kolkata, Mohanpur, West Bengal 741252, India

¹³⁵Scuola Normale Superiore, Piazza dei Cavalier' 7, 1-56126 Pisa, Italy

¹³⁵Scuola Normale Supersité de Lyon, F-69361 Lyon, France

¹³⁷Hobart and William Smi

¹⁵⁰Station de Radioastronomie de Nançay, Observatoire de Paris, CNRS/INSU, F-18330 Nançay, France
¹⁵¹INAF—Osservatorio Astronomico di Cagliari, via della Scienza 5, 09047 Selargius, Italy
¹⁵²Max-Planck-Institut für

Manchester M13 9PL, United Kingdom
¹⁵⁶International Centre for Radio Astronomy Research, Curtin University, Bentley, Western Australia 6101, Australia

† Deceased.