

The Diurnal Cycles of Cloud Profiles and Water Vapor over Land and Ocean seen by the CATS Spaceborne Lidar and Megha-Tropiques Sensors

Vincent Noël, Hélène Chepfer, Marjolaine Chiriaco, Hélène Brogniez, John Yorks, Patrick Raberanto

▶ To cite this version:

Vincent Noël, Hélène Chepfer, Marjolaine Chiriaco, Hélène Brogniez, John Yorks, et al.. The Diurnal Cycles of Cloud Profiles and Water Vapor over Land and Ocean seen by the CATS Spaceborne Lidar and Megha-Tropiques Sensors. ATMOS 2018, Nov 2018, Salzburg, Austria. insu-01979657

HAL Id: insu-01979657 https://insu.hal.science/insu-01979657v1

Submitted on 28 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Diurnal Cycles of Cloud Profiles and Water Vapor seen by the CATS Lidar and Megha-Tropiques Spaceborne Sensors


- V. Noel LA, CNRS
- H. Chepfer LMD, UPMC
- M. Chiriaco LATMOS, UVSQ
- H. Brogniez LATMOS, UVSQ
- J. Yorks NASA Goddard
- P. Raberanto LMD, CNRS

Spaceborne lidars and clouds

- Since 2006 the spaceborne lidar CALIPSO maps the 3D distribution of clouds on a near-global scale with a high vertical resolution
- Many important and useful results
- CALIPSO and almost all spaceborne lidars -ADM-Aeolus, IceSAT 1 and 2, Earth-CARE...
 - follow a sun-synchronous orbit
 - always sample a given location at the same local time
 - capture a single moment of the diurnal cycle
- CATS is an exception

CATS The Cloud-Aerosol Transport System


- The CATS lidar operated on the ISS from Feb 2015 to Oct 2017 (NASA mission)
- ISS orbits ±51° latitudes = samples 78% of the Earth
- data:mostly 1064nm
- 350m between profiles, 60m vertical resolution, 5kHz



From single overpasses to the diurnal cycle

By aggregating overpasses along local time, we can reconstruct the diurnal cycle with vertical variability


JJA 2015-2017 ±51°



From single overpasses to the diurnal cycle

By aggregating overpasses along local time, we can reconstruct the diurnal cycle with vertical variability


JJA 2015-2017 ±51°




From single overpasses to the diurnal cycle

By aggregating overpasses along local time, we can reconstruct the diurnal cycle with vertical variability

JJA 2015-2017 ±51°




Diurnal cycles in smaller regions



Local Time


Local Time

Implications of CATS results for CALIPSO & sun-synchronous missions




- CATS at 0130PM ~ CF daily min
 CALIPSO daytime overpass
- CATS at 0130AM ~ CF daily max
 CALIPSO nighttime overpass
- CALIPSO = daily extremes of the cloud fraction profiles over ocean, especially below 10km
 ASL

Implications of CATS results for CALIPSO & sun-synchronous missions



- CATS at 0130PM ~ CF daily min
 = CALIPSO daytime overpass
- CATS at 0130AM ~ CF daily max
 = CALIPSO nighttime overpass
- CALIPSO = daily extremes of the cloud fraction profiles over ocean, especially below 10km ASL
- Also works for Earth-CARE
 (0200LT) & A-Train instruments


Implications of CATS results for CALIPSO & sun-synchronous missions



- CATS at 06AM/PM : intermediate CF profiles
 - = ADM-Aeolus overpasses

diurnal cycles: clouds and water vapour

LAND - w500 < 0 (convection) - JJA 2015-2017



Takeaway Points


- CATS and Megha-Tropiques can describe the diurnal cycle of vertical profiles of clouds and water vapor in the Tropics
- CATS data shows CALIPSO, Earth-CARE data document extremes of Cloud Fraction profiles over ocean
- Other factors will impact agreement between instruments, eg wavelength — cf. talk from M. Chiriaco, next

Noel, V., H. Chepfer, M. Chiriaco, and J. Yorks, 2018: The diurnal cycle of cloud profiles over land and ocean between 51°S and 51°N, seen by the CATS spaceborne lidar from the International Space Station, **Atmos. Chem. Phys. 18**, 9457–9473

Chepfer, H., H. Brogniez, V. Noel, P. Raberanto, 2018: Influence of water vapor and cloud profiles on temperature diurnal variation across the tropics. submitted.

backup slides

CATS at CALIOP local times




CALIOP overpasses any location at 0130AM or 0130PM local times

CATS overpasses at those local times are consistent with those derived from CALIPSO

Comparisons of CATS with groundbased lidar/radar data and ISCCP are also consistent (cf. Noel et al. 2018)


CATS clouds are reliable

JJA Results by latitude band




- low/high separation everywhere
- different cycles at different latitudes

JJA Results by latitude band




PBL clouds sun-activated

JJA Results by latitude band




JJA Results by latitude band




JJA Results by latitude band

Southern

Ocean





Even though some profiles get fully attenuated by atmospheric components, sampling is better than 80% at most altitudes, and ~100% above 7km except late in the afternoon

