



HAL
open science

Tentative integration of paleoseismic data from lake sediments and from nearby trenches: The central section of the Boconó Fault (northern Venezuela)

Christian Beck, Eduardo Carrillo, Franck Audemard, Aurelien van Welden,
Jean-Robert Disnar

► To cite this version:

Christian Beck, Eduardo Carrillo, Franck Audemard, Aurelien van Welden, Jean-Robert Disnar. Tentative integration of paleoseismic data from lake sediments and from nearby trenches: The central section of the Boconó Fault (northern Venezuela). *Journal of South American Earth Sciences*, 2019, 92, pp.646-657. <10.1016/j.jsames.2019.03.028>. <insu-02092447>

HAL Id: insu-02092447

<https://insu.hal.science/insu-02092447v1>

Submitted on 10 Jul 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Accepted Manuscript

Tentative integration of paleoseismic data from lake sediments and from nearby trenches: The central section of the Boconó Fault (northern Venezuela)

Christian Beck, Eduardo Carrillo, Franck Audemard, Aurélien van Welden, Jean-Robert Disnar



PII: S0895-9811(18)30110-X

DOI: <https://doi.org/10.1016/j.jsames.2019.03.028>

Reference: SAMES 2153

To appear in: *Journal of South American Earth Sciences*

Received Date: 30 March 2018

Revised Date: 29 March 2019

Accepted Date: 29 March 2019

Please cite this article as: Beck, C., Carrillo, E., Audemard, F., van Welden, Auré., Disnar, J.-R., Tentative integration of paleoseismic data from lake sediments and from nearby trenches: The central section of the Boconó Fault (northern Venezuela), *Journal of South American Earth Sciences* (2019), doi: <https://doi.org/10.1016/j.jsames.2019.03.028>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Ref.** SAMES_2018_99 Revised Version

2 **Title:** Tentative integration of paleoseismic data from lake sediments and from nearby
3 trenches: the central section of the Boconó Fault (northern Venezuela).

4 **Authors:**

5 Christian Beck¹, Eduardo Carrillo², Franck Audemard³, Aurélien van Welden^{1*}, Jean-Robert
6 Disnar⁴

7 **Adresses:**

8 (1) CNRS ISTERre, Université Savoie-Mont-Blanc, 73 376 Le Bourget du Lac, France.

9 beck@univ-smb.fr

10 (2) Instituto de Ciencias de la Tierra, Universidad Central de Venezuela, Caracas.

11 (3) Fundación Venezolana de Investigaciones Sismológicas FUNVISIS, El Llanito, Caracas
12 1073, Venezuela.

13 (4) CNRS ISTO, Campus Géosciences, Université d'Orléans, 45071 Orléans Cedex 2, France.

14 (*) Presently at CGGVeritas, OH Bangs, Vei 70, NH 1323 HØVIK, Norway.

15

16 **Abstract**

17 The right-lateral strike slip Boconó Fault (Mérida Andes, northern Venezuela)
18 accommodates an important part of the South-American Plate northern transform boundary.
19 Along its central portion, preserved post-LMG lake fills are intersected by two surface-
20 reaching active traces which could be trenched just beside. Outcropping and cored lacustrine
21 sedimentary archives are combined with trench data in order to achieve a Holocene
22 paleoseismicity record for a 7 km-long segment. For lakes sediments, several types of
23 sedimentary “events” were taken into account as co-seismic: mass wasting on deltaic foreset,
24 liquefaction and slumping, reflected tsunami effects, re-suspension, abrupt change in
25 sedimentary dynamics and sources, abrupt emptying and lake surface changes. Time
26 coincidences between two lacustrine archives and two trenches can be proposed for the last 10
27 kyr BP. Among a total of 24 events, 13 events are detected in two sites, 3 events in 3 sites. 9
28 possible correlations concern separate traces while 4 concern the same trace; a relay between
29 the activity of the two traces is also deduced. This combination of surveys both reinforces and
30 completes the trenches results, leading to a better knowledge of local to regional seismic
31 hazard. Nevertheless, the total information results probably incomplete and/or biased. The co-
32 seismic origin of lacustrine fills disturbances evidenced but the associated archive is
33 incomplete and/or biased due to: changing recording potential through time, possible impacts
34 by strong distant earthquakes. Trenches data appear to fill lacustrine “gaps” but with a number

35 of events possibly overestimated if all ruptures and associated ^{14}C data are considered as
36 representing separated earthquakes.

37

38 **Key-words:** paleosismicity, lake sediments, trenches, Last Glacial Maximum, Holocene,
39 Mérida Andes, Boconó Fault.

40

41 **1. Introduction:**

42 Since pioneer investigations and discoveries on sedimentary recording of seismic
43 shocks (e.g. Sims, 1973; Adams, 1990), “subaqueous” paleosismology widely developed both
44 for marine and for lacustrine accumulations. Meanwhile, paleoseismology based on field
45 survey, which began several decades before, largely benefited from the development and
46 improvement of trenches analyses (*in* McCalpin and Nelson, 2009; Weldon et al, 2009).
47 Beside, major active faults, as the Dead Sea Fault and the North-Anatolian Fault, have been
48 surveyed through their coeval sedimentation (e.g. Marco and Agnon, 1995; McHugh et al.,
49 2006, 2014; Gorsline et al., 2000; Beck et al., 2007, 2015). For lacustrine or marine
50 sediments, the most frequent inferred processes leading to earthquake recording belong to
51 three groups: 1) *in situ* disturbances related to rheological contrast between successive layers
52 (e.g. Moretti et al., 1999; Rodriguez-Pascua et al, 2003; Wetzler et al, 2005), 2) simple gravity
53 reworking (turbidites, Mass Transport Deposits/MTDs; e.g. Strasser et al., 2006; Goldfinger
54 et al., 20012), and, 3) gravity reworking with additional coeval specific particles re-settling
55 (combination with tsunami/seiche effect) (e.g. Chapron et al., 1999; Beck, 2009; Beck et al.,
56 2007; McHugh et al., 2011; Campos et al., 2013). For small-sized lakes, possible impacts on
57 watershed and basin geometry may be recorded through sedimentary feeding and/or
58 depositional dynamic (e.g. Carrillo et al., 2008; Avşar et al., 2015; Aguilar et al., 2015).

59 To our knowledge, the combination of trench data and lacustrine sedimentary archives,
60 both related to the same active seismogenic fault, has scarcely been attempted. The here-
61 presented work concerns a favourable site for such combined approaches: the central part of
62 the Mérida Andes (North-western Venezuela) crosscut by the Boconó Fault. Within the frame
63 of the same research program, we could achieve: outcrop analyses, lake sediments coring, and
64 trenching. The present article presents the last phase of these investigations: an attempt to
65 integrate trenches results published in 1999 and 2008, and lacustrine archives data published
66 in 2006 and 2008. The here-discussed data combined these results with new additional
67 investigations dedicated: 1) to lacustrine sediments depositional processes and chronology, 2)
68 to deep-seated gravitational slope deformations bounding the Boconó Fault (Audemard et al.,

69 2010), 3) to the chronology of the last deglaciation using cosmogenic isotopes (Carcaillet et
70 al., 2013). Thus, for the following, we present as new data: a final interpretation of
71 sedimentological investigations, including structural positions, and a completed chronology of
72 all compared events.

73

74 **2. Investigated area: the central portion of the Boconó Fault**

75 **2.1. Geodynamic and tectonic setting**

76 The southern limit of the Caribbean Plate - the northern limit of the South-America
77 Plate - is a complex transform system with some convergence component (Fig. 1-B;
78 simplified from Stéphan et al., 1990; Audemard et al., 2000; DeMets et al., 2000; Pérez et al.,
79 2001; Weber et al., 2001; Symithe et al., 2015). In north-western Venezuela, part of the 2
80 cm/yr global displacement is partitioned into major faults bounding the so-called “Santa
81 Marta Triangular Block” or “Maracaibo Triangular Block”. The Mérida Andes (Fig. 1-A)
82 represent the active relieves accommodating part of this relative plate displacement and
83 associated stress field; they culminate at 4998 m a.s.l.. The Boconó Fault - dextral strike slip -
84 is roughly following the backbone of this SW-NE trending chain. In a review of available
85 Global Navigation Satellite System surveys data applied to northern Venezuela, Reinoza et al.
86 (2015) mention a 9-to-11 mm/y horizontal slip rate and a 1 mm/y convergence orthogonal to
87 the Mérida Andes. Previous estimations of the slip rate along the Boconó Fault, based on field
88 surveys, concluded to a 7-to-10 mm/y (Audemard et al., 1999). More recent investigations
89 combining ^{10}Be and high resolution satellite imagery concluded to similar values (up to 11,2
90 mm/yr) for the north-eastern strand (Pousse-Beltran et al., 2017).

91 As other major strike slip faults in northern Venezuela (El Pilar F., San Sebastian F.,
92 Fig. 1-B), the Boconó Fault was soon recognized by Rod (1956) as responsible for large
93 historical earthquakes. Later works (Cluff and Hansen, 1969; Aggarwal, 1983; *in* Audemard
94 et al. (2010)) attributed several major events (e.g. 1610, 1812, 1644, 1875, 1950) to the
95 Boconó Fault activity; 6.5 to 7.4 Mw magnitudes have been estimated (Palme et al., 2005, and
96 *in* Audemard et al. (2008)). More recent paleoseismic investigations through trenching, and
97 seismotectonic approaches, permitted to precise the distribution of major earthquakes along
98 different segments of the fault, and, by mean, to better estimate seismic hazards (Audemard,
99 1997, 2003, 2005; Pousse-Beltran et al., 2018). Results from the trenches performed in the
100 investigated area (white rectangle on Fig. 1B) will be summarized in § 4. The two concerned
101 trenches (Morro de los Hoyos and Mesa del Caballo; Audemard et al., 1999, 2008) are
102 precisely located on Figure 2.

103 In the investigated area, the highest relieves consist of Late Proterozoic igneous and
104 metamorphic rocks (orthogneiss, paragneiss, amphibolites, granitic pegmatites, aplites). The
105 Boconó Fault crosscuts this basement and also the Late Quaternary sediments developed
106 during the last climatic cycles, and directly resting on the Precambrian formations. Gneiss and
107 pegmatite fragments are the main components of Pleistocene coarse deposits, and especially
108 of the well-preserved morainic systems related to the last glaciation.

109

110 **2.2. Imprint of the last climatic cycles in central Mérida Andes**

111 Although located in tropical zone, due to the altitude of their central part, the Mérida
112 Andes, and especially the studied area, underwent a neat impact of the last glacio-eustatic
113 cycles. Several well-preserved morainic systems (Fig. 2) have been attributed to the Last
114 Glacial Maximum (equivalent for MIS2) here called “Mérida Glaciation” (Schubert, 1974;
115 Salgado-Labouriau and Schubert, 1976). Glacigenic sediments and erosion surfaces
116 developed above 2500 m with associated moraine-dammed lacustrine fills and peat-bogs. The
117 different LGM and post-LGM formations have been the focus of paleoecological and
118 paleoclimatic reconstructions (Salgado-Labouriau et al, 1977, 1992; Bradley et al., 1985;
119 Weingarten et al., 1990; Rull, 1996; Mahaney et al., 1997, 2000; Stansell et al., 2005). Beside
120 the well-defined LGM remnants, few tills bodies and organic-rich lacustrine sediments have
121 been attributed to a pre-LGM glacial stage, based on their relationships with the LGM
122 glacigenic sets and ^{14}C dating (Rull, 1996; Mahaney et al., 1997, 2000). More recent
123 cosmogenic isotopes analyses permitted to estimate erosion rates (Wesnousky et al., 2012)
124 and to precise the pattern and velocity of the post-LGM glacial retreat (Angel et al., 2013;
125 Carcaillet et al., 2013). The impact of the Little Ice Age has also been tested in the studied
126 area (Polissar et al., 2006). For the here-presented area, a set of 16 ^{10}Be measurements were
127 performed: on morainic complex crests boulders, and on free striated surfaces (Angel et al.,
128 2013; Carcaillet et al., 2013). The maximum advance is dated back to 18 kyr BP for the
129 highest site (Mucubají), which is in agreement with ^{14}C ages previously found for associated
130 lacustrine sediments (Carrillo et al, 2008).

131

132 **2.3. The Boconó Fault across the post-LGM lacustrine basins**

133 Beside their high paleoclimatic interest, the Mérida Andes' glacigenic and post-glacial
134 sedimentary bodies were soon used as precious tectonic benchmarks to quantify the Boconó
135 Fault activity (Schubert, 1982). Most of them are crosscut by surface-reaching active traces,
136 and horizontally displaced (Fig. 2). We focused our work on two of these morainic systems

137 and their associated lacustrine sedimentary fills: the Los Zerpa site and the Mucubají site. A
138 major south-eastern active trace and a less visible north-western trace were chosen for
139 structural and paleoseismic analyses in trenches (Audemard et al., 1999, 2008). The south-
140 eastern active trace just bounds the Los Zerpa lacustrine sediments (Fig. 3, section AA') while
141 it crosscuts Lake Mucubají (locally named "Laguna de Mucubají") (Fig. 3, section BB'). A
142 minor transtensional component also characterizes the studied portion of the Boconó Fault,
143 indicated by: normal ruptures within moraines, "micro pullapart" depressions or sagponds,
144 and low angle NE-dipping slickendides. In the Los Zerpa/Mucubají área, a large north-
145 westwards landslide displaces the whole Late Quaternary pile on the southeastern side of the
146 fault (Audemard et al., 2010). This gravitational deformation is also related to the local
147 transtensional component and inferred to be activated by strongest earthquakes (Audemard et
148 al., 2010; and this work).

149

150 **3. Summary and interpretation of paleoseismic results deduced from lacustrine** 151 **sedimentary archives**

152 Our investigations for subaqueous paleoseismic purpose were dedicated to the Los Zerpa
153 and the Mucubají lacustrine sediments accumulated within morainic systems. We used
154 outcrops for the Los Zerpa site (Carrillo et al., 2006), and cores and outcrops for the Mucubají
155 site (Carrillo et al., 2008). For both sedimentary fills, we performed different analyses and
156 interpretative approaches aiming to detect and characterize specific layers and sedimentary
157 features induced by seismic shaking, as proposed for alpine lakes (Chapron et al., 1999; Beck,
158 2009). The here-discussed results are based on the data published in 2006 and 2008 (Carrillo
159 et al., 2006, 2008) and on new data and interpretations:

- 160 1) additional field observations (Los Zerpa lacustrine and deltaic sediments and Mucubají
161 Lake's watershed for surface structural features);
- 162 2) additional laboratory analysis on Lake Mucubají cores (sediments textures, Organic
163 Matter characterization through Rock-Eval pyrolysis, new ^{14}C measurements and
164 calibrations, Table 1).

165 Within the frame of the same investigation program, two trenches had been achieved and
166 analysed: respectively at Morro de Los Hoyos and Mesa del Caballo sites (Audemard et al.
167 1999; 2008). For the following comparison we used their complete raw chronological results.

168

169 **3.1. The Los Zerpa paleolake**

170 This well-preserved simple glacial system (Fig. 2-A and Fig. 4) displays numerous
171 fractures affecting the lateral moraines, and the major Boconó Fault active trace which offsets
172 the frontal morainic arc (see also Schubert, 1981). Microtectonic observations on the major
173 trace (30° dipping slickensides on fine-grained lacustrine sediments) indicate a normal/right
174 lateral movement. Successive abandoned outlets with slightly decreasing altitudes can be
175 observed.

176 Imbricated terraces are related to episodes of emptying of the lake which have been
177 attributed to major co-seismic displacement leading to rupturing of the frontal moraine
178 (Carrillo et al., 2006). Between the lake genesis and its definitive emptying, the sedimentary
179 infilling was interrupted at least three times. The rather simple depositional system, related to
180 a unique tributary, can be divided into a lacustrine part with fine-grained laminated
181 bottomsets, and a complex deltaic part. Carrillo et al. (2006) investigated the downstream part
182 where they could recognize both specific deposits synchronous with local fracturing and post-
183 depositional disturbances (slumping above liquefied layers). Our new works were dedicated
184 to the upstream deltaic part in order to check a possible influence of the large landslide
185 bounding the whole Quaternary deposits (Audemard et al., 2010) (Fig. 3-A). Deformation
186 within the upstream part are mainly related to slumping; 3 major sliding surfaces were
187 observed (outcrops close up in Carrillo et al, 2006, Fig.6 and 7). Beside highly variable dips
188 of foreset layers, several discontinuities appear followed by coarse-grained “onlapping” layers
189 with upstream fining. We interpret these layers as due to temporary powerful upslope currents
190 with transport of coarse material (up to few cm gravels). At the top of the different foreset
191 units, a centimetric to decimetric fine-grained laminated episode is draping the coarser
192 sediments. The topmost millimetres, enriched in organic matter, provided 6 almost regularly
193 spaced ¹⁴C dates between 500 and 2000 yr cal. BP. A chronology of the different Los Zerpa
194 deposits is discussed in Carrillo et al. (2006).

195 The upstream prolongation of the deltaic set is more complex and highly deformed,
196 with extensional ruptures. The uppermost outcropping layers are downstream-dipping with
197 different angles and fractured unconformities. We associate all the above-mentioned features
198 to major co-seismic displacements of the Boconó Fault, along a portion which, at least,
199 includes the segment crossing the Los Zerpa paleo-lake. A detailed model for one of these
200 events, similar for Los Zerpa and Mucubají sites, is presented here-after (§ 3.3 and §3.4).

201

202 **3.2. Lake Mucubají**

203 About 5 km South-West of Los Zerpa paleolake, Lake Mucubají (or “Laguna de
204 Mucubají”), is a 16 m-deep, still active, lacustrine basin at 3560 m a.s.l.. Short gravity cores
205 and longer piston core (up to 8 m) were retrieved close to one of the two fault traces (Fig. 5).
206 Isolated outcrops have been attributed to an initial much larger lake (Salgado-Labouriau et al.,
207 1977; Carrillo et al, 2008). Although strongly depending on the seismo-tectonic setting, this
208 sedimentary archive permitted to also detect the global climatic episodes (Carrillo et al.,
209 2008). The southern part of the “paleo-lake Mucubají” was cut by a large deep-seated
210 landslide separating the quaternary sediments from their rocky Precambrian basement
211 (Audemard et al., 2010) (Fig. 3-B and white arrows on Fig. 5). The bottom of core MUCL-02-
212 02 is dated back to the cal. ^{14}C 16 340/15 390 yr BP interval. For detailed description and
213 analyses performed on cores and outcrops, we refer to Carrillo et al. (2008). In order to detect
214 and characterize specific layers for paleoseismic investigation, different parameters were used
215 for sediments composition and texture. Several abrupt changes in organic and mineral
216 sources, and particular depositional processes, may argue for the co-seismic local origin of
217 several major sedimentary “events”. As for the Los Zerpa paleolake, these events appear
218 recorded through gravity reworking and reflected “tsunami” effects. We selected one of these
219 events - considered as related to co-seismic rupturing of the Boconó Fault across the
220 investigated lakes - to illustrate hereafter both Los Zerpa and Mucubají sedimentary archives.

222 3.3.. Sedimentological characterization of a single major event

223 Although the selected event (dated back to around 13 kyr BP) was detected and analysed
224 in Core MUCL0202 from Mucubají site (Fig. 6), all evidenced sedimentary processes have
225 also been observed in Los Zerpa site. The 41 cm-thick complex succession of layers is
226 intercalated between two intervals of slow (“normal” or “background”) sedimentation.

227 The earthquake-related “event” (core close up on Fig. 6) consists in a succession of layers
228 which we related to different phases of reworking and subsequent re-settling. Basically, it
229 resembles a turbidite in a very broad sense, but several neat particularities have to be
230 underlined:

- 231 1) a poorly sorted coarse base, lacking transition with overlying deposit,
 - 232 2) reworked mudclasts (indicating microfracturing of previously deposited coherent mud),
 - 233 3) indication of two opposite transport directions;
 - 234 4) sharp basal limit of the final homogeneous fine-grained settling.
- 235 Furthermore, the end of the fine-grained plume settling shows: i) a specific chemical and
236 mineralogical concentration (vivianite), possible consequence of strong P release from bottom

237 sediments during shaking, ii) concentration of particulate O.M. which was used for ^{14}C dating
238 of the event. The association of the above-mentioned features corresponds to the
239 “turbidite+homogenite” model as defined in lakes and marine semi-closed basins (Chapron et
240 al., 1999; McHugh et al., 2006, 2011, 2014; Beck, 2009; Beck et al., 2007; Campos et al.,
241 2013) and attributed to major earthquake-induced gravity reworking, *in situ* disturbances, and
242 possible seiche effect.

243 In addition, sediments underlying and overlying this event are significantly different.
244 Among different parameters (layering, grain size, mineralogy and chemistry) we selected the
245 characteristics of Organic Matter content to illustrate a major and abrupt change synchronous
246 with this event. ROCK-EVAL pyrolysis results (Fig. 6) indicated that O.M. content and
247 characteristics also changed. With respect to underlying sediments, we noticed: increased
248 COT, increased HI, decreased and stabilised OI. The whole evidences an abrupt increase of
249 *in situ* lacustrine OM and more reducing conditions.

250

251 **3.4. Interpretative model for major sedimentary events: impact of *in situ* co-seismic** 252 **rupturing.**

253 Figure 7 depicted two steps of the structural evolution of Lake Mucubají while Figure 8 is
254 a tentative 2D model illustrating the successive sedimentary processes involved in one single
255 major event in Los Zerpa paleo-lake. This mechanism is also entirely available for the
256 sedimentary events observed in Lake Mucubají, especially the one presented and depicted in §
257 3.3 (Fig. 6). For both sites, we assumed that, in addition to the Boconó Fault main strand,
258 other rupturing occurred within, and at the base of, the morainic sediments pile. The 12 650-
259 13 150 yr BP event (Fig. 7-A) is considered as responsible for a major morphological change
260 (size and depth) and the coeval sedimentary event; a similar more recent event (Fig. 7-B) was
261 also depicted (Carrillo et al., 2008).

262 On Figure 8, depicting the details of the sedimentation during one single major event in
263 Los Zerpa site, step B corresponds to the tectonic *s.s.* triggering: the unconsolidated sediments
264 (lacustrine deposits and bounding moraines) undergo fracturing with partial to total separation
265 from underlying units. The vertical component of the offset is inferred to play a major role for
266 the water mass mobilization. Step B and C depict the effect of “backwash” responsible for
267 removing sediments upstream; it can be compared to a small “channelized” tsunami. With
268 steps C to E, we explain the different features mentioned by Carrillo et al. (2006) in the distal
269 (bottomsets) lacustrine deposits. Step F should represent the quiet final settling of re-
270 suspended fraction of the reworked material, including lighter organic particles (cf. Fig. 6

271 close up). Each unit defined within the deltaic foreset is believed to correspond to such a
272 scenario. The whole process may be applied to Lake Mucubají evolution, with slight
273 differences only concerning the position of involved fracturing.

274

275 **4. Discussion: comparison with Mesa del Caballo y Morro de Los Hoyos trenches data.**

276 The two lacustrine archives provided paleoseismic records at about 5 km distance from
277 each other, along the same active fault trace. In order to compare these data with results
278 yielded by the two neighbouring trenches (Fig. 2), we need to ensure their relationships with
279 local co-seismic rupturing, i.e. with displacements along the Boconó Fault strands crossing
280 the two lakes/paleolakes. For several major events - as the 12 650-13 150 yr BP one (Fig. 7-
281 A) - this relationship may be assumed. Conversely, we cannot rule out the impacts: i) of
282 remote strong earthquakes along farther strand of the Boconó Fault, ii) of other active faults
283 within or bounding the Mérida Andes as the Valera fault. At least for the historical seismicity
284 (Audemard, 1997; Audemard et al., 1999; Pousse-Beltran et al., 2018), none of the major
285 events known elsewhere along the Boconó Fault (south-westward or north-eastward) appear
286 within our results. Thus, we propose to consider the two lacustrine archives as representing
287 local events records. If comparing with historical seismicity the here-involved portion of the
288 Boconó Fault main strand most probably represents only a part of the total rupturing segments
289 with respect to estimated magnitudes which rather require few tens of kilometres (Audemard,
290 1997; Audemard et al., 1999; Pousse-Beltran et al., 2018).

291 The Morro de los Hoyos and the Mesa del Caballo trenches were respectively dug across
292 the south-eastern and the north-western traces (Fig. 2) (Audemard et al., 1999, 2008). To
293 attempt correlations between the different paleoseismic registers, a precise and reliable
294 chronology is requested, a point discussed hereafter.

295

296 **4.1. Chronological uncertainties**

297 With respect to the types of discussed archives and the differences between the two
298 lacustrine fills, tentative correlations of separate events, between the four sites, may only rely
299 on coincidences of radiocarbon dates. Thus, the actual significance of the obtained ages and
300 their precision have to be discussed. For all samples, a correction was applied to conventional
301 AMS ^{14}C ages using OxCal v4.3 software (Bronk Ramsey and Lee, 2013) considering only
302 the atmospheric CO_2 variations (IntCal123 curve, Reimer et al, 2013). The analysis of
303 Organic Matter (O.M.), using Rock-Eval parameters (Espitalié et al., 1985) for Lake
304 Mucubají cores (Fig. 6), indicates dominant terrestrial O.M. and minor lacustrine production.

305 Thus, and regarding the reduced sizes and depths of the lake basins, no reservoir effect
306 correction was applied to the lacustrine samples.

307 For trenches data, the dated O.M. matter comes from soils or sag ponds vegetal
308 accumulation. These measurements probably give a mean age for mixed organic debris which
309 represent a time interval difficult to estimate. Most of the ages obtained from the two lakes
310 represent fine-grained particulate O.M. settling at the end of reworking events (Carrillo et al,
311 2008). In this case, dated samples have a more precise position with respect to the earthquake-
312 induced layer, but the reworked thickness from previous deposits may also induce a mixing of
313 not contemporaneous O.M. particles. Table 1 concerns AMS ^{14}C data previously published
314 and new results; for the trenches, we used already published results (Audemard et al, 1999,
315 2008) and the same re-calibration was applied to all values. Tentative correlations between
316 the events from the four sites (Fig. 9) are based on the complete ^{14}C error bars (2σ , 95.4 %
317 prob.). The last 10 000 years BP (almost the entire Holocene) could be checked.

318

319 **4.2. Tentative correlations between the two trenches and the two lacustrine records**

320 For the following discussion, we first assume that all fracturing episodes detected and
321 dated in trenches have to be taken into account (Fig. 9-A and Fig. 10). Among a total of 24
322 inferred seismic events (Fig. 9-B), only 3 may be *correlated for three sites*, including one or
323 two lake fills and one or two trenches. 3 other *correlations between two sites - one lake fill*
324 *and one trench* - may be added. An overall mean 400 yr return interval may be deduced;
325 published historical “frequency” is higher but it concerns to a much longer portion of the
326 Boconó Fault (§ 2.1; Audemard, 1997). If comparing with the north-east portion of the
327 Boconó Fault recently surveyed (Pousse-Beltran et al., 2018) no historical mutual event
328 appears.

329 For *Lake Mucubají*, only few events were detected, but they all have a possible equivalent
330 in other sites. This may indicate that, during part of the last 10 kyr BP, sedimentation was not
331 favourable for shock-induced reworking (e.g. slow rate with cohesive deposits) and that only
332 *the strongest and local events* were registered. The about 9 kyr BP event (Fig. 7-A) illustrates
333 this case. The results for the *Los Zerpa paleo-lake* appears partly different as the “quiet”
334 interval between 2 kyr and 7.5 kyr BP is due to a gap in sedimentary infilling (see § 3.1).

335 According to *trench data*, many possible correlations appear between the two traces prior
336 to 2.5 kyr BP. No more events are detected on north-western trace, whilst several are detected
337 on the south-eastern trace; furthermore, they have possible equivalents in Los Zerpa lacustrine
338 archive. As noticed on Figure 9-A, ending of the north-western trace may be proposed, with a

339 continuation on the south-eastern trace. An alternate hypothesis is to consider the north-
340 western trace as a more temporary, or a secondary, one (Audemard et al., 2010). Our initial
341 assessment - to consider all rupturing phases dated in trenches as major earthquake marks -
342 appears questionable.

343 However, the observed deformations only concern few hundreds of meters of
344 unconsolidated sediments. These surficial structures may not exactly reflect the geometry of
345 the Boconó Fault activity within the underlying basement, and the huge landslide (white line
346 on Fig. 10; *cf.* also Audemard et al. (2010)) possibly plays a major role in co-seismic
347 fracturing distribution. Future investigation (through trenching ?) should investigate this
348 major structure.

349

350 **Conclusions**

351 Although we considered the studied area as particularly adequate to test the
352 complementarity of lacustrine sedimentary archives and trench dating, possible correlations
353 between very close sites could be *established only for part of the detected events*. The
354 lacustrine records may partly confirm the results of trenching across the same active traces. At
355 the difference, *large gaps* in studied lacustrine archives appear, at least partly filled by trench
356 data. As the four sites are concentrated within a small area, it is unlikely to put forward the
357 *distance* between the sedimentary accumulations and the *epicentral areas* as a limiting factor
358 for lacustrine recording. Furthermore, for major events (approx. 9 kyr and 13 kyr BP, Fig. 7-A
359 and 7-B), combined structural and sedimentological observations (§ 3.3 and 3.4) clearly
360 evidence the association of “*intra-lake*” *co-seismic rupturing and sedimentary reworking*. In
361 the here-studied case, a weak intrinsic potential for earthquake recording better explains the
362 sedimentary “events” scarcity. In particular, shock-induced reworking may lack during
363 periods of clayey cohesive slow sedimentation; for a deltaic foreset accumulation, a major
364 reworking may sweep off a major amount of unconsolidated sediment and prevent new
365 reworking during a time interval (in Beck, 2009).

366 More generally, these results also show that a unique sedimentary archive or a unique
367 trench, *even for a well-known fault and within a small area*, may be an *incomplete or biased*
368 record of local seismicity, and deduced seismic hazard assessment. The present attempt to
369 combine lacustrine sedimentary archives and trench data points out two “opposite” biases:
370 “incomplete” recording by sedimentation, and possible “overestimation” of co-seismic
371 rupturing events deduced from trench analysis.

372

373 **Acknowledgements:**

374 The authors thank the Venezuelan Foundation for Seismological Research, the
375 Venezuelan National Foundation for Science and Technology (FONACIT grants
376 20001002492 and 2003000090), and the French National Council for Scientific Research
377 (UMR CNRS 5025), for funding and logistical support. The French Ministry of Foreign
378 Affairs, through EGIDE and the French Embassy in Caracas (Technological and Cultural
379 Cooperation), supported the initial phase of project building. The Latin-American/French
380 scientific cooperation ECOS-Nord program contributed to these researches through the
381 V04O01 2004-2007 and V10U01 2010-2013 grants; thank you to ECOS-Nord Committee.
382 Mrs. Maura Elena Remiro (Instituto Nacional de Parques), and Mrs. Ana Elisa Osorio
383 Granado, Minister of Environment and Natural Resources, provided permission to sample
384 inside the Sierra Nevada and La Culata National Parks. E. Carrillo's research PhD work and
385 stay in LGCA were supported by the Scientific and Humanistic Development Centre of the
386 Central University of Venezuela. A. van Welden's PhD research was supported by a Thesis
387 grant from French National Education Ministry. Great thanks to the whole Piva family and the
388 Santo Domingo hotel staff for friendship and logistical help during all field surveys.

389 Our initial manuscript greatly benefited from reviewing by Pr. Dr. Carlos Costa, who
390 we sincerely acknowledge for suggested improvements.

391

392 **References:**

- 393 Adams J. (1990) Paleoseismicity of the Cascadian subduction zone: evidence from turbidites
394 off the Oregon-Washington margin. *Tectonics*, vol. 9, 4:569-583.
- 395 Aguilar, I., Beck, C., Audemard, F., Davelle, A.-L., Boussafir, M., Campos, C., Crouzet, C..
396 2015. Last millennium sedimentation in the Gulf of Cariaco (NE Venezuela): evidence
397 for morphological changes of Gulf entrance and possible relations with large
398 earthquakes. *C. R. Geoscience*, 348 (2016) 70–79.
- 399 Angel, I., Carrillo, E., Carcaillet, Audemard, F., & Beck, C., 2013. Geocronologia con el
400 isótopo cosmogénico ^{10}Be , aplicación para el estudio de la dinámica glaciár en la
401 región central de los Andes de Mérida. *GEOS*, 44: 73-82.
- 402 Audemard, F.A., 1997. Holocene and historical earthquakes on the Boconó Fault System,
403 southern Venezuelan Andes: trench confirmation. *Journal of Geodynamics*, 24:183-
404 200.
- 405 Audemard, F.A., 2003. Geomorphic and geologic evidence of ongoing uplift and deformation
406 in the Mérida Andes, Venezuela. *Quaternary International*, 101/102C: 43-65.

- 407 Audemard, F.A., 2005. Paleoseismology in Venezuela: objectives, methods, applications,
408 limitations and perspectives. *Tectonophysics*, 408:29-61.
- 409 Audemard, F.A., Beck, C., Carrillo, E., 2010. Deep-seated gravitational slope deformations
410 along the active Boconó Fault in the central portion of the Mérida Andes, western
411 Venezuela. *Geomorphology*, 124:164–177, doi:10.1016/j.geomorph.2010.04.020
- 412 Audemard, F. A., Machette, M.N., Cox, J.W., Dart, R.L., and Haller, K.M., 2000. Map of
413 Quaternary faults of Venezuela, USGS Open-File report 00–0018.
- 414 Audemard, F.A., Ollarves, R., Bechtold, M., Díaz, G., Beck, C., Carrillo, E., Pantosti, D.,
415 Diederix, H., 2008. Trench investigation on the main strand of the Boconó fault in its
416 central section, at Mesa del Caballo, Mérida Andes, Venezuela. *Tectonophysics*,
417 459:38–53.
- 418 Audemard, F.A., Pantosti, D., Machette, M., Costa, C., Okumura, K., Cowan, H., Diederix,
419 H., Ferrer, C., and SAWOP Participants, 1999. Trench investigation along the Merida
420 section of the Boconó fault (central Venezuelan Andes). *Tectonophysics*, 308:1-21.
- 421 Avşar, U., Hubert-Ferrari, A., De Batist, M., Schmidt, S., Fagel, N., 2015. Sedimentary
422 records of past earthquakes in Boraboy Lake during the last ca 600 years (North
423 Anatolian Fault, Turkey). *Palaeogeography, Palaeoclimatology, Palaeoecology*,
424 433:1-9.
- 425 Beck, C., 2009. Late Quaternary lacustrine paleo-seismic archives in north-western Alps:
426 Examples of earthquake-origin assessment of sedimentary disturbances. *Earth-Science*
427 *Reviews*, 96:327–344.
- 428 Beck, C., Campos, C., Eriş, K.K., Çağatay, N., Mercier de Lépinay, B., and Jouanne, F., 2015.
429 Estimation of successive coseismic vertical offsets using coeval sedimentary events –
430 application to the southwestern limit of the Sea of Marmara’s Central Basin (North
431 Anatolian Fault). *Natural Hazards and Earth System Sciences*, 15:247–259,
432 doi:10.5194/nhess-15-247-2015.
- 433 Beck, C., Mercier de Lépinay, B., Schneider, J.-L., Cremer, M., Çağatay, N., Wendenbaum,
434 E., Boutareaud, S., Ménot, G., Schmidt, S., Weber, O., Eris, K., Armijo, R., Meyer, B.,
435 Pondard, N., Gutscher, M.-A., and the MARMACORE Cruise Party, J.-L. Turon, L.
436 Labeyrie, E. Cortijo, Y. Gallet, H. Bouquerel, N. Gorur, A. Gervais, M.-H. Castera, L.
437 Londeix, A. de Rességuier, A. Jaouen, 2007. Late Quaternary co-seismic
438 sedimentation in the Sea of Marmara's deep basins. *Sedimentary Geology*, 199:65–89.
- 439 Bradley, R.S., Yuretich, R., Salgado-Labouriau, M.L., Weingarten, B., 1985. Late Quaternary

- 440 paleoenvironmental reconstruction using lake sediments from the Venezuelan Andes:
441 preliminary results. *Zeitschrift für Gletscherkunde und Glazialgeologie*, 21:97–106.
- 442 Bronk Ramsey, C, Lee, S. 2013. Recent and Planned Developments of the Program OxCal.
443 *Radiocarbon*, 55(2-3):720–730.
- 444 Campos, C., Beck, C., Crouzet, C., Demory, F., Van Welden, A., Eris, K.. 2013. Deciphering
445 hemipelagites from homogenites through anisotropy of magnetic susceptibility.
446 Paleoseismic implications (Sea of Marmara and Gulf of Corinth). *Sedimentary*
447 *Geology*, 292:1–14.
- 448 Carcaillet, J., Angel, I., Carrillo, E., Audemard, F.A., Beck, C., 2013. Timing of the last
449 deglaciation in the Sierra Nevada of the Mérida Andes, Venezuela. *Quaternary*
450 *Research*, 80:482–494, doi:10.1016/j.yqres.2013.08.001
- 451 Carrillo, E., Audemard, F., Beck, C., Cousin, M., Jouanne, F., Cano, V., Castilla, R., Melo,
452 L., and Villemin, T., 2006. A Late Pleistocene natural seismograph along the Boconò
453 Fault (Mérida Andes, Venezuela): the moraine-dammed Los Zerpa paleo-lake. *Bulletin*
454 *of the French Geological Society*, t. 177, 1:3-17.
- 455 Carrillo, E., Beck, C., Audemard, F.A., Moreno, E., Ollarves, R., 2008. Disentangling Late
456 Quaternary climatic and seismo-tectonic controls on Lake Mucubají sedimentation
457 (Mérida Andes, Venezuela). *Palaeogeogr. Palaeoclim. Palaeoecol.*, 259: 284-300.
- 458 Chapron, E., Beck, C., Pourchet, M. and Deconinck, J.-F., 1999. 1822 AD earthquake-
459 triggered homogenite in Lake Le Bourget (NW Alps). *Terra Nova* 11, 86-92.
- 460 DeMets, C., Jansma, P.E., Mattioli, G.S., Dixon, T.H., Farina, F., Bilham, R., Calais, E.,
461 and Mann, P., (2000). GPS geodetic constraints on Caribbean-North America plate
462 motion, *Geophysical Research Letters*, 27(3):437–440.
- 463 Espitalié, J, Deroo, G., and Marquis, F., 1985. La pyrolyse Rock-Eval et ses applications.
464 Deuxième partie, *Oil & Gas Science and Technology*, 40: 755-84.
- 465 Garrity, C., Hackley, P., and Urbani, F., 2004. *Digital shaded-relief map of Venezuela*.
466 <http://pubs.usgs.gov/of/2004/1322>.
- 467 Goldfinger, C., Nelson, C.H., Morey, A., Johnson, J.E., Gutierrez-Pastor, J., Eriksson, A.T.,
468 Karabanov, E., Patton, E., Grácia, E., Enkin, E., Dallimore, A., Dunhill, G., and
469 Vallier, T., 2012. Turbidite event history: methods and implications for Holocene
470 Cascadian Subduction Zone. U.S. Geological Survey Professionnal Paper, 1662-F, 184
471 pp..
- 472 Gorsline, D.S., De Diego, T., Nava-Sanchez, E.H., 2000. Seismically triggered turbidites in
473 small margin basins: Alfonso Basin, Western Gulf of California and Santa Monica

- 474 Basin, California Borderland. *Sedimentary Geology*, 135:21–35.
- 475 Mahaney, W., Kalm, V., Bezada, M., 1997. Estratigrafía del Cuaternario tardío de un
476 ambiente proglacial en el area de Mucubají, Mucuchache, El Pedregal. Andes
477 Venezolanos. Memorias Ier Congreso Latinoamericano de Sedimentología, Sociedad
478 Venezolana de Geólogo, Tomo I, p. 417-424.
- 479 Mahaney, W.C., Milner, M.W., Voros, J. Kalm, V. Hütt, G., Bezada, M., Hancock, R.G.V.
480 Autreiter, S., 2000. Stratotype for the Mérida Glaciation at Pueblo Llano in the Northern
481 Venezuela Andes. *Journal of South American Earth Sciences*. 13:761-774.
- 482 McCalpin, J.P., and Nelson, A.R., 2009. Introduction to paleoseismology. Chapter 1 in
483 “Paleoseismology”, (J.P. McCalpin Edr.), Academic Press, ISBN 978-0-12-373576-8.
- 484 McHugh, C., Seeber, L., Cormier, M.-H., Dutton, J., Çağatay, N., Polonia, A., Ryan, W.B.F.,
485 Gorur, N., 2006. Submarine earthquake geology along the North Anatolian fault in the
486 Marmara Sea: a model for transform basin sedimentation. *Earth and Planetary Sciences
487 Letters*, 248:661-684.
- 488 McHugh, C., Seeber, L., Braudy, N., Cormier, M.-H., Davis, M.B., Diebold, J.B., Dieudonné,
489 N., Douilly, R., Gulik, S.P.S., Hornbach, M.J., Johnson III, H.E., Ryan Miskin, K.,
490 Sorlien, C., Steckler, M., Symithe, S.J., Templeton, J., 2011. Offshore sedimentary
491 effects of the 12 January 2010 Haïti earthquake. *Geology*, 39(8):723-726.
- 492 McHugh, C., Braudy, N., Çağatay, N., Sorlien, C., Cormier, M.-H., Seeber, L., Henry, P.,
493 2014. Seafloor fault ruptures along the North Anatolia Fault in the Marmara Sea,
494 Turkey: Link with the adjacent basin turbidite record. *Marine Geology*, 353:65-83.
- 495 Marco, S., Agnon, A., 1995. Prehistoric earthquake deformations near Masada, Dead Sea
496 Graben. *Geology*, vol. 23, 8:695–698.
- 497 Moretti, M., Alfaro, P., Caselles, O., Canas, J.A., 1999. Modelling seismites with a digital
498 shaking table. *Tectonophysics*, 304:369–383.
- 499 Palme, C., Morandi, M., Choy, J., 2005. Re-evaluación de las intensidades de los grandes
500 sismos históricos de la región de la cordillera de Mérida utilizando el método de
501 Bakun & Wentworth. *Revista Geográfica Venezolana*, 233–253 (número especial).
- 502 Perez, O.J., Bilham, R., Bendick, R., Velandia, J.R., Hernandez, N., Moncayo, C., Hoyer, M.,
503 and Kozuch, M., 2001. Velocity field across the southern Caribbean plate boundary
504 and estimates of Caribbean/South-American plate motion using GPS geodesy 1994-
505 2000. *Geophysical Research Letters*, 28:2987-2990.
- 506 Polissar, P.J., Abbott, M.B., Wolfe, A.P., Bezada, M., Rull, V., and Bradley, R.S., 2006. Solar

- 507 modulation of Little Ice Age climate in the tropical Andes. *Proceeding of the National*
508 *Academy of Sciences*, U.S.A., Vol. 103, 24:8937-8942.
- 509 Pousse Beltran, L., R. Vassallo, F. Audemard, F. Jouanne, J. Carcaillet, E. Pathier, M. Volat,
510 2017. Pleistocene slip rates on the Boconó Fault along the North Andean Block plate
511 boundary, Venezuela. *Tectonics*, 36, 1207–1231, doi:10.1002/2016TC004305.
- 512 Pousse-Beltran L., R. Vassallo, F. Audemard, F. Jouanne, J. Oropeza, S. Garambois, J. Aray,
513 2018. Earthquake geology of the last millennium along the Boconó Fault, Venezuela,
514 *Tectonophysics*, 747-748, 40-53, doi.org/10.1016/j.tecto.2018.09.010.
- 515 Reimer, P.J., Bard, E., Bayliss, A., Beck, J. W., Blackwell, P.J., Bronk Ramsey, C., Buck, C.
516 , Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P.,
517 Haflidason, H., Irka Hajdas, Hatté, C., Heaton, T.J., Dirk L Hoffmann, D.L., Hogg,
518 A.G, Hughen, K.G., Kaiser, K.F., Kromer, B., Manning, S.W., Niu, M., Reimer, R.W.,
519 Richards, D.A., Scott, E.M., Southon, J.R., Staff, R.A., Turney, C.S.M., van der Plicht,
520 J., 2013. IntCal13 and Marine13 Radiocarbon Age Calibration Curves 0–50,000 Years
521 cal BP.. *Radiocarbon*, Vol 55, No 4. IntCal13 Special Issue.
- 522 Reinoza, C., Audemard, F.A., Jouanne, F., and Beck, C., 2015. An overview of the GNSS
523 geodetic measurements applied to geodynamic studies in Venezuela. In “The northern
524 limit of the South-American Plate - lithospheric structure from surface to the mantle”
525 (M., Schmitz, F. Audemard, F. Urbani, F., Eds.) Editorial Inovación Tecnológica,
526 Universidad Central de Venezuela, 378 pp..
- 527 Rod, E., 1956. Earthquakes of Venezuela related to strike slip fault ? *American Association of*
528 *Petroleum Geologists Bulletin*, 40:2509-2512.
- 529 Rodriguez-Pascua, M.A., De Vicente, G., Calvo, J.P., Pérez-López, R., 2003. Similarities
530 between recent seismic activity and palæoseismites during the Late Miocene in the
531 external Betic Chain (Spain): relationship between “b” value and the fractal
532 dimension. *Journal of Structural Geology*, 25:749–763.
- 533 Rull, V. 1996. Late Pleistocene and Holocene climates of Venezuela. *Quaternary*
534 *International*.
535 31:85-94.
- 536 Salgado-Labouriau, M.-L., Schubert, C., Valastro S.J., 1977. Paleoecologic analysis of a Late
537 Quaternary terrace from Mucubají, Venezuelan Andes. *Journal of Biogeography*,
538 4:313–325.
- 539 Salgado-Labouriau, M.-L., Schubert, C., 1976. Palynology of Holocene peat bogs from the

- 540 central Venezuelan Andes. *Palaeogeography, Palaeoclimatology,*
541 *Palaeoecology*,19:147–156.
- 542 Salgado-Labouriau, M.-L., Bradley, R.S., Yuretich, R., Weingarten, B., 1992. Paleocological
543 analysis of the sediments of Lake Mucubají, Venezuelan Andes. *Journal of*
544 *Biogeography*: 4:313–325.
- 545 Schubert, C. (1974). Late Pleistocene Merida glaciation, Venezuelan Andes. *Boreas*, 3:147-
546 152.
- 547 Schubert, C., 1981. Evolución post-glacial de un valle morrénico, Andes Merideños. *Acta*
548 *Científica Venezolana*, 32:151-158.
- 549 Schubert, C., 1982. Neotectonics of Boconó Fault, western Venezuela. *Tectonophysics*, 85:
550 205-220.
- 551 Schubert, C., Henneberg, H.G., 1975. Geological and geodetic investigations on the
552 movement along the Boconó fault, Venezuelan Andes. *Tectonophysics*, 29(1/4):199-
553 207.
- 554 Sims, J., 1973. Earthquake-induced structures in sediments of Van Norman Lake, San
555 Fernando, California. *Science*, 182:161–163.
- 556 Stansell N. D., Abbott M. B., Polissar P. J., Wolfe A. P., Bezada M., Rull V. 2005 Late
557 Quaternary deglacial history of the Merida Andes, Venezuela. *Journal of Quaternary*
558 *Science*, 20(7-8):801-812.
- 559 Stéphan, J.-F., Mercier de Lépinay, B., Calais, E., Tardy, M., Beck, C., Carfantan, J.-C.,
560 Olivet, J.-L., Vila, J.-M., Bouysse, P., Mauffret, A., Bourgois, J., Théry, J.-M.,
561 Tournon, J., Blanchet, R. & Dercourt, J., 1990. Paleogeodynamic maps of the
562 Caribbean: 14 steps from Lias to Present. *Bulletin of the French Geological Society*, 8,
563 VI, p. 915-919. 14 appendices.
- 564 Strasser, M., Anselmetti, F.A., Fäh, D., Giardini, D., and Schnellmann, M., (2006).
565 Magnitudes and source areas of large prehistoric northern Alpine earthquakes revealed
566 by slope failures in lakes. *Geology*, vol. 34, 12:1005-1008.
- 567 Symithe, S., Calais, E., de Chabaliere, JB, Robertson, R., and Higgins, M., 2015. Current
568 Block Motions and Strain Accumulation on Active Faults in the Caribbean. *Journal of*
569 *Geophysical Research: Solid Earth*, 120, 3748–3774, doi:10.1002/2014JB011779.
- 570 Weber, J., Dixon, T., DeMets, C., Ambeh, W., Jansma, P., Mattioli, G., Saleh, J., Sella, G.,
571 Bilham, R. & Pérez, O., 2001. GPS estimate of relative motion between the Caribbean
572 and South American plates and geologic implications for Trinidad and Venezuela.
573 *Geology*, vol. 29, 1:75-78.

- 574 Weingarten, B., Yuretich, R.F., Bradley, R.S., Salgado-Labouriau, M.-L., 1990.
 575 Characteristics of sediments in an altitudinal sequence of lakes in the Venezuelan
 576 Andes: climatic implications. *Journal of South American Earth Sciences*, 3:113–124.
- 577 Weldon II, R.J. II, McCalpin, J.P., and Rockwell, T.K., 2009. Paleoseismology in Strike-Slip
 578 Tectonic Environments: Introduction. Chapter 6 in “Paleoseismology”, (J.P. McCalpin
 579 Edr.), Academic Press, ISBN 978-0-12-373576-8.
- 580 Wesnousky, S.G., Aranguren, R., Rengifo, M., Owen, L.A., Caffee, M.W., Krishna Murari,
 581 M., Pérez, O.J., 2012. Toward quantifying geomorphic rates of crustal displacement,
 582 landscape development, and the age of glaciation in the Venezuelan Andes.
 583 *Geomorphology*, 141–142, 99–113.
- 584 Wetzler, N., Marco, S., Heifetz, H., 2010. Quantitative analysis of seismogenic shear-induced
 585 turbulence in lake sediments. *Geology*, vol. 38, 4:303-306.

586
 587 **Figures captions:**

- 588
 589 Figure 1.- Geodynamic and morphological settings of the Boconó Fault. A: *Morphology of*
 590 *the*
 591 *Merida Andes and location of the studied area. Shade relief from Garrity et al. (2004).*
 592 *Epicentral location of several major historical earthquakes in central Merida*
 593 *Andes, from Palme et al. (2005) and Audemard (2005). Insert B: South-Caribbean*
 594 *geodynamics simplified from Audemard et al. (2000), Weber et al. (2001), Pérez et al.*
 595 *(2001), Symithe et al. (2015).*
- 596 Figure 2.- Detailed morphology of the surveyed area, and location of the investigated sites. A
 597 *set of morainic systems are crosscut and deformed by an active trace of the right*
 598 *lateral strike slip Boconó Fault; several associated moraine-dammed lakes are also*
 599 *affected by the fault activity. For the Los Zerpa site, the sedimentary fill is preserved*
 600 *as outcrops; for the Mucubají site, part of the fill is still in sub-lacustrine position.*
 601 *Aerial photos from Instituto Geográfico “Simón Bolívar” de Venezuela. (North*
 602 *direction towards the left-bottom corner of the picture for better relief visibility). (AA’*
 603 *and BB’: cross section displayed on Figure 3).*
- 604 Figure 3.- Geometry and Structural setting of Los Zerpa and Mucubají Late Quaternary
 605 morainic and lacustrine deposits. (*Location of sections AA’ and BB’ on Figure 2*). 1)
 606 *Precambrian metamorphic and igneous basement (gneisses, amphibolites, granites);*
 607 *2) LGM morainic deposits; 3) post-LGM lacustrine deposits (for the Los Zerpa site,*

608 *they are represented at their altitude as viewed through lateral moraines). NB: scales*
 609 *are different for the two sections.*

610 Figure 4.- Location of the different types of disturbances in the Los Zerpa paleo-lake fill
 611 remnants. *Lacustrine sediments trapped within the Los Zerpa morainic system appear*
 612 *as terraces. Earthquake-impacts are either direct fracturing (Gilbert-delta and bottom*
 613 *sets) or gravity reworking and liquefaction (delta foresets and bottomsets). Aerial*
 614 *photos from Instituto Geográfico “Simón Bolívar” de Venezuela. (North direction*
 615 *towards the left-bottom corner of the picture for better relief visibility).*

616 Figure 5.- The Mucubají Lake and paleo-lake: morphological setting and location of coring
 617 sites. (isobaths curves interval: 2 m). *Light grey area: outcropping lacustrine*
 618 *sediments. White arrows underline the limit of a deep-seated landslide following*
 619 *Audemard et al. (2013). Aerial photos from Instituto Geográfico “Simón Bolívar” de*
 620 *Venezuela. (North direction towards the left-bottom corner of the picture for better*
 621 *relief visibility)*

622 Figure 6.- Example of major sedimentary event in Lake Mucubají. *Layering, texture,*
 623 *and composition (left), and deduced sedimentary processes (right), depict the*
 624 *combined effect of gravity reworking and water mass oscillation. This complex event*
 625 *corresponds to a “homogenite+turbidite” association. The chosen event exactly*
 626 *precedes a major change of the lake size and sedimentation (see Fig. 8-A) and is thus*
 627 *considered as co-seismic with respect to an offset of the main strand of the Boconó*
 628 *fault across the lake.*

629 Figure 7.- Reconstruction of two inferred co-seismic modifications of Lake Mucubají in
 630 relation with the Boconó Fault. *The scenario represented in A corresponds to the main*
 631 *event displayed on Figure 6.*

632 Figure 8.- Reconstruction of a co-seismic sedimentary event related to an offset of the Boconó
 633 Fault across the Los Zerpa paleolake. *This scenario is based on the sedimentary*
 634 *structures and the specific layering observed in the different parts of the lacustrine fill*
 635 *(detailed in Carrillo et al. 2006).*

636 Figure 9.- Synthesis of paleoseismic data from the Los Zerpa/Mucubají area, combining
 637 trench data and lacustrine sedimentary archives. A) Tentative time correlations. *They*
 638 *are based on overlapping of error bars corresponding to 2 σ intervals of Calibrated*
 639 *¹⁴C ages. Correlations concern the 10 last kyr BP. Trench data provide a more*
 640 *regularly distributed set of data, while lacustrine archives give two periods with*
 641 *recorded events (2 kyr to Present, 10 to 8 kyr BP); several possible correlations*

642 *between trenches and lacustrine fills may be proposed for these two periods.*
643 *Combined data indicate a possible dis-activation of the North-West trace with a*
644 *coeval continuation of the South-East trace. Insert B) Summary of trenches and*
645 *lacustrine data for the last 10 kyr BP.*


646 Figure 10.- Geographical distribution and summary of paleoseismic data sources in the Los
647 Zerpa/Mucubají Area.


648 Table 1: ^{14}C dating results for the lacustrine sedimentary archives. *Compilation of Carrillo et*
649 *al. (2006, 2008)'s data and additional unpublished measurements. LZAF: Los Zerpa*
650 *paleo-lake; MUCL: Lake Mucubají cores samples.*


651


652


Sample	Conventional ¹⁴C age yrs BP	Calibrated BP age ; 95.4 % probability yrs BP
MUCL-02-01-60	8180 ± 60	9000/9400
MUCL0201-02A30	9620 ± 50	10750/11170
MUCL-02-01-176	10490 ± 50	12050/12900
MUCL-02-01-203	10910 ± 60	12650/13150
MUCL-02-01-269	11750 ± 70	13350/14150
MUCL-02-0- 453	13200 ± 80	14850/16450
MUCL-02-02-150	2630 ± 50	2700/2860
MUCL-02-023A90	7890 ± 50	8580/8990
MUCL-02-02-590	10800 ± 90	12600/13150
MUCL-02-02-800	13200 ± 120	14850/16450
LDMU-04-3	750 ± 40	640/740
LZAF-04 1	760 ± 40	650/760
LZAF-04 2	1280 ± 40	1080/1290
LZAF-04 3	1550 ± 50	1330/1540
LZAF-04 4	1290 ± 40	1120/1300
LZAF-04 5	1800 ± 40	1600/1830
LZA-6 C1	1755 ± 40	1550/1740
LZA-4-4MO1	8500 ± 50	9430/9550
LZA-4-4MO2	8590 ± 50	9480/9700
LZA-AL-SU-1	6870 ± 60	7580/7830

























Along an active seismogenic strike slip fault - the Boconó Fault, Mérida Andes, Northwestern Venezuela - a 7 km long strand offers a unique situation to combine, for paleoseismic purpose, two lacustrine sedimentary records and two trenches. For a 10 kyr record, tentative correlations are presented: among a total of 24 inferred seismic events, only 3 may be correlated for three sites, including one or two lake fills and one or two trenches. 3 other correlations between two sites - one lake fill and one trench - may be added. An overall mean 400 yr return interval may be deduced, which appears longer than historical events return period. These results show that a unique sedimentary archive or a unique trench, even for a well-known fault and within a small area, may be an incomplete or biased record of local seismicity, and deduced seismic hazard assessment. The present attempt to combine lacustrine sedimentary archives and trench data points out two “opposite” biases: “incomplete” recording by sedimentation, and possible “overestimation” of co-seismic rupturing events deduced from trench analysis.