



**HAL**  
open science

# Chemistry in Titan's ionosphere: contribution of dissociative recombination to an ion- neutral coupled model

Sylvain Plessis, P. Pernot, Nathalie Carrasco

► **To cite this version:**

Sylvain Plessis, P. Pernot, Nathalie Carrasco. Chemistry in Titan's ionosphere: contribution of dissociative recombination to an ion- neutral coupled model. European Planetary Science Congress (EPSC2009), Sep 2009, Potsdam, Germany. pp.EPSC2009-148. insu-02896805

**HAL Id: insu-02896805**

**<https://insu.hal.science/insu-02896805v1>**

Submitted on 10 Jul 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Chemistry in Titan's ionosphere: contribution of dissociative recombination to an ion-neutral coupled model

S. Plessis (1,2), P. Pernot (1,2) and N. Carrasco (3)

(1) Laboratoire de Chimie Physique, CNRS, UMR 8000, Orsay, F-91405, (2) Université Paris-Sud 11, Orsay, F-91405, (3) Service d'Aéronomie, CNRS, IPSL, Université de Versailles Saint Quentin, BP 3, Verrières le Buisson, F-91371, (sylvain.plessis@lcp.u-psud.fr / Fax: +33 1 69 15 44 47)

Modelling the chemical composition of Titan's ionosphere is a very challenging issue. Latest works perform either inversion of CASSINI's INMS mass spectra (neutral[1] or ion[2]), or design coupled ion-neutral chemistry models[3].

Coupling ionic and neutral chemistry has been reported to be an essential feature of accurate modelling[3]. Electron Dissociative Recombination (EDR), where free electrons recombine with positive ions to produce neutral species, is a key component of ion-neutral coupling.

Experimental databases of EDR are incomplete. Indeed, for heavy hydrocarbon ions ( $C_nH_y^+$ ,  $n \geq 3$ ), final neutral molecules distribution is unknown. Data are available only over the carbon composition of fragments, which leaves a deterministic approach irrelevant.

Thanks to a novel stochastic description of the EDR chemical reactions, we were able to calculate neutral species fluxes due to EDR processes and compare them to fluxes due to neutral chemistry processes.

Some species' fluxes, like nitrile molecules, are mainly described by EDR processes (Fig. 1). Although hydrocarbon fluxes are dominated by the neutral chemistry contribution, EDR contribution grows with the molecules' mass (Fig. 2) and is therefore of importance for heavy molecules coupling.

### References

- [1] J. Cui, R.V. Yelle, V. Vuitton, J.H. Waite Jr., W.T. Kasprzak, D.A. Gell, H.B. Niemann, I.C.F. Miller-Wodarg, N. Borggren, G.G. Fletcher, E.L. Patrick, E. Raen, and B.A. Magee. *Icarus*, 200(2):581–615, 2008.
- [2] V. Vuitton, R. V. Yelle, and M.J. McEwan. *Icarus*, 191:722–742, 2007.
- [3] V. De La Haye, J.H. Waite Jr., T.E. Cravens, I.P. Robertson, and S. Lebonnois. *Icarus*, 197(1):110 – 136, 2008.




Figure 1: Nitrile species fluxes; blue: EDR processes flux; green: neutral chemistry processes fluxes




Figure 2: Global fluxes for hydrocarbon molecules, sorted by mass; blue: EDR processes flux; green: neutral chemistry processes fluxes