



Effects of organic matter–goethite interactions on reactive transport of nalidixic acid: column study and modeling

Wei Cheng, Lian Zhou, Remi Marsac, Jean-François Boily, Khalil Hanna

► To cite this version:

Wei Cheng, Lian Zhou, Remi Marsac, Jean-François Boily, Khalil Hanna. Effects of organic matter–goethite interactions on reactive transport of nalidixic acid: column study and modeling. *Environmental Research*, 2020, 191, pp.110187. 10.1016/j.envres.2020.110187 . insu-02940049

HAL Id: insu-02940049

<https://insu.hal.science/insu-02940049>

Submitted on 16 Sep 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Journal Pre-proof

Effects of organic matter–goethite interactions on reactive transport of nalidixic acid: column study and modeling

Wei Cheng, Lian Zhou, Rémi Marsac, Jean-François Boily, Khalil Hanna



PII: S0013-9351(20)31084-7

DOI: <https://doi.org/10.1016/j.envres.2020.110187>

Reference: YENRS 110187

To appear in: *Environmental Research*

Received Date: 25 May 2020

Revised Date: 25 August 2020

Accepted Date: 7 September 2020

Please cite this article as: Cheng, W., Zhou, L., Marsac, R., Boily, J.-F., Hanna, K., Effects of organic matter–goethite interactions on reactive transport of nalidixic acid: column study and modeling, *Environmental Research*, <https://doi.org/10.1016/j.envres.2020.110187>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Elsevier Inc. All rights reserved.

Effects of organic matter–goethite interactions on reactive transport of nalidixic acid: column study and modeling

Wei Cheng,^{a,b} Lian Zhou,^b Rémi Marsac,^c Jean-François Boily^d and Khalil Hanna^{*b,e}

^a *College of Resources and Environmental Science, South-Central University for Nationalities, Wuhan, China.*

^b *Univ Rennes, École Nationale Supérieure de Chimie de Rennes, CNRS, ISCR – UMR6226, F-35000 Rennes, France*

^c *Univ Rennes, CNRS, Géosciences Rennes - UMR 6118, F-35000 Rennes, France*

^d *Department of Chemistry, Umeå University, SE-901 87 Umeå, Sweden*

^e *Institut Universitaire de France (IUF), MESRI, 1 rue Descartes, 75231 Paris, France.*


*Corresponding author: Tel.: +33 2 23 23 80 27; Fax : +33 2 23 23 81 20

E-mail address: khalil.hanna@ensc-rennes.fr (K. Hanna)

A revised manuscript submitted to *Environmental Research*

August 2020

Graphical Abstract



Abstract

The fractionation of natural organic matter (NOM) and its impact on the binding of quinolones to mineral surfaces and transport behaviour under flow-through conditions have been scarcely investigated. In this study, the sorption and transport of a widely used quinolone antibiotic, Nalidixic acid (NA), were investigated in goethite-coated sand (GCS) columns over a wide concentration range (5 – 50 mg/L) of Leonardite humic acid (LHA), a representative NOM. Simultaneous injection of NA and LHA in GCS columns mutually alter transport of each other, *i.e.* NA mobility and LHA molecular fractionation. Preloading of GCS column with LHA dramatically facilitated the transport behaviour of NA, where nonspecific interactions with LHA-covered goethite surfaces controlled NA mobility. Simulations using a two-site nonequilibrium model showed that a modified sorption rate constant was required to accurately describe the breakthrough curves of NA under these conditions. This altered rate constant suggests that nonspecific interactions of NA on bound LHA may take place as an additional binding mechanism affecting adsorption kinetics. NOM fractionation alters sorption mechanisms and kinetics of quinolone antibiotics, which in turn affect their fractionation. These results may have important implications for an accurate assessment of the fate of these types of antibiotics in aquatic environments.

Keywords: Quinolones; Adsorption; Column; Reactive transport; Modeling.

1. Introduction

Quinolone antibiotics are broad-spectra antimicrobial agents that are widely used in human and veterinary medicine to treat and prevent infectious bacterial diseases (Oliphant and Green, 2002). Due to their extensive uses and incomplete metabolisms *in vivo*, they can be disseminated into terrestrial and aquatic environments at concentrations as high as several hundred ng per L (Fatta-Kassinos et al., 2011). Adsorption to immobile soil/sediment minerals and/or mobile colloids is one of the key processes governing the fate and mobility of quinolones in environment (Marsac et al., 2016; Xu et al., 2017a; Yu et al., 2019). This process is strongly related to the nature and relative abundance of mineral phases of common occurrence in the Earth's near-surface environment (Cornell and Schwertmann, 2003). Among these reactive mineral surfaces, goethite (α -FeOOH) is the most common and stable iron oxyhydroxide mineral with high specific surface area and high reactivity (Liu et al., 2014), and generally found as coatings on less reactive soil particles, such as silica sand (Stumm et al., 1992), in soils and sediments.

Quinolones binding on minerals can however be strongly influenced by Natural Organic Matter (NOM) (Gu and Karthikeyan, 2008; Yan et al., 2012; Peng et al., 2015; Martínez-Mejía et al., 2017; Qin et al., 2018; Cheng et al., 2018). Some studies have suggested that the presence of NOM can provide additional sorption sites for quinolones due to hydrophobic interactions, π - π interactions, ligand exchange, and hydrogen bonding between NOM and quinolones, thus promoting the adsorption of quinolones (Peng et al., 2015; Martínez-Mejía et al., 2017; Qin et al., 2018). However, other studies have reported that NOM will suppress quinolones binding because of the competition for sorption sites on minerals (Cheng et al., 2018), or electrostatic repulsions between the coated

NOM and quinolones (Gu and Karthikeyan, 2008; Yan et al., 2012). Although NOM may interact strongly with both minerals (Vindedahl et al., 2016) and antibiotics (Aristilde and Sposito, 2013; Richter et al., 2009), little is known about these co-occurring molecular interactions in natural settings where water migrates through soils and sediments. This is complicated by the complex nature of NOM — which consists of a polydisperse mixture of organic molecules of varying molecular size and chemical composition — and by its molecular fractionation resulting from its association to mineral surfaces. Currently available studies on the subject (Cheng et al., 2018; Kulshrestha et al., 2004; Qin et al., 2018; Yan et al., 2012) were limited to batch experiments, and little is known about the influence of NOM on binding and transport of antibiotics under flow-through conditions. Accounting for dynamic flow on nonequilibrium sorption of quinolones is particularly relevant when multiple mechanisms (*e.g.* binding vs co-binding and NOM molecular fractionation) concurrently influence breakthrough.

In this study, we addressed these concerns by studying the mobility of Nalidixic acid (NA), a model quinolone antibiotic, over a wide concentration range (5 – 50 mg/L) of Leonardite humic acid (LHA), a representative hydrophilic NOM. The NOM concentration range was chosen in order to cover a wide variety of dissolved organic carbon found in shallow groundwater, soil pore waters and surface waters (Degueldre et al., 1996; McKnight et al., 1992). Transport was studied under water saturated flow conditions, which are chosen to emulate contaminant migration and transformation in natural environments such as aquifer and sediments. Columns were packed with goethite-coated sand, a structurally stable and hydraulically conductive porous medium, to mimic natural mineral assemblages (Hanna et al., 2014). Impacts of LHA on NA transport was investigated in two scenarios:

(i) NOM initially present in contaminated water and then interact with co-occurring compound to mineral surfaces or (ii) NOM found as organic coatings on the mineral surface as in natural sedimentary materials (McKnight et al., 1992). In both cases, molecular fractionation of LHA and NA breakthrough were monitored by High Performance Liquid Chromatography (HPLC), UltraViolet Visible (UV-Vis) spectrophotometry and Total Organic Carbon (TOC) analyses of column effluents. A transport model that accounts for adsorption kinetics was used to help us in understanding the modifications in adsorption and desorption fronts of the breakthrough behavior of NA over a wide range of conditions. Changes in sorption kinetics can result from modifications in binding mechanisms and thus explain how NA transportation is facilitated in the presence of LHA.

2. Materials and methods

2.1. Materials

All reagents except Leonardite humic acid were purchased from Sigma-Aldrich and were used without further purification. All solutions were prepared with ultrapure water. Leonardite Humic Acid Standard (1S104H) was purchased from the International Humic Substance Society. A LHA stock solution (2 g/L, 1276 mg C/L) was prepared by dissolving 2 g LHA in 100 mL of 1 M NaOH, then diluted to 1 L with ultrapure water. A 1 mM stock solution of Nalidixic acid (NA) (purity >99%) was prepared by dissolving 1 mmole NA in 20 mL of 1 M NaOH, then diluted to 1 L with ultrapure water.

2.2. Synthesis of goethite-coated sand (GCS)

Goethite was prepared as described in previous studies (Marsac et al., 2016; Hanna et al., 2014),

and coated onto cleaned Fontainebleau quartz sand (100-150 μm) as previously detailed (Scheidegger et al., 1993; Hanna et al., 2014). All sample preparation and characterization procedures are in the Supplementary Material. All synthetic solids were washed with doubly distilled deionized water to remove soluble Fe and electrolyte ions. The solids were then stored in an anaerobic $\text{N}_2(\text{g})$ chamber at ambient temperature.

Previous work (Hanna et al., 2010, 2014) from our group confirmed that the coating procedure did not alter goethite particles and no silicates were released from quartz sand. The goethite content deposited on the sand surface, measured by acid digestion analysis, was 0.99 g/100 g of sand.

2.3. Breakthrough Column Experiments

Breakthrough column experiments were conducted at constant flow rate and under water-saturated conditions. pH conditions (i.e. pH 5) were specifically selected to ensure stable goethite coatings on sand and high adsorption of NA and LHA on goethite. Briefly, 15 g of dry goethite coated sand was packed into a glass chromatographic column of 1.6 cm internal diameter to give a porous bed length of 4.7 cm. After packing to a uniform bulk density ($1.59 \pm 0.05 \text{ g/cm}^3$), the column was wetted upward with a 0.01 M NaCl solution at pH_{in} 5 and 0.5 mL/min. The pore volume (V_{p}) was estimated by weight differences between the saturated and dry column. Once the column became water saturated, the flow characteristics of the porous bed were determined by a nonreactive tracer (bromide) experiment and described by the classic advection dispersion equation (ADE). The fitting parameters of the bromide elution confirmed the flow homogeneity and predominance of a convective regime in the column.

10 μM NA and different concentrations of LHA (0, 5, 10, 50 mg/L) were equilibrated in 0.01 M

NaCl at pH_{in} 5 and then injected simultaneously into column at 0.5 mL/min. LHA and NA concentrations in the effluents were measured. After total breakthrough of NA/LHA, desorption was initiated by injecting 0.01 M NaCl at pH_{in} 5 into column at 0.5 mL/min. After the desorption of NA/LHA, the solid was transferred to 0.1 M NaOH and shaken for 1 week. Then the supernatants were filtered (0.2 μm), and analyzed by UV-Vis spectrometry and TOC analyzer.

In another experimental series, a LHA solution (5, 10 or 50 mg/L) in 0.01 M NaCl at pH_{in} 5 was thereafter injected into the column with a continuous mode at 0.5 mL/min. The effluent of the column was collected and stored at 4°C for further analyses. After the complete breakthrough of LHA, 10 μM NA with the corresponding LHA (5, 10, 50 mg/L) in 0.01 M NaCl at pH_{in} 5 was injected in the same column at the same constant flow rate. NA concentrations in the collected fractions were measured by HPLC/UV.

The concentration of LHA in the effluent solution was determined by a total organic carbon (TOC) analyzer (Shimadzu TOC-VCSH). The UV absorbance of LHA samples were recorded in the range of 200 - 800 nm with a UV-visible spectrophotometer using 1 cm quartz cell. The following specific UV absorbance parameters were calculated as indicators of NOM fractionation. The relative absorbance ratio E_2/E_3 is defined as the ratio of absorbance at 250 nm and 365 nm. E_2/E_3 has been reported to inversely correlate with the molecular weight (Mw) and aromaticity of NOM (Li et al., 2009; Peuravuori and Pihlaja, 1997; Qin et al., 2015). Specific ultraviolet absorbance of LHA at 280 nm (SUVA_{280}) was used to estimate NOM aromaticity by dividing the molar absorptivity at 280 nm (m^{-1}) by the TOC concentration (mg/L) in the solution (Qin et al., 2015; Hur et al., 2003). The parameter S_R (slope ratio) is the ratio of the slope of the shorter wavelength region (275 nm – 295 nm)

to that of the longer wavelength region (350 nm – 400 nm) (Helms et al., 2008). S_R is reported to be negatively correlated with the M_w (Helms et al., 2008; Valencia et al., 2013).

Aqueous NA concentrations were determined by high performance liquid chromatography (Waters 600 Controller) using a UV detector (Waters 2489) and a reversed-phase C18 column (250 mm×4.6 mm i.d., 5 μ m). The mobile phase (1 mL/min) was a mixture of acetonitrile/water (60:40 v/v) contained 0.1% formic acid. The detector was set to 258 nm for NA.

Column experiments were conducted in duplicates and showed a good reproducibility within 5 % of relative standard deviation.

2.4. Model approach

The transport of NA both in the presence and absence of LHA through the columns was simulated using a one-dimensional advection-dispersion model coupled with two-site nonequilibrium adsorption model (van Genuchten and Wagenet, 1989). The two-site nonequilibrium model is based on the assumption that two types of sorption sites exist, an equilibrium site and a kinetic site. Sorption is instantaneous on the equilibrium site and described by a sorption isotherm (Type 1, equilibrium), and time dependent on the kinetics site and follows first-order kinetics (Type 2, kinetic) (van Genuchten and Wagenet, 1989). The final governing equations of the model are given by:

$$\frac{\partial C}{\partial t} + \left(\frac{\rho}{\theta}\right) \left(\frac{\partial S_1}{\partial t} + \frac{\partial S_2}{\partial t}\right) = D \frac{\partial^2 C}{\partial x^2} - v \frac{\partial C}{\partial x} \quad (1)$$

$$S_1 = K_d C^\beta \quad (2)$$

$$\frac{\partial S_2}{\partial t} = \alpha [(1 - f) K_d C^\beta - S_2] \quad (3)$$

In these equations, C is the volume-averaged solution concentration of the adsorbate (NA) (μ M), t is time (h), ρ is the bulk density of goethite-coated sand in the column (g/cm^3), θ is the volumetric water

content (cm^3/cm^3), and it equals to the porosity of the saturated column, S_1 and S_2 are sorbed-phase concentrations of NA on the equilibrium and kinetic nonequilibrium sites ($\mu\text{mol/g}$), respectively, D is the dispersion coefficient (cm^2/h), x is distance (cm), v is the average pore water velocity (cm/h), f is the fraction of equilibrium sites (Type 1), α is the first-order rate coefficient associated with the kinetic site (h^{-1}), and K_d is the Freundlich isotherm adsorption coefficient (cm^3/g) related to the sorption capacity, and β is the Freundlich exponent that characterizes the degree of nonlinearity.

Hydrus-1D program was used to obtain the model parameters for the two-site nonequilibrium model. The values of θ and D were obtained by fitting the breakthrough data of the Br^- tracer, and the values of β , K_d , f and α were obtained by fitting the transport data of NA. The best fitted f value to describe the mobility of NA alone lied at 0.6. This value was then kept constant for all simulations, which allows to reduce the number of estimated parameters. A complete sensitivity analysis, including the modeling approach used in the present work, is provided in the Supplementary Material.

3. Results and discussion

3.1. Mutual effects of NA sorption and LHA molecular fractionation

Simultaneous injection of NA and LHA in goethite-coated sand column alters NA mobility. This can be seen through the increased steepness of asymmetrical breakthrough curve of NA at larger LHA concentrations while the breakthrough point remained constant (Fig. 1). The latter was shifted down only with the highest concentration of LHA (i.e. 50 mg/L), leading to a rapid breakthrough of NA. While the NA retention did not vary the outflow pH, probably due to the low NA loading ($10 \mu\text{M}$), the presence of LHA caused pH variations in the initial stage. The increase in pH suggested that proton

co-sorption is involved when the COO⁻ groups of LHA replaced the surface –OH on goethite due to ligand exchange mechanism (Gu et al., 1994). At higher LHA loading, more COO⁻ groups replaced surface –OH group resulting in more significant increase in pH. When all –OH groups are consumed, the effluent pH decreased and finally approached the influent pH after reaching total breakthrough (Fig. 1).

Since both NA and LHA can bind to goethite through ligand-exchange and hydrogen bonding (Cheng et al., 2019; Xu et al., 2017a), competing for surface sorption sites could occur during simultaneous injection. Oxygenated polycyclic aromatics and carboxylic compounds, which generally contain in DOM of high oxidation state and high aromaticity, tend to have higher affinities for iron oxyhydroxides than alcohols, ethers and aliphatics (Cheng et al., 2019; Lv et al., 2016, 2018; Coward et al., 2019). As LHA exhibits high chemical heterogeneity with higher aromaticity (i.e. carbon aromaticity of 0.58) (Thorn and Cox, 2009) and lower aliphatic carbons contents, we expect a molecular fractionation caused by selective adsorption of LHA to goethite (Coward et al., 2019; Kleber et al., 2007). This falls in line with the breakthrough behavior of LHA in goethite-coated sand column, particularly in terms of the discrepancy between TOC and UV (Fig. 2 for LHA without NA).

Higher TOC values with respect to UV absorbance at 254 nm were observed during the first adsorption stage, before both of the two parameters followed each other to reach their inflow values. This can also be seen by the low SUVA₂₈₀ values observed at the first stage (lower PV values), suggesting that aromatics were primarily and preferentially adsorbed to goethite in the column. In addition, E₂/E₃ (Abs_{250nm}/Abs_{365nm}) and S_R of outflow solution were first lower than the initial value of

LHA and then increased sharply, and finally decreased to reach the initial value (Fig. 2c,d). This confirms that LHA compounds with smaller molecular size were preferentially retained, leading to higher Mw fractions in the column effluent (lower E_2/E_3 and S_R than the initial LHA). The increase in both parameters above the initial value suggested that outflow solution contained lower Mw fractions with higher aromaticity, while higher Mw fractions were adsorbed in the column. Note that different reports have contrast results for humic acids fractionation onto Fe-oxides, which is likely related to the HA nature/source (Kang and Xing, 2008; Qin et al., 2015, 2012; Seders Dietrich et al., 2013; Zhou et al., 2001). Higher molecular weight were preferentially adsorbed and then replaced by lower ones in solution for commercial HA (Sinopharm Chemical Reagent Co. Ltd. China.) to goethite (Qin et al., 2015), while in other reports lower Mw molecules of NOM are firstly adsorbed onto iron oxides and successively replaced by higher Mw fractions (Kang and Xing, 2008; Qin et al., 2012; Seders Dietrich et al., 2013; Zhou et al., 2001). The preferential adsorption of relatively small size fractions can be explained by their fast diffusion to the mineral surface, while the bigger HA compounds containing more reactive groups can exhibit stronger overall binding, and thus replace the lower Mw fractions during the adsorption process (Weng et al., 2007). The primary adsorption of lower Mw of LHA aromatic components were also confirmed under batch conditions, where fast and simultaneous decrease in TOC and UV was observed over the first 15 min of contact time (Fig. S1a). Higher TOC values relative to UV were observed during adsorption kinetics, while E_2/E_3 and S_R values tend to exceed the initial value after 6 h of contact time (Fig. S1b), suggesting that the lower Mw of LHA components first adsorbed may be replaced by the larger LHA components on the goethite surface. This is consistent with a recent kinetic study (Coward et al., 2019) showing a rapid primary phase

adsorption of aromatic and polycyclic aromatic compounds to goethite at the first stage, followed by lignin-like and aliphatic compounds.

On the other hand, LHA fractionation under simultaneous injection with NA is illustrated using the relative UV absorbance at 254 nm (ΔA_{250}) and 400 nm (ΔA_{400}) of outflow solutions for the three LHA concentrations (empty symbols in Fig. 3). Except at the highest LHA loading (50 mg/L), the breakthrough curves display a first step where high absorbance values were observed following by a classical breakthrough step, whereby ΔA_{400} gradually increased from zero and tended to 1 (complete breakthrough). The beginning of a second step coincides with the NA breakthrough, suggesting that NA adsorption at the first step may influence LHA retention and thus mobilize some LHA compounds. Compounds that are flushed out early have greater A_{400}/A_{254} ratios than in raw LHA (Fig. 3 and Fig. S2). The greater absorbance at 400 nm is attributed to $\pi - \pi^*$ transitions of polycyclic aromatic compounds (Chen et al., 2002). This suggests that the smaller aromatic compounds are preferentially adsorbed than multiple benzene ring structures at the first stage of breakthrough. This result is also consistent with the fractionation of LHA in column where lower Mw fractions were preferentially adsorbed (lower E_2/E_3 and S_R than the initial LHA in Fig. 2).

It is worth noting that this early breakthrough of aromatics is not observed in LHA transport experiments in the absence of NA (See UV absorbance at 400 nm in Fig. S3), confirming the impact of NA retention on the LHA molecular fractionation in column. NA likely increased the mobility of some LHA compounds through competitive binding with goethite surfaces. The full breakthrough behavior of LHA in goethite-coated sand columns was also found affected by the co-existing NA (See Breakthrough Curve (BTC) of LHA with and without NA in Fig. S4).

258

259 At low LHA loadings (5 or 10 mg/L), both NA and LHA compounds bind to surface sites of goethite
260 and the breakthrough of NA was only altered in the adsorption front (Fig. 1). In contrast, at higher
261 LHA loading (50 mg/L), NA breakthrough was altered both in the initial breakthrough point and
262 adsorption front. This rapid breakthrough of NA is likely ascribed to strong competitive adsorption of
263 LHA compounds to goethite, as well as intermolecular interactions between NA and unbound LHA
264 that decreased NA retention. This was confirmed further by a fluorescence test shown in Fig. S6.

265 This breakthrough behavior can be understood in terms of the competitive binding of NA and
266 carboxylic/aromatics compounds of LHA to the same surface sites of goethite (Cheng et al., 2019; Gu
267 et al., 1994). In particular, ligand exchange reaction with singly-coordinate -OH sites of goethite
268 (Cheng et al., 2019) is likely to drive the first step in the breakthrough curve as our previous
269 spectroscopic work (Xu et al., 2017b) shows that NA binds both as metal- and hydrogen-bonded
270 complexes on these sites. Therefore, competitive binding of NA and carboxylic/aromatics compounds
271 of LHA on -OH sites occur at the first step, followed probably by further adsorption of NA to LHA
272 covered goethite. We note that low amount of reactive phase (i.e. goethite $\sim 1\text{wt } \%$) in coated sand
273 material as well as the low amount of adsorbed NA ($0.06\text{-}0.24 \mu\text{mol} / \text{m}^2$) hinder our ability to directly
274 assess possible modifications in binding mechanisms in the column system. In addition, the great
275 molecular complexity of LHA components and the strong overlapping of vibrational bands of NA and
276 LHA compounds explain why our attempt to get molecular-level details in the ternary
277 NA/LHA/goethite system failed. Instead, we used a macroscopic sorption model as a mean to examine
278 the possibility of different mechanisms involved during the simultaneous injection of NA and LHA. In

particular, a model integrating a sorption kinetic term was applied as a theoretical tool to describe the influence of LHA on NA adsorption behavior. The model adequately predicted NA adsorption ($r^2 \geq 0.9823$, Table 1) using a lower K_d but a higher α at high LHA concentrations, thus suggesting the modification of sorption mechanisms. The lower K_d indicates smaller NA surface loadings, whereas the total amount of NA adsorbed was positively correlated with K_d (Fig. S7). Moreover, the higher α value indicates faster kinetics, as suggested by the less tailing of NA in the presence of LHA (Fig. 1). The long tailing observed in the breakthrough curve of NA without LHA was likely caused by chemical kinetic limitations, as previously reported (Hanna et al., 2012, 2010), and confirmed here for NA by the tailing mitigation obtained when a lower flow rate (0.1 mL/min) was applied (See Fig. S8). To test whether modification of sorption kinetics takes place when NA interacts with LHA-covered goethite, injection of NA in LHA preloaded columns was investigated, as detailed in the following section.

3.2. Sorption of NA in LHA preloaded columns

Preloading of goethite coated sand columns with LHA dramatically changed the transport behavior of NA in terms of both retardation factor and breakthrough curve shape (Fig. 4). Asymmetrical curve shape with an extensive tailing became sigmoidal when NA was injected in LHA preloaded columns. Increasing in LHA concentrations (from 0 to 50 mg/L) advanced the breakthrough point, yet the latter remained very close for the three tested LHA concentrations. Overall, the preliminary loading of column with LHA considerably facilitated the transport of NA, with a very similar breakthrough behavior over the concentration range of LHA (5-50 mg/L).

As LHA ligand exchange predominantly involves ~ 3.3 sites/nm² of the reactive

singly-coordinated –OH groups of goethite (Cheng et al., 2019), it becomes useful to express breakthrough results in terms of surface loadings. Based on our previous work (Cheng et al., 2019), goethite surfaces expose a mixture of non-reacted –OH groups and bound LHA at C/Fe ratios below 0.1 (namely, 26 C/nm²). All -OH groups are, on the other hand, consumed above this ratio (Cheng et al., 2019). Using the TOC of column effluents (Fig. 2a), the integrated area above the BTC shows that the total LHA adsorbed in the column was ~75 mg C/g goethite, which is equivalent to ~42 C atoms/nm² and therefore ~3 times the total crystallographic density of all surface oxygens (~15 sites/nm²) (Boily and Felmy, 2008; Cheng et al., 2019).

Under these conditions, NA retention was mainly driven by weak van der Waals-type hydrophobic interactions, and/or through π – π electron coupling interactions with LHA-covered goethite surfaces. These interactions could govern the transport behavior in preloaded columns, independently on the LHA preloading concentration. This hypothesis was further confirmed in the simulation results using a two-site nonequilibrium model of the breakthrough curves of NA in LHA-preloaded columns. Similar adsorption parameters (K_d , α and β) described the breakthrough curve of NA for the three preloading LHA concentration (Fig. 4, Table 1). The first-order rate coefficient associated with the kinetically limited site (α) increased when column is previously loaded with LHA, suggesting the modification of NA sorption kinetics. This phenomenon is also observed under batch conditions where the presence of LHA affected the adsorption kinetics of NA (Fig. S10).

3.3. Impact of LHA on the breakthrough behavior of NA

Dynamic adsorption experiments in column provided relevant data on the dual binding mechanisms of NA, *i.e.* adsorption to (i) goethite in competition with carboxylic/aromatics LHA

compounds, and (ii) LHA-covered goethite through nonspecific interactions, together with the molecular fractionation of LHA. To highlight the impact of LHA on the NA mobility/transport in the column system, full breakthrough curve consisting of both adsorption/desorption fronts was determined for NA alone and NA in presence of 50 mg/L LHA.

As shown in Fig. 5a, the full breakthrough curve of NA is asymmetrical, and exhibits tailing both in adsorption front and elution-wave, confirming the rate-limited sorption/desorption process of NA on goethite surfaces. Interestingly, the desorption front can be well predicted with the same parameters used for the adsorption, suggesting a reversible sorption process. As expected from previous column tests, the presence of LHA alters the desorption front, making the full breakthrough curve of NA more symmetrical, with less tailing in both adsorption and desorption fronts (Fig. 5b and Fig. S11). As it is difficult to directly access molecular-level information in the NA/LHA/goethite system, we have used the same model as for the previous column adsorption experiments, to describe the influence of LHA on NA mobility. If the same adsorption parameters (i.e. α , K_d , β) were used for full BTC description, the calculated curve failed to well describe the desorption tailing (Simulation#1 in Fig. 5b). Attempt to describe better the latter by changing α (see Table S1) provided a curve where adsorption front is shifted down with respect to the experimental data (Simulation#2). An accurate description of the full breakthrough curve required two different values of α for adsorption and desorption fronts (Simulation#3, See Table S1). Indeed, lower α for desorption than adsorption was needed, thereby underscoring possible modification of surface reactions. Due to the very slow and incomplete desorption of LHA compounds in column, the amount of LHA desorbed in the effluent solution becomes below detection limit after 5 PV (Fig. S12), and extension up to 800 PV had no influence on

the amount of LHA desorbed (data not shown). Solid-phase extraction conducted on the goethite coated sand solid after termination of the column experiment revealed that at least 80 % of total sorbed LHA are still present on the solid (total LHA adsorbed in the column was ~ 75 mg C/g goethite or ~ 42 C atoms/nm²). This can also be observed visually since the solid kept its brown color (due to the attachment of LHA), confirming the incomplete LHA desorption under our experimental conditions and over the whole breakthrough time. A desorption batch experiment showed that only 40 % of initial bound LHA (based on TOC measurement) were removed after 3 days and even under vigorous mixing conditions (Fig. S13). This is consistent with previous works (Gu et al., 1994; Joo et al., 2008) where the slow desorption kinetics and low desorption extent of NOM at oxide surfaces have been attributed to the complex nature of NOM compounds (though different NOM were used in these studies) and their multiple interactions involving multiple binding sites with oxide surfaces (Gu et al., 1994; Joo et al., 2008). Therefore, experimental observations and simulation results suggested that nonspecific interactions between NA and bound LHA compounds may influence desorption process. While NA desorption is almost complete, most of sorbed LHA compounds remained attached to goethite surfaces, resulting in modification in desorption kinetics.

4. Conclusions

NOM, one of the most widespread and ubiquitous components in aquatic and terrestrial environments, can affect the mobility and fate of emerging contaminants. Here, we have considered two situations where target contaminants (i) are present together with NOM in groundwater and surface waters, then bind to mineral surfaces, and (ii) where they interact with NOM coatings on

minerals. In both scenarios, NOM strongly affected quinolone binding to goethite, and thus transport behavior under flow through conditions. We notably demonstrated that LHA facilitated NA transport in columns but this effect depends on how organic matter is contacted to mineral surfaces. When LHA was present as a mineral coating, van der Waals-type hydrophobic interactions governed NA transport. However, when both LHA and NA were initially present in the influent solution, the breakthrough was characterized first by a primary step involving competitive adsorption of NA and carboxylic/aromatics compounds, and a second step involving nonspecific adsorption of NA to LHA-covered goethite. This two-step process controls both NA retention and LHA molecular fractionation in column. NOM fractionation alters sorption mechanisms and kinetics of quinolone antibiotics, which in turn affect their fractionation. Furthermore, the low and incomplete desorption of LHA compounds from mineral surfaces also alters the desorption and mobility of NA under flow through conditions.

Our work thereby shows how molecular interactions between the NOM and soil materials control the migration of contaminants within surface and subsurface environments. This calls for in-depth consideration of molecular characteristics of binding mechanisms in assessment studies of contaminant fate. These findings can have important implications on the prediction of transport of quinolone antibiotics, and of their ecological impacts in the environment.

Acknowledgements

This work was supported by the Centre National de la Recherche Scientifique (PICS CNRS 2018-2020), the Swedish Research Council (2016-03808), and the French National Research Agency

384 via the C-FACTOR project (ANR-18-CE01-0008-01).

385

386 **Supplementary Material**

387 Details of the synthesis and characterization of goethite and goethite coated sand; kinetics experiment
388 conditions and results; experimental and calculated breakthrough curves; UV spectra; Fluorescence
389 results; Sensitivity tests.

Journal Pre-proof

References

- Aristilde, L., Sposito, G., 2013. Complexes of the antimicrobial ciprofloxacin with soil, peat, and aquatic humic substances. *Environ. Toxicol. Chem.* 32, 1467–1478. <https://doi.org/10.1002/etc.2214>.
- Boily, J.-F., Felmy, A.R., 2008. On the protonation of oxo- and hydroxo-groups of the goethite (α -FeOOH) surface: A FTIR spectroscopic investigation of surface O–H stretching vibrations. *Geochim. Cosmochim. Acta* 72, 3338–3357. <https://doi.org/10.1016/j.gca.2008.04.022>.
- Chen, J., Gu, B., LeBoeuf, E.J., Pan, H., Dai, S., 2002. Spectroscopic characterization of the structural and functional properties of natural organic matter fractions. *Chemosphere* 48, 59–68. [https://doi.org/10.1016/S0045-6535\(02\)00041-3](https://doi.org/10.1016/S0045-6535(02)00041-3).
- Cheng, W., Hanna, K., Boily, J.-F., 2019. Water Vapor Binding on Organic Matter-Coated Minerals. *Environ. Sci. Technol.* <https://doi.org/10.1021/acs.est.8b05134>.
- Cheng, W., Marsac, R., Hanna, K., 2018. Influence of Magnetite Stoichiometry on the Binding of Emerging Organic Contaminants. *Environ. Sci. Technol.* 52, 467–473. <https://doi.org/10.1021/acs.est.7b04849>.
- Cornell, R.M., Schwertmann, U., 2003. The iron oxides structure, properties, reactions, occurrences and uses. Wiley-VCH, Weinheim; Cambridge.
- Coward, E.K., Ohno, T., Sparks, D.L., 2019. Direct Evidence for Temporal Molecular Fractionation of Dissolved Organic Matter at the Iron Oxyhydroxide Interface. *Environ. Sci. Technol.* 53, 642–650. <https://doi.org/10.1021/acs.est.8b04687>.
- Deguelldre, C., Pfeiffer, H.-R., Alexander, W., Wernli, B., Bruetsch, R., 1996. Colloid properties in

- 411 granitic groundwater systems. I: Sampling and characterisation. *Appl. Geochem.* 11, 677–695.
412 [https://doi.org/10.1016/S0883-2927\(96\)00036-4](https://doi.org/10.1016/S0883-2927(96)00036-4).
- 413 Fatta-Kassinos, D., Meric, S., Nikolaou, A., 2011. Pharmaceutical residues in environmental waters
414 and wastewater: current state of knowledge and future research. *Anal. Bioanal. Chem.* 399, 251–
415 275. <https://doi.org/10.1007/s00216-010-4300-9>.
- 416 Gaboriaud, F., Ehrhardt, J.-J., 2003. Effects of different crystal faces on the surface charge of colloidal
417 goethite (α -FeOOH) particles: an experimental and modeling study. *Geochim. Cosmochim. Acta* 67,
418 967–983. [https://doi.org/10.1016/S0016-7037\(02\)00988-2](https://doi.org/10.1016/S0016-7037(02)00988-2).
- 419 Gu, Baohua., Schmitt, Juergen., Chen, Zhihong., Liang, Liyuan., McCarthy, J.F., 1994. Adsorption and
420 desorption of natural organic matter on iron oxide: mechanisms and models. *Environ. Sci. Technol.*
421 28, 38–46. <https://doi.org/10.1021/es00050a007>.
- 422 Gu, C., Karthikeyan, K.G., 2008. Sorption of the Antibiotic Tetracycline to Humic-Mineral
423 Complexes. *J. Environ. Qual.* 37, 704–711. <https://doi.org/10.2134/jeq2007.0030>.
- 424 Hanna, K., Boily, J.-F., 2010. Sorption of Two Naphthoic Acids to Goethite Surface under Flow
425 through Conditions. *Environ. Sci. Technol.* 44, 8863–8869. <https://doi.org/10.1021/es102903n>.
- 426 Hanna, K., Lassabatere, L., Bechet, B., 2012. Transport of two naphthoic acids and salicylic acid in
427 soil: experimental study and empirical modeling. *Water Res.* 46, 4457–4467.
428 <https://doi.org/10.1016/j.watres.2012.04.037>.
- 429 Hanna, K., Martin, S., Quilès, F., Boily, J.-F., 2014. Sorption of Phthalic Acid at Goethite Surfaces
430 under Flow-Through Conditions. *Langmuir* 30, 6800–6807. <https://doi.org/10.1021/la4049715>.
- 431 Hanna, K., Rusch, B., Lassabatere, L., Hofmann, A., Humbert, B., 2010. Reactive transport of gentisic

- acid in a hematite-coated sand column: Experimental study and modeling. *Geochim. Cosmochim. Acta* 74, 3351–3366. <https://doi.org/10.1016/j.gca.2010.03.022>.
- Helms, J.R., Stubbins, A., Ritchie, J.D., Minor, E.C., Kieber, D.J., Mopper, K., 2008. Absorption spectral slopes and slope ratios as indicators of molecular weight, source, and photobleaching of chromophoric dissolved organic matter. *Limnol. Oceanogr.* 53, 955–969. <https://doi.org/10.4319/lo.2008.53.3.0955>.
- Hur, J., Schlautman, M.A., 2003. Using Selected Operational Descriptors to Examine the Heterogeneity within a Bulk Humic Substance. *Environ. Sci. Technol.* 37, 880–887. <https://doi.org/10.1021/es0260824>.
- Joo, J.C., Shackelford, C.D., Reardon, K.F., 2008. Association of humic acid with metal (hydr)oxide-coated sands at solid–water interfaces. *J. Colloid Interface Sci.* 317, 424–433. <https://doi.org/10.1016/j.jcis.2007.09.061>.
- Kang, S., Xing, B., 2008. Humic Acid Fractionation upon Sequential Adsorption onto Goethite. *Langmuir* 24, 2525–2531. <https://doi.org/10.1021/la702914q>.
- Kleber, M., Sollins, P., Sutton, R., 2007. A conceptual model of organo-mineral interactions in soils: self-assembly of organic molecular fragments into zonal structures on mineral surfaces. *Biogeochemistry* 85, 9–24. <https://doi.org/10.1007/s10533-007-9103-5>.
- Kulshrestha, P., Giese, R.F., Aga, D.S., 2004. Investigating the Molecular Interactions of Oxytetracycline in Clay and Organic Matter: Insights on Factors Affecting Its Mobility in Soil. *Environ. Sci. Technol.* 38, 4097–4105. <https://doi.org/10.1021/es034856q>.
- Li, A., Hu, J., Li, W., Zhang, W., Wang, X., 2009. Polarity based fractionation of fulvic acids.

- 453 Chemosphere 77, 1419–1426. <https://doi.org/10.1016/j.chemosphere.2009.09.002>.
- 454 Liu, H., Chen, T., Frost, R.L., 2014. An overview of the role of goethite surfaces in the environment.
- 455 Chemosphere 103, 1–11. <https://doi.org/10.1016/j.chemosphere.2013.11.065>.
- 456 Lv, J., Miao, Y., Huang, Z., Han, R., Zhang, S., 2018. Facet-Mediated Adsorption and Molecular
- 457 Fractionation of Humic Substances on Hematite Surfaces. *Environ. Sci. Technol.* *acs.est.8b03940*.
- 458 <https://doi.org/10.1021/acs.est.8b03940>.
- 459 Lv, J., Zhang, S., Wang, S., Luo, L., Cao, D., Christie, P., 2016. Molecular-Scale Investigation with
- 460 ESI-FT-ICR-MS on Fractionation of Dissolved Organic Matter Induced by Adsorption on Iron
- 461 Oxyhydroxides. *Environ. Sci. Technol.* 50, 2328–2336. <https://doi.org/10.1021/acs.est.5b04996>.
- 462 Marsac, R., Martin, S., Boily, J.-F., Hanna, K., 2016. Oxolinic Acid Binding at Goethite and
- 463 Akaganéite Surfaces: Experimental Study and Modeling. *Environ. Sci. Technol.* 50, 660–668.
- 464 <https://doi.org/10.1021/acs.est.5b04940>.
- 465 Martínez-Mejía, M.J., Sato, I., Rath, S., 2017. Sorption mechanism of enrofloxacin on humic acids
- 466 extracted from Brazilian soils. *Environ. Sci. Pollut. Res.* 24, 15995–16006.
- 467 <https://doi.org/10.1007/s11356-017-9210-3>.
- 468 McKnight, D.M., Bencala, K.E., Zellweger, G.W., Aiken, G.R., Feder, G.L., Thorn, K.A., 1992.
- 469 Sorption of dissolved organic carbon by hydrous aluminum and iron oxides occurring at the
- 470 confluence of Deer Creek with the Snake River, Summit County, Colorado. *Environ. Sci. Technol.*
- 471 26, 1388–1396. <https://doi.org/10.1021/es00031a017>.
- 472 Oliphant, C.M., Green, G.M., 2002. Quinolones: a comprehensive review. *Am. Fam. Physician* 65,
- 473 455–464.

- 474 Peng, H., Liang, N., Li, H., Chen, F., Zhang, D., Pan, B., Xing, B., 2015. Contribution of coated
 475 humic acids calculated through their surface coverage on nano iron oxides for ofloxacin and
 476 norfloxacin sorption. *Environ. Pollut.* 204, 191–198.
 477 <https://doi.org/10.1016/j.envpol.2015.04.029>.
- 478 Peuravuori, J., Pihlaja, K., 1997. Molecular size distribution and spectroscopic properties of aquatic
 479 humic substances. *Anal. Chim. Acta* 337, 133–149.
 480 [https://doi.org/10.1016/S0003-2670\(96\)00412-6](https://doi.org/10.1016/S0003-2670(96)00412-6).
- 481 Qin, X., Du, P., Chen, J., Liu, F., Wang, G., Weng, L., 2018. Effects of natural organic matter with
 482 different properties on levofloxacin adsorption to goethite: Experiments and modeling. *Chem.*
 483 *Eng. J.* 345, 425–431. <https://doi.org/10.1016/j.cej.2018.03.125>.
- 484 Qin, X., Liu, F., Wang, G., 2012. Fractionation and kinetic processes of humic acid upon adsorption on
 485 colloidal hematite in aqueous solution with phosphate. *Chem. Eng. J.* 209, 458–463.
 486 <https://doi.org/10.1016/j.cej.2012.08.026>.
- 487 Qin, X., Liu, F., Wang, G., Hou, H., Li, F., Weng, L., 2015. Fractionation of humic acid upon
 488 adsorption to goethite: Batch and column studies. *Chem. Eng. J.* 269, 272–278.
 489 <https://doi.org/10.1016/j.cej.2015.01.124>.
- 490 Richter, M.K., Sander, M., Krauss, M., Christl, I., Dahinden, M.G., Schneider, M.K., Schwarzenbach,
 491 R.P., 2009. Cation Binding of Antimicrobial Sulfathiazole to Leonardite Humic Acid. *Environ.*
 492 *Sci. Technol.* 43, 6632–6638. <https://doi.org/10.1021/es900946u>.
- 493 Scheidegger, A., Borkovec, M., Sticher, H., 1993. Coating of silica sand with goethite: preparation and
 494 analytical identification. *Geoderma* 58, 43–65. [https://doi.org/10.1016/0016-7061\(93\)90084-X](https://doi.org/10.1016/0016-7061(93)90084-X).

- 495 Seders Dietrich, L.A., McInnis, D.P., Bolster, D., Maurice, P.A., 2013. Effect of polydispersity on
 496 natural organic matter transport. *Water Res.* 47, 2231–2240.
 497 <https://doi.org/10.1016/j.watres.2013.01.053>.
- 498 Stumm, W., Sigg, L., Sulzberger, B., 1992. Chemistry of the solid-water interface: processes at the
 499 mineral-water and particle-water interface in natural systems. Wiley, New York.
- 500 Thorn, K.A., Cox, L.G., 2009. N-15 NMR spectra of naturally abundant nitrogen in soil and aquatic
 501 natural organic matter samples of the International Humic Substances Society. *Org. Geochem.* 40,
 502 484–499. <https://doi.org/10.1016/j.orggeochem.2009.01.007>.
- 503 Valencia, S., Marín, J.M., Restrepo, G., Frimmel, F.H., 2013. Application of excitation–emission
 504 fluorescence matrices and UV/Vis absorption to monitoring the photocatalytic degradation of
 505 commercial humic acid. *Sci. Total Environ.* 442, 207–214.
 506 <https://doi.org/10.1016/j.scitotenv.2012.10.058>.
- 507 van Genuchten, M.Th., Wagenet, R.J., 1989. Two-Site/Two-Region Models for Pesticide Transport
 508 and Degradation: Theoretical Development and Analytical Solutions. *Soil Sci. Soc. Am. J.* 53,
 509 1303. <https://doi.org/10.2136/sssaj1989.03615995005300050001x>.
- 510 Vindedahl, A.M., Stemig, M.S., Arnold, W.A., Penn, R.L., 2016. Character of Humic Substances as a
 511 Predictor for Goethite Nanoparticle Reactivity and Aggregation. *Environ. Sci. Technol.* 50, 1200–
 512 1208. <https://doi.org/10.1021/acs.est.5b04136>.
- 513 Weng, L., Van Riemsdijk, W.H., Hiemstra, T., 2007. Adsorption of humic acids onto goethite: Effects
 514 of molar mass, pH and ionic strength. *J. Colloid Interface Sci.* 314, 107–118.
 515 <https://doi.org/10.1016/j.jcis.2007.05.039>.

- 516 Xu, J., Marsac, R., Costa, D., Cheng, W., Wu, F., Boily, J.-F., Hanna, K., 2017a. Co-Binding of
517 Pharmaceutical Compounds at Mineral Surfaces: Molecular Investigations of Dimer Formation at
518 Goethite/Water Interfaces. *Environ. Sci. Technol.* 51, 8343–8349.
519 <https://doi.org/10.1021/acs.est.7b02835>.
- 520 Xu, J., Marsac, R., Wei, C., Wu, F., Boily, J.-F., Hanna, K., 2017b. Cobinding of Pharmaceutical
521 Compounds at Mineral Surfaces: Mechanistic Modeling of Binding and Cobinding of Nalidixic
522 Acid and Niflumic Acid at Goethite Surfaces. *Environ. Sci. Technol.* 51, 11617–11624.
523 <https://doi.org/10.1021/acs.est.7b02900>.
- 524 Yan, W., Hu, S., Jing, C., 2012. Enrofloxacin sorption on smectite clays: Effects of pH, cations, and
525 humic acid. *J. Colloid Interface Sci.* 372, 141–147. <https://doi.org/10.1016/j.jcis.2012.01.016>.
- 526 Yu, C., Bahashi, J., Bi, E., 2019. Mechanisms and quantification of adsorption of three
527 anti-inflammatory pharmaceuticals onto goethite with/without surface-bound organic acids.
528 *Chemosphere* 222, 593–602. <https://doi.org/10.1016/j.chemosphere.2019.01.155>.
- 529 Zhou, Q., Maurice, P.A., Cabaniss, S.E., 2001. Size fractionation upon adsorption of fulvic acid on
530 goethite: equilibrium and kinetic studies. *Geochim. Cosmochim. Acta* 65, 803–812.
531 [https://doi.org/10.1016/S0016-7037\(00\)00536-6](https://doi.org/10.1016/S0016-7037(00)00536-6).

- NOM compounds with higher aromaticity and lower Mw are preferentially adsorbed.
- NOM fractionation alters sorption mechanisms and kinetics of quinolones.
- Multiple mechanisms influence NA mobility when NOM are present together with NA.
- Nonspecific interactions govern NA mobility when NOM are found as mineral coatings.
- A transport model is developed to interpret the modifications in binding mechanisms.

Credit Author Statement:

WC, RM and KH designed the study. WC carried out the experiments and wrote the manuscript. LZ performed the calculations. JFB and KH: Wrote, reviewed and edited the manuscript. All authors discussed the results and commented on the manuscript.

Declaration of interests

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: