

Paleotemperature investigation of the Variscan southern external domain: the case of the Montagne Noire (France)

Clément Montmartin, Michel Faure, Hugues Raimbourg

► To cite this version:

Clément Montmartin, Michel Faure, Hugues Raimbourg. Paleotemperature investigation of the Variscan southern external domain: the case of the Montagne Noire (France). Bulletin de la Société Géologique de France, 2020, 10.1051/bsgf/2020043. insu-03086940v1

HAL Id: insu-03086940 https://insu.hal.science/insu-03086940v1

Submitted on 23 Dec 2020 (v1), last revised 31 Mar 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Paleotemperature investigation of the Variscan southern external

- 2 domain: the case of the Montagne Noire (France)
- 3 Étude de paléotempérature dans le domaine méridional externe varisque: le cas de la
- 4 Montagne Noire (France)
- 5 Clément Montmartin*, Michel Faure, Hugues Raimbourg
- 6 Institut des Sciences de la Terre d'Orléans (ISTO), UMR 7327, Université d'Orléans-CNRS-
- 7 BRGM, Campus Géosciences, 1A Rue de la Férollerie 45071 Orléans Cedex 2, France
- 8 * Corresponding author, <u>clement.montmartin@univ-orleans.fr</u>, 02 38 49 25 73

9 Keywords:

- 10 Raman Spectroscopy of Carbonaceous Material (RSCM) Spectrométrie Raman sur Matière
- 11 Carbonée (RSCM)
- 12 Granite-migmatite dome Dôme granite-migmatite
- 13 Montagne Noire Montagne noire
- 14 Variscan orogen Orogenèse Varisque
- 15 French Massif Central Massif Central Français
- 16 Recumbent folds Plis couchés

17 Résumé

18 La Montagne Noire, située dans la partie sud du Massif Central français, représente la partie

19 nord de l'avant-pays varisque. La zone est divisée en trois parties. Le dôme granite-

- 20 migmatite de la zone axiale est entouré de séries sédimentaires paléozoïques pas ou
- 21 faiblement métamorphisées. Les flancs nord et sud du dôme de la Montagne Noire sont
- 22 déformés par des plis d'échelle kilométrique, déversés vers le sud sud-est. La méthode de
- 23 spectrométrie Raman sur matière carbonée (RSCM), réalisée dans les roches de bas grade
- 24 métamorphique du flanc sud de la Montagne Noire, a donné des températures comprises
- 25 entre 400°C près du dôme et 230°C dans le domaine sud. Trois géothermomètres Raman

ont été utilisés pour couvrir cette gamme de température. Ces températures RSCM sont
qualitativement conformes aux estimations précédentes basées sur la cristallinité de l'illite,
l'altération de la couleur des conodontes et les inclusions fluides effectuées dans la même
zone, qui démontrent une augmentation de la température vers le dôme.

Les isothermes traversent les différents contacts de nappe et sont orientées
parallèlement à la marge sud de la zone axiale. Cette distribution de température suggère
que la structure thermique a été acquise lors de la mise en place du dôme de la zone axiale.
La structure thermique acquise lors de la mise en place des plis couchés et de
l'enfouissement des séries sédimentaires est ainsi totalement effacé par le dôme.

De plus, dans un domaine relativement éloigné du dôme de la zone axiale, les mesures RSCM ont donné des températures significativement plus élevées que la cristallinité illite. Cette divergence indique une sensibilité plus élevée du RSCM face à la cristallinité illite aux événements thermiques de courte durée, probablement en raison d'une cinétique plus efficace de la réaction de carbonisation. D'autre part, de fortes températures RSCM analysées loin de la Zone Axiale, entre 300°C et 360°C, pourraient être expliquées par la présence de plutons granitiques sous le bassin d'avant-pays.

42 Abstract

The Montagne Noire located in the southern part of the French Massif Central represents the 43 northern part of the South-Variscan Foreland. It is subdivided into three parts. The granite-44 migmatite Axial Zone dome is surrounded by non- or weakly metamorphosed Paleozoic 45 sedimentary series. Both northern and southern flanks of the Montagne Noire dome are 46 deformed by km-scale, south to southeast facing recumbent folds and thrusts sheets. The 47 Raman Spectroscopy of Carbonaceous Material (RSCM) method, carried out in the low-48 49 grade metamorphic rocks of the southern flank of the Montagne Noire, yielded temperatures comprised between 400°C near the dome, and 230°C in the southern domain. Three Raman 50 geothermometers were used to cover this temperature range. RSCM temperatures comply 51

qualitatively with previous estimates based on illite crystallinity, conodont colour alteration,
and fluid inclusions carried out in the same area, which document a metamorphic
temperature increase towards the dome.

The isotherms cut across the different nappe contacts and are oriented parallel to the southern margin of the Axial Zone. This temperature distribution supports the idea that the thermal structure was acquired during the Axial Zone dome emplacement. The thermal structure acquired during the recumbent folds emplacement and burial of the sedimentary series is totally overprinted by the doming event.

In addition, in a domain relatively remote from the Axial Zone dome, the RSCM measurements yielded significantly higher temperatures than illite crystallinity. This discrepancy points to a higher sensitivity of RSCM to short-lived thermal events than illite crystallinity, possibly because of more efficient kinetics of the carbonization reaction. On the other hand, high RSCM temperatures analysed far from the Axial Zone, between 300°C and 360°C could be explained by the presence of granitic plutons under the foreland basin.

66

1. Introduction

67 The knowledge of quantitative constraints, such as pressure, temperature, duration of 68 heating, strain, strain rate, exhumation rate, and uplift rate, is essential to understand the formation and evolution of a mountain belt. In recent years, many studies (e.g Beyssac et al., 69 70 2002; Gerya and Stöckhert, 2002) dealt with these issues thanks to significant advances in petrology and geochronology. In the inner domain of an orogen, temperature and pressure 71 conditions experienced by metamorphic rocks can be approached by petrological 72 investigations based on mineral parageneses. On the contrary, in the outer domain, where 73 74 the deformation and associated metamorphism are usually less developed than in the inner 75 domain, the evaluation of the thermo-barometric conditions is difficult, since the size of syntectonic metamorphic minerals is generally small and conditions of thermodynamic equilibria 76 are often not achieved (Frey, 1987; Merriman and Frey, 1999; Merriman and Peacor, 1999). 77

The measurement of low temperature or low pressure values is therefore delicate, and often 78 79 bears a large error. However, several methods have been developed to overcome this 80 problem and to reach a quantitative knowledge of the conditions of low-grade 81 metamorphism, including illite crystallinity, conodont alteration colour, calcite-dolomite 82 thermo-barometer (e.g. Kübler, 1968; Dunoyer de Segonzac et al., 1968; Epstein et al., 1977; Frey, 1987). More recently, the Raman Spectroscopy on Carbonaceous Material 83 (RSCM) method based on carbonaceous material crystallinity has been developed for the 84 85 temperature range 330-600°C (Beyssac et al., 2002). Its applicability has been further extended towards low temperatures (Rahl et al., 2005; Lahfid et al., 2010; Kouketsu et al., 86 87 2014), using different calibrations of the Raman spectra based on other geothermometers, such as vitrinite reflectance, fluid inclusions microthermometry or illite crystallinity (Rahn et 88 89 al., 1995, Hara et al., 2013) or indirect data on temperature provided by fission-tracks and U-Th/He method (Rahl et al., 2005). 90

The applicability of the RSCM method is a priori restricted to the conditions of its 91 calibration. In all the different studies that focused on RSCM at low temperature, the case 92 93 studies were chosen in subduction or collision settings such as Sanbagawa Belt in Japan (Kouketsu et al., 2014) and the Glarus Alps in Switzerland (Lahfid et al., 2010), providing 94 examples of metamorphic gradients. Therefore, in these cases, the thermal imprint on the 95 96 rocks is related to burial then exhumation within a subduction zone or a collision belt, i.e. the 97 thermal event lasts for at least one or several million years. Heating by a magmatic intrusion 98 is potentially much shorter-lived and the inter-compatibility of the different geothermometers in such a case is not guaranteed (Velde and Lanson, 1993). 99

When compared to other geothermometers, the biases linked to the kinetics of the thermal recording by the carbonaceous material with respect to other mineral reactions should be considered (Velde and Lanson, 1993; Mullis et al., 2017). It appears that duration of heating heavily impacts maturity processes and is a key parameter that controls the measured temperature (Wada et al., 1994; Mählmann, 2001; Le Bayon et al., 2011; Mori et al., 2017). Moreover, the geothermometers based on carbonaceous material differ from fluid inclusion or metamorphic mineral ones mostly because no water is needed for the maturation
of carbonaceous material (Velde and Lanson, 1993; Le Bayon et al., 2011). Thus, even if the
whole rock experienced the same thermal history, disparities might be observed between the
different methods.

110 Further complexity arises when dealing with multi-metamorphic events, as in the case of post-orogenic, large-scale heating and pluton emplacement postdating collisional 111 112 metamorphism, for instance in the Jebilet Massif in Moroccan Variscan Belt (Delchini et al., 113 2016). A contact metamorphism linked to a magmatic intrusion may overprint the crystallinity of carbonaceous material inherited from earlier stages, as it has already been documented in 114 several studies. For instance, Mori et al. (2017), measured temperatures imposed by thin 115 (50m) magmatic bodies such as the Great Whin Sill in the UK, overprinting the regional low-116 117 grade metamorphism (<300°C). On a larger scale, Aoya et al. (2010) focussed on contact metamorphic aureoles around granitic plutons in Japan within two poorly metamorphosed 118 119 accretionary complexes (<400°C). Hilchie et al. (2014) have studied the South Mountain Batholith intrusion thermicity of Nova Scotia emplaced under a regional greenschist facies 120 121 metamorphism. In each case, RSCM temperatures decrease with the distance to the intrusive bodies. Finally, shear heating between structural units could also influence the 122 organic matter organization (e.g. Mori et al., 2015). The interpretation of RSCM data in 123 124 domains with a poly-stage metamorphic evolution is therefore often ambiguous, as the 125 RSCM measurements combine the contribution of regional syn-tectonic metamorphism 126 coeval with crustal thickening due to nappe stacking and the later thermal overprint related to 127 a magmatic event.

The present study aims to decipher the thermal evolution recorded in a cold foreland domain of a collisional orogen overprinted by a hot metamorphic dome (e. g. Gèze, 1949; Schuiling, 1960; Demange, 1982; Echtler and Malavieille, 1990; Soula et al., 2001; Alabouvette et al., 2003; Franke et al., 2011). The study area is located in the southern outer zone of the Variscan belt, in the Montagne Noire of the French Massif Central (Fig.1). The bulk paleotemperature field is derived from 73 RSCM measurements of graphite crystallinity

134	on weakly deformed silts and pelites from Paleozoic formations that constitute the southern
135	Variscan fold-and-thrust belt of the Massif Central. These new results are discussed in the
136	tectonic framework of this segment of the Variscan orogen.

137

7 2. Geological Background

The Variscan belt develops from South Spain to PolandThe French Massif Central is a stack 138 of nappes formed through a long and complex polyorogenic evolution (for details see Faure 139 140 et al., 2005, 2009; 2017). The SE part of the Massif exposes a succession of thrusts and nappes formed from Visean to Bashkirian during the late collisional intracontinental evolution 141 142 (the D3 event of Faure et al., 2009). In the Montagne Noire area, the southernmost part of 143 Massif Central, weakly metamorphosed sedimentary series develop in a southeast-verging 144 fold-and-thrust belt (Fig. 1). There, kilometre-scale recumbent folds affect the entire 145 sedimentary series, including the Visean-Serpukhovian flysch. Over the whole belt, the 146 crustal thickening resulting from the successive emplacement from North to South of large-147 scale nappes (UGU, LGU, PAU), was followed by pervasive melting affecting the tectonic pile. Middle to Late Carboniferous migmatites and granites are widespread in the Massif 148 149 Central (Fig. 1).

150

3. The outer Variscan domain in the Montagne Noire

151 3.1. General structure

152 The Montagne Noire is subdivided into southern and northern flanks separated by the Axial Zone (e.g. Gèze, 1949; Arthaud, 1970; Fig. 2). The former two zones consist of Cambrian, 153 154 Ordovician, Devonian and Carboniferous sedimentary rocks folded and thrust to the south 155 from the Visean to the Serpukhovian. The Axial Zone is a granite-migmatite dome emplaced within the folded series. Paleozoic formations are well exposed in the Montagne Noire 156 southern flank (Fig. 3). Detailed lithological descriptions are available in Gèze (1949); Feist 157 and Galtier (1985); Álvaro et al. (1998); Vizcaïno and Álvaro (2001) and Alabouvette et al. 158 (2003). The Visean-Serpukhovian series, well exposed on the southern flank, consists of 159

syn-tectonic gravity flow deposits such as greywacke turbidites and olistostromes, which 160 161 recycle pre-Visean sedimentary rocks of the advancing nappe pile (Engel et al., 1978, 1980-1981; Feist and Galtier, 1985; Poty et al., 2002; Vachard and Aretz, 2004; Cózar et al., 2017; 162 163 Vachard et al., 2017). During the Carboniferous, the progressive deepening of the sedimentary basin argues for a foreland trough, filled by syntectonic deposits, and coeval 164 with the formation of recumbent folds. The entire Paleozoic series has been deformed into 165 several km-scale, southeast-verging, thrust sheets and recumbent folds (Arthaud 1970; 166 Echtler and Malavieille, 1990). 167

168 3.2. Tectonic subdivisions

169 3.2.1. The Southern Flank

170 The upper recumbent fold, or Pardailhan unit, is the highest tectonic unit of the southern 171 flank of the Montagne Noire. It is composed of Cambrian to Devonian sedimentary rocks with an inverted stratigraphic order, i.e. the geometrically upper part consists of Early Cambrian 172 sandstone, and the lower part is formed by Devonian limestone boudins that delimits the 173 basal shear zone separating the upper recumbent fold from the lower one. From north to 174 175 south, perpendicularly to the fold axes, the inverted limb of the Pardailhan recumbent fold develops along 12 km. This inverted sedimentary series is subdivided into three km-scale 176 recumbent anticlines overturned to the south (Gèze, 1949; Arthaud, 1970; Alabouvette et al., 177 178 1982; 2003).

179 Structurally below the upper recumbent fold, the lower one, also named Mt-Peyroux 180 unit, can be observed both in the eastern and western sides of the upper recumbent fold (Fig. 2). The western part of the lower recumbent fold is called the Minervois unit. The 181 182 highest part of this recumbent fold is occupied by Ordovician flysch while Visean turbidites 183 crop out in the lower part, indicating a stratigraphic inversion. NE-SW upright antiforms and synforms deform the inverted series of the entire nappe stack. The poorly exposed contact 184 between the lower recumbent fold and the foreland basin led to controversial interpretations. 185 According to Arthaud (1970), and Alabouvette et al. (1982, 2003) the inverted limb is in 186

contact with Visean-Serpukhovian turbidites. However, according to (Engel et al., 1978, 187 188 1980-1981), the flysch sequence is part of the lower recumbent (Mt. Peyroux) fold. The normal and inverted limbs of that fold nappe are connected by a D1 fold hinge. Whatever the 189 190 structural interpretation, this does not change the RSCM results presented below, and their 191 interpretations. The presence of Paleozoic olistoliths in the Visean-Serpukhovian turbidites basin in which the recumbent folds are emplaced documents the syn-sedimentary character 192 193 of this event. It provides an important time constraint for the tectonic evolution of the 194 southern flank: the nappe thrusting is contemporary or older than 318Ma (Engel et al., 1978, 195 1980-1981; Alabouvette et al., 1982, 2003; Vachard et al., 2017).

196 Below the eastern part of the lower (Mt Peyroux) recumbent fold, characterized by overturned Ordovician to Early Carboniferous series, the Mts de Faugères unit consists of 197 198 Devonian and Early Carboniferous sedimentary rocks deformed by southeast-verging recumbent folds. The Para-autochthon unit is composed of Devonian to Middle 199 200 Carboniferous sedimentary rocks exposed in the normal stratigraphic order. This domain is observed between the lower recumbent fold and the Axial Zone metamorphic rocks. Close to 201 202 the Axial Zone, for instance near Saint-Pons (Fig. 2), the Para-autochthonous unit consists of Devonian marbles and micaschists (Engel et al., 1980-1981; Feist and Galtier, 1985; Poty et 203 al., 2002; Vachard and Aretz, 2004; Cózar et al., 2017; Vachard et al., 2017). 204

The autochthonous turbiditic basin represents the foreland basin into which the Mt-Peyroux and Mts-de-Faugères recumbent folds were emplaced. The basin substratum is unknown but might probably be similar to that observed in the northern para-autochthonous unit. To the west, the basin underlies the stack of recumbent folds, and to the south, it is hidden below the Cenozoic formations.

210 3.2.2. The Axial Zone

The Axial Zone is composed of orthogneiss, paragneiss, amphibolite, micaschist, and rare marble that experienced a partial melting giving rise to migmatites and granites. The Axial Zone presents a dome architecture of about 90 km long, and 20 km wide, with a N70E long

axis (Gèze, 1949; Faure and Cottereau, 1988; Echtler and Malavieille, 1990; Matte et al., 214 215 1998; Van den Driessche and Brun, 1992; Demange, 1993; Alabouvette et al., 2003). The 216 Eocene Pyrenean Mazamet fault divides the western part of the dome in two parts, the Agout 217 and Nore massifs in the North and South, respectively. The orthogneiss yields zircon U-Pb Ordovician to Devonian ages at 472 ± 2.8 Ma , 456 ± 3 Ma, 450 ± 6 Ma, 455 ± 2 Ma, and 416218 \pm 5 Ma interpreted as those of the granite protolith (Roger et al., 2004; Cocherie et al., 2005; 219 Franke et al., 2011; Pitra et al., 2012; Trap et al., 2017). The country rocks protoliths of these 220 221 orthogneiss are interpreted as Neoproterozoic to Cambrian or Ordovician (Alabouvette et al., 222 2003). Several geochronological ages on the migmatite and anatectic granites range between 333 Ma and 294 Ma (Hamet and Allègre, 1976; Faure et al., 2010; Franke et al., 223 2011; Poilvet et al., 2011; Roger et al., 2015; Trap et al., 2017) with pegmatite until 282 Ma 224 225 (Doublier et al., 2015). The abundance of granitoids and migmatite argue for widespread 226 thermal event. The Axial Zone experienced two metamorphic and tectonic events. The early 227 one is recorded only in rare mafic eclogitic restites enclosed in the migmatite. Though zircon in eclogite yields a ca. 315 Ma age (Faure et al., 2014; Whitney et al., 2015), this can hardly 228 229 be interpreted as the age of the high pressure event since the enclosing migmatite yields ages older than the eclogite blocks. A hydrothermal event might be responsible for zircon 230 recrystallization. The second, and main one is coeval with the dome formation (Schuilling, 231 1960; Bard and Rambeloson, 1973; Thompson and Bard, 1982; Soula et al., 2001; Faure et 232 233 al., 2014; Fréville et al., 2016; Trap et al., 2017).

The metasedimentary rocks that form the outer envelope of the granite-migmatite dome underwent a pervasive high temperature/medium pressure (HT/MP) metamorphism coeval with the dome formation (Soula et al., 2001; Faure et al., 2014; Fréville et al., 2016). From South to North, chlorite, biotite, garnet, andalousite, staurolite, and sillimanite isograds appear successively. As already pointed out, chlorite, biotite, and muscovite belonging to the low temperature part of the HT/MP metamorphism develops in the various lithotectonic units close to the Axial Zone, such as the para-autochthonous, Mts-de-Faugères, Mt-Peyroux and Pardailhan units (Arthaud, 1970; Franke et al., 2011; Fréville et al., 2016). The Axial Zone
HT/MP metamorphism started after the emplacement of the thrusts and recumbent folds but
continued still afterwards into the Permian (Alabouvette et al., 2003; Franke et al., 2011; Pitra
et al., 2012).

245 3.2.3. The Northern Flank

246 The Montagne Noire northern flank is composed of Late Neoproterozoic (Ediacaran) to Silurian sedimentary formations subdivided into several south-directed folds and thrusts 247 considered as equivalent to those of the southern flank. The deformation age is estimated of 248 249 Visean age (342-333 Ma; Gèze, 1949; Alabouvette et al., 2003; Doublier et al., 2006). The 250 southern part of the northern flank, east and SW of the Lacaune fault (Fig. 2), exposes 251 biotite, garnet, and alousite micaschists comparable to the metapelites of the southern envelope of the Axial Zone (Demange, 1982). This metamorphism that superimposes to the 252 fold-and thrust structure was coeval with the HT/MP event related to the Axial Zone doming 253 254 (Doublier et al., 2006). The contact between the northern flank and Axial Zone is the Mts-de-255 Lacaune NE-dipping ductile normal fault that accommodated the dome emplacement (Van den Driessche and Brun, 1989). Furthermore, a brittle normal fault controlled the opening of 256 257 the Late Carboniferous (Gzhelian) Graissessac coal basin. In the following, the northern flank will not be considered. 258

3.3. The tectonic outline of the southern flank

The structural analysis of the Montagne Noire southern flank revealed two main tectonic events (Arthaud, 1970). The first deformation stage corresponds to the emplacement of the south verging recumbent folds (F1) and thrusts. The bedding and cleavage relationships demonstrate the south to southeastward vergence of the F1 folds (Arthaud, 1970; Fig. 4a). A recent study proposes a nappe thrusting towards the northeast (Chardon et al., 2020). Whatever the tectonic interpretation, in spite of 15 km of subhorizontal displacement, and the ca 1.5 km overload due to the stratigraphic inversion, the Paleozoic series poorly exhibit a ductile syn-metamorphic deformation. A slaty cleavage is only visible in the hinges of F1
folds (Arthaud, 1970). The metamorphic grade reached by the upside down Paleozoic series
in the Upper (Pardailhan) and Lower (Mt-Peyroux) recumbent folds is below the biotite grade,
which firstly appear near the dome, in the para-autochthonous unit.

The second deformation event is an upright folding phase (F2). Field observations 271 document the refolding of the F1 recumbent folds by the F2 upright ones with NE-SW striking 272 axes (Arthaud, 1970; Alabouvette et al., 2003). Close to the Axial Zone, a subvertical axial 273 274 planar cleavage develops in the F2 folds, whereas away from the Axial Zone dome, the F2 folds are rather open and devoid of cleavage (Fig. 4b). The F2 upright folding corresponds to 275 a NW-SE shortening that formed large- scale antiforms and synforms including the Axial 276 277 Zone dome. A N70 mineral and crenulation lineation appears in the Carboniferous formation 278 at about 5 km south of the dome and northward (Harris et al., 1983). Sericite is also recognized north of this point (close to sample H11 - Fig. 2). However, Franke et al. (2011) 279 280 interpreted this lineation as an extension direction linked to the pull apart emplacement of the Axial Zone. Anyhow, these metamorphic and structural features argue for a syntectonic 281 282 metamorphism linked to the doming (Soula et al., 2001; Franke et al., 2011).

283

4. The RSCM method

The Raman spectroscopy on carbonaceous material (RSCM) method was carried out to estimate the paleotemperature field of the low grade, weakly deformed, sedimentary rocks of the southern flank of the Montagne Noire. The method is based on the irreversible transformation of carbonaceous matter towards graphite structure, using empirical correlations to correlate Raman spectra of carbonaceous matter into maximum temperature of heating, in the range 150-650°C (Beyssac et al., 2002, Lahfid et al., 2010).

290 4.1. Analytical settings

The Raman measurements were carried out on a Renishaw inVia reflex system belonging to the ISTO-BRGM analytic platform. The Wire 3.4 software was used for the data acquisition. 293 The argon-ion laser source excitation of 514.5nm was set at a power of about 5% of its 294 capacity (2.5mW). The monochromatic ray was coupled to a reflection microscope with an 295 x100 objective. Before each series of measurement, the spectrometer was calibrated using 296 an internal silica standard for the wavenumber (520,4 cm⁻¹) and the signal intensity (at least 297 30 000 counts per second). Fifteen carbonaceous matter spectra were systematically acquired for each sample within the thin section in order to obtain a representative and 298 reliable temperature. Only organic particles located below transparent quartz, a few µm 299 300 under the thin section surface, were analysed, to avoid any mechanical damaging of their 301 crystalline structure due to thin section preparation and polishing. The acquisition duration 302 was set to at least 60s and adapted depending on the quality of the spectrum.

303 4.2. RSCM data processing

Several empirical calibrations were proposed to correlate the carbonaceous matter Raman spectra to temperature (Beyssac et al., 2002; Rahl et al., 2005; Aoya et al., 2010; Lahfid et al., 2010; Kouketsu et al., 2014). Carbonaceous matter presents characteristic Raman bands in the wavenumber range between 1100-1800 cm⁻¹ and 2500-3100 cm⁻¹. The first order region of the carbonaceous matter spectrum for 1100-1800 cm⁻¹ provided the data of this study.

310 Depending on its crystallinity (hence on the temperature experienced), carbonaceous matter presents a variable number of Raman peaks, which can be used to discriminate 311 between the different calibration methods. Since the sedimentary rocks of the Montagne 312 Noire southern flank are weakly to un-metamorphosed, most of the Raman spectra were 313 processed with the Lahfid et al. (2010) method. Accordingly, each spectrum was 314 315 decomposed into five bands (Lorentzian functions), with a well-constrained position: G (1590 cm⁻¹) D1 (1350 cm⁻¹), D2 (1620 cm⁻¹), D3 (1515 cm⁻¹) and D4 (1250 cm⁻¹) (Lahfid et al., 2010; 316 Sadezky et al., 2005). From this decomposition, the RSCM temperature was computed using 317 the formula proposed by Lahfid et al. (2010) which is valid between 200°C and 320°C, with 318 an accuracy of $\pm 30^{\circ}$ C (Fig. 5a): 319

320
$$T^{\circ}C = \frac{RA1 - 0.3758}{0.0008}$$
 with $RA1 = \frac{D1 + D4}{D1 + D2 + D3 + D4 + G}$

Samples with higher crystallinity have fewer Raman peaks and were processed accordingly using the method proposed by Beyssac et al. (2002) that allows an estimation of temperature in the range between 330 up to 650°C, with a \pm 50°C uncertainties (Fig. 5b)

324
$$T^{\circ}C = (-445 * R2 + 641)$$
 with $R2 = \frac{D1}{D1 + D2 + G}$

Considering the low metamorphic grade of the analysed samples, only the low range 325 of this method was used (<400°C). Since the Beyssac et al. (2002) and Lahfid et al. (2010) 326 327 methods do not exactly overlap with each another; we also used a third thermometer 328 (Kouketsu et al., 2014). This method does not correspond to peaks area ratios but focuses on the peak full width at half maximal (FWHM) of the D1 and D2 band. Because of the 329 difficulty to distinguish the D2 band from the G band, only the formula about the former band 330 331 was retained. According to this method, the temperature is derived from Raman peaks as: $T^{\circ}C = -2.15 * (FWHM - D1) + 478$ 332

The Kouketsu et al. (2014) method leads to temperatures comprised between 150 and

334 400°C with a \pm 30°C uncertainty.

4.3. Uncertainty in RSCM temperatures

336 The different RSCM methods are calibrated against several geothermometers. The higher 337 range temperatures method proposed by Beyssac et al. (2002) is mostly based on mineral assemblages. However, vitrinite reflectance, illite crystallinity, chlorite geothermometers were 338 compared with RSCM spectra both in Kouketsu et al. (2014) and Lahfid et al. (2010) studies. 339 Furthermore, thermal modelling, garnet-chlorite geothermometers were also used for 340 calibration in Kouketsu et al. (2014) while Lahfid et al. (2010) used calcite-dolomite 341 thermometry, quartz-chlorite isotopic thermometry and fluid inclusion. These calibrations 342 were carried out on domains that were affected by a single metamorphic event. 343 344 The maximum error (1σ) on R2 ratio obtained by Beyssac et al. (2002) is ± 0.08 corresponding to a ± 36°C intra-sample variability. Adding the uncertainty linked to the 345

calibration of RSCM with reference temperatures, the authors considered that the uncertainty on the estimation of temperature was about $\pm 50^{\circ}$ C.

In the low temperature range, Lahfid et al. (2010) method uses the ratio of Raman peak spectra area, while Kouketsu et al. (2014) method uses the width of the peaks, to derive the paleotemperature. Kouketsu et al. (2014) estimated that the difference in estimated temperature due to the use of the two different methods is of the order of 50°C.

352 4.4. RSCM acquisition

353 The temperatures derived from the RSCM method were obtained with a minimum of 10 to 15 spectra for each sample in order to ensure their validity. Some of the grains of carbonaceous 354 355 matter present within a sediment might originate from the erosion of a sediment that has already been affected by a prior metamorphic event. Since the organic matter only records 356 357 the highest temperature, these inherited grains might yield higher conditions of temperature than their host sediments, if they come from higher-grade material. This case is often 358 suspected when a few CM grains give much higher temperatures than most grains in a 359 360 sample. In such a case, we disregarded the few anomalous grains and estimated the 361 temperature from the statistical distribution concentrating most of the grains (Beyssac et al., 2002). 362

The 122 samples analysed cover the major part of the Montagne Noire southern flank 363 in order to get a general picture of the paleotemperature field reached by the Paleozoic 364 365 series. Almost all lithologies were tested for the RSCM analysis. Several samples were 366 polished at 35µm for optical microscope observation (Fig. 6) but most of the measurements were carried out on 200-300µm-thick thin sections. The thin sections were prepared 367 regardless of the mineral preferred orientation of the carbonaceous matter because of the 368 369 poorly developed carbonization in these low metamorphic grade series (Aoya et al., 2010). The RSCM method reliability depends on the richness of the carbonaceous matter 370 371 enclosed in the sample. In the field, the dark lithologies, i.e. the Early Ordovician and Visean-372 Serpukhovian turbidites, turned out to be the most appropriate lithologies (Fig. 6a, b, d, f). On the contrary, the Early Cambrian Marcory green sandstone, and the Early Cambrian and
Devonian carbonates did not yield sufficient organic matter (Fig. 6c, e), and thus, were
avoided for sampling.

4.5. Sampling Strategy

The main goal of the present study was to establish the paleotemperature field associated with the regional tectonic-metamorphic events. Several hypotheses may be formulated.,

379 The temperature development was coeval with the recumbent folding and thrusting event for this purpose, samples were collected in each structural unit except for the South 380 Minervois and Mts-de-Faugères units. Higher temperatures can be expected at deeper 381 tectonic levels. Furthermore, areas distant from the Axial Zone dome where the metamorphic 382 383 and structural imprint of the Axial Zone dome upon the recumbent folds seems absent 384 (Doublier et al., 2006) represent attractive sites. For instance, in the Ordovician sandstonesiltstone formation of the Upper (Pardailhan) recumbent fold, only the S1 slaty cleavage is 385 386 observed (Fig. 4a).

The thermal effect was related to the late-stage emplacement of the migmatites and 387 plutons in the Axial Zone dome. The strategy to assess the dome effect, as suggested by 388 Wiederer et al. (2002) and Franke et al. (2011) studies, was totally different. In the southern 389 flank, upright folds coeval with a steep S2 cleavage (Fig. 4b) structurally represents this 390 391 event. The occurrence of a N70E striking mineral lineation marked by white mica (sericite) becomes more and more difficult to recognize south of the dome. Consequently, high 392 temperatures should be recorded near the dome while they must decrease away from it, 393 southwards. Nord-south sections were sampled in order to test this hypothesis. 394

Finally, a last possibility can be considered. The axial dome exhumation itself results of a regional horizontal flow that affected the entire crust. Thus, a hot crust could be located below the nappes piles. Beside the dome thermal impact, a high temperature event overprinting the nappe stacking can be expected far from the Axial Zone. 399

400

401 **5. Results**

402 Despite the low-grade metamorphism of the area, both high and low temperature methods 403 were used during this study. The results are well distributed in the 230 to 400°C range 404 covered by both methods (Fig. 5 and 7). Seventy-three temperature values were calculated 405 in the Montagne Noire southern flank. Forty-eight RSCM spectra were processed using 406 Lahfid et al. (2010) method, while 25 spectra were computed through the Beyssac et al. 407 (2002) method (Table. 1). Thirty-six temperatures values were also reassessed by the FWHD-D1-linked thermometer (Kouketsu et al., 2014; Table. 2). The results, plotted in Fig.8, 408 409 reveal a decreasing of maximum temperatures towards the south. It appears that the 410 metamorphic zonation cuts across the boundaries between the tectonic units and the limbs of recumbent folds close the dome and a more homogenous in further distance. In contrast 411 to one of our working hypotheses, the RSCM temperatures do not indicate any thermal gap 412 on both sides of the tectonic contacts. 413

414 **6.** Discussion

6.1. Temperature field acquired by RSCM method

416 In this study, the entire Montagne Noire southern flank has been investigated by the RSCM 417 method. Most of the data set obtained from the sedimentary rocks records a gradient consistent over the whole investigated area, visible in both normal and inverted sedimentary 418 419 series (Fig. 8, 9 cross sections 1, 2, 3, 4, 5). Globally, the temperature decreases from the 420 southern edge of the Axial Zone dome towards the southeast in the Paleozoic sedimentary rocks (Fig. 10). The maximum temperature is located in the micaschists series that form the 421 422 dome envelope with a value close to 400°C. The lowest temperature, about 220°C, was 423 obtained in the Visean turbidites. In the western part of the southern flank, the gradient is

424 less apparent that in the eastern part. It decreases from 400°C to 315°C towards the
425 Cenozoic sedimentary rocks.

The RSCM method was also used within the dome micaschist envelope (Fréville et 426 427 al., 2016). The measured temperatures, ranging between 450°C and 600°C are higher than 428 those obtained in the sedimentary rocks of the southern flank (Fig. 11). The temperature gradient is steeper in the dome northern edge than in the micaschist series of the eastern 429 430 area. Indeed, the deformation is more intense in the dome northern edge, as also shown by 431 the tight arrangement of the metamorphic isograds (Alabouvette et al., 1993; Fréville et al., 432 2016). This is probably a consequence of a brittle shearing that poste-dates the dome 433 exhumation (Thompson and Bard, 1982). The data set obtained via the RSCM method documents a continuous evolution from the dome core southward with a temperature 434 435 decrease from 580°C to less than 300°C (Fig. 11). Since the dome foliation is plunging under the sedimentary series, samples horizontally away from the dome are also vertically distant 436 from the dome. 437

In case of a thermal field predating the emplacement of the recumbent folds, a 438 439 temperature discontinuity should coincide with the contacts between different units. In case of a thermal event coeval with nappe thrusting and tectonic thickening, the gradient should 440 be consistent with the structural position in the tectonic pile. Indeed, it has been shown that 441 442 temperatures recorded in terrigenous sedimentary series deposited in foreland basins 443 involved in fold-and-thrust belts of orogens are related to tectonic thickening. For instance, in 444 the northern Variscan foreland basin of the Ardennes massif, the temperature of 200 to 445 300°C, measured by illite crystallinity, vitrinite reflectance, conodont alteration index, and 446 fluid inclusion methods, has been linked to the burial of sedimentary series below the 447 Variscan thrust front (Fielitz and Mansy, 1999).

The RSCM temperature map of the Montagne Noire southern flank (Fig. 10) is clearly incompatible with these assumptions. Since the isotherms cut across the different fold units irrespective of their contacts, the bulk thermal structure of the Montagne Noire southern flank cannot be linked to the pre-doming thickening stage. Moreover, the RSCM temperature is not 452 correlated with the structural position of the analyzed samples since the highest nappes of
453 the tectonic pile record-temperatures similar to those measured in the para-autochthonous
454 units (Fig. 10). Thus, the link of the temperature field with nappe emplacement, and tectonic
455 burial is unlikely. The measured temperature is also not in agreement with shear heating
456 along major contacts, as that observed in Japan (Mori et al., 2015).

In contrast, there is a clear temperature gradient decreasing with the distance to the 457 458 dome. Whatever the thrusting direction of the nappe, southeastward (Arthaud, 1970) or 459 northeastward (Chardon et al., 2020), the interpretation of a thermal effect related to the Axial Zone dome is therefore preferred to explain the temperature field as suggested by 460 previous authors (Wiederer et al., 2002; Doublier et al., 2005; Franke et al., 2011). At first 461 order, the RSCM method records the HT metamorphic event already recognized close to the 462 Axial Zone over a very broad area including its entire southern flank. It has been shown that 463 RSCM investigations around an intrusion provide high temperatures away from the 464 metamorphic aureole (e. g. Hilchie et al., 2014; Beyssac et al., 2019). 465

However, it is quite unlikely that the heat propagation away from the dome may develop temperatures above 300°C as those observed south of St-Pons at ca 10km in the southernmost part of the upper recumbent fold. We suggest that the temperature field is linked with an hypothetic hot crust below the recumbent folds.

470 6.2. Anomalies in the temperature field

Two temperature anomalies are visible within the N-S temperature gradient (Fig. 8). To the east of the study area, within the Visean-Serpukhovian turbiditic basin, several samples present relatively high temperatures around 360°C, similar to those found near the dome micaschists (Figs. 6, 9 cross section 5). However, in this area, the Axial Zone dome is not exposed. The other temperature anomaly is located in the southern part of the Pardailhan upper recumbent fold (Fig. 9 cross section 3). The gradient is decreasing southward from ca 380°C to 270°C, but at the very end of the exposed Paleozoic series, the temperature increases to reach ca 320°C. Since this area is located about 15 km away from the Axial
Zone dome south margin, it is unlikely that these high temperatures result from the direct
dome thermal influence.

481 Three hypotheses might explain these singularities.

First, the effect of a late fault might be taken in account as it could reorganize the distribution of paleotemperatures. It is especially noticeable for the anomaly within the upper recumbent fold (Pardailhan unit). The N-S striking sinistral fault at the east of the points H25 an H26 Fig. 2) separates high temperature in the west >315°C (H27-H28-H29) from lower temperatures in the east <290°C (H36-H39-H40; Fig. 8; Fig. 9). This effect is also shown by the bending of the 300°C isotherm (Fig. 10).

Second, the circulation of fluids, related to the nappes thickening or the doming may be considered. Several quartz veins, developed during these circulations, appear within the basin and close to the tectonic contacts (Guiraud et al., 1981) or are linked with the micaschist /sedimentary rocks contact as observed in Salsigne (Lescuyer et al., 1993).

Finally, at the regional scale, the district is riddled by Late Carboniferous granitic 492 493 intrusions, such as the Sidobre or Folat plutons (Fig. 2), thus the presence of a hidden pluton below the sedimentary rocks cannot be discarded. Granitic plutons emplaced in an orogen 494 495 external zone and in gneissic domes have been already documented in the Nappe and 496 External zones of Sardinia (Carmignani et al., 1994; Carosi et al., 1998) and more recently 497 by RSCM investigation in Morocco (Delchini et al., 2016). Thus, the existence of several 498 hidden gneiss-granite-migmatite massifs underlying the southern flank stack of recumbent folds and the turbiditic basin must be considered. Moreover, a pegmatite dyke, 2 km north of 499 500 the anomaly in the basin, dated at 282 Ma (Doublier et al., 2015), could be the witness of one of these intrusions. This hypothesis is closely related with the presence of a hot crust which 501 502 can explain the isotherm pattern in the easternmost part of the southern flank, more than 503 15km away from the dome (Fig. 10).

504 6.3. Comparison of the RSCM results with other thermometers

505 6.3.1. General pattern of the temperature field

The illite crystallinity has been widely studied in the eastern part of the Montagne Noire 506 507 southern flank to unravel low-grade metamorphic gradients. In our study area, two previous 508 works (Engel et al., 1980-1981; Doublier et al., 2015) documented a north to south decreasing thermal gradient (Fig. 12). The latter study used the Árkai index (Árkai, 1991; 509 Guggenheim et al., 2002) and the Kübler index (Kübler, 1964) to estimate a relative indicator 510 of temperature. Furthermore, Wiederer et al. (2002) compared the results obtained via the 511 Weber index (Weber 1972) in Engel et al. (1980-1981) to the diagenetic domains defined in 512 513 Frey and Robinson, (1999). In agreement with our own results, these works show the same N-S variations as those documented by the RSCM method. 514

In the same area, the conodont color alteration has been used to estimate the paleotemperature repartition (Wiederer et al., 2002). The method is based on the color of apatite crystals that form the conodonts (Epstein et al., 1977). During a metamorphic event, the crystallinity of organic matter trapped within the apatite grains increases, allowing a possible correlation with the metamorphic grade experienced by the fossil. As shown in Fig. 13, the dome thermal effect is also visible by this method. The intensity of the metamorphism is globally higher to the northwest and decreases away to the southeast (Fig. 13).

A fluid inclusion study of quartz veins (see location on Fig. 11) has been carried out 522 (Guiraud et al., 1981). The analyzed rocks lie along the basal tectonic contact of the 523 524 Pardailhan upper recumbent fold, called "queue de cochon" (pig tail) contact (Gèze, 1949) 525 composed of Devonian limestone boudins surrounded by Ordovician sandstone and pelite. These 1- to 10-cm sized quartz lenses were inferred to be related to fluid circulation coeval 526 527 with the emplacement of the upper recumbent fold (Guiraud et al., 1981). Analyses show a temperature around 275±25°C, which is in agreement with our results at ca 300°C, but 528 529 higher than those provided by illite crystallinity at 200°C (Doublier et al., 2015). This 300°C temperature has been related to the thickening event (Guiraud et al., 1981), however, fluid 530 circulation might have occurred also during the doming. The tectonic contact between the 531

upper and lower recumbent folds has been reworked during doming, as indicated by E-W
striking slickenlines on flat lying surfaces, and N70E striking fibers infilling tension gashes
(Arthaud, 1970; Sauniac, 1980; Harris et al., 1983). Therefore, the deformation observed
along the contact suggests that the ca 300°C temperature might be related to the Axial Zone
dome thermal effect or even to a late event as documented by Aerden (1998) and Franke et
al. (2011).

538 6.3.2. Discrepancy in the low T range between RSCM temperature and other thermometers

For the sake of comparison between the different geothermometers, both conodont 539 540 color alteration index and illite crystallinity data were converted in temperatures using to the 541 tables provided by Merriman and Frey (1999) and Wiederer et al. (2002) for low grade metamorphism and the equation (1) from Zhu et al. (2016) work. Several studies have shown 542 a good correlation between illite crystallinity (IC, estimated from Kübler Index, KI) and 543 544 organic geothermometers such as vitrinite reflectance (VR) (Underwood et al., 1993; 545 Mukoyoshi et al., 2009; Fukuchi et al., 2014). This is true in basinal settings (Baludikay et al., 2018) but also in collision/subduction settings (Rahn et al., 1995). In the Montagne Noire, 546 547 and despite qualitatively convergent trends indicating a north-south gradient from the Axial 548 Zone dome to the sedimentary southern flank, the measured temperatures vary significantly, 549 depending on the method considered.

As example of these differences, the temperature range estimated using the 550 conodont approach is the widest, with temperatures comprised between 75°C and 475°C 551 552 against 85°C to 300°C for IC and 230°C to 400°C for RSCM. Furthermore, for a given location, the results are sometimes different between the two illite crystallinity studies by 553 Engel et al. (1980-1981) and Doublier et al. (2015). This is remarkable at the contact 554 between the lower recumbent fold and the Visean basin where the data from Engel et al. 555 556 (1980-1981) are 50°C higher than those from Doublier et al. (2015). It can be explained by 557 the use of a standardized and more precise procedure in the study of Doublier et al. (2015) while Engel et al. (1980-1981) employed an early version of the method known to have 558

interlaboratory standardization issues (Merriman and Peacor, 1999). Comparing with RSCM 559 560 result, other geothermometers provide temperature lower by at least 70°C. Baludikay et al. (2018) reported a general overestimation of RSCM temperatures (using Kouketsu et al., 561 562 2014 calibration) with respect to other geothermometer in low-grade sediments (T<200°C) from intra-cratonic basins. The discrepancy is here much higher, as illite crystallinity data 563 from Doublier et al. (2015) and RSCM data from this study show a difference up to 175°C 564 (Fig. 14). A first possible source of these differences lies in the large uncertainty in the 565 566 conversion of any of the metamorphic/organic indicators into temperatures. A second source of error is related to the complex geological history of the domain treated here, involving 567 568 several tectonic and heating stages, which might have affected to a variable extent the 569 organic and mineral signals (García-López et al., 2001).

570 Nonetheless, irrespective of the comparison between temperatures themselves, there 571 are also large differences in the temperature gradients derived from the different 572 geothermometers. Between 1 to 6 km away from the dome, the illite crystallinity seems to reach a plateau around 275°C (dashed line in Fig. 14). This plateau could correspond to the 573 574 temperature above which small muscovite or sericite have completely replaced the original clays and would be the upper limit of the method, while carbonaceous matter crystallinity 575 (hence its Raman spectra) continue its evolution at higher temperature. Hence, for 576 comparing the RSCM temperature with illite crystallinity temperature, we shall exclude the 577 578 highest temperature domain in the vicinity of the dome and consider the section at a distance 579 from 5 km to 15 km from the dome. In this distance range, the gap between RSCM temperatures and illite crystallinity temperatures increases with the distance to the dome 580 (Fig. 14). This divergence is apparently irrespective of the RSCM calibration used (Fig. 15a, 581 582 b) or the calibration to convert KI in temperature (T):

-(i) With Merriman and Frey (1999) table: the difference between KI-derived and
 RSCM-derived temperatures varies from ~75°C to ~150°C (Lahfid et al., 2010) and from
 ~80°C to ~120°C (Kouketsu et al., 2014) at 5 km and 14 km from the dome, respectively.

-(ii) With Zhu et al. (2016) equation: the T difference varies from ~75°C to ~175°C
(Lahfid et al., 2010) and from ~80°C to ~150°C (Kouketsu et al., 2014) in the same section.
Therefore, these disparities suggest that organic matter record the dome exhumation thermal
impact on a much broader area than illite crystallinity.

To interpret these discrepancies, one has to consider the nature of the physical and 590 chemical processes involved in carbonaceous matter and illite maturation. Velde and Lanson 591 (1993), Belmar et al. (2002) and Mählmann et al. (2012) suggested that the carbonaceous 592 593 matter records short thermal episodes, such as magmatic intrusions, more easily than clays. 594 This contrasted record could be linked to the fact that organic matter maturation beyond its low-T cracking is principally isochemical, while illite and other clays evolution requires 595 596 chemical reactions involving elemental exchanges (Velde and Lanson, 1993). The samples 597 used to calibrate RSCM with illite crystallinity (Kouketsu et al., 2014) were collected in an accretionary complex in SW Japan (Hara et al., 2013). In this regional metamorphism 598 599 context, both illite and carbonaceous material had sufficient time to maturate and tend 600 towards some "equilibrium" state. With a duration longer than 10Ma, several 601 geothermometers based on various mineralogical assemblages yield the same temperature (Le Bayon et al., 2011; Mullis et al., 2017). 602

Hence, the disparities between illite crystallinity and RSCM in Montagne Noire suggest that the dome emplacement, which controls IC and RSCM evolution, was a relatively short event in terms of heat source. Accordingly, it left a more significant imprint on carbonaceous matter than on illite.

One can note that the much smaller-scale T anomaly in the basin, recorded by RSCM but not by illite crystallinity, can be similarly explained by hidden magmatic bodies that provided a local heat source for a short period of time. We therefore propose that a regional high heatflow linked to a hot crust is present over a long period 330-295Ma, as recorded by Illite and CAI. However, local events resulting from this regional heat such as plutonic intrusion and on a larger scale the migmatization mostly affected the carbonaceous material. Moreover, the high RSCM, illite and condont temperatures observed in the basin far from the eastern terminaison of the dome suggest that a sinistral movement occurred during the dome emplacement. It is then possible that the tectonic origin of the dome emplacement took place during a sinistral transpression as suggested by Chardon et al. (2020). This view is also in agreement with a N-S shortening responsible for the double anticline present within the Axial Zone dome (Matte et al., 1988; Malavieille, 2010).

619 **7. Conclusion**

620 The 75 RSCM measurements acquired in this study combined with the results of several previous studies, derived from different methods, allow us to reconstruct the thermal history 621 of the Montagne Noire southern flank. The RSCM measurements, processed either by Lahfid 622 623 et al. (2010) or Beyssac et al. (2002) methods, cover the whole metamorphic range present 624 in the sedimentary rocks. In agreement with the RSCM results, the conodont color alteration, and the illite crystallinity geothermometers revealed a temperature gradient within the 625 Paleozoic formations decreasing from the Axial Zone dome toward the SE. This RSCM 626 627 temperature gradient represents the effect of the Axial Zone dome overprint on the 628 sedimentary rocks as suggested by previous studies on a smaller area (Engel et al., 1980-1981; Wiederer et al., 2002; Doublier et al., 2015) and also observed in the Montagne Noire 629 northern side (Doublier et al., 2006). 630

Moreover, an underneath heating by the hot crust might also explain the relatively high temperatures, >300°C, recorded by RSCM, IC and CAI in the very southern part of the upper recumbent fold and in the east of the southern flank.

In spite of a common North to South decreasing temperature gradient, the different thermometers present quantitative discrepancies in the estimated temperature. IC temperatures decrease away from the dome to temperatures below 150°C, while RSCM temperatures remain above 250°C. In this case, the fast kinetics of carbonaceous material crystalline evolution, with respect to illite evolution, could then account for the record of the heating event over a larger area by RSCM than by IC. The link between the bulk temperatures and the recumbent folding is not supported by this study. Furthermore, anomalies are observed within the thermal gradient. Since these anomalies are not documented by the IC method, the most likely explanation would be that hidden bodies such as granitic plutons, would have emplaced beneath the Paleozoic sediments after the dome exhumation. In addition, the presence of a hot crust underlying the Paleozoic series cannot be discarded.

Finally, the temperatures obtained in the southern part of the Upper recumbent fold, alike those measured by fluid inclusion study (Guiraud et al., 1981), are compatible with the Axial Zone doming. The RSCM approach suggests that the thermal effects related to the thickening event has been totally overprinted by the dome thermal effect.

As already been investigated in the Rheno-Hercynian zone in the northern Variscan
branch (e.g. Fielitz and Mansy, 1999; Doublier et al., 2012), the study of thermicity in
foreland basins can be extended to other areas within the Variscan belt. For instance,
Mouthoumet massif (Kretschmer et al., 2015), Balearic or southern Sardinia-areas are
suitable places to get a general picture of the Variscan thermicity.

655 Acknowledgments

This study has been funded by the Labex VOLTAIRE, and the Observatoire des Sciences de
l'Univers en Région Centre (OSUC). Jean-Gabriel Badin, Sylvain Janiec, and Ida di Carlo are
thanked for their assistance during the thin section preparation and Raman spectroscopy
measurements. We aslo thank Stanislas Sizaret for his help during sampling session.
Constructive comments by J. Malavieille, R. Carosi, and an anonymous reviewer are deeply
acknowledged.

663 **References**

Aerden DG. 1998. Tectonic evolution of the Montagne Noire and a possible orogenic model 664 for syncollisional exhumation of deep rocks, Variscan belt, France. Tectonics, 17(1), 665 62-79. 666 667 668 Alabouvette B, Arthaud F, Bambier A, Freytet P, Paloc H. 1982. Notice explicative de la carte géologique au 1/ 50 000 de Saint-Chinian, n°1014. Bureau de Recherches 669 Géologiques et Minières, Orléans, France. 1-44. 670 671 Alabouvette B, Demange M, Sauvel C, Vautrelle C. 1993. Notice explicative de la feuille Saint-672 Pons à 1/50 000. Édition du Bureau de Recherches Géologiques et Minières, Orléans. 673 674 Alabouvette B, Demange M, Guérangé-Lozes J, Ambert P. 2003. Notice explicative de la carte 675 géologique au 1:250000 de Montpellier. Bureau de Recherches Géologiques et 676 677 Minières, Orléans, France. 678 Álvaro JJ, Courjault-Radé P, Chauvel JJ, Dabard M P, Debrenne F, Feist R, Pillola GL, Vennin 679 E, Vizcaïno D. 1998. Nouveau découpage stratigraphique des séries cambriennes des 680 nappes de Pardailhan et du Minervois (versant sud de la Montagne Noire, France). 681 Géologie de la France, 2, 3-12. 682 683 Aoya M, Kouketsu Y, Endo S, Shimizu H, Mizukami T, Nakamura D, Wallis S. 2010. Extending 684 the applicability of the Raman carbonaceous - material geothermometer using data 685 from contact metamorphic rocks. Journal of Metamorphic Geology, 28, 895-914 686 687

688	Árkai P. 1991. Chlorite crystallinity: an empirical approach and correlation with illite
689	crystallinity, coal rank and mineral facies as exemplified by Palaeozoic and Mesozoic
690	rocks of northeast Hungary. Journal of Metamorphic Geology, 9, 723–734.
691	
692	Arthaud F. 1970. Etude tectonique et microtectonique comparée de deux domaines
693	hercyniens : les nappes de la Montagne Noire (France) et l'anticlinorium de
694	l'Iglesiente (Sardaigne). Université des Sciences et Techniques du Languedoc, 175
695	pp.
696	
697	Baludikay BK, François C, Sforna MC, Beghin J, Cornet Y, Storme JY, Fagel N, Fontaine F,
698	Littke R, Baudet D, Delvaux D, Javaux EJ. 2018. Raman microspectroscopy, bitumen
699	reflectance and illite crystallinity scale: comparison of different geothermometry
700	methods on fossiliferous Proterozoic sedimentary basins (DR Congo, Mauritania and
701	Australia). International Journal of Coal Geology, 191, 80-94.
702	
703	Bard JP, Rambeloson R. (1973). Métamorphisme plurifacial et sens de variation du degré
704	géothermique durant la tectogenèse polyphasée hercynienne dans la partie orientale
705	de la zone axiale de la Montagne Noire (Massif du Caroux, Sud du Massif Central
706	français). Bulletin de la Société géologique de France, 7(5-6), 579-586.
707	
708	Berger G, Alabouvette B, Guérangé-Lozes J, Demange M, Ambert P. 2001. Carte géol. France
709	(1/250 000), feuille Montpellier (38). Orléans : BRGM. Notice explicative par B.
710	Alabouvette M, Démange J, Guérangé-Lozes P. Ambert (2003), 164 pp.
711	
712	Beyssac O, Goffé B, Chopin C, Rouzaud JN. 2002. Raman spectra of carbonaceous material
713	in metasediments: a new geothermometer. Journal of metamorphic Geology, 20, 859-
714	871.
715	

716	Beyssac O, Pattison DRM, Bourdelle F. 2019. Contrasting degrees of recrystallization of
717	carbonaceous material in the Nelson aureole, British Columbia and Ballachulish
718	aureole, Scotland, with implications for thermometry based on Raman spectroscopy of
719	carbonaceous material, J. Metamorphic Geol., 37, 71-95.
720	
721	Belmar M, Schmidt ST, Frey M, Ferreiro-Mählmann R, Mullis J, Stern W. 2002. Diagenesis,
722	low-grade and contact metamorphism in the Triassic-Jurassic of the Vichuquén-
723	Tilicura and Hualané-Gualleco Basins, Coastal Range of Chile. Schweizeriche
724	Mineralogische und Petrographische Mitteilungen, 82, 375-392.
725	
726	Carmignani L, Carosi R, Di Pisa A, Gattiglio M, Musumeci G, Oggiano G, Carlo Pertusati P.
727	1994. The hercynian chain in Sardinia (Italy). Geodinamica Acta, 7(1), 31-47.
728	
729	Carosi R, Perillo M, Pertusati PC. 1998. Structural evolution of the Southern Sulcis
730	Metamorphic Complex (SW Sardinia, Italy). C. R. Acad. Sci. Paris, Sciences de la terre
731	et des planètes/Earth & Planetary Sciences, 326, 505-512, Paris, Elsevier Gauthier-
732	Villars, 1998.
733	
734	Chardon D, Aretz M, Roques D. 2020. Reappraisal of Variscan tectonics in the southern
735	French Massif Central. Tectonophysics, 787.
736	
737	Cocherie A, Baudin T, Autran A, Guerrot C, Fanning CM, Laumonier B. 2005. U-Pb zircon (ID-
738	TIMS and SHRIMP) evidence for the early Ordovician intrusion of metagranites in the
739	late Proterozoic Canaveilles Group of the Pyrenees, and the Montagne Noire (France).
740	Bull. Société géologique de France, 176, 269-282.
741	

742	Cózar P, Izart A, Vachard D, Coronado I. 2017 A mid-Tournaisian-late Viséan carbonate ramp
743	reconstructed from nappes and olistolites in the southern Montagne Noire (France).
744	Sedimentary Geology, 358, 148-175.
745	
746	Delchini S, Lahfid A, Plunder A, Michard A. 2016. Applicability of the RSCM geothermometry
747	approach in a complex tectono-metamorphic context: The Jebilet massif case study
748	(Variscan Belt, Morocco). Lithos, 256, 1-12.
749	
750	Demange M. 1982. Etude géologique du massif de l'Agoût, Montagne Noire, France. Thèse
751	d'Etat, Paris 6, 1050p.
752	Demange M. 1993. Que signifie la faille des Monts de Lacaune (Montagne Noire, France) ?
753	Implications quant au problème de la patrie des nappes. Comptes rendus de
754	l'Académie des sciences. Série 2, 317, 411-418.
755	
756	Doublier MP, Potel S, Wemmer K. 2006. Age and grade of metamorphism in the eastern Monts
757	de Lacaune-implications for the collisional accretion in Variscan externides (French
758	Massif Central). Geodinamica Acta, 19, 391-407.
759	
760	Doublier MP, Potel S, Franke W, Roache T. 2012. Very low-grade metamorphism of
761	Rhenohercynian allochthons (Variscides, Germany): facts and tectonic
762	consequences. International J. Earth Sciences, 101, 1229–1252.
763	
764	Doublier MP, Potel S, Wemmer K. 2015. The tectono-metamorphic evolution of the very low-
765	grade hangingwall constrains two-stage gneiss dome formation in the Montagne Noire
766	(Southern France). Journal of Metamorphic Geology, 33, 71-89.
767	
768	Dunoyer de Segonzac G, Ferrero J, Kübler B. 1968. Sur la cristallinité de l'illite dans la
769	diagenèse et l'anchimétamorphisme. Sedimentology, 10, 137-143

770	
771	Echtler H, Malavieille J. 1990. Extensional tectonics, basement uplift and Stephano-Permian
772	collapse basin in a late Variscan metamorphic core complex (Montagne Noire,
773	Southern Massif Central). Tectonophysics, 177, 125-138.
774	
775	Engel W, Feist R, Franke W. 1978. Synorogenic gravitational transport in the Carboniferous of
776	the Montagne Noire (S-France). Zeitschrift der deutschen geologischen Gesellschaft,
777	461-472.
778	Engel W, Feist R, Franke W. 1980-81. Le Carbonifère anté-stéphanien de la Montagne Noire:
779	rapports entre mise en place des nappes et sédimentation. Bull. B.R.G.M. 2, 341-389.
780	
781	Epstein AG, Epstein JB, Harris LD. 1977. Conodont color alteration: an index to organic
782	metamorphism. Geological Survey of America, Professional paper 995, 1-27.
783	
784	Faure M, Cottereau N. 1988. Données cinématiques sur la mise en place du dôme migmatique
785	carbonifère moyen de la zone axiale, de la Montagne Noire (Massif Central, France).
786	Comptes rendus Acad. Sci. Série 2, 307, 1787-1794.
787	
788	Faure M. 1995. Late orogenic carboniferous extensions in the Variscan French Massif Central.
789	Tectonics, 14, 132-153.
790	
791	Faure M, Bé Mézème E, Duguet M, Cartier C, Talbot JY. 2005. Paleozoic tectonic evolution of
792	medio-Europa from the example of the French Massif Central and Massif Armoricain.
793	Journal of the virtual Explorer, 19, 1-25.
794	
795	Faure M, Lardeaux JM, Ledru P. 2009. A review of the pre-Permian geology of the Variscan
796	French Massif Central. Comptes Rendus Geoscience, 341, 202-213.
797	

Faure M, Cocherie A. Bé Mézème E, Charles N, Rossi P. 2010. Middle Carboniferous crustal
melting in the Variscan belt: New insights from U–Th–Pb tot monazite and U–PB zircon
ages of the Montagne Noire Axial Zone (southern French Massif Central). Gondwana
Res. 18, 653–673.

802

Faure M, Cocherie A, Gaché J, Esnault C, Guerrot C, Rossi P, Lin W, Qiuli L. 2014. Middle
Carboniferous intracontinental subduction in the Outer Zone of the Variscan belt
(Montagne Noire Axial Zone, French Massif Central): multimethod geochronological
approach of polyphase metamorphism. Multimethod geochronological approach of
polyphase metamorphism. Geological Society, London, Special Publications, 405(1),
289-311.

809

Faure M, Li XH, Lin W. 2017. The northwest-directed "Bretonian phase" in the French Variscan
Belt (Massif Central and Massif Armoricain): a consequence of the Early Carboniferous
Gondwana-Laurussia collision. C. R Géoscience, 349, 126-136.

813

Feist R, Galtier J. 1985. Découverte de flores d'âge namurien probable dans le flysch à
olistolites de Cabrières (Hérault). Implication sur la durée de la sédimentation
synorogénique dans la Montagne Noire (France méridionale). Comptes Rendus Acad.
Sciences. Série 2, 300, 207-212.

818

Fielitz W, Mansy JL. 1999. Pre-and synorogenic burial metamorphism in the Ardenne and
neighbouring areas (Rhenohercynian zone, central European Variscides).
Tectonophysics, 309, 227-256.

822

Franke W, Doublier MP, Klama K, Potel S, Wemmer K. 2011. Hot metamorphic core complex
in a cold foreland. International Journal of Earth Sciences, 100, 753-785.

826	Fréville K, Cenki-Tok B, Trap P, Rabin M, Leyreloup A, Régnier JL, Whitney DL. 2016. Thermal
827	interaction of middle and upper crust during gneiss dome formation: Example from the
828	Montagne Noire (French Massif Central). Journal of Metamorphic Geology, 34, 447-
829	462.
830	
831	Frey M. Very low grade metamorphism of clastic sedimentary rock. In : Frey M, Ed. Low
832	temperature metamorphism. Chapman and Hall, New York, 1987, pp. 9-58.
833	
834	Frey, M., Robinson, D., (eds) 1999, Low-grade metamorphism. Blackwell, 313 pp.
835	
836	Fukuchi R, Fujimoto K, Kameda J, Hamahashi M, Yamaguchi A, Kimura G, Hamada Y,
837	Hashimoto Y, Kitamura Y, Saito S. 2014. Changes in illite crystallinity within an ancient
838	tectonic boundary thrust caused by thermal, mechanical, and hydrothermal effects: an
839	example from the Nobeoka Thrust, southwest Japan, Earth, Planets and Space,
840	66:116, 1-12.
841	
842	García-López S, Bastida F, Aller J, Sanz-López J. 2001. Geothermal palaeogradients and
843	metamorphic zonation from the conodont colour alteration index (CAI). Terra Nova,
844	13(2), 79-83.
845	
846	Gerya T, Stöckhert B. 2002. Exhumation rates of high-pressure metamorphic rocks in
847	subduction channels: The effect of Rheology. Geophysical Research Letters, 29, 1261-
848	1264.
849	
850	Gèze B. 1949. Etude géologique de la Montagne Noire et des Cévennes méridionales.
851	Soc.Géol. Fr. Mém. 62, 1–125.
852	

853	Guggenheim S, Bain DC, Bergaya F, Brigatti MF, Drits VA, Eberl DD, Formoso MLL, Galán
854	E, Merriman RJ, Peacor DR, Stanjek H, Watanabe T. 2002. Report of the Association
855	Internationale pour l'Etude des Argiles (AIPEA) Nomenclature Committee for 2001:
856	Order, disorder and crystallinity in phyllosilicates and the use of the 'Crystallinity
857	Index'. Clay Minerals, 37, 389-393.
858	
859	Guiraud M, Sauniac S, Burg JP. 1981. Précision sur les conditions Pression-Température lors
860	de la mise en place de la nappe de Pardailhan (Montagne Noire), par la détermination
861	des inclusions fluides. Comptes-rendus de l'Académie des Sciences, Série II, 292, 229-
862	232.
863	
864	Hamet J, Allègre CJ. 1976. Hercynian orogeny in the Montagne Noire (France): Application
865	of Rb87-Sr87 systematics. Geological Society of America Bulletin, 87, 1429-1442.
866	
867	Hara H, Kurihara T, Mori H. 2013. Tectono-stratigraphy and low-grade metamorphism of
868	Late Permian and Early Jurassic accretionary complexes within the Kurosegawa belt,
869	Southwest Japan: Implications for mechanisms of crustal displacement within active
870	continental margin. Tectonophysics, 592, 80-93.
871	
872	Harris LB, Burg JP, Sauniac S. 1983. Strain distribution within the Pardailhan nappe
873	(Montagne Noire, France), and structure of its basal thrust zone: implications for
874	events associated with nappe emplacement. J. Struct. Geol. 5, 431-440.
875	
876	Hilchie LJ, Jamieson RA. 2014. Graphite thermometry in a low-pressure contact aureole,
877	Halifax, Nova Scotia. Lithos, 208, 21-33.
878	

879	Kouketsu Y, Mizukami T, Mori H, Endo S, Aoya M, Hara H, Nakamura D, Wallis S. 2014. A
880	new approach to develop the Raman carbonaceous material geothermometer for low-
881	grade metamorphism using peak width. Island Arc, 23, 33–50.
882	
883	Kretschmer S, Franke W, Wemmer K, Köngigshof P, Gerdes A. 2015. Tectono-metamorphic
884	evolution of the Mouthoumet Massif (Variscides, S-France): Interference of orogenic
885	accretion and crustal extension: Géologie de la France, 1, 78.
886	
887	Kübler B. 1964. Les argiles, indicateurs de métamorphisme. Revue Institut de la Française
888	de Pétrole, 19, 1093–1112.
889	
890	Kübler B. 1968. Evaluation quantitative du metamorphism par la cristallinite de l'Illite. Bull.
891	Centre Rech. Pau-SNPA, 2, 385-397.
892	
893	Lahfid A, Beyssac O, Deville E, Negro F, Chopin C, Goffé B. 2010. Evolution of the Raman
894	spectrum of carbonaceous material in low-grade metasediments of the Glarus Alps
895	(Switzerland). Terra Nova, 22, 354–360.
896	
897	Le Bayon R, Brey GP, Ernst WG, Mählmann RF. 2011. Experimental kinetic study of organic
898	matter maturation: time and pressure effects on vitrinite reflectance at 400 C. Organic
899	Geochemistry, 42, 340-355.
900	
901	Lescuyer JL, Bouchot V, Cassard D, Feybesse JL, Marcoux E, Moine B, Piantone P, Teygey
902	M, Tollon F. Le gisement aurifère de Salsigne (Aude, France) : Une concentration
903	syntectonique tardivarisque dans les sédiments détritiques et carbonatés de la
904	Montagne Noire: Chronique de la Recherche Minière, n° 512, (1993) pp 3–73.
905	

Mählmann RF. 2001. Correlation of very low-grade data to calibrate a thermal maturity model 906 in a nappe tectonic setting, a case study from the Alps. Tectonophysics, 334, 1-33. 907 908 Mählmann RF, Bozkaya Ö, Potel S, Le Bayon R, Šegvić B, Nieto F. 2012. The pioneer work 909 of Bernard Kübler and Martin Frey in very low-grade metamorphic terranes: paleo-910 geothermal potential of variation in Kübler-Index/organic matter reflectance 911 correlations. A review. Swiss Journal of Geosciences, 105(2), 121-152. 912 913 Malavieille J. 2010. Impact of erosion, sedimentation and structural heritage on the structure 914 and kinematics of orogenic wedges: analog models and case studies. GSA Today, 20, 915 916 4–10. 917 Mattauer M, Laurent P, Matte P. 1996. Plissement hercynien synschisteux post-nappe et 918 étirement subhorizontal dans le versant sud de la Montagne noire (Sud du Massif 919 920 central, France). Comptes rendus de l'Acad. des Sciences. Série 2, 322, 309-315. 921 922 Matte P, Lancelot J, Mattauer M. 1998. La zone axiale hercynienne de la Montagne Noire n'est pas un "metamorphic core complex" extensif mais un anticlinal post-nappe à cœur 923 anatectique. Geodinamica Acta, 11, 13-22. 924 925 Merriman R, Frey M. Patterns of very low-grade metamorphism in metapelitic rocks. In : 926 927 Robinson D, Frey M, Eds. Low-grade metamorphism, 1999, pp. 61-108. 928 Merriman R, Peacor DR. Very low-grade metapelites: mineralogy, microfabrics and measuring 929 reaction progress. In : Robinson D, Frey, M Eds. Low-grade metamorphism, 1999, pp. 930 10-60. 931 932

933	Mori H, Wallis S, Fujimoto K, Shigematsu N. 2015. Recognition of shear heating on a long-
934	lived major fault using Raman carbonaceous material thermometry: implications for
935	strength and displacement history of the MTL, SW Japan. Island Arc, 24, 425-446.
936	
937	Mori H, Mori N, Wallis S, Westaway R, Annen C. 2017. The importance of heating duration
938	for Raman CM thermometry: evidence from contact metamorphism around the Great
939	Whin Sill intrusion, UK. Journal of Metamorphic Geology, 35, 165-180.
940	
941	Mukoyoshi H, Hirono T, Hara H, Sekine K, Tsuchiya N, Sakaguchi A, Soh W. 2009. Style of
942	fluid flow and deformation in and around an ancient out-of-sequence thrust: An example
943	from the Nobeoka Tectonic Line in the Shimanto accretionary complex, southwest
944	Japan, Island Arc, 18, 333-351.
945	
946	Mullis J, Mählmann RF, Wolf M. 2017. Fluid inclusion microthermometry to calibrate vitrinite
947	reflectance (between 50 and 270° C), illite Kübler-Index data and the
948	diagenesis/anchizone boundary in the external part of the Central Alps. Applied Clay
949	Science, 143, 307-319.
950	
951	Pitra P, Poujol M, Van Den Driessche J, Poilvet JC, Paquette JL. (2012). Early Permian
952	extensional shearing of an ordovician granite: The saint-eutrope "c/s-like" orthogneiss
953	(montagne noire, French massif central). Comptes Rendus Géoscience, 344(8), 377-
954	384.
955	
956	Poilvet JC, Poujol M, Pitra P, Van Den Driessche J, Paquette JL. (2011). The Montalet granite,
957	Montagne Noire, France: An Early Permian syn-extensional pluton as evidenced by
958	new U-Th-Pb data on zircon and monazite. Comptes Rendus Géoscience, 343(7), 454-
959	461.

Poty É, Aretz M, Barchy L. 2002, Stratigraphie et sédimentologie des «calcaires à Productus»
du Carbonifère inférieur de la Montagne noire (Massif central, France). Comptes
Rendus Geoscience, 334, 843-848.

963

Rahl JM, Anderson KM, Brandon MT, Fassoulas C. 2005. Raman spectroscopic carbonaceous
 material thermometry of low-grade metamorphic rocks: calibration and application to
 tectonic exhumation in Crete, Greece. Earth and Planetary Science Letters, 240, 339 354.

968

Rahn M, Mullis J, Erdelbrock K, Frey M. 1995. Alpine metamorphism in the north Helvetic
flysch of the Glarus-Alps, Switzerland. Eclogae Geologicae Helvetiae, 88, 157-178.

971

Proger F, Respaut JP, Brunel M, Matte P, Paquette JL. 2004. Première datation U-Pb des
orthogneiss œillés de la zone axiale de la Montagne noire (Sud du Massif central):
nouveaux témoins du magmatisme ordovicien dans la chaîne Varisque. Comptes
Rendus Geoscience, 336, 19-28.

976

877 Roger F, Teyssier C, Respaut JP, Rey PF, Jolivet M, Whitney DL, Brunel M. 2015. Timing of
878 formation and exhumation of the Montagne Noire double dome, French Massif Central.
879 Tectonophysics, 640, 53-69.

980

981 Sadezky A, Muckenhube H, Grothe H, Niessner R, Pöschl U. 2005. Raman

982 microspectroscopy of soot and related carbonaceous materials: spectral analysis and 983 structural information. Carbon, 43, 1731-1742.

984

Sauniac S. 1980. Utilisation des exsudats de quartz come critères de reconnaissance d'un
régime cisaillant: exemple de la base de la nappe de Pardailhan (versant sud de la
Montagne Noire). Rev. Géol. Dyn. Geog. Phys, 22, 177-186.

988	
989	Schuiling R. 1960. Le dôme gneissique de l'Agoût (Tarn et Hérault). Mémoire de la
990	Société Géologique de France, 91.
991	
992	Soula JC, Debat P, Brusset S, Bessière G, Christophoul F, Déramond J. 2001. Thrust-
993	related, diapiric, and extensional doming in a frontal orogenic wedge: example of the
994	Montagne Noire, Southern French Hercynian Belt. Journal of Structural Geology, 23,
995	1677-1699.
996	
997	Thompson PH, Bard JP. 1982. Isograds and mineral assemblages in the eastern Axial Zone,
998	Montagne Noire (France): implications for temperature gradients and P–T history.
999	Canadian Journal of Earth Sciences, 19, 129-143.
1000	
1001	Trap P, Roger F, Cenki-Tok B, Paquette JL. 2017. Timing and duration of partial melting and
1002	magmatism in the Variscan Montagne Noire gneiss dome (French Massif Central).
1003	International Journal of Earth Sciences, 106, 453-476.
1004	
1005	Underwood M, Byrne T, Hibbard JP, DiTullio L. A comparison among organic and inorganic
1006	indicators of diagenesis and low-temperature metamorphism, Tertiary Shimanto Belt,
1007	Shikoku, Japan. In : Underwood M, Ed. Thermal evolution of the Tertiary Shimanto
1008	Belt, southwest Japan: an example of ridge-trench interaction, 1993, pp. 45-61.
1009	
1010	Vachard D, Aretz M. 2004. Biostratigraphical precisions on the Early Serpukhovian (Late
1011	Mississippian), by means of a carbonate algal microflora (cyanobacteria, algae and
1012	pseudo-algae) from La Serre (Montagne Noire, France). Geobios, 37, 643-666.
1013	

1014	Vachard D, Izart A, Cózar P. 2017. Mississippian (middle Tournaisian-late Serpukhovian)
1015	lithostratigraphic and tectonosedimentary units of the southeastern Montagne Noire
1016	(Hérault, France). Géologie de la France, 1, 47-88
1017	
1018	Van Den Driessche J, Brun JP. (1989). Un modèle cinématique de l'extension paléozoïque
1019	supérieur dans le Sud du Massif Central. Comptes rendus de l'Académie des
1020	sciences. Série 2, Mécanique, Physique, Chimie, Sciences de l'univers, Sciences de
1021	la Terre, 309(16), 1607-1613.
1022	
1023	Van Den Driessche J, Brun JP. 1992. Tectonic evolution of the Montagne Noire (French Massif
1024	Central): a model of extensional gneiss dome. Geodinamica Acta, 5, 85-97.
1025	
1026	Velde B, Lanson B. 1993. Comparison of I/S transformation and maturity of organic matter at
1027	elevated temperatures. Clays and Clay Minerals, 41, 178-178.
1028	
1029	Vizcaïno D, Álvaro JJ. 2001. The Cambrian and lower Ordovician of the southern Montagne
1030	Noire: a synthesis for the beginning of the new century. Ann. Soc. géol.Nord (2° sér.)
1031	8, 185–242.
1032	
1033	Wada H, Tomita T, Matsuura K, Tuchi K, Ito M, Morikiyo T. 1994. Graphitization of
1034	carbonaceous matter during metamorphism with references to carbonate and pelitic
1035	rocks of contact and regional metamorphisms, Japan. Contributions to Mineralogy
1036	and Petrology, 118, 217-228.
1037	
1038	Weber K. 1972. Notes on the determination of illite crystallinity. Neues Jahrbuch für
1039	Mineralogie, Monatshefte, 6, 267-276.
1040	

1041	Whitney DL, Roger F, Teyssier C, Rey PF, Respaut JP. 2015. Syn-collapse eclogite
1042	metamorphism and exhumation of deep crust in a migmatite dome: the P–T–t record
1043	of the youngest Variscan eclogite (Montagne Noire, French Massif Central). Earth
1044	and Planetary Science Letters, 430, 224-234.
1045	
1046	Wiederer U, Koenigshof P, Feist R, Franke W, Doublier MP. (2002). Low-grade metamorphism
1047	in the Montagne Noire (S-France): Conodont Alteration Index (CAI) in Palaeozoic
1048	carbonates and implications for the exhumation of a hot metamorphic core complex.
1049	Schweizerische Mineralogische und Petrographische Mitteilungen, 82, 393–407.
1050	
1051	Zhu C, Rao S, Hu S. 2016. Application of illite crystallinity for paleo - temperature
1052	reconstruction: A case study in the western Sichuan basin, SW China. Carpathian
1053	Journal of Earth and Environmental Sciences, 11(2), 599-608.
1054	
1055	

1056 Figure captions

Fig. 1. Variscan massifs map with location of the Visean-Serpukhovian migmatites. The red rectangle represents the study area (modified from Faure et al., 2010).

1059 Fig. 2. Simplified structural map of the Montagne Noire showing the granite-migmatite dome

1060 of the Axial Zone, and the sedimentary northern and southern flanks (modified from Gèze,

1061 1949; Arthaud, 1970; Faure et al., 2010). The recumbent folding in the south area is coeval

1062 with the sedimentation in the Visean-Serpukhovian basin. The stratigraphy presents a

1063 reverse order in most part of these series. The figure was made from the vectorised map

1064 250 000 of Montpellier by the BRGM (Berger et al., 2001).

Fig. 3. Paleozoic stratigraphic log, with the abundance in organic matter shown on the left
column (modified from Arthaud, 1970; Engel et al. 1980-1981; Álvaro and Vizcaïno, 1998;
and Vizcaïno and Álvaro, 2001).

Fig. 4. Pictures showing the two main deformation phases in the Montagne Noire southern flank. (a). Steep inverted limb of a S-verging fold coeval with the formation of the km-scale recumbent folds (F1), near Ferrals-les-Montagnes. Sample H52 was picked up for RSCM study. (b). Upright fold with axial planar cleavage (S2) coeval with the Axial Zone doming (F2) and location of the sample H41, close to S^t-Pons. The folded layers are S0-S1 surface where (S1) is coeval with recumbent folding (F1).

Fig. 5. Peakfitting of samples H52 (a), and H75 (b) by low temperature (Lahfid et al., 2010)
Sand high temperature method (Beyssac et al., 2002), respectively.

Fig. 6. Thin sections observed with optical microscope in polarized non-analysed light showing the lithology diversity, (a). Mudstone in the Visean Turbidite (sample G8), (b). Siltstone in the Early Ordovician rocks (sample G1), (c). Early Cambrian green sandstone (Marcory formation, sample H32), (d). Mudstone in the Early Ordovician rocks (sample G2), (e). Middle Cambrian fine-grained sandstone (sample H22), (f). Early Ordovician mudstone (sample G3). Fig. 7. Graph showing the repartition of our data with respect to the two methods used. The
right column shows the evolution of the spectra aspect and the deconvolution chosen with
D1 band (blue), D2 band (red), D3 band (yellow), D4 band (orange) and G band (green).

Fig. 8. Geological map of the Montagne Noire with the Tmax measured in the southern flank.
Totally, 74 samples from the recumbent folds and turbiditic basin have been analysed by the
RSCM method in the present study.

Fig. 9. Projections of the analyzed samples on five cross sections (located in figure 8) of the
southern flank of the Montagne Noire. Temperatures are globally increasing toward the Axial
Zone dome.

Fig. 10. Temperature map determined by the RSCM Tmax results of this study. The isotherms crosscut the tectonic units of the southern flank. Roughly, the temperature decreases from the NW to the SE away from the Axial Zone dome. The isotherms cut the micaschists between Salsigne and St-Pons because of a lack of data.

Fig. 11. Comparison of temperatures in °C acquired via RSCM method in this study and
Fréville et al. (2015) and Guiraud et al. (1981) temperature from fluid inclusion.

Fig. 12. Illite crystallinity data from previous studies (Engel et al., 1981; Doublier et al., 2015) based on the crystalline organization of illite analysed via X-ray diffraction converted from the Kübler index into °C (Merriman and Frey, 1999; Zhu et al., 2016). The Fig. 14 is based on the data from the red rectangle.

Fig. 13. Conodont index alteration method from previous study (Wiederer et al., 2002). The
colour of apatite crystals indicates its metamorphic degree. Wiederer et al. (2002) have
studied more than three hundred conodonts.

Fig. 14. Comparison of Lahfid et al. (2010) and Kouketsu et al. (2014) thermometers, arrowsrepresent the major linear trends.

- 1105 **Fig. 15.** (a). Plot of the variation of measured RSCM temperatures using different methods
- 1106 with respect to the distance to the dome. (b). Histograms of frequencies of the full width at
- 1107 half maximum of the band D1 (FWHM) acquired using the Kouketsu et al. (2014) method.

1108 Table captions

- 1109 **Table. 1.** Tmax results obtained by RSCM according to Beyssac et al. (2002) and Lahfid et al.
- 1110 (2010) methods
- 1111 **Table. 2.** Tmax results obtained by RSCM according to Kouketsu et al. (2014) method