



HAL
open science

Sulfur monoxide dimer chemistry as a possible source of polysulfur in the upper atmosphere of Venus

Joseph P. Pinto, Jiazheng Li, Franklin P. Mills, Emmanuel Marcq, Daria Evdokimova, Denis Belyaev, Yuk L. Yung

► To cite this version:

Joseph P. Pinto, Jiazheng Li, Franklin P. Mills, Emmanuel Marcq, Daria Evdokimova, et al.. Sulfur monoxide dimer chemistry as a possible source of polysulfur in the upper atmosphere of Venus. *Nature Communications*, 2021, 12 (1), pp.175. 10.1038/s41467-020-20451-2 . insu-03108861

HAL Id: insu-03108861

<https://insu.hal.science/insu-03108861>

Submitted on 13 Jan 2021







HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Sulfur monoxide dimer chemistry as a possible source of polysulfur in the upper atmosphere of Venus

Joseph P. Pinto¹, Jiazheng Li ^{2✉}, Franklin P. Mills ^{3,4}, Emmanuel Marcq ⁵, Daria Evdokimova ^{5,6}, Denis Belyaev ⁶ & Yuk L. Yung ^{2,7}

The abundance of SO dimers (SO)₂ in the upper atmosphere of Venus and their implications for the enigmatic ultraviolet absorption has been investigated in several studies over the past few years. However, the photochemistry of sulfur species in the upper atmosphere of Venus is still not well understood and the identity of the missing ultraviolet absorber(s) remains unknown. Here we update an existing photochemical model of Venus' upper atmosphere by including the photochemistry of SO dimers. Although the spectral absorption profile of SO dimers fits the unknown absorber, their abundance is found to be too low for them to contribute significantly to the absorption. It is more likely that their photolysis and/or reaction products could contribute more substantively. Reactions of SO dimers are found to be important sources of S₂O, and possibly higher order S_nO species and polysulfur, S_n. All of these species absorb in the critical ultraviolet region and are expected to be found in both the aerosol and gas phase, indicating that in-situ high resolution aerosol mass spectrometry might be a useful technique for identifying the ultraviolet absorber on Venus.

¹University of North Carolina at Chapel Hill, Chapel Hill, NC, USA. ²Division of Geological and Planetary Science, California Institute of Technology, Pasadena, CA, USA. ³Australian National University, Canberra, ACT, Australia. ⁴Space Science Institute, Boulder, CO, USA. ⁵LATMOS/CNRS/Sorbonne Université/UVSQ, Paris, France. ⁶Space Research Institute of the Russian Academy of Sciences (IKI), Moscow, Russia. ⁷Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA. ✉email: jjazheng@caltech.edu

The photochemistry of sulfur species in the upper atmosphere of Venus is not well understood and is poorly quantified. The known parent molecules upwelling from the deep atmosphere are SO₂ and OCS. Near and above the cloud tops, these molecules are photolyzed, leading to the production of S_n ($n = 1-7$), S_x ($x = 8+$), SO, SO₃, and H₂SO₄¹. Significant amounts of sulfur aerosol were predicted in the lower and middle cloud layers by ref. ² and also ref. ³, which assumes a particle radius of 0.5 μm. Ref. ⁴ suggested that the dimers of sulfur monoxide (SO), (SO)₂, formed from the self-recombination of SO, could contribute substantially to the ultraviolet (UV) absorption found on Venus. In addition, production of S_nO could also occur, as suggested by ref. ⁵.

There are three main isomeric forms of the SO dimer. Reference ⁴ concluded that SO recombination yields ~49% as *cis*-OSSO, ~49% as *trans*-OSSO isomers and no more than 2% as the trigonal isomer (S=SO₂) which is the lowest energy isomer. Reference ⁶ considered only the formation of the trigonal form, photolysis of which leads to the formation of S and SO₂. Reference ³, based on photochemical modeling, suggested that the abundance of OSSO is too low by two orders of magnitude to explain the near UV absorption of Venus. Photolysis of SO dimer is expected to yield mainly two SO radicals⁴, but we will consider other possible reaction paths. Reference ⁷ found evidence for the formation of the *cis*-OSSO, *trans*-OSSO, and C1-S₂O₂ (cyclic OS (=O)S) dimers, along with several other isomeric forms in much lower abundance, following condensation into a solid matrix with N₂ and subsequent irradiation. Production of S₂ and SO₂ was also observed, but the trigonal (S=SO₂) dimer form was beneath detection in the observations of ref. ⁷. S₂ was formed initially from dissociation of the ethylene episulfoxide used as source of SO; further irradiation at 365 nm depleted the 370 nm band, which they ascribed to destruction of *syn*-OSSO (*cis*-OSSO). This depletion occurred simultaneously with an increase in absorption in the 287 nm band of S₂. Formation of S₂ requires only 1 eV above that for dissociation of dimer back to SO and could account for the very strong absorption and characteristic vibrational structure seen at

287 nm in their experiments. We suggest that the photochemistry of (SO)₂ could provide a significant source of S₂ on Venus based on the results of ref. ⁷, leading ultimately to the production of polysulfur, a candidate for the unidentified UV absorber on Venus¹ and possibly also to production of polysulfur oxides⁵.

In this work, we use a photochemical model to assess the contributions of the proposed dimer chemistry to the production of polysulfur. We find that the SO dimers are more likely to be important intermediaries in the formation of more complex S species that could be responsible for the UV absorption. Our model chemistry is also applied to interpret data from the European Space Agency's Venus Express (VEx) mission that was recently processed and analyzed and to provide input to plans for future missions.

Results

Figure 1 presents the mixing ratios of SO₂, OCS, and SO in the middle atmosphere using an SO₂ mixing ratio of 3.0 ppm, which we have adopted for our standard model at the lower boundary (58 km), based on ref. ⁸ and 0.3 ppm for OCS, based on ref. ⁹. The profiles of the rates of the reactions in Supplementary Table 1 are shown in Figure Supplementary Fig. 1. Comparison of the model with several observational datasets at higher altitudes is also shown in Fig. 1. As can be seen (curve a), the model is in reasonable agreement with SO₂ mixing ratios at 70 km during the first four years (2006 through 2009) of VEx measurements¹⁰. Calculated values are slightly higher than the interquartile range and the mean of VEx measurements within ±20° latitude; but the data include a large number of spikes, as indicated by the difference between the median and mean observations. As an indication of the model sensitivity to the choice of SO₂ at the lower boundary, we also show the model-measurement comparison using an SO₂ mixing ratio of 0.3 ppm at the lower boundary in Supplementary Fig. 2. This lower boundary condition could also be seen as reflecting in some ways the lower cloud top SO₂ mixing ratios seen from the start of 2010 through 2014. Calculated values are within the interquartile range and are quite close to the mean of VEx measurements within ±20° latitude. As is the case for the earlier period in the record, the mean differs significantly from the median because of the existence of a large number of concentration spikes, which are roughly three orders of magnitude greater than median levels.

Modeled SO₂ (thin solid blue line) is also consistent with Hubble Space Telescope (HST) data (dashes b) of ref. ¹¹ and with the upper limit derived from ground-based submillimeter observations (dashes d) of ref. ¹² from 85 to 100 km and is in reasonable agreement with Spectroscopy for Investigation of Characteristics of the Atmosphere of Venus/Solar Occultation at Infrared (SPICAV/SOIR) measurements from 90 to 100 km (curve c) of ref. ¹³ collected from 2006 through 2014. Note that the upper level SO₂ mixing ratios (curve c) obtained by ref. ¹³ from 2006 through 2009 tend to be higher than those from 2010 through 2014 in a manner similar to SO₂ observed at 70 km by ref. ¹⁰. Both ground-based submillimeter spectroscopy¹² and solar and stellar occultations¹³ with one standard deviation (curve c and e) show higher SO₂ mixing ratio at high altitudes ($z > \sim 85$ km) compared to lower altitudes. There are two possible sources of SO₂ to be considered at these altitudes, meteorite ablation and photolysis of H₂SO₄. We have included ablation of meteoritic material, which is a very minor source of S to the upper atmosphere of Venus based on current estimates of meteoric input¹⁴ and S content¹⁵. The calculated profile of H₂SO₄ is within limits obtained by ref. ¹⁶. Photolysis of H₂SO₄ is also included based on calculations of its absorption spectrum by ref. ¹⁷, which provided evidence for the existence of a long-wavelength tail that




Fig. 1 Modeled profiles of SO₂, OCS, and SO. Comparison of modeled profiles of SO₂, OCS and SO with observations for SO₂ = 3.0 ppm and OCS = 0.3 ppm at 58 km, the lower boundary of the model. Model profiles are shown as thin solid lines: (blue) SO₂, (green) OCS, (red) SO. Data sources: **a** SO₂, interquartile range from 2006 through 2009, diamond shows mean of distribution, ref. ¹⁰; **b** SO₂, ref. ¹¹; **c** SO₂, ref. ¹³ solar occultation with 1-σ error bars; **d** SO₂, ref. ¹²; **e** SO₂, ref. ¹³ stellar occultation with 1-σ error bars; **f** OCS, ref. ³⁵; **g** SO, ref. ¹¹; **h** SO, ref. ¹⁴; **i** SO, ref. ¹³; **j** SO, ref. ¹².




Fig. 2 Model results. **a** Modeled profiles of *cis*-(SO)₂, *trans*-(SO)₂, and trigonal-(SO)₂. **b** Modeled profiles of S_n species. **c** Modeled profiles of S₂O, S₃O and S₄O. **d** Comparison of production of aerosol S versus recycling of S back to SO.

substantially increases the photolysis rate of H₂SO₄ in the current model.

The reasons for the differences between observed and calculated SO₂ and OCS for some datasets shown in Fig. 1 are not entirely clear, but some of the applicable datasets share the common feature that they were obtained on relatively short time scales. The SO₂ record at 70 km exhibits dramatic spatial and temporal variability with SO₂ mixing ratios in the equatorial region spanning three orders of magnitude from ppb to ppm levels. Data for some datasets could have been obtained during the brief periods when vertical transport was greatly enhanced compared to the mean values used here. Examining variability in transport as a source of variability in mixing ratios of SO₂ and OCS is beyond the scope of this paper and is best accomplished using a multidimensional model incorporating radiative, chemical and dynamical feedbacks.

Reasonably good agreement of modeled SO with Hubble Space Telescope observations of ref. ¹¹ and the mean SO abundance determined by submillimeter observations of ref. ¹² at higher altitudes is found using a model with 3.0 ppm SO₂ at the lower boundary. Larger model-observation differences are found for the SO mixing ratios observed by ref. ¹⁸ and ref. ¹⁹ beneath an altitude of about 80 km. Reference ¹⁸ concluded that IUE data were best fit with an SO mixing ratio of 20 ± 10 ppb above ~70 km with

no SO beneath that altitude; ref. ¹⁹ derived a best fit to their data with a constant mixing ratio of SO of 12 ± 5 ppb for $z \geq 64$ km and falling off sharply beneath 64 km. As can be seen from Fig. 1, our calculated mixing ratio of SO is 0.14 ppb at 64 km, increasing with height. As can also be seen from Fig. 1, our calculated mixing ratio of SO is ~1 ppb at 70 km.

The SO₂ profiles shown in Supplementary Fig. 2 (solid blue line) calculated using 0.3 ppm at the lower boundary consistently underpredict the higher altitude observations. As seen in Supplementary Fig. 2, a mixing ratio of 0.3 ppm for SO₂ applied at the lower boundary leads to an SO mixing ratio that is slightly too low compared to observations (dashes g, h, i, j). However, much of the disagreement may simply be due to the sparse nature of observations of SO that are not able to capture the extent of spatial and temporal variability as was possible for SO₂.

The mixing ratios of the three isomers of (SO)₂, *c*-(SO)₂, *t*-(SO)₂, and *r*-(SO)₂, are shown in Fig. 2a. Our values are consistent with the model of ref. ³ but are far lower than those calculated by ref. ⁴, particularly in the crucial layer beneath 70 km. The reason is that ref. ⁴ fixed the mixing ratio of SO to 12 ppb at 64 km based on the model of ref. ¹⁹ while decreasing it to 3 ppb at 70 ppb based on the microwave observations of ref. ¹² and then increasing it to 150 ppb at 96 km based on ref. ²⁰. The altitude profile of SO derived by ref. ¹⁹ should be reevaluated in the




Fig. 3 Correlation between S_x and SO_2 . Scatterplot showing imaginary index of refraction at 250 nm versus SO_2 at 70 km from Venus Express¹⁰ (blue dots) compared to the scatterplot of S_x versus SO_2 at 70 km (red diamonds) and 72 km (green dots) obtained from our model. The mixing ratios of S_x in our model are multiplied by 2×10^5 in order to compare with the imaginary index of refraction derived from ref. ¹⁰.

context of the shape of the modeled profile, which shows an increase in the mixing ratio of SO with height. This is a more likely shape for the SO mixing ratio profile based on measurements of ref. ¹¹, ref. ¹² and current understanding of photochemistry. It must also be kept in mind that conditions on Venus are highly spatially and temporally variable, so short-term measurements might not represent conditions at other times. The mixing ratios of S_n ($n = 1-7$) are shown in Fig. 2b and S_nO ($n = 2-4$) are shown in Fig. 2c.

There is competition between production of reduced aerosol and recycling of oxidized S. It is clear from Fig. 2d, that the reaction, $S + O_2 \rightarrow SO + O$, is the primary sink for S throughout the modeling domain, except in a few thin layers at about 64 km and below. This demonstrates the difficulty of producing S_n via S atoms, which are rapidly converted to SO in the presence of O_2 . In our current model, production of S_2 does not directly involve S atoms, as it is derived from alternative paths involving SO dimer photochemistry and as seen below from catalytic cycles involving Cl, thus bypassing the rapid recycling of S to SO by O_2 . Sensitivity studies carried out by varying the rate coefficients in Supplementary Table 1 uniformly upward and downward by a factor of ten indicate that species concentrations are typically within a factor of two of their values shown in Supplementary Table 1.

Cl has been shown to facilitate the stability of CO^{21-23} Cl also exerts strong influence on the production of S_x . The column production rate of aerosol (on an S basis) is $8.3 \times 10^{11} \text{ cm}^{-2} \text{ s}^{-1}$ in our standard model ($SO_2 = 3.0 \text{ ppm}$, $HCl = 0.4 \text{ ppm}$) but it is reduced to $6.4 \times 10^{11} \text{ cm}^{-2} \text{ s}^{-1}$ for $HCl = 0.2 \text{ ppm}$. At the lower SO_2 mixing ratio ($SO_2 = 0.3 \text{ ppm}$) used at 58 km, the results are even more dramatic. Column production rates of aerosol drop from $2.9 \times 10^{11} \text{ cm}^{-2} \text{ s}^{-1}$ to $1.1 \times 10^{11} \text{ cm}^{-2} \text{ s}^{-1}$ for a decrease in HCl mixing ratio from 0.4 to 0.2 ppm. Although changes in major species such as SO_2 associated with changes in HCl are relatively minor, order of magnitude changes in S and Cl-S species are seen. This strong sensitivity results in part from Cl reactions depleting O_2 , the major sink for S, through recombination of CO_2 . This is similar to what was found for production of S_2 via chlorosulfanes²⁴.

There are a number of intriguing clues linking SO_2 photochemistry to the unknown absorber. As shown in Fig. 2d, the maximum production rate of absorbing aerosol in our model


occurs in the lower half of the upper cloud deck, but noticeable depletion of SO_2 occurs only at higher altitudes. Our results are consistent with VEx observations showing an inverse relation between SO_2 at 70 km and UV absorption at 250 nm¹⁰, as shown in Fig. 3. The data points in Fig. 3 are obtained from the time steps towards equilibrium when running our model and are meant to provide an indication of the adequacy of the model's production rate of absorbing aerosol. The density of S_x is found to be negatively correlated with the local SO_2 mixing ratio, which is consistent with the observations of ref. ¹⁰. However, as noted by ref. ²⁵, the lifetime of the UV absorber is much longer than that of SO_2 , and as a result, the ratio of their abundances is strongly affected by transient atmospheric dynamics, in particular convective activity. Upward transport in the ascending branch of the Hadley Cell could bring absorber upward from where it is formed in the lower half of the upper cloud deck. Events such as these would reduce the strength of any correlation between SO_2 and S_x . Since our one-dimensional model is meant to simulate mean conditions at low latitudes, it cannot encompass the full range of conditions sampled by *Venus Express*. It therefore cannot simulate the behavior of SO_2 and the UV absorber undergoing poleward transport; a detailed simulation of their latitudinal behavior requires a two- or three-dimensional model.

Discussion

The SO dimers are more likely to be important intermediaries in the formation of more complex S species that could be responsible for the UV absorption, rather than being the UV absorbers themselves. Cl is an important modulator of important chemical processes maintaining the stability of CO_2 and the production of condensable sulfur species. These species include S_x and S_n , which have been the subject of several studies (see e.g. ref. ²⁴ and references therein), and possibly S_nO ⁵. In our current model, which only includes condensational loss to existing particles and does not consider new particle formation, S_2O is the major condensable species followed by S_2 and S_4 . Although it's still not clear exactly what is driving the overall temporal behavior of the albedo at 365 nm, a possible explanation is that it is driven by the variability in SO_2 levels, in particular through the formation of some of the UV absorbers discussed in the literature (e.g. ref. ²⁶) and/or from polysulfur and polysulfur oxide species. The implications of our model are that more complex sulfur compounds, beyond OCS, SO, SO_2 , SO_3 , and H_2SO_4 , that are also produced by SO_2 photochemistry could contribute to the absorption of UV in the enigmatic 320–400 nm range. Gas phase abundances of polysulfur and S_nO by themselves are not capable of accounting for the NUV absorption but a combination of their condensed and gas phase forms (or in addition to $FeCl_3$) might.

Our base modeling provides reasonable fits to available data, although the significance of these results is not entirely clear given the sparsity of data. The sporadic nature of observations of key species precludes making definitive conclusions regarding the major mechanisms involved in production of the enigmatic UV absorber and for understanding the photochemistry occurring in the upper atmosphere of Venus, especially given the large spatial and temporal variability of species that has been observed, e.g. by *Venus Express*. Rather, a more concentrated approach measuring a number of interacting species simultaneously would provide much needed input and hence greater confidence in our understanding of major chemical processes. In situ measurements using high resolution aerosol mass spectrometry (e.g. ref. ²⁷) might be a way to identify the UV absorber on a future probe. Such measurements would be able to supply information not only identifying the major elements (S, O, C, Cl, N, and H) that may be present in the aerosol, but also their isotopic composition. Long

polymerization via reactions, such as



perhaps producing higher order S_nO , in a manner similar to that for producing S_n .

Data availability

The model data and observational data that support the findings of this study are available from the corresponding author on reasonable request.

Code availability

The code for the photochemical model written in Fortran and the post processing of the data written in Matlab is available from the corresponding author on reasonable request.

Received: 11 May 2020; Accepted: 27 November 2020;

Published online: 08 January 2021

References

- Mills, F. P., Esposito, L. W., & Yung, Y. L. In *Exploring Venus as a Terrestrial Planet* (eds. Esposito, L. W., Stofan, E. & Cravens, T.) 73–100 (American Geophysical Union, Washington, DC, 2007).
- Krasnopolsky, V. A. Sulfur aerosol in the clouds of Venus. *Icarus* **274**, 33–36 (2016).
- Krasnopolsky, V. A. Disulfur dioxide and its near-UV absorption in the photochemical model of Venus atmosphere. *Icarus* **299**, 294–299 (2018).
- Frandsen, B. N., Wennberg, P. O. & Kjaergaard, H. G. Identification of OSSO as a near-UV absorber in the Venusian atmosphere. *Geophys. Res. Lett.* **43**, 11–146 (2016).
- Hapke, B. & Graham, F. Spectral properties of condensed phases of disulfur monoxide, polysulfur oxide, and irradiated sulfur. *Icarus* **79**, 47–55 (1989).
- Krasnopolsky, V. A. A photochemical model for the Venus atmosphere at 47–112 km. *Icarus* **218**, 230–246 (2012).
- Wu, Z. et al. The near-UV absorber OSSO and its isomers. *Chem. Commun.* **54**, 4517–4520 (2018).
- Zhang, X. et al. Sulfur chemistry in the middle atmosphere of Venus. *Icarus* **217**, 714–739 (2012).
- Yung, Y. L. et al. Evidence for carbonyl sulfide (OCS) conversion to CO in the lower atmosphere of Venus. *J. Geophys. Res.* **114**(E5), 113368–113377 (2009).
- Marcq, E. et al. Climatology of SO₂ and UV absorber at Venus' cloud top from SPICAV-UV nadir dataset. *Icarus* **335**, 113368–113377 (2020).
- Jessup, K. L. et al. Coordinated Hubble Space Telescope and Venus Express observations of Venus' upper cloud deck. *Icarus* **258**, 309–336 (2015).
- Sandor, B. J. et al. Sulfur chemistry in the Venus mesosphere from SO₂ and SO microwave spectra. *Icarus* **208**, 49–60 (2010).
- Belyaev, D. A. et al. Night side distribution of SO₂ content in Venus' upper mesosphere. *Icarus* **294**, 58–71 (2017).
- Plane, J. M. C. Cosmic dust in the earth's atmosphere. *Chem. Soc. Rev.* **41**, 6507–6518 (2012).
- Labidi, J. et al. Mass independent sulfur isotope signatures in CMs: implications for sulfur chemistry in the early solar system. *Geochim. Cosmochim. Acta* **196**, 326–350 (2017).
- Sandor, B. J., Clancy, R. T. & Moriarty-Schieven, G. Upper limits for H₂SO₄ in the mesosphere of Venus. *Icarus* **217**, 839–844 (2012).
- Farahani, S. et al. Simulated electronic absorption spectra of sulfur-containing molecules present in Earth's atmosphere. *J. Phys. Chem. A* **123**, 6605–6617 (2019).
- Na, C. Y., Esposito, L. W. & Skinner, T. E. International ultraviolet explorer observation of Venus SO₂ and SO. *J. Geophys. Res.* **95**, 7485–7491 (1990).
- Na, C. Y. et al. Sulfur dioxide in the atmosphere of Venus: II. Modeling results. *Icarus* **112**, 389–395 (1994).
- Belyaev, D. A. et al. Vertical profiling of SO₂ and SO above Venus' clouds by SPICAV/SOIR solar occultations. *Icarus* **217**, 740–751 (2012).
- Yung, Y. L. & DeMore, W. B. Photochemistry of the stratosphere of Venus: implications for atmospheric evolution. *Icarus* **51**, 199–247 (1982).
- Krasnopolsky, V. A. & Parshev, V. A. Chemical composition of the atmosphere of Venus. *Nature* **292**, 610–613 (1981).
- Pernice, H. et al. Laboratory evidence for a key intermediate in the Venus atmosphere: peroxychloroformyl radical. *Proc. Natl Acad. Sci. USA* **101**, 14007–14010 (2004).
- Mills, F. P. & Allen, M. A review of selected issues concerning the chemistry in Venus' middle atmosphere. *Planet. Space Sci.* **55**, 1729–1740 (2007).
- Jessup, K. L. et al. On Venus' cloud top chemistry, convective activity and topography: a perspective from HST. *Icarus* **335**, 113372 (2020).
- Pérez-Hoyos, S. et al. Venus upper clouds and the UV absorber from MESSENGER/MASCS observations. *J. Geophys. Res.* **123**, 145–162 (2018).
- Jayne, J. T. et al. Development of an aerosol mass spectrometer for size and composition analysis of submicron particles. *Aerosol Sci. Technol.* **33**, 49–70 (2000).
- Herron, J. T. & Huie, R. E. rate constant at 298 K for the reactions SO + SO + M → (SO)₂ + M and SO + (SO)₂ → SO₂ + S₂O. *Chem. Phys. Lett.* **76**, 322–324 (1980).
- Zhang, X. et al. Photolysis of sulphuric acid as the source of sulphur oxides in the mesosphere of Venus. *Nat. Geosci.* **3**, 834–837 (2010).
- Atkinson, R. et al. Evaluated kinetic and photochemical data for atmospheric chemistry: volume I-gas phase reactions of Ox, HOx, NOx and SOx species. *Atmos. Chem. Phys.* **4**, 1461–1738 (2004).
- Singleton, D. L. & Cvetanović, R. J. Evaluated chemical kinetic data for the reactions of atomic oxygen O (3P) with sulfur containing compounds. *J. Phys. Chem. Ref. Data* **17**, 1377–1437 (1988).
- Sander, S. P. et al. Chemical kinetics and photochemical data for use in atmospheric studies, evaluation number 14. *JPL Publ.* **02**, 334 (2003).
- Moses, J. I., Zolotov, M. Y. & Fegley, B. Jr. Photochemistry of a volcanically driven atmosphere on Io: Sulfur and oxygen species from a Pele-type eruption. *Icarus* **156**, 76–106 (2002).
- Krasnopolsky, V. A. Chemical kinetic model for the lower atmosphere of Venus. *Icarus* **191**, 25–37 (2007).
- Krasnopolsky, V. A. Spatially-resolved high-resolution spectroscopy of Venus 2. Variations of HDO, OCS, and SO₂ at the cloud tops. *Icarus* **209**, 314–322 (2010).

Acknowledgements

This research was supported by NASA Grant P1969079 under subcontract to the Space Science Institute (SSI) and NASA grant NNX16AN03G to SSI. D.E. and D.B. (IKI) acknowledge funding from the Russian government (topic VENERA). D.E. acknowledges the support by the French Government Scholarship Vernadski for PhD students.

Author contributions

J.P.P., J.L., F.P.M., and Y.L.Y. contributed to the paper writing. J.P.P., J.L., and F.P.M. carried out the modelling work. E.M., D.E., and D.B. provided the data from Venus Express. Y.L.Y. supervised the research.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information is available for this paper at <https://doi.org/10.1038/s41467-020-20451-2>.

Correspondence and requests for materials should be addressed to J.L.

Peer review information *Nature Communications* thanks Vladimir Krasnopolsky and the other, anonymous, reviewer for their contribution to the peer review of this work. Peer reviewer reports are available.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021