



**HAL**  
open science

## The variety of subaerial active salt deformations in the Kuqa fold-thrust belt (China) constrained by InSAR

Cindy Colón, A. Alexander G. Webb, Cécile Lasserre, Marie-Pierre Doin, Francois Renard, Rowena Lohman, Jianghai Li, Patrick F. Baudoin

### ► To cite this version:

Cindy Colón, A. Alexander G. Webb, Cécile Lasserre, Marie-Pierre Doin, Francois Renard, et al.. The variety of subaerial active salt deformations in the Kuqa fold-thrust belt (China) constrained by InSAR. *Earth and Planetary Science Letters*, 2016, 450, pp.83-95. 10.1016/j.epsl.2016.06.009 . insu-03188547

**HAL Id: insu-03188547**

**<https://insu.hal.science/insu-03188547v1>**

Submitted on 2 Apr 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **The variety of subaerial active salt deformations in the Kuqa fold-thrust belt**  
2 **(China) constrained by InSAR**

3

4 Cindy Colón<sup>1\*</sup>, A. Alexander G. Webb<sup>2</sup>, Cécile Lasserre<sup>3,4</sup>, Marie-Pierre Doin<sup>3,4</sup>,  
5 François Renard<sup>3,4,5</sup>, Rowena Lohman<sup>6</sup>, Jianghai Li<sup>7</sup>, Patrick F. Baudoin<sup>1</sup>

6

7

8 <sup>1</sup>*Department of Geology and Geophysics, Louisiana State University, Baton Rouge,*  
9 *Louisiana, 70803, USA*

10 <sup>2</sup>*Department of Earth Sciences, University of Hong Kong, Hong Kong, China*

11 <sup>3</sup>*ISTerre, University Grenoble Alpes, BP53, 38041, Grenoble, Cedex 09, France*

12 <sup>4</sup>*ISTerre, CNRS, BP53, 38041, Grenoble, Cedex 09, France*

13 <sup>5</sup>*PGP, Dept. Geosciences, University of Oslo, box 1048, Blindern, 0316 Oslo, Norway*

14 <sup>6</sup>*Earth and Atmospheric Sciences, Cornell University, Ithaca, New York, 14850, USA*

15 <sup>7</sup>*The Key Laboratory of Orogenic Belts and Crustal Evolution, Ministry of Education,*  
16 *School of Earth and Space Sciences, Peking University, Beijing 100871, China*

17

18

19

20 \*Corresponding author: [cinolon@gmail.com](mailto:cinolon@gmail.com)

21

22

23 Submitted to: Earth and Planetary Science Letters

24 Submission Date: December 2, 2015

25 Resubmission Date: May 8, 2016

26 **ABSTRACT**

27 **Surface salt bodies in the western Kuqa fold-thrust belt of northwestern**  
28 **China allow study of subaerial salt kinematics and its possible correlations with**  
29 **weather variations. Ephemeral subaerial salt exposure during the evolution of a salt**  
30 **structure can greatly impact the subsequent development and deformation of its**  
31 **tectonic setting. Here, we present a quantitative time-lapse survey of surface salt**  
32 **deformation measured from interferometric synthetic aperture radar (InSAR) using**  
33 **Envisat radar imagery acquired between 2003 and 2010. Time series analysis and**  
34 **inspection of individual interferograms confirm that the majority of the salt bodies**  
35 **in western Kuqa are active, with significant InSAR observable displacements at 3 of**  
36 **4 structures studied in the region. Subaerial salt motion toward and away from the**  
37 **satellite at rates up to 5 mm/yr with respect to local references. Rainfall**  
38 **measurements from the Tropical Rainfall Measuring Mission (TRMM) and**  
39 **temperature from a local weather station are used to test the relationship between**  
40 **seasonality and surface salt motion. We observe decoupling between surface salt**  
41 **motion and seasonality and interpret these observations to indicate that regional**  
42 **and local structural regimes exert primary control on surface salt displacement**  
43 **rates.**

44

45 **Key words:** Kuqa fold-thrust belt, salt deformation, namakiers, InSAR

46

47

48

49 **1. INTRODUCTION**

50 Analysis of subaerial salt behavior is limited because allochthonous salt outcrops  
51 are rarely preserved on the Earth's surface, making subsurface imaging the primary  
52 technique for observing salt deposits (Talbot and Pohjola, 2009). Few examples of salt  
53 bodies outcrop at the surface today (Zagros, Iran; Salt Range, Pakistan), not including  
54 submarine salt exposures. Interpretations of many subsurface salt bodies throughout the  
55 world have indicate subaerial exposure during structural evolution (e.g., Canerot et al.,  
56 2005). This subaerial exposure may be short-lived and yet have key impacts on the long-  
57 term evolution because salt bodies interacting with surface processes may display sharply  
58 distinct kinematics versus subsurface salt bodies. As a result, it is important for the  
59 overall exploration of crustal salt to study the few available subaerial salt bodies. Many  
60 structural systems dominated by salt deformation are commonly associated with major  
61 hydrocarbon reservoirs. Additional compelling reasons to explore subaerial salt bodies  
62 are (1) distinctive mechanical properties of salt make it one of the few solid Earth  
63 materials able to flow at the surface, and (2) these represent Earth-surface deformation  
64 systems that can be highly responsive to weather conditions (Talbot and Rogers, 1980).

65 Much of our current understanding of surface salt kinematics stems from  
66 namakier (extrusive salt "glaciers", also called "salt fountains") studies predominantly in  
67 the Alborz and Zagros Mountains in northern and southern Iran, respectively (e.g., Kent,  
68 1979). Over one hundred Hormuz salt diapirs, fed from pipe-like conduits (point-  
69 sourced), are known to exist in the Zagros Mountains (Kent, 1979). Of these salt bodies,  
70 surface motion has been observed on twenty diapirs using a satellite-based remote  
71 sensing technique called interferometry of synthetic aperture radar (InSAR) capable of

72 measuring millimeter-scale vertical displacements (Barnhart and Lohman, 2012). The  
73 sheet-like Garmsar salt nappe of the Alborz Mountains extruded along a frontal thrust  
74 before undergoing open-toed advance and is presently inactive; the motion detected by  
75 InSAR is attributed to dissolution (Baikpour et al., 2010). Furthermore, InSAR has also  
76 been used to detect continuous vertical, upward flow rates between 5.5-8.3 mm/yr across  
77 the line-sourced Sedom diapir in the Dead Sea basin (Weinberger et al., 2006).

78         Recently recognized namakiers in the western Kuqa fold-thrust belt in  
79 northwestern China provide a new natural laboratory to observe and constrain subaerial  
80 salt kinematics (Li et al., 2014). Four active point-sourced namakiers exist in western  
81 Kuqa fold-thrust belt (**Figure 1**). An active line-sourced diapir, the Tuzimazha salt wall,  
82 is comparable to the Sedom diapir with vertical, upward flow. Quele namakier, exposed  
83 in western Kuqa fold-thrust belt, is an active open-toed, line-sourced subaerial salt  
84 structure extruded from a surface-breaching thrust fault on Earth's surface. Kuqa fold-  
85 thrust belt provides an invaluable opportunity to study the behavior of a variety of  
86 subaerial salt bodies within the same geologic setting. Furthermore, the Kuqa fold-thrust  
87 belt is more accessible than other examples of subaerial salt extrusion like the Salt Range  
88 of Pakistan.

89         Existing work concludes that the presence of meteoric water and fluctuating  
90 temperatures are the principal factors enabling salt recrystallization and/or dissolution  
91 after surface extrusion (Talbot and Rogers, 1980; Urai et al., 1986; Desbois et al., 2010).  
92 In the present study, we use InSAR to measure the spatial and temporal deformation  
93 patterns of namakiers in Kuqa fold-thrust belt. We compare these observations to rainfall

94 and temperature measurements in order to explore potential relationships between the  
95 active surface salt kinematics and local weather conditions.

## 96 **2. BACKGROUND**

### 97 **2.1 Geologic setting**

98         The Kuqa foreland fold-thrust belt actively accommodates ~4-7 mm/yr of north-  
99 south tectonic shortening between the Tian Shan to the north and the Tarim basin to the  
100 south across a ~400 km long, ~20-65 km wide region (e.g., Allen et al., 1991).  
101 Contractional tectonics have formed structural hydrocarbon traps across Kuqa fold-thrust  
102 belt where the deformational style is controlled by the presence and varying thickness of  
103 salt layers (e.g. Tang et al., 2004; Chen et al., 2004). The subsalt sequence (Upper  
104 Permian-Cretaceous) includes Triassic-Jurassic coal and lacustrine mudstones that  
105 matured into regional hydrocarbon source rocks (Hendrix et al., 1992; Tang et al., 2004;  
106 Chen et al., 2004). Two salt dominated units control deformation across Kuqa (Li et al.,  
107 2012): the Miocene Jidike Formation and Paleocene-Eocene Kumugeliemu Group  
108 concentrated in eastern and western Kuqa, respectively. The Kumugeliemu Group  
109 consists of halite and gypsum interbedded with mudstone, sandstone, conglomerate and  
110 marl (Hendrix et al., 1992; Chen et al., 2004). These strata had an original thickness of  
111 ~1100 m and the range of current thicknesses spans 110-3000 m (Hendrix et al., 1992;  
112 Chen et al., 2004; Yixin and Pengwan, 2009). This dramatic thickness range testifies to  
113 the importance of salt flow, which thinned and thickened the unit laterally. Where the salt  
114 units thicken toward the south, deformation is concentrated in the suprasalt sequence.  
115 From west to east, salt décollements in Kuqa fold-thrust belt have developed into salt  
116 nappes exposed at the surface, triangle zones and pop-up structures (Chen et al., 2004;

117 Wang et al., 2011, Li et al., 2012). The suprasalt sequence (Oligocene-Quaternary)  
118 reaches a thickness of ~7 km (Tang et al., 2004; Li et al., 2014). The salt-dominated  
119 Kumugeliemu Group in western Kuqa fold-thrust belt (**Figure 1**) outcrops at line-sourced  
120 and point-sourced salt structures. These surface salt structures display structural and  
121 stratigraphic records of active vertical motion, exhumation and surface flow (Li et al.,  
122 2014).

123 Point-sourced salt structures, like Awate namakier, Bozidun namakier and an  
124 unspecified structure referred to hereafter as Daxiagu namakier, are fed from a salt source  
125 via pipe-like feeders (i.e., stems) (e.g., Jackson and Talbot, 1986). Awate namakier  
126 occurs at the western extent of the Quilitage anticline system, divided by a south-flowing  
127 river. The eastern portion of Awate namakier is approximately 2.5 km long and 1.5 km  
128 wide. The western portion may have the same salt source (Li et al., 2014). Salt mining at  
129 the southern edge of Awate exposes freshly cut outcrop surfaces and reveals a mud-rich,  
130 1-3 m thick carapace above the namakier. Bozidun namakier is located ~17 km north of  
131 Awate with a length of ~2.2 km and 1.4 km width. Lastly, Daxiagu namakier is the  
132 westernmost surface salt structure in Kuqa, located ~12 km west-northwest of Awate.  
133 Daxiagu is ~4.5 km long and 3.4 km wide and we interpret it to be a point-sourced based  
134 on its circumscribed surface geometry.

135 Line-sourced salt structures such as Quele namakier and Tuzimazha salt wall have  
136 relatively thin, elongated salt feeders (e.g., Jackson and Talbot, 1986). The Quele salt  
137 thrust propagates southward and the salt décollement is exhumed where the thrust fault  
138 intersects the surface, forming the Quele namakier. Quele namakier is exposed across  
139 ~32 km E-W with a vertical thickness generally ranging from ~50-200 m. The namakier

140 thins and locally pinches out toward the west and the salt thrust extends to a north-  
141 trending, salt-lubricated strike-slip fault that appears to transfer slip to another E-W  
142 trending thrust fault further north (Zhong and Xia, 1998a; Li et al., 2014). Quele  
143 namakier broadens at its eastern extent and covers an area ~5km wide, called the  
144 Chaerhan namakier, which also features an isolated salt dome (Li et al., 2014). Large,  
145 siliciclastic rock bodies, of varying sizes reaching up to 2.5 km length, occur across  
146 Quele namakier. These rock bodies may be considered rafts if that have been dislodged  
147 from surrounding sedimentary units and incorporated into the namakier as it flows. These  
148 sedimentary bodies may also be the more brittle, siliciclastic rocks that make up the  
149 Kumugeliemu Group. Toward the north in the hinterland of Kuqa, the subvertical  
150 Tuzimazha salt wall trends east-southeast and maintains an ~50 m thickness along its ~10  
151 km long surface exposure.

## 152 **2.2 Subaerial salt deformation and the impact of weather conditions**

153       Relatively low viscosity and density as well as a strong resistance to compression  
154 allows rock salt to flow as a non-Newtonian fluid that can pierce through surrounding  
155 sedimentary layers in the Earth's upper crust (e.g., Weijermars et al., 1993).  
156 Microstructural analysis shows that salt flow converges and accelerates as it moves from  
157 a buried source layer into a diapiric feeder, or stem (Desbois et al., 2010). The principal  
158 deformation mechanisms observed at the stem of the diapir are grain-boundary migration  
159 (GBM) and sub-grain rotation (SGR) caused by increased differential stress (Desbois et  
160 al., 2010). Exhumation of salt initially forms salt domes that later develop into salt  
161 fountains as circumferential lateral spreading propagates away from the orifice (e.g.,  
162 Kehle, 1988). Dominant deformation mechanisms shift from GBM and SGR to pressure-

163 solution accommodated grain-boundary sliding (GBS) from the top to the distal part of  
164 the salt fountain (Desbois et al., 2010). This shift can be explained by the influx of  
165 meteoric water and the decreased differential stress as salt flow divergences radially away  
166 from the orifice (i.e., Jackson, 1985; Desbois et al., 2010). A ground-based geodetic  
167 study, with hourly to daily sampling, of the Kuh-e-Namak namakier in southern Iran  
168 proposed that the presence of meteoric water causes plastic deformation of the salt while  
169 daily temperature variations cause elastic deformation (Talbot and Rogers, 1980).  
170 Temporary surface salt flow rate of ~500 mm/day was observed at Kuh-e-Namak  
171 namakier shortly after a large rain event (~330 mm average rainfall from October to  
172 February) (Talbot and Rogers, 1980). Furthermore, it was observed that during dry  
173 periods the surface salt shrunk, retreating and losing some of lateral extent attained  
174 during the wet period (Talbot and Rogers, 1980).

175         Given the typical low-pressure stress conditions of subaerial salt bodies, the  
176 namakier porosity is stimulated by joints formed as a result of diurnal/seasonal thermal  
177 expansion and contraction (Merriam et al., 1962; Talbot and Aftabi, 2004; Desbois et al.,  
178 2010). Vertical joints in subaerial salt bodies dilate as the salt carapace spreads in  
179 response to continual growth of the viscous salt fountain, which permits infiltration of  
180 meteoric water (Desbois et al., 2010). The carapace of a namakier is typically composed  
181 of granular clastic material left behind after the shallower layers of salt have dissolved  
182 (e.g., Walker 1973). An annual temperature range of -0.5°C to 37.8°C was enough to  
183 facilitate the plastic deformation observed at Kuh-e-Namak (Talbot and Rogers, 1980).  
184 However, surface motion during dry periods not only depends on temperature but also the  
185 thermal gradient in the salt, which is dependent on the wetness of the salt (Talbot and

186 Rogers, 1980). The semi-arid climate of the Kuqa fold-thrust belt allows the preservation  
187 of subaerial salt exposures. Here, annual rainfall averages ~74 mm (April-October) and  
188 average temperatures can range from -13.3°C to 31.2°C.

### 189 **2.3 Models, predictions, and tests**

190 The rapid kinematic response of surface salt structures to temporary stimuli  
191 requires subaerial salt evolution analysis to consider short-term (days to years)  
192 deformation kinematics. The rates of salt supply, salt dissolution, sediment accumulation  
193 of the overburden, erosion and regional tectonic strain influence the long-term evolution  
194 of a namakier (i.e., Koyi, 1998; McGuinness and Hossack, 1993). Crustal shortening may  
195 impede diapirism if lateral compression results in an obstruction of the salt feeder (Koyi,  
196 1998). In other circumstances the compression of a diapir feeder may physically separate  
197 the diapir from the source layer. This can produce a temporary pulse of increased salt  
198 extrusion rates that would mark the onset of the final phases of namakier evolution (Koyi,  
199 1998).

200 Extreme daily temperature variations in semi-arid climates, like Kuqa fold-thrust  
201 belt, lead to oscillating inflation and deflation of a subaerial salt body on a sub-hourly to  
202 daily scale (Talbot and Rogers 1980). We would expect the entire surface of the  
203 namakiers to uplift during daylight hours, when surface temperatures can be very high,  
204 under the influence of thermal expansion (Merriam et al., 1962). Similarly, we expect  
205 subsidence across the namakier surfaces, as a result of thermal contraction, when  
206 temperatures drop well below freezing. Hourly sampling, at minimum, of temperature  
207 and surface motion would be necessary to detect the rate and degree of daily oscillations  
208 across the namakier surfaces at Kuqa fold-thrust belt. Similar inflation and deflation

209 cycles in response to seasonal variations may also be detectable. However all these  
210 variations due to the thermal expansion or contraction of the salt body are likely small  
211 because the coefficient of the very low thermal expansion of NaCl, in the range of  $5.10^{-6}$   
212  $^{\circ}\text{C}^{-1}$  (White, 1965). For example, in the idealized case of a homogeneous variation of  
213  $40^{\circ}\text{C}$  on a 1 meter thick salt layer, these thermal effects would result in a vertical  
214 displacement of the order of 200 micrometers.

215         The short-term deformation patterns of namakiers must consider moisture at the  
216 surface. When sufficient quantities of meteoric water make contact with/infiltrate  
217 namakiers, pressure-solution and grain boundary sliding allow accelerated salt flow. We  
218 expect namakiers at Kuqa fold-thrust belt to flow faster within days after storm and rain  
219 events. As salt flow accelerates the leading edge of the surface salt bodies should extend  
220 further laterally. As a namakier spreads laterally, the surface should appear to subside.  
221 This can be visualized for a static namakier by considering how a drop of molasses would  
222 behave on a tabletop. As the drop spreads slowly in every direction, the surface will  
223 flatten, bringing the top of the molasses closer to the table.

224         Two models suggest conflicting roles of the carapace in surface salt kinematics  
225 during dry periods. One scenario is the dry-inflation model that suggests the carapace that  
226 develops on a surface salt body inhibits lateral spreading and facilitates swelling at the  
227 dome (above the orifice) as the salt extrudes from the feeder (Talbot and Pohjola, 2009).  
228 If this model holds for Kuqa fold-thrust belt then we should see a relative surface uplift  
229 above the orifice during dry periods. Alternatively, dry-deflation model suggests that the  
230 carapace hinders uplift of the central dome and instead promotes lateral spreading (Aftabi  
231 et al., 2010). With a constant salt supply rate, this model suggests a continual surface

232 subsidence of the namakier as it spreads laterally. To test these models we must record  
233 time-lapse surface motions of the Kuqa fold-thrust belt namakiers during wet and dry  
234 periods as well as document the occurrences of rain events.

235         The resistance supplied by a load overlying the orifice of a namakier is another  
236 factor that can potentially impact subaerial salt. Emergent, surface breaching, diapirs rise  
237 faster than pre-emergent diapirs because of the reduced confinement and load when the  
238 overburden is removed (e.g., Weinberg, 1993). Strain rates from microstructural analysis  
239 for pre-emergent diapirs range from  $10^{-15}$  to  $10^{-13}$  s<sup>-1</sup> while higher strain rates from  $10^{-11}$   
240 to  $10^{-9}$  s<sup>-1</sup> have been observed for emergent diapirs (Talbot and Jackson, 1987). This  
241 aligns with flow velocity estimates that show pre-emergent diapirs in Germany rise ~0.3  
242 mm/yr while emergent diapirs in Iran can exceed 10 mm/yr (e.g., Trusheim, 1960; Talbot  
243 and Aftabi, 2004). Strain and velocity rates of pre-emergent and emergent diapirs  
244 reasonably suggest that the rate of salt extrusion can increase if enough material above  
245 the orifice of a namakier is removed. The salt above the orifice is a load, and this load is  
246 reduced by salt spreading away from the orifice at the surface.

247         Removal of material above the orifice can occur via erosion and salt dissolution.  
248 During periodic large storm events, the shallow, relatively uncompact layers of the  
249 namakier carapace can erode considerably (Wenkert, 1979). Furthermore, significant salt  
250 dissolution can occur during storm events if the carapace thickness allows sufficient  
251 meteoric water to infiltrate the namakier. In the event that the removal of material above  
252 the orifice is large enough, we expect acceleration of salt extrusion resulting in an overall  
253 rise of the namakier surface. Conversely, if the load above the namakier orifice controls

254 the salt extrusion rate, a thickening carapace would cause a deceleration of salt extrusion  
255 that would also lead to a deceleration of lateral spreading.

### 256 **3. METHODS**

257 To better understand the complex set of processes that govern deformation in salt  
258 bodies, we examine several different bodies using a diverse set of data, including space-  
259 based constraints on ground deformation and weather information from both ground  
260 stations and remote sensing. Surface salt deformation is accessed using a remote sensing  
261 technique called interferometry of synthetic aperture. To perform the InSAR analysis we  
262 utilize Envisat ASAR (Advanced Synthetic Aperture Radar) imagery acquired between  
263 June 2003 and October 2010. To access the relationship between subaerial salt motions  
264 and weather conditions we use Tropical Rainfall Measuring Mission (TRMM) data and  
265 local weather station data during the same period of Envisat acquisitions. The temporal  
266 resolution of the Envisat imagery available must be less than the temporal range of  
267 surface salt motion to detect the salts' kinematic responses to weather variations (**Figure**  
268 **3**). Because of this it is unlikely that our Envisat dataset could resolve the impact of  
269 individual storm events as our best temporal resolution is 36 days. However, it is possible  
270 to detect the impact of seasonal weather variations. The combination of these disparate  
271 datasets, which have very different spatial and temporal resolutions and degrees of  
272 uncertainty, presents some challenges, but also allows a more complete analysis of the  
273 setting.

#### 274 **3.1 Weather estimates**

275 The TRMM satellite has utilized microwave and radar instruments since late 1997  
276 with a spatial coverage between latitude 50°S to 50° (Huffman, 2011). The Daily TRMM

277 Rainfall Estimate (3B42 v7) used for our analysis is a product derived from rain gauge  
278 and multiple TRMM satellite measurements. First, independent precipitation estimates  
279 from various microwave instruments are calibrated to TRMM Combined Instrument  
280 (TCI) precipitation estimates (Huffman, 2011). The 3-hourly multi-satellite  
281 measurements are combined with the accumulated monthly Global Precipitation  
282 Climatology Centre (GPCC) rain gauge analysis using inverse-error-variance weighting  
283 to form a monthly best-estimate precipitation rate (Huffman, 2011). Lastly, the 3-hourly  
284 estimates are scaled for the month then summed to the monthly value for each individual  
285 grid box (Huffman, 2011). The final data set has precipitation estimates with a 3-hourly  
286 temporal resolution and a  $0.25^{\circ} \times 0.25^{\circ}$  spatial resolution (Huffman, 2011). This temporal  
287 resolution is sufficient to observe storms that can occur in the span of a few hours across  
288 the surface salt structures in Kuqa fold-thrust belt.

289 Another dataset we have used includes daily humidity, temperature, and rainfall  
290 measurements from a meteorological weather station in Kuqa, Xinjiang Province, China  
291 (Tutiempo Network). We utilize this station, despite its location over 150 km away,  
292 because it is the closest ground truth. When we compare the rainfall estimates from the  
293 weather station and the TRMM dataset we look at the cumulative measurements for the  
294 time periods between each Envisat image acquisition. Similarly, we compare the average  
295 surface temperature and humidity during each time period between image acquisitions.

### 296 **3.2 Interferometry of synthetic aperture radar (InSAR)**

297 We constructed an InSAR time series to constrain the surface displacements of  
298 active extruded salt structures in western Kuqa fold-thrust belt as well as several surface  
299 displacements associated with hydrocarbon withdrawal and subsurface fluid injection.

300 InSAR is a remote sensing technique that uses radar imagery to provide spatially dense  
301 measurements of surface displacements in the satellite line of sight (LOS) with millimeter  
302 to centimeter accuracy (e.g., Zebker et al., 1994; Massonnet and Feigl, 1998; Bürgmann  
303 et al., 2000). Multiple SAR images are used to generate sets of interferograms to form a  
304 time series after a joint inversion (e.g., Berardino et al., 2002). InSAR time series analysis  
305 helps reduce the impact of several noise sources (decorrelation, orbital and DEM errors,  
306 atmospheric delays, phase unwrapping errors) on displacement rates estimates during the  
307 time period spanned by the full dataset (e.g., Grandin et al., 2012) with an accuracy for  
308 surface displacement velocity at the mm/yr scale. The time series analysis also allows  
309 identification of time variable deformation including seasonal signals and response to  
310 earthquakes or anthropogenic activity.

311 We use 40 ESA ASAR C-band radar images acquired by the Envisat satellite  
312 between June 2003 and October 2010 from Track 291 (along descending orbits). A small-  
313 baseline approach (e.g., Berardino et al., 2002) was used to process interferograms and  
314 invert for average displacement rates and evolution through time with the New Small  
315 Baseline Algorithm Subset (NSBAS) chain, as described in detail by Doin et al. (2011)  
316 and Jolivet et al. (2013). 85 individual interferograms were generated using a modified  
317 version of the Caltech/JPL Repeat Orbit Interferometry Package (ROI\_PAC, Rosen et al.,  
318 2004) and the STRM 90-global DEM. Interferograms were corrected from DEM errors  
319 (Ducret et al., 2013) and stratified tropospheric delays (based on the ERA-Interim global  
320 atmospheric model, Doin et al., 2009; Jolivet et al., 2011), multi-looked by a factor of 4  
321 in the range direction and 20 in the azimuth direction, flattened and filtered before  
322 unwrapping. Individual interferogram coherence was used to determine the optimal

323 unwrapping path for each interferograms. The phase unwrapping was achieved  
324 iteratively, beginning in areas of high coherence then incrementally reducing the  
325 coherence threshold for each iteration (Grandin et al., 2012). The InSAR inversion  
326 process does not allow for a direct measurement of the error margin of the LOS  
327 displacement measurements, however we estimate the noise level is on the order of ~1  
328 mm/year. Radar amplitude images were used to interpret the namakiers' boundaries,  
329 outlined on the velocity maps.

330         We present average velocity (mm to cm/yr) and displacement maps with a spatial  
331 resolution of 80 x 80 m. The velocity and displacement maps are displayed in line of  
332 sight (LOS) with positive (red) and negative (blue) values corresponding to movement  
333 toward and away from the sensor that we furthermore interpret as uplift and subsidence,  
334 respectively, due to the dominant sensitivity of InSAR to vertical motion. Differential  
335 incremental displacement maps and time series are used to track the temporal phase  
336 evolution. To isolate relative displacements, we generate these differential maps and time  
337 series computing the difference between time series from points within deforming areas  
338 and times series from local reference points. This helps minimizing the effect of spatially  
339 coherent noise. The remaining, relative, correlated temporal fluctuations observed and  
340 comparisons to weather conditions are discussed in the following section.

341         The present study uses three datasets that were processed to obtain a time-series  
342 that cover the period June 2003 to October 2010 where InSAR data provide estimates of  
343 surface displacement at a pixel size of ~30 m and time resolution of 36 to 210 days, with  
344 an average of 75 days. From the nearest weather station, average temperatures and  
345 precipitations are obtained. Finally, the TRMM data provide the same time-lapse series as

346 the weather station, at a pixel size resolution of 4 km and for the same periods at the  
347 InSAR data. These three data sets are then processed to 1) analyze the surface  
348 deformation of salt bodies, 2) correlate these deformations with weather conditions.

## 349 **4. RESULTS**

### 350 **4.1 Kinematic observations**

351 The overall distribution of LOS velocity at Bozidun namakier is relatively  
352 uniform. From the average velocity map, minor subsidence is observed on the western  
353 margin and localized uplift near the north and east (**Figure 4A**). The velocity distribution  
354 across Bozidun namakier is nearly indistinguishable from its immediate surroundings  
355 supporting field observations that suggest Bozidun namakier is not actively extruding  
356 material to the surface, as shown in the time series of the **Figure 4**.

357 The distribution of positive and negative LOS velocities at Daxiagu is  
358 asymmetrical and ranges from -4.5 to 5 mm/yr (**Figure 4B**). An area of uplift is observed  
359 on the northeastern side of the structure, oriented toward the northwest. Areas of overall  
360 subsidence occur predominately on the southwestern side of the structure as well as the  
361 northwestern. The incremental displacements maps highlight deformation patterns that do  
362 not reflect the long-term deformation patterns measured by the average displacement  
363 velocity maps (**Figure 5**). Each incremental displacement map of Daxiagu displays  
364 relatively uniform deformation across the namakier with fluctuations between surface  
365 uplift and subsidence at different time intervals. Time series analysis of east and west  
366 Daxiagu namakier inversely mirror each other; the east is uplifting at approximately the  
367 same rate (~4.5 mm/yr) as the west is subsiding. The ribbed pattern that appears in both  
368 the average velocity map and the incremental displacements maps do not correlate to

369 topographical variations but instead may reflect noisiness in the data likely due to the  
370 namakiers' rugged surface.

371 Displacement velocities at Awate namakier are distributed asymmetrically  
372 (**Figure 4C**). Awate namakier is divided in two by a major river with stronger subsidence  
373 observed on the eastern segment, slight subsidence on the west, and uplift at the river  
374 channel in between. On the eastern Awate namakier the displacement velocities decrease  
375 radially away from the center and the maximum rate of displacement approaches -5  
376 mm/yr. Incremental displacement maps show that the east Awate namakier has  
377 experienced periods of uplift despite overall subsidence suggesting it is an active  
378 structure. The areas of displacement at east Awate namakier have consistent shapes in the  
379 incremental displacement maps for periods of a year to three, but gradually shift from a  
380 central block to a more variegated area. These areas all subside and uplift more  
381 dramatically than the surroundings. Minor subsidence and unvarying incremental  
382 displacements are observed at west Awate namakier. Localized uplift at the river channel  
383 approaches 4 mm/yr. The zone of uplift is broader than the river channel, extending  
384 farther to the west. Incremental displacement maps show uplift at the riverbed, dividing  
385 Awate namakier, initiated toward the western riverbank then propagated toward the  
386 center. The time series analysis at the river channel is notably less noisy and fluctuating  
387 than the time series from areas across the namakier, especially for east Awate namakier  
388 (**Figure 4**). This observation indicates that uplift at the river channel is relatively regular  
389 and constant while displacement of the namakier is relatively irregular.

390 The distribution of average displacement velocities across Quele namakier is  
391 predominately uniform (**Figure 4D**). Relative subsidence is measured along the exhumed

392 thrust sheet but strong, localized subsidence is observed at the central segment of Quele  
393 namakier where the average displacement velocity approaches  $\sim 4$  mm/yr. Velocities  
394 measured at Chaerhan namakier and the isolated salt dome at the eastern extent of Quele  
395 are within the noise level of background velocity measurements. A few periods of minor  
396 uplift are observed in the incremental displacement maps at the central segment despite  
397 the overall subsidence suggesting an actively extruding structure (**Figure 4**). Time series  
398 analysis shows the displacement trends of two areas in the central segment partially  
399 mirror each other, from 2003 to 2006. After 2006, higher rates of subsidence are  
400 concentrated toward the east suggesting heterogeneities along the salt thrust.

#### 401 **4.2 Correlation between kinematics and weather conditions**

402 To test the relationship between weather conditions and surface displacements of  
403 namakiers in Kuqa fold-thrust belt we compare the incremental displacements between  
404 each image acquisition to the average humidity, average temperature, and cumulative  
405 rainfall during the respective time intervals (**Figure 6**). There is no clear correlation  
406 between average humidity estimates and surface displacements for any of the studied  
407 namakiers. Similarly, no correlation is observed between displacements and average  
408 temperature for Quele namakier. Furthermore, correlation coefficients between surface  
409 displacement measurements on across the 4 namakiers studied and average temperature  
410 and humidity estimates yield a maximum correlation of 0.2. However, at Awate, Bozidun  
411 and Daxiagu namakiers we see temperature extremes are associated with more surface  
412 salt motion than intermediate temperatures. When we compare the displacements  
413 observed at the namakiers to the cumulative rainfall values estimated by TRMM as well  
414 as the distantly located weather station, we find a positive correlation. There is an

415 increased occurrence of surface salt displacements during time intervals that experienced  
416 lower rainfall estimates suggesting that with less rainfall there is more surface motion.  
417 The correlation observed is not simply an artifact of having more data during periods of  
418 lower rainfall events because there are equal amounts of temporal coverage during the  
419 rainy (April-October) and dry seasons (October-April).

420 Another way to test the correlation between observed surface displacements and  
421 rainfall is to look at how the displacement residuals correlate to rainfall estimates. The  
422 residuals of a dataset are the observed values minus the predicted (or average) values.  
423 Residuals highlight the data points' deviation from the linear average, which are used to  
424 test the correlation between rainfall and surface salt displacement response. When we  
425 calculate the correlation coefficient between the residual time series displacements,  
426 showing the deviation of each displacement data point from the average, at the namakiers  
427 of interest and TRMM rainfall we find no correlation greater than 0.1. The low  
428 correlation coefficient is likely due to the occurrence of both positive and negative  
429 motion with hot and cold temperatures.

## 430 **5. DISCUSSION**

431 InSAR analysis uncovers a variety of ground motions across the subaerial salt  
432 structures of the Kuqa fold-thrust belt. While the majority of the namakiers studied show  
433 surface motion, InSAR estimates of surface motion at Bozidun and Chaerhan namakiers  
434 are relatively indistinct from the surrounding area. The lack of observable deformation at  
435 the Tuzimazha salt wall could mean that (1) the salt wall has recently become inactive,  
436 (2) it is deforming at rates below our detection threshold ( $<1$  mm/yr), or (3) the activity is  
437 confined to a small area that is more challenging to resolve. We favor the third

438 interpretation because the thickness of the Tuzimazha salt wall (~50 m) cannot be  
439 adequately resolved by the 80 x 80 m spatial resolution of our InSAR dataset (Figure S4).  
440 The namakiers exhibit asymmetrical spatial displacement patterns with fluctuating  
441 deformation rates over varying temporal scales (weeks to months). Average velocity  
442 estimates in addition to the incremental displacement steps allow us to explore the  
443 seasonality of subaerial salt deformation and its link to weather conditions.

#### 444 **5.1 Daxiagu namakier**

445         The asymmetric displacement patterns observed at Daxiagu namakier (**Figure**  
446 **7A**) do not conform to the expected radial flow pattern of a salt fountain. The irregular  
447 shape and flow kinematics is likely controlled by the surrounding topography and sub-  
448 salt morphology beyond the feeder, beneath the namakier. A river east of Daxiagu has  
449 eroded steep channel walls, and likely rapidly dissolves salt that flows toward the east.

450         Because surface uplift is expected directly above the feeder (orifice) of a point-  
451 source namakier, we interpret the area of overall uplift, on the eastern side of the  
452 namakier, to indicate the location of the salt feeder. Long-term average uplift also  
453 suggests that the rate of salt extrusion into the namakier exceeds the rate of salt  
454 dissolution/erosion. The similarity in magnitude of surface displacement on the east and  
455 west may indicate the two sides of the namakier are not mutually exclusive (**Figure 4**).  
456 Diminutive subsidence at the northwestern margin could also indicate partial flow toward  
457 the northwest from the namakiers' feeder. Eastward flow is expected, where a river  
458 passes, but the material that flows toward the river is likely dissolved relatively rapidly.  
459 The subsidence is interpreted to reflect an approximate volume balance as the namakier  
460 spreads farther out. This interpretation suggests that the dominant direction of flow is

461 toward the west and that lateral spreading is faster than the rate of supply. The  
462 southwestern lobe of Daxiagu is subsiding on average, but does not appear in the time-  
463 lapse data in any one time-step. However, we find no statistical correlation to suggest  
464 weather conditions control the surface motion patterns measured. Furthermore, this  
465 observation is not likely to be climatically driven because we do not see it in incremental  
466 displacement maps. The north-south tectonic shortening (~4-7 mm/yr) accommodated  
467 across the Kuqa fold-thrust belt is likely to have a greater influence on the exhumation  
468 rates and subsequent surface displacement patterns (e.g., Allen et al., 1991; Zhong and  
469 Xia, 1998; Yang et al., 2010; Zubovich et al., 2010). However, to positively discriminate  
470 the precise influence of regional shortening on subaerial salt motion, spatially dense GPS  
471 observations are necessary across the area.

## 472 **5.2 Awate namakier**

473 Subsidence is concentrated at the center of the eastern Awate namakier (**Figure**  
474 **7B**). The broad, long-term deformation pattern of centralized subsidence is indicative of  
475 a mature namakier that is experiencing a dissolution / erosion rate which exceeds the rate  
476 at which salt is fed into the structure. Distinct anthropogenic forcing may play a role: salt  
477 mining operations spray water along the southwestern margin of the eastern Awate  
478 namakier branch, which may influence the observed long-term subsidence by locally  
479 increasing salt dissolution.

480 Incremental displacement maps do not entirely correspond with observations from  
481 the average displacement velocity map, thereby revealing different deformation patterns  
482 at different temporal scales. Unlike the long-term displacement pattern, four intervals of  
483 incremental surface uplift at eastern Awate namakier occur between January and March

484 (colder/drier months) confirming that new material (salt) is being fed into the namakier.  
485 We observe uplift when our temperature-kinematic salt models predict Awate would  
486 experience overall surface subsidence during colder periods in response to thermal  
487 contraction. Calculated correlation coefficients between incremental displacements and  
488 temperature variations show no statistical correlation. Temperature variations do not  
489 appear to control periods of wide spread uplift and subsidence. The intervals of uplift do,  
490 however, support the dry-inflation model. The dry-inflation kinematic salt model predicts  
491 a namakier will inflate during dry periods because the carapace inhibits lateral spreading.  
492 At Awate namakier, we observe uplift above a central location (dome) during periods of  
493 low rainfall implying that the carapace restricts lateral spreading and prefers dome  
494 inflation during dry periods. Furthermore, periods of increased rainfall do not correspond  
495 to increased surface salt uplift.

496         The average, long-term (~7.3 years) uplift signal measured at the river channel  
497 between the east and west segments is not notably observed in the incremental  
498 displacement maps. If river deposition were responsible for the rise of the river channel  
499 surface between the Awate namakier branches, the area of uplift would not be as  
500 localized as observed. Instead, the rise would continue as the river does, toward the south.  
501 It is more likely that the concentrated uplift observed at the river channel is due the rise  
502 of salt from the source layer through the stem of the diapir that feeds Awate namakier. If  
503 the river cuts off the salt feeder that sources Awate, the observed uplift could be the result  
504 of accumulating strain in the overburden of a partially buried, active salt feeder.  
505 However, the estimated long-term rate of uplift here is not likely to persist for time scales  
506 of >100 years because that would significantly perturb the long-term equilibrium

507 topography by developing a hill in the very place there is a river. It is possible to perturb  
508 the equilibrium topography if the diapir is new or suddenly changing character, but it is  
509 not new because the namakier is well established. Thus, we interpret the equilibrium to  
510 be controlled by the regular frequency of large flooding events allowing this area to  
511 neither rise nor fall with respect to regional baseline for longer periods than ~7 years.  
512 Alternatively, the surface of the river channel in between the Awate namakier can be  
513 controlled by a natural equilibrium between the diapir extrusion rate and river erosion. A  
514 study of climatic conditions from 1951 to 2009 in Xinjiang, China reveals that the  
515 average precipitation has increased by 15 mm since 1990 (Kong and Pang, 2012). To our  
516 knowledge, a detailed study of this particular river does not exist, however, work done on  
517 the proximal Tailan River, west of Awate, found that glacial melt-water accounts for 60%  
518 of its  $7.5 \times 10^8 \text{ m}^3$  annual river discharge (Zhao et al., 2015). The combination of  
519 increased rainfall and possible glacial melt-water, due to the  $1^\circ\text{C}$  raise in average annual  
520 temperature, would facilitate flash-floods every 10 to 100 years that removes a lot of  
521 material through erosion, maintaining a topographic equilibrium (Kong and Pang, 2012).

### 522 **5.3 Quele namakier**

523         Quele namakier (**Figure 7C**) is the exhumed portion of a significant salt thrust  
524 and its surface kinematic patterns vary from typical salt fountains. Because it is a line-  
525 source structure, we do not expect to see radial flow comparable to a point-sourced  
526 structure. Unlike a salt fountain, if the exhumation rate of Quele namakier increases the  
527 surface is not likely to uplift significantly because it is confined by its overburden, the  
528 hanging wall of the Quele salt thrust. Instead we would see increased subsidence due to  
529 amplified southward spreading. Decreased exhumation would result in stagnant to

530 minimal lateral spreading. The overall subsidence observed across Quele namakier is  
531 within the range of the data resolution. However, one area of the structure has  
532 experienced a higher rate of subsidence. Weather conditions are unlikely to be  
533 responsible because we see no statistical relationship between surface motion and  
534 weather variations (temperature and moisture). Furthermore, it is unlikely that weather  
535 conditions would exclusively and consistently affect only one spot along a relatively  
536 smooth hill-front. Alternatively, the isolated subsidence at Quele namakier may be  
537 structurally controlled. The subsidence could be the result of a pinched salt source  
538 because analog models by Cotton and Koyi (2000) show that an overriding hanging wall  
539 can pinch a salt décollement during shortening. A possible indicator of a pinched salt  
540 décollement includes the left-lateral tear fault that occurs in the hanging wall, coincident  
541 with the area of concentrated subsidence (**Figure 7E**). The tear fault could have resulted  
542 from lateral variations along the salt décollement, such as thickness, which could indicate  
543 a pinched salt source for Quele namakier. Alternatively, the tear fault may be the result of  
544 differential shortening of the thrust sheet. A large siliciclastic rock body, which appears  
545 to be a raft, is located along Quele namakier and south of the tear fault (**Figure 7E**). If  
546 this siliciclastic rock body acted as a structural barrier, limiting southward propagation of  
547 the exposed Quele salt décollement (namakier), then it may have caused sufficient  
548 differential shortening rates to generate a tear fault. This alternative suggests that the  
549 localized, increased subsidence at Quele is the result of differential shortening across one  
550 portion of the Quele thrust.

#### 551 **5.4 Limitations**

552           Across Kuqa fold-thrust belt, we discovered that subaerial salt motion is  
553 decoupled from weather conditions. The sparse temporal sampling of this Envisat dataset  
554 is significantly limited when compared to the daily sampling rate of temperature and  
555 rainfall from spaceborne satellites or ground-based weather stations. Additionally, the  
556 restricted spatial rainfall resolution of ~4 km estimates from spaceborne satellites may  
557 also preclude correlation with meter-scale InSAR sampling resolution. Looking forward,  
558 the association of the European Space Agency's Sentinel 1A and 1B could provide  
559 improved temporal coverage after several years of observation and better quantify the  
560 temporal fluctuations that may reveal the weather dependence we expected

## 561 **6. CONCLUSIONS**

562           In this study we present quantitative estimates of sub-aerial salt displacement  
563 rates of surface salt bodies in Kuqa fold-thrust belt of northwest China. We predicted  
564 strong dependence of surface salt motion on weather conditions based on previous  
565 surface salt studies but discovered weather conditions do not control surface kinematics  
566 here. Although the namakiers occur in the same fold-thrust belt, none of the salt bodies  
567 share the same short-term kinematic displacement pattern, which suggests contrasting  
568 subsurface geometries between the salt bodies. We measure the dominant vertical  
569 displacement and identify asymmetric deformation patterns of uplift and subsidence at  
570 Daxiagu and Awate, likely controlled by local tectonics and topography. At Quele  
571 namakier we identify an area of concentrated subsidence that is coincident with  
572 surrounding structural features that may indicate a pinched salt thrust. Li et al., 2014  
573 detailed the geometries of the Awate, Bozidun, Quele and Daxiagu namakiers. Our study

574 furthers our understanding of the Kuqa fold-thrust belt namakiers by providing a  
575 kinematic description of the active subaerial salt motion.

576

577 **Acknowledgements**

578 We thank two anonymous reviewers for their helpful comments. This work was  
579 supported by the Marathon Oil GeoDE Graduate Assistantship at LSU, LA-AMP Bridge  
580 to Doctorate Fellowship at LSU, Université Grenoble-Alpes – Institut des Sciences de la  
581 Terre (ISTerre), Tarim Oil Company (a subsidiary of the China National Petroleum  
582 Corporation), grants from the American Chemical Society’s Petroleum Research Fund,  
583 U.S. National Science Foundation (EAR-1322033), and Cornell University Earth and  
584 Atmospheric Sciences Department. I would especially like to thank Matthieu Volat and  
585 Simon Daout at ISTerre for their technical support. The European Space Agency (ESA)  
586 provided the Envisat ASAR imagery used for this study (Dragon project 10686).

587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636

## REFERENCES

- Aftabi, P., Roustaie, M., Alsop, G. I., and Talbot, C. J., 2010, InSAR mapping and modelling of an active Iranian salt extrusion: *Journal of Geological Society – London*, v. 167. P. 155-170.
- Allen, M. B., Windley, B. F., Chi, Z., Zhong-Yan, Z., and Guang-Rei, W., 1991, Basin evolution within and adjacent to the Tien Shan Range, NW China: *Journal of the Geological Society*, v. 148(2), 369-378.
- Baikpour, S., Zulauf, G., Dehghani, M., and Bahroudi, A., 2010, InSAR maps and time series observations of surface displacements of rock salt extruded near Garmsar, northern Iran: *Journal of the Geological Society, London*, v. 167, p. 171-181.
- Barnhart, W.D., and Lohman, R.B., 2012, Regional trends in active diapirism revealed by mountain range-scale InSAR time series: *Geophysical Research Letters*, v.39, p. 1-5.
- Berardino, P., Fornaro, G. and Lanari, R., 2002, A new algorithm for surface deformation monitoring based on small baseline differential SAR interferograms: *IEEE Transactions on Geoscience and Remote Sensing*, v. 40(11), p. 2375-2383.
- Bürgmann, R., Rosen, P.A., and Fielding, E. J., 2000, Synthetic aperture radar interferometry to measure Earth's surface topography and its deformation: *Annual Review of Earth and Planetary Sciences* v. 28.1 p. 169-209.
- Canerot, J., Hudec, M. R., and Rockenbauch, K., 2005, Mesozoic diapirism in the Pyrenean orogen: Salt tectonics on a transform plate boundary: *AAPG Bulletin*, v. 89(2), p. 211-229.
- Chen, S., Tang, L., Jin, Z., Jia, C., and Pi, X., 2004, Thrust and fold tectonics and the role of evaporites in deformation in the Western Kuqa Foreland of Tarim Basin, Northwest China: *Marine and Petroleum Geology*, v. 21, p. 1027–1042.
- Cotton, J.T. and Koyi, H.A., 2000, Modeling of thrust fronts above ductile and frictional detachments: Application to structures in the Salt Range and Potwar Plateau, Pakistan: *Geological Society of America Bulletin*, v. 112(3), p. 351-363.
- Desbois, G., Závada, P., Schléder, Z., and Urai, J. L., 2010, Deformation and recrystallization mechanisms in actively extruding salt fountain: Microstructural evidence for a switch in deformation mechanisms with increased availability of meteoric water and decreased grain size (Qum Kuh, central Iran): *Journal of Structural Geology*, v. 32.
- Doin, M. P., Lasserre, C., Peltzer, G., Cavalié, O., and Doubre, C., 2009, Corrections of stratified tropospheric delays in SAR interferometry: Validation with global atmospheric models. *Journal of Applied Geophysics*, 69(1), 35-50.
- Doin, M-P, Lodge, F., Guillaso, S., Jolivet, R., Lasserre, C., Ducret, G., Grandin, R., Pathier, E., and Pinel, V., 2011, Presentation of the small baseline NSBAS processing chain on a case example: the Etna deformation monitoring from 2003 to 2010 using Envisat Data Proceedings from Fringe 2011 Workshop, Frascati, Italy, 19-23 September 2011(ESA SP-697, January 2012).
- Ducret, G., Doin, M. P., Grandin, R., Lasserre, C., and Guillaso, S., 2014, DEM corrections before unwrapping in a small baseline strategy for InSAR time series analysis. *Geoscience and Remote Sensing Letters, IEEE*, 11(3), 696-700.
- Grandin, R., Doin, M. P., Bollinger, L., Pinel-Puységur, B., Ducret, G., Jolivet, R., and Sapkota, S. N., 2012, Long-term growth of the Himalaya inferred from interseismic InSAR measurement. *Geology*, 40(12), 1059-1062.
- Hendrix M. S., Graham, S. A., Carroll A. R., Sobel, E.R., McKnight, C. L., Schulein, B. J., and Wang, Z., 1992, Sedimentary record and climatic implications of recurrent deformation in the Tian Shan: Evidence from Mesozoic strata of the north Tarim, south Junggar, and Turpan basins, northwest China: *Geological Society of America Bulletin*, v. 104 (1), p.

637 53-79.

638 Hudec M. R., and Jackson, M. P.A., 2007, Terra infirma: Understading salt tectonics: Earth-  
639 Science Reviews, v. 82, p. 1-28.

640 Huffman, G., 2011, Global Change Master Directory - Daily TRMM and Others Rainfall  
641 Estimate:[http://gcmd.gsfc.nasa.gov/KeywordSearch/Metadata.do?Portal=GCMD&MetadataType=0&MetadataView=Full&KeywordPath=&EntryId=GES\\_DISC\\_TRMM\\_3B42\\_](http://gcmd.gsfc.nasa.gov/KeywordSearch/Metadata.do?Portal=GCMD&MetadataType=0&MetadataView=Full&KeywordPath=&EntryId=GES_DISC_TRMM_3B42_daily_V7)  
642 [daily\\_V7](http://gcmd.gsfc.nasa.gov/KeywordSearch/Metadata.do?Portal=GCMD&MetadataType=0&MetadataView=Full&KeywordPath=&EntryId=GES_DISC_TRMM_3B42_daily_V7), Accessed: 07/2015.

643

644 Koyi, H., 1998, The shaping of salt diapirs: *Journal of Structural Geology*, v. 20(4), p. 321-338.

645 Jackson, M.P.A., 1985. Natural Strain in Diapiric and Glacial Rocksalt, with Emphasis  
646 on Oakwood Dome, East Texas. Bureau of Economic Geology, The University of  
647 Texas at Austin, Texas

648 Jackson, M. P. A., and Talbot, C. J., 1986, External shapes, strain rates, and dynamics of salt  
649 structures: *Geological Society of America Bulletin*, v. 97, p. 305-323.

650 Jolivet, R., Grandin, R., Lasserre, C., Doin, M.-P., and Peltzer, G., 2011, Systematic InSAR  
651 tropospheric phase delay corrections from global meteorological reanalysis data:  
652 *Geophysical Research Letters*, v. 38.

653 Jolivet, R., Lasserre, C., Doin, M. P., Peltzer, G., Avouac, J. P., Sun, J., and Dailu, R., 2013,  
654 Spatio-temporal evolution of aseismic slip along the Haiyuan fault, China: Implications  
655 for fault frictional properties. *Earth and Planetary Science Letters*, 377, 23-33.

656 Kehle, R.O., 1988, The origin of salt structures. In: Schreiber, B.C. (Ed.), *Evaporites and*  
657 *Hydrocarbons*. Columbia University Press, New York, pp. 345–404.

658 Kent, P. E., 1979, The emergent Hormuz salt plugs of southern Iran: *Journal of Petroleum*  
659 *Geology*, v.2: p.117–144.

660 Kong, Y., and Pang, Z., 2012, Evaluating the sensitivity of glacier rivers to climate change based  
661 on hydrograph separation of discharge: *Journal of Hydrology*, in press.

662 Li, S., Wang, X., and Suppe, J., 2012, Compressional salt tectonics and synkinematic strata of the  
663 western Kuqa basin, southern Tian Shan, China: *Basin Research*, v. 23, p. 1-23.

664 Li, J., Webb, A. A. G., Mao, X., Eckhoff, I., Colón, C., Zhang, K., Wang, H., Li, A., He, D., 2014,  
665 Active surface salt structures of the western Kuqa fold-thrust belt, Northwestern China:  
666 *Geosphere*, v. 10 (6).

667 Massonnet, D., and Feigl, K. L., 1998 Radar interferometry and its application to changes in the  
668 Earth's surface: *Reviews of Geophysics*, v. 36(4), p. 441-500.

669 Merriam, M. F., Smoluchowski, R., and Wiegand, D. A., 1962, High-temperature thermal  
670 expansion of rocksalt: *American Physical Society*, v. 125

671 McGuinness, D.B. and Hossack, J.R., 1993, The development of allochthonous salt sheets as  
672 controlled by the rates of extension, sedimentation, and salt supply: In *Rates of*  
673 *geological processes: 14th Annual Gulf Coast Section, SEPM Foundation Bob F. Perkins*  
674 *Research Conference* p. 127-139).

675 Talbot, C. J., and Aftabi, P., 2004, Geology and models of salt extrusion at Qum Kuh, central  
676 Iran: *Journal of Geological Society – London*, v. 161. P. 321-334.

677 Talbot, C. J., and Jackson, M. P. A., 1987, Internal kinematics of salt diapirs: *Bulletin of*  
678 *American Association of Petroleum Geologists*, v. 71(9). p. 1068-1093.

679 Talbot, C. J., and Pohjola, V., 2009, Subaerial salt extrusions in Iran as analogues of ice sheets,  
680 streams and glaciers: *Earth Science Reviews*, v. 97(1-4), p. 155-183.

681 Talbot, C. J., and Rogers, E. A., 1980, Seasonal movements in a salt glacier in Iran. *Science*, v.  
682 208(4442), p. 395-397.

683 Tang, L.-J., Jia, C.-Z., Jin, Z.-J., Chen, S.-P., Pi, X.-J., and Xie, H.-W., 2004, Salt tectonic  
684 evolution and hydrocarbon accumulation of Kuqa fold belt, Tarim Basin, NW China:  
685 *Journal of Petroleum Science and Engineering*, v. 41, p. 97-108.

686 Trusheim, F., 1960, Mechanism of salt migration in northern Germany: *Bulletin of the American*  
687 *Association of Petroleum Geologists*, v.A(9) p. 1519-1540.

688 Tutiempo Network, S.L., "Climate KUQA - Climate Data 1956 – 2015." *TuTiempo.net*.  
689 November 2015 (updated). Accessed: December 2014. URL: [en.tutiempo.net/climate/ws-](http://en.tutiempo.net/climate/ws-516440.html)  
690 [516440.html](http://en.tutiempo.net/climate/ws-516440.html).  
691 Urai, J. L., C. J. Spiers, H. J. Zwart, and G. S. Lister, 1986, Weakening of rock salt by water  
692 during long-term creep: *Nature*, v. 324, p. 554-557.  
693 Walker, C. W., 1973, Nature and Origin of Caprock Overlying Gulf Coast Salt Domes: Fourth  
694 International Symposium on Salt–Northern Ohio Geological Society. p. 169-195.  
695 Wang, X., Suppe, J., Guan, S., Hubert-Ferrari, A., Gonzalez-Mieres, R., and Jia, C., 2011,  
696 Cenozoic structure and tectonic evolution of the Kuqa fold belt, southern Tianshan,  
697 China, in K. McClay, J. H. Shaw, and J. Suppe, eds., Thrust fault-related folding: AAPG  
698 Memoir 94, p. 215-243.  
699 Weijermars, R., Jackson, M.P.A., and Vendeville, B., 1993, Rheological and tectonic modeling  
700 of salt provinces: *Tectonophysics*, p. 143–174.  
701 Weinberg, R. F., 1993, The upward transport of inclusions in Newtonian and power-law salt  
702 diapirs: *Tectonophysics*, v. 228. p. 141-150.  
703 Weinberger, R., Lyakhovsky, V., Baer, G., and Begin, Z. B., 2006, Mechanical modeling and  
704 InSAR measurements of Mount Sedom uplift, Dead Sea basin: Implications for effective  
705 viscosity of rock salt: *Geochemistry, Geophysics, Geosystems*, v. 7(5).  
706 Wenkert, D., 1979, The flow of salt glaciers: *Geophysical Research Letters*, v. 6(6).  
707 Wu, Z., Yin, H., Wang, X., Zhao, B., and Jia, D., 2013, Characteristics and deformation  
708 mechanism of salt-related structures in the western Kuqa depression, Tarim basin:  
709 Insights from scaled sandbox modeling: *Tectonophysics*, v. 613-614, p. 81-96.  
710 Yang, S., Li, J., and Wang, Q., 2010, The deformation pattern and fault rate in the Tianshan  
711 Mountains inferred from GPS observations: *Science in China Series D: Earth Sciences*, v.  
712 51(8), p. 1064-1080.  
713 Yixin, Y. and Pengwan, W., 2009, Balanced Cross-sections of Salt Structures in Kuqa Foreland  
714 Thrust Belt in Northern Part of Tarim Basin: *Journal of Marine Origin Petroleum*  
715 *Geology*, p.012.  
716 Zebker, H. A., Rosen, P. A., Goldstein, R. M., Gabriel, A., and Werner, C., 1994, On the  
717 derivation of coseismic displacement fields using differential radar interferometry: the  
718 Landers earthquake: *Journal of Geophysical Research*, v. 99(B10), p. 19,617-19,634.  
719 Zhong, D., and Xia, W. S., 1998a, The investigation report of Mesozoic–Cenozoic strata,  
720 structure, sedimentary faces, and petroleum potential of Kuche foreland basin outcrop  
721 area (with 1:100,000 Kuche foreland basin geologic map): Research report of Tarim  
722 Oilfield Company, Xinjiang, p. 462.  
723 Zubovich, A. V., Wang, X. Q., Scherba, Y. G., Schelochkov, G. G., Reilinger, R., Reigber, C.,  
724 and Beisenbaev, R. T., 2010, GPS velocity field for the Tien Shan and surrounding  
725 regions: *Tectonics*, v. 29(6).

726 **FIGURE CAPTIONS**

727 **Colón et al., Figure 1** (Top) Geologic map of the western Kuqa fold-thrust belt, adapted  
728 from Li et al., 2014, located south of Tian Shan Mountain Range and northern Tarim  
729 Basin. Blue and green units toward the north represent Tian Shan bedrock. Yellow and  
730 orange units represent Cenozoic basin deposits. (Bottom) Cross section (adapted from Li  
731 et al., 2012) from the southern Tian Shan through the Kuqa fold-thrust belt, toward the  
732 Tarim basin and dissects the Quele salt nappe and the Misikantage anticline. The  
733 schematic cross section of Quele namakier is more representative of the central portion of  
734 Quele namakier.

735

736 **Colón et al., Figure 2** Chart highlights variables that may control subaerial salt motion,  
737 their temporal and spatial ranges, and an illustration and description of the predicted  
738 kinematic impact. White arrows represent dominant salt flow direction. Black arrows  
739 represent surface displacement direction. Swelling of the salt in hot weather due to  
740 thermal expansion and deflation during cold weather because of thermal contraction may  
741 occur however the range of motion may be very small. During and shortly after rainfall  
742 events surface salt flow is predicted to accelerate significantly albeit short lived. During  
743 dry periods, 2 opposing models explain surface salt behavior: Dryness (a) predicts the  
744 strong carapace that develops during dry periods inhibits lateral spreading and facilitates  
745 swelling at the dome (above the orifice) as the salt extrudes from the feeder. Dryness (b)  
746 predicts that the thick carapace hinders uplift of the central dome and instead promotes  
747 lateral spreading. The presence of a thin weak overburden/carapace can potentially

748 reduce overburden stress and accelerate salt extrusion. A thick, strong  
749 overburden/carapace may apply enough overburden stress to reduce salt extrusion rates.  
750  
751 **Colón et al., Figure 3** Logarithmic plot with spatial range in meters over temporal range  
752 in days. The yellow, green and blue regions represent the range of temporal and spatial  
753 scales that deformation in response to respective variables could be observed. The purple  
754 highlights possible GPS sampling resolution. The dashed black rectangle represents the  
755 range of spatial and temporal sample of our Envisat dataset. Satellites like the Sentinel  
756 1A and 1B have potential to reach temporal resolution of up to 5 days. The dashed gray  
757 rectangle represents the maximum Tropical Rainfall Measuring Mission (TRMM)  
758 resolution used in this study.

759

760 **Colón et al., Figure 4 (Top)** Average surface velocity maps for Bozidun namakier (A),  
761 Daxiagu namakier (B), Awate namakier (C), and Quele namakier (D) displayed in  
762 satellite line of sight (LOS) and saturated to the color scale 0.5 to -0.5 cm/yr. Surface  
763 movement toward (uplift) and away (subsidence) from the satellite is represented by  
764 warm and cool colors, respectively. (Bottom) Differential time series analysis of select  
765 points on Bozidun, Daxiagu, Awate, and Quele namakiers (locations are highlighted by  
766 black circles on the velocity maps above) is displayed in LOS displacement (cm) across  
767 time (from June 2003 to October 2010). The displacement measurements are relative to  
768 local references represented by the label “Ref”.

769

770 **Colón et al., Figure 5** Line-of-sight (LOS) differential incremental displacement maps of  
771 Daxiagu (1A-33A), Awate (1B-33B), and Quele namakiers (1C-33C), saturated to the  
772 color scale 5 to -5 cm. Displacement values of the local reference points (labeled in  
773 Figure 4) were removed from each incremental displacement map (*Ref1* was used for  
774 Quele namakier) to reduce spatially coherent noise and highlight relative motion. Each  
775 incremental map spans the time between 2 successive acquisitions, the temporal coverage  
776 is represented by the shaded red area on each timeline.

777


778 **Colón et al., Figure 6** Comparison between surface displacements at Kuqa fold-thrust  
779 belt and weather conditions. We evaluate the dependence of incremental LOS  
780 displacement rates (cm) from each structure (Bozidun, Daxiagu, Awate, and Quele  
781 namakiers) to daily humidity (left), temperature (center left), and rainfall/snowmelt  
782 measurements (center right) at a meteorological weather station in Kuqa (Xinjiang  
783 Province, China) from Tutiempo Network and daily cumulative rainfall estimates from  
784 the Tropical Rainfall Measuring Mission (right).

785


786 **Colón et al., Figure 7 (Top)** Schematic cross-section and overview of displacement  
787 patterns across Daxiagu, Awate and Quele namakiers and kinematic interpretations.  
788 Daxiagu is interpreted to have an inflating dome that actively extrudes from salt feeder on  
789 the eastern side. Similarly, Awate is interpreted to flow predominantly toward the east to  
790 southeast. Awate is interpreted to have a salt feeder on the eastern segment that may be  
791 partially buried by the adjacent river channel causing the localized uplift. Unlike Awate  
792 and Bozidun namakiers, Quele namakier does not have a central dome because it is an

793 exhumed salt thrust décollement. We interpret the dominant flow direction of Quele  
794 namakier to be toward the south. (Bottom) The average displacement map (displayed in  
795 radians) overlaid on top of Google Earth. The concentrated area subsidence at Quele  
796 corresponds precisely to a large raft immediately to the east of the area of deformation.  
797 The local left-lateral tear fault may also have created a structure barrier between this part  
798 and the rest of the namakier.

# FIGURES









(B) Kuqa fold-thrust cross-section



Schematic cross-section of Quele namakier (central segment)


Colón et al., Figure 1

### Factors impacting subaerial salt structure motion

Variable	Temporal range	Spatial flow range	Kinematic response to external condition
Hot (surface temp.)	hourly, daily, seasonal	<mm to cm	 <p>Thermal expansion may cause inflation and overall relative surface uplift of the namakiers when surface temperatures are high</p>
Cold (surface temp.)	hourly, daily, seasonal	<mm to cm	 <p>Thermal contraction may cause deflation and overall relative surface subsidence of the namakier when temperatures are low</p>
Wetness	daily, seasonal	<mm to meters	 <p>Meteoric waters facilitates pressure-solution GBS which accelerates salt flow. As salt spreads faster and further laterally, the namakiers' surface appears to subside</p>
Dryness (a)	seasonal	<mm to cm	 <p>Dry-inflation scenario suggests the carapace, developed during dry periods, inhibits lateral spreading and facilitates swelling above the orifice resulting in relative surface uplift above the orifice</p>
Dryness (b)	seasonal	<mm to cm	 <p>Dry-deflation scenario suggests the carapace, developed during dry periods, hinders uplift of the central dome and promotes lateral spreading resulting in surface subsidence of the namakier as it spreads laterally</p>
Thin Overburden	seasonal, annually	<mm to cm	 <p>If enough material is removed from the top of a namakier, thereby reducing the overburden above the orifice, and causing an increase salt extrusion rate which results in surface uplift above the orifice</p>
Thick Overburden	>annually	<mm to cm	 <p>If a namakier carapace thickens significantly then the additional load from may cause enough resistance to decrease the salt extrusion rate resulting in reduced namakier growth</p>


↓ = vertical surface motion    ↘ = dominant salt flow direction

Colón et al., Figure 2




Colón et al., Figure 3

**Average surface displacement rate maps**




**Time series analysis of surface salt structures**




Colón et al., Figure 4


Colón et al., Figure 5



Colón et al., Figure 6



## InSAR surface displacement rates and kinematic interpretations



Colón et al., Figure 7