



Dissolved Oxygen stable isotopes in groundwater: a new tracer of biogeochemical reactivity for the underground Critical Zone

Ivan-David Osorio-Leon, Eliot Chatton, Marlene Dordoni, Davit Piatka, Johannes Barth, Tanguy Le Borgne, Camille Bouchez

► To cite this version:

Ivan-David Osorio-Leon, Eliot Chatton, Marlene Dordoni, Davit Piatka, Johannes Barth, et al.. Dissolved Oxygen stable isotopes in groundwater: a new tracer of biogeochemical reactivity for the underground Critical Zone. Goldschmidt Conference 2021, European Association of Geochemistry & Geochemical Society., Jul 2021, online, United States. <insu-03286671>

HAL Id: insu-03286671

<https://insu.hal.science/insu-03286671v1>

Submitted on 15 Jul 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Dissolved Oxygen stable isotopes in groundwater: a new tracer of biogeochemical reactivity for the underground Critical Zone

Ivan-David Osorio-Leon ¹, Eliot Chatton ¹, Marlene Dordoni ², Davit Piatka ², Johannes Barth ², Tanguy Le Borgne¹, Camille Bouchez ¹

¹ University of Rennes 1, CNRS, Géosciences Rennes, UMR 6118, 35000 Rennes, France

² GeoZentrum Nordbayern – Friedrich-Alexander University, Germany

The persistence of dissolved oxygen (DO) in deep groundwater sustains microbial life and biogeochemical reactivity, with potential impacts on large scale nutrient cycling. In aquifers, dissolved oxygen is often heterogeneous and intermittent, but the driving factors of the spatial and temporal DO distributions remain poorly identified. This study aims to identify the reactive processes controlling DO dynamics in a fractured-bedrock catchment in Brittany, characterized by strong surface-groundwater connectivity, rapid hydrologic response and deep microbial hot-spots.

Oxygen stable isotopes of DO and of the water molecule, major gases, CFC concentrations and major cations and anions were analyzed on rainwater and on groundwater sampled from 10 150m-deep piezometers located in recharge and discharge zones of the catchment. We show that DO concentrations decrease with increasing water residence time while the $\delta^{18}O - DO$ signatures show two different trends depending on DO concentrations. DO concentrations ranging from 10 - 1.5 mg/L are associated with an unexpected depletion of $\delta^{18}O - DO$ suggesting that the DO pool enriches in ^{16}O as it is consumed. Conversely, DO concentrations below 1.5 mg/L (the so-called microaerobic range) are associated with an increase of the $\delta^{18}O - DO$ consistent with an enrichment fractionation of 0.9950 – 0.9980. Our results reveal a shift in the reactive processes controlling DO dynamics and $\delta^{18}O - DO$ at catchment scale.

Water chemical data as well as laboratory reactivity batch experiments indicate that, at high DO concentrations, the abiotic oxidation of dissolved Fe(II) is the main reactive process consuming DO but is not responsible for the observed $\delta^{18}O - DO$ depletion. Isotopic equilibrium exchanges of ^{18}O between DO and water, which $\delta^{18}O - H_2O$ compositions range from -5 to -5.2 ‰, could however explain the light DO isotopic signatures observed despite short (below 50 years) residence times. On the other hand, the observed enrichment in $\delta^{18}O - DO$ at low DO concentrations reflects a clear oxygen consumption by the activity of microaerobic iron-oxidizing bacteria (FeOB).

The present study shows a spatial heterogeneity of biogeochemical reactive processes controlled by oxygen concentrations. DO stable isotopes proved useful to unravel biotic and abiotic processes at stake in the deep critical zone with possible applications for the assessment of primary production of deep biomass.

Keywords: isotopic fractionation, dissolved oxygen isotopes, biogeochemical reactivity, Critical Zone, groundwater.