



**HAL**  
open science

## Distribution of RV- and transiting exoplanets by masses and orbital periods taking into account observational selection

Vladislava Ananyeva, Alexander Tavrov, Oleg Korablev, Jean-Loup Bertaux

### ► To cite this version:

Vladislava Ananyeva, Alexander Tavrov, Oleg Korablev, Jean-Loup Bertaux. Distribution of RV- and transiting exoplanets by masses and orbital periods taking into account observational selection. European Planetary Science Congress. EPSC 2021, Sep 2021, Virtual Meeting, Germany. 10.5194/epsc2021-802 . insu-03312828

**HAL Id: insu-03312828**

**<https://insu.hal.science/insu-03312828>**

Submitted on 2 Aug 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



## Distribution of RV- and transiting exoplanets by masses and orbital periods taking into account observational selection

Vladislava Ananyeva<sup>1</sup>, Alexander Tavrov<sup>1</sup>, Oleg Korablev<sup>1</sup>, and Jean-Loup Bertaux<sup>2</sup>

<sup>1</sup>Space Research Institute of the Russian Academy of Sciences (IKI), Lab 501, Moscow, Russian Federation (a-lada@yandex.ru)

<sup>2</sup>LATMOS/IPSL/CNRS/UVSQ, 78280, Guyancourt, France

More than 95% of the known exoplanets were discovered by the transit- and radial velocity techniques. However, the observed distributions of planets by their masses and by their orbital periods are significantly distorted by numerous observational selections, different for these techniques and different surveys.

We found and studied the de-biased statistical distributions of exoplanets by masses and by orbital periods for three groups of exoplanets: I. transiting planets discovered by Kepler ST, whose masses were measured by the follow-up radial velocity technique, II. transiting planets discovered by ground-based surveys (SuperWASP, HATNet, NGTS, XO, KELT, etc.), III. planets discovered by the radial velocity technique. The synthetic projective-mass distribution of RV planets obeys the piecewise power law with the breakpoints at  $\sim 0.14\text{MJ}$  and  $\sim 1.7\text{MJ}$ . The distribution of RV-planets with  $m = (0.011-0.087)\text{MJ}$  (or  $(3.5-28)\text{ME}$ ) accurately obeys the power law with an exponent of  $-2$ ,  $dN/dm \propto m^{-2}$ . The distribution of RV planets with  $m = (0.21-1.7)\text{MJ}$  follows the power law with an exponent ranging from  $-0.7$  to  $-0.8$ ,  $dN/dm \propto m^{(-0.7\dots-0.8)}$ . The distribution of RV planets with  $m = (1.7-13)\text{MJ}$  is fitted by a power law with an exponent ranging from  $-1.7$  to  $-2.0$ ,  $dN/dm \propto m^{(-1.7\dots-2.0)}$ . In general, the synthetic projective-mass distribution of RV planets well agrees with the predictions of the population synthesis theory (Mordasini, 2018) and includes more detailed features to be discussed.

The exoplanets distribution by periods generally follows a power law with an exponent of  $-0.75$  and indicates a predominant (averaged) structuring of the planetary systems.

De-biased mass distribution of RV-planets with orbital periods of 2-58 days is well consistent with a similar distribution of the Kepler planets. The mass distribution of the transit exoplanets detected by ground-based surveys is consistent with the similar distribution of RV planets with periods of 1-20 days.