



**HAL**  
open science

## **An atmospheric origin for HCN-derived polymers on Titan**

Zoé Perrin, Nathalie Carrasco, Audrey Chatain, Lora Jovanovic, Nathalie Ruscassier, Ludovic Vettier, Guy Cernogora

### ► **To cite this version:**

Zoé Perrin, Nathalie Carrasco, Audrey Chatain, Lora Jovanovic, Nathalie Ruscassier, et al.. An atmospheric origin for HCN-derived polymers on Titan. European Planetary Science Congress (EPSC 2021), Sep 2021, Online, France. pp.491, <10.5194/epsc2021-491>. <insu-03324109>

**HAL Id: insu-03324109**

**<https://insu.hal.science/insu-03324109v1>**

Submitted on 23 Aug 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License



## An atmospheric origin for HCN-derived polymers on Titan

Zoé Perrin<sup>1</sup>, Nathalie Carrasco<sup>1</sup>, Audrey Chatain<sup>1</sup>, Lora Jovanovic<sup>1</sup>, Nathalie Ruscassier<sup>2</sup>, Ludovic Vettier<sup>1</sup>, and Guy Cernogora<sup>1</sup>

<sup>1</sup>CNRS/LATMOS, Université Paris-Saclay/UVSQ, Guyancourt, France

<sup>2</sup>LGPM, Centrale-Supélec, Gif-sur-Yvette, France

### ▪ Introduction

Titan, Saturn's largest moon, has a thick nitrogen-based atmosphere, where during its flyby, the IRIS infrared spectrometer aboard the Voyager 1 spacecraft detected nitriles, such as HCN, formed by EUV photochemistry based on methane CH<sub>4</sub> and molecular nitrogen N<sub>2</sub>. Titan is also surrounded by an organic photochemical haze, according to the observations of the Cassini-Huygens and Voyager 1 missions.

In-situ measurements by the Huygens spacecraft [1], and laboratory experiments synthesizing Titan analog aerosols (called tholins), have revealed that HCN is one of the major chemical signatures extracted from the aerosols, and their laboratory analogues.

Laboratory experiments were conducted using a powder plasma reactor mimicking Titan's ionosphere, replicating the formation of Titan-like organic aerosols, and the associated chemistry. In this work, we simultaneously study the temporal evolution of HCN present in the gas phase, and the formation and growth of Titan tholins.

### ▪ 1 - Aerosol production

Analogues of Titan aerosols are formed using the experiment PAMPRE [4]. PAMPRE is a reaction chamber, where a cold radiofrequency plasma capacitively coupled is ignited at low pressure (~0.9 mbar). A gas mixture of 95% N<sub>2</sub> and 5% CH<sub>4</sub> is introduced into the reactor. Subsequently, a 12W discharge is generated between two electrodes, which ionizes the N<sub>2</sub> and CH<sub>4</sub> present.


In this plasma, the radicals and products formed by ionization of methane and nitrogen, will combine by several reaction pathways, to form more complex organic particles and nitriles such as HCN, in the same way as in the ionosphere of Titan.

In this study, the gas flow rate was optimized, to increase the residence time of the gas mixture in the reactor as much as possible. The chemical growth of the solid particles is thus favored, allowing to follow simultaneously the formation and the evolution of the particles, as well as the co-evolution

of the gas mixture composition.

## ▪ 2 - Morphology analysis of tholins by scanning electron microscopy

The formed samples were observed by scanning electron microscopy (SEM field emission gun).



*Figure 1 - SEM picture of Titan tholins obtained after 160 s in the discharge*

The images show primary nanometer monomers coagulated and formed aggregates (Figure 1). These aggregates continue to evolve to form single spherical particles of  $\sim 1.5 \mu\text{m}$  in diameter (Figure 1). These experimental results are in agreement with the evolution predicted by the model [2] on aerosols residing in the atmosphere of Titan.

## ▪ 3 - Chemical analysis of tholins by infrared spectroscopy

Absorbance measurement is performed using an FTIR, on pellets composed of  $\sim 99.6\%$  KBr and  $\sim 0.4\%$  tholins. Three absorption bands stand out on the mid-infrared absorption spectrum. In particular, the intense bands at  $1560 \text{ cm}^{-1}$  and  $1630 \text{ cm}^{-1}$  (Figure 2) which correspond to different functional groups, such as aromatic or aliphatic  $-\text{NH}_2$ ,  $\text{C}=\text{N}$ ,  $\text{C}=\text{C}$ , aromatic or heteroaromatic groups, difficult to distinguish from each other. The presence of these nitrogenous aromatic compounds can be promoted by HCN [3] [5].




Figure 2 - Normalized IR spectra showing the aromatic or aliphatic groups bands

#### ▪ 4 - Conclusions and Perspectives

The growth of the particles could be observed by the SEM analysis. Moreover, the IR analysis shows the presence of nitrogenous chemical bonds, corresponding to aromatic compounds or nitrile. This growth may be due to the incorporation of HCN chemistry on the surface of the tholins. Indeed, MS analysis of the gas phase shows a progressive consumption of HCN during the formation and growth of Titan tholins. Our experimental results are consistent with numerical models based on Cassini-Huygens observations: A formation of atmospheric HCN on Titan, and its efficient incorporation in Titan's haze to produce large organic polymers.

#### References

- [1] Israël G. et al., Nature 438 : 796-99 (2005).
- [2] Lavvas P. et al., The Astrophysical Journal (2011).
- [3] Imanaka H. et al., Icarus, 168: 344-66 (2004).
- [4] Szopa C. et al., Planetary and Space Science 54 (2006).
- [5] Gautier T. et al., Icarus, 221, 320-327 (2012).