



**HAL**  
open science

## The influence of landscape spatial configuration on nitrogen and phosphorus exports in agricultural catchments

Antoine Casquin, Rémi Dupas, Sen Gu, Ewan Couic, Gérard Gruau, Patrick Durand

► **To cite this version:**

Antoine Casquin, Rémi Dupas, Sen Gu, Ewan Couic, Gérard Gruau, et al.. The influence of landscape spatial configuration on nitrogen and phosphorus exports in agricultural catchments. *Landscape Ecology*, 2021, 36 (12), pp.3383-3399. 10.1007/s10980-021-01308-5. insu-03325139

**HAL Id: insu-03325139**

**<https://insu.hal.science/insu-03325139>**

Submitted on 17 Sep 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Landscape Ecology

## The influence of landscape spatial configuration on nitrogen and phosphorus exports in agricultural catchments --Manuscript Draft--

<b>Manuscript Number:</b>	LAND-D-21-00075	
<b>Full Title:</b>	The influence of landscape spatial configuration on nitrogen and phosphorus exports in agricultural catchments	
<b>Article Type:</b>	Original research	
<b>Keywords:</b>	Eutrophication; landscape pattern; landscape organisation; landscape composition; heterogeneity; scaling	
<b>Corresponding Author:</b>	Antoine Casquin, M.S. Institut National de Recherche pour l'Agriculture l'Alimentation et l'Environnement Rennes, Bretagne FRANCE	
<b>Corresponding Author Secondary Information:</b>		
<b>Corresponding Author's Institution:</b>	Institut National de Recherche pour l'Agriculture l'Alimentation et l'Environnement	
<b>Corresponding Author's Secondary Institution:</b>		
<b>First Author:</b>	Antoine Casquin, M.S.	
<b>First Author Secondary Information:</b>		
<b>Order of Authors:</b>	Antoine Casquin, M.S. Rémi Dupas, PhD Sen Gu Ewan Couic, PhD Gérard Gruau, PhD Patrick Durand, PhD	
<b>Order of Authors Secondary Information:</b>		
<b>Funding Information:</b>	Interreg (Channel Payments for Ecosystem Services)	Dr. Rémi Dupas Dr Gérard Gruau Dr Patrick Durand
<b>Abstract:</b>	<p>Nitrogen (N) and phosphorus (P) exports from rural landscapes can cause eutrophication of inland and coastal waters. Few studies have investigated the influence of the spatial configuration of nutrient sources – i.e. the spatial arrangement of agricultural fields in headwater catchments – on N and P exports. This study aimed to 1) assess the influence of the spatial configuration of nutrient sources on nitrate (NO<sub>3</sub><sup>-</sup>) and total phosphorus (TP) exports at the catchment scale, and 2) investigate how relationships between landscape composition and landscape configuration vary depending on catchment size. We analysed NO<sub>3</sub><sup>-</sup> and TP in 19 headwaters (1-14 km<sup>2</sup>, Western France) every two weeks for 17 months. The headwater catchments had similar soil types, climate, and farming systems but differed in landscape composition (percentage agricultural land-use) and spatial configuration. We developed a landscape configuration index (LCI) describing the spatial configuration of nutrient sources as a function of their hydrological distance to streams and flow accumulation zones. We calibrated the LCI's two parameters to maximise the rank correlation with median concentrations of TP and NO<sub>3</sub><sup>-</sup>. We found that NO<sub>3</sub><sup>-</sup> exports were controlled by landscape composition, whereas TP exports were controlled by landscape configuration. For a given landscape composition, landscape spatial configuration was heterogeneous at small scales (&lt;10 km<sup>2</sup>) but homogeneous at larger scales (&gt;50 km<sup>2</sup>).</p>	

	<p>The spatial configuration of nutrient sources influences TP but not NO<sub>3</sub>- exports. An ideal placement of mitigation options to limit TP export should consider both the hydrological distance to streams and flow accumulation zones.</p>
<p><b>Suggested Reviewers:</b></p>	<p>Sarah E Godsey, PhD  PI, Idaho State University  godsey@isu.edu  Relevant work on landscape pattern and biogeochemistry of headwater streams</p> <hr/> <p>Erin Peterson, PhD  PI, Queensland University of Technology  erin.peterson@qut.edu.au  Research Interests:  Environmental monitoring assessment and reporting  Spatial networks  Spatial statistics</p> <p>Her work on spatially explicit metrics and water quality</p> <hr/> <p>Jay Zarnetske, PhD  Associate professor, Michigan State University  jtz@msu.edu  Research interest : how catchment hydrology and groundwater - surface water interactions affect aquatic ecosystems through solute generation, fate, and transport</p> <hr/> <p>Ian A Thomas, PhD  Post Doc, University College Dublin  ian.thomas@ucd.ie  Specialist of critical source areas ;agricultural diffuse pollution; hydrological modelling</p> <hr/> <p>Philip Haygarth, PhD  Professor, Lancaster Environment Centre  p.haygarth@lancaster.ac.uk  Professor Phil Haygarth conducts research on the interface between soils and freshwaters, with a focus on diffuse (particularly phosphorus) pollution and runoff control in a catchment context.</p> <p>Numerous works on scaling the phosphorus signal from land to aquatic ecosystems.</p>

[Click here to view linked References](#)

## 1 **Title**

2 The influence of landscape spatial configuration on nitrogen and phosphorus exports in  
3 agricultural catchments

## 4 **Authors**

5 Antoine Casquin. INRAE, UMR SAS 1069 / L'Institut Agro, 65 rue de St Brieuc, 35200  
6 Rennes, France. [antoine.casquin@inrae.fr](mailto:antoine.casquin@inrae.fr) / [a.casquin@gmail.com](mailto:a.casquin@gmail.com). [https://orcid.org/0000-](https://orcid.org/0000-0001-6897-0183)  
7 0001-6897-0183

8 Rémi Dupas. INRAE, UMR SAS 1069 / L'Institut Agro, 65 rue de St Brieuc, 35200 Rennes,  
9 France. [remi.dupas@inrae.fr](mailto:remi.dupas@inrae.fr). <https://orcid.org/0000-0002-5932-4056>

10 Sen Gu. OSUR, Géosciences Rennes, CNRS, UMR 6118, Campus de Beaulieu, 35042  
11 Rennes, France / Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan 430072,  
12 China. [gusen@ihb.ac.cn](mailto:gusen@ihb.ac.cn)

13 Ewan Couic. OSUR, Géosciences Rennes, CNRS, UMR 6118, Campus de Beaulieu, 35042  
14 Rennes, France. [ewan.couic@univ-rennes1.fr](mailto:ewan.couic@univ-rennes1.fr). <https://orcid.org/0000-0003-4223-2726>

15 Gérard Gruau. OSUR, Géosciences Rennes, CNRS, UMR 6118, Campus de Beaulieu,  
16 35042 Rennes, France. [gerard.gruau@univ-rennes1.fr](mailto:gerard.gruau@univ-rennes1.fr)

17 Patrick Durand. INRAE, UMR SAS 1069 / L'Institut Agro, 65 rue de St Brieuc, 35200  
18 Rennes, France. [patrick.durand@inrae.fr](mailto:patrick.durand@inrae.fr). <https://orcid.org/0000-0002-0984-693X>

## 19 **Correspondent author:**

20 Antoine Casquin. INRAE, UMR SAS 1069 / L'Institut Agro, 65 rue de St Brieuc, 35200  
21 Rennes, France. [antoine.casquin@inrae.fr](mailto:antoine.casquin@inrae.fr) (institutional) / [a.casquin@gmail.com](mailto:a.casquin@gmail.com)  
22 (permanent). Tel.: +33(0)2 23 48 54 22. Fax: + 33(0)2 23 48 54 30.

## 23 **Abstract**

### 24 **Context**

25 Nitrogen (N) and phosphorus (P) exports from rural landscapes can cause eutrophication of  
26 inland and coastal waters. Few studies have investigated the influence of the spatial  
27 configuration of nutrient sources – i.e. the spatial arrangement of agricultural fields in  
28 headwater catchments – on N and P exports.

### 29 **Objectives**

30 This study aimed to 1) assess the influence of the spatial configuration of nutrient sources on  
31 nitrate (NO<sub>3</sub><sup>-</sup>) and total phosphorus (TP) exports at the catchment scale, and 2) investigate

32 how relationships between landscape composition and landscape configuration vary  
33 depending on catchment size.

#### 34 **Methods**

35 We analysed  $\text{NO}_3^-$  and TP in 19 headwaters (1-14 km<sup>2</sup>, Western France) every two weeks for  
36 17 months. The headwater catchments had similar soil types, climate, and farming systems  
37 but differed in landscape composition (percentage agricultural land-use) and spatial  
38 configuration. We developed a landscape configuration index (LCI) describing the spatial  
39 configuration of nutrient sources as a function of their hydrological distance to streams and  
40 flow accumulation zones. We calibrated the LCI's two parameters to maximise the rank  
41 correlation with median concentrations of TP and  $\text{NO}_3^-$ .

#### 42 **Results**

43 We found that  $\text{NO}_3^-$  exports were controlled by landscape composition, whereas TP exports  
44 were controlled by landscape configuration. For a given landscape composition, landscape  
45 spatial configuration was highly heterogeneous at small scales (<10 km<sup>2</sup>) but became  
46 homogeneous at larger scales (>50 km<sup>2</sup>).

#### 47 **Conclusions**

48 The spatial configuration of nutrient sources influences TP but not  $\text{NO}_3^-$  exports. An ideal  
49 placement of mitigation options to limit TP export should consider both the hydrological  
50 distance to streams and flow accumulation zones

51

#### 52 **Keywords**

53 Eutrophication; landscape pattern; landscape organisation; landscape composition;  
54 heterogeneity; scaling

55

## 56 **1. Introduction**

57 Excessive loads of nitrogen (N) and phosphorus (P) cause the eutrophication of marine and  
58 freshwater ecosystems (Dodds and Smith 2016; Le Moal et al. 2018), which threatens  
59 biodiversity and human activities (Steffen et al. 2015). In streams and rivers that drain  
60 agricultural landscapes, N and P originate mainly from fertilised agricultural fields. Water  
61 quality mitigation measures often result in reduced yields or increased production costs;  
62 hence, they decrease profitability for farmers (Withers et al. 2014). It is therefore necessary  
63 to locate mitigation options such as set-aside areas and buffer zones where they will be most  
64 effective (Cole et al. 2020). Mechanistic models (Casal et al. 2019; McDowell et al. 2014)  
65 and methods based on statistical analysis of landscape properties (Doody et al. 2012;  
66 Hashemi et al. 2018) have been used to optimise locations of mitigation practices.

67 Most N and P enter the hydrographic network in headwater streams (Dodds and Oakes  
68 2008), which represent 90% of global stream length (Downing 2012) but are rarely  
69 monitored, except for research purposes (Bishop et al. 2008). At this scale, nutrient exports  
70 vary greatly in space (Abbott et al. 2018; Temnerud and Bishop 2005). While in large (>100  
71 km<sup>2</sup>) catchments, both N and P exports can be statistically related to proxies of agricultural  
72 pressure intensity, such as percentage of agricultural land-use or agricultural surplus (Dupas  
73 et al. 2015; Goyette et al. 2018), these relationships break down at the scale of headwater  
74 catchments (<20 km<sup>2</sup>) (Bol et al. 2018; Burt and Pinay 2005). The reason why the  
75 relationship between landscape composition and nutrient exports is scale-dependent is  
76 unclear. Identifying the factor(s) responsible for this loss of relationship, and the  
77 characteristic spatial scale at which it occurs, would help guide water quality mitigation  
78 measures at the catchment scale. In this article, we explore the hypothesis that the spatial  
79 configuration of nutrient sources – i.e. the spatial arrangement/distribution/organisation of  
80 agricultural fields, hereafter called “landscape configuration”, –influences N and P exports at  
81 the headwater catchment scale. For a given region, landscape configuration can be highly  
82 heterogeneous among headwater catchments but homogeneous among larger catchments  
83 (Abbott et al. 2018; Temnerud and Bishop 2005). If verified, these two hypotheses could  
84 explain the breakdown of the relationship between agricultural pressure intensity and nutrient  
85 exports at the headwater catchment scale.

86 Several authors have reviewed the influence of landscape spatial configuration on nutrient  
87 loads at multiple scales (Lintern et al. 2018; Uemaa et al. 2007). Metrics used in landscape  
88 ecology are based on the area/density/edge, shape, isolation, interspersion, and connectivity  
89 of patches, and they have been applied to predict water quality parameters at the catchment  
90 scale (Liu et al. 2020; Shi et al. 2013; Xiao et al. 2016). However, these approaches often  
91 rely on regressions of several landscape metrics and water quality parameters, which risks

92 over-fitting certain relationships and/or spurious correlations. Landscape spatial configuration  
93 can also be understood as the overlap and proximity of features such as streams,  
94 depressions, and flow accumulation areas to land-use patches (e.g. crops, forest fragments,  
95 urban fabric). In line with this concept, Peterson et al. (2011)) related parameters of stream  
96 ecological conditions to multiple spatially explicit landscape “topological” metrics and found  
97 that these spatially explicit methods clearly had more predictive power than landscape  
98 composition metrics. Staponites et al. (2019)), using similar metrics, suggested that the  
99 spatial organisation of nutrient sources (i.e. landscape configuration) influenced the transfer  
100 of reactive or labile water quality parameters (e.g. TP,  $\text{NO}_2^-$  and  $\text{PO}_4^{3-}$ ), whereas percentages  
101 of land use (i.e. landscape composition) have more predictive power for more temporally  
102 stable water quality parameters (e.g.  $\text{Ca}^{2+}$ ,  $\text{NO}_3^-$ , conductivity). These results are consistent  
103 with current knowledge on transfer pathways of nutrients in catchments, with deeper  
104 flowpaths for N species than for P species (Strohmer et al. 2020), and explains why  
105 landscape features had less influence on  $\text{NO}_3^-$  than P (Thomas and Abbott 2018). These  
106 landscape metrics, however, are relatively rigid, as topological influences (e.g. hydraulic  
107 distance to stream, surface flow accumulation, slope), whether considered in the metrics or  
108 not, cannot be weighted. In addition, certain topological features vary over a much wider  
109 range of values than others, which may obscure other features when no weighting coefficient  
110 is included in the landscape metric. For example, flow accumulation (which can have large  
111 values) can overshadow the influence of slope (which varies over a narrow range of values)  
112 in an index that considers both without weighting them.


113 The idea that landscape configuration influences N and P loads is the basis for the concept  
114 of critical source area (CSA), i.e. the idea that a small percentage of the agricultural area  
115 (e.g. <20%) contributes disproportionately to the nutrients transferred to streams (e.g.  
116 >80%). CSAs are defined as the intersection of nutrient sources and hydrologically sensitive  
117 areas (Gburek and Sharpley 1998; Pionke et al. 2000). Initially defined at the sub-field scale,  
118 the concept of CSA was extended to larger scales; entire fields or subcatchments can also  
119 be CSAs (Buchanan et al. 2013b; Page et al. 2005; Reaney et al. 2019; Sharpley et al. 2011;  
120 Srinivasan and McDowell 2009). Substantial uncertainties remain, however, in their  
121 delineation at all scales (Doody et al. 2012). Validation at the sub-field scale can be based on  
122 observing erosion marks (Reaney et al. 2019) or tracers (Collins et al. 2012). However,  
123 studies that validate CSA delineation based on N or P concentrations in streams and rivers  
124 are rare and are based on only a few hydrologically contrasting headwater catchments  
125 (McDowell and Srinivasan 2009; Shore et al. 2014; Thomas et al. 2016a) or larger  
126 catchments for which land-use composition is already a good predictor (Giri et al. 2018).

127 To address the limits of “expert-based” delineation of CSAs and the rigidity of spatially  
128 explicit landscape metrics, we developed a stochastic, data-driven approach based on 30  
129 synoptic samplings of 19 agricultural headwater catchments to answer two questions: i) Does  
130 landscape configuration influence N and P exports?; and ii) Does the location of CSAs  
131 depend on the hydrological distance of the nutrient source to the hydrological network, their  
132 overlap with flow accumulation areas, or both? We then investigated why some relationships  
133 between water quality parameters and landscape composition metrics break down below a  
134 certain catchment size. For this, we studied how the relationship between landscape  
135 composition and configuration varied as a function of catchment size in 500 randomly  
136 selected subcatchments in the study area.

## 137 **2. Methods**

### 138 **2.1. Study site**

139 The Yvel catchment is a 375 km<sup>2</sup> agricultural catchment of Strahler order 5 in Brittany,  
140 western France (Fig. 1). The Yvel River is the main tributary of a three million m<sup>3</sup> water  
141 reservoir that has been subjected to cyanobacteria blooms since the 1970s (ODEM 2012),  
142 for which N and P are deemed responsible (Shatwell and Köhler 2018).



143

144 **Fig. 1** Monitored subcatchments, land use in 2018, hydrography and roads in the Yvel  
 145 catchment. The inset shows the location of Brittany, France, in western Europe

146 The climate is temperate oceanic with mean annual precipitation (1998-2017) of 777 mm  
 147 (sd=132 mm), mean annual temperature of 11.7°C (sd=0.5°C), and mean annual runoff of  
 148 254 mm (sd=143 mm). The Yvel River's discharge is monitored for 300 km<sup>2</sup> of its 375 km<sup>2</sup>,  
 149 and monthly mean discharge varies from 157 L.s<sup>-1</sup> in August to 5,600 L.s<sup>-1</sup> in February. The  
 150 hydrology is controlled by the dynamics of the shallow groundwater within unconsolidated  
 151 weathered material that caps impervious schist bedrock (Casquin et al. 2020). The land use  
 152 is dominated by arable fields (maize and winter cereals), which cover 54% of the catchment  
 153 (Fig. 1). Grasslands (21%, mainly leys in rotation), forests (18%), and urban areas (6%)  
 154 comprise the rest of the catchment area. Hedgerow density is 71 m.ha<sup>-1</sup>. Soils in the  
 155 catchment are generally shallow (<100 cm), are well-drained in the upland part of the  
 156 hillslope, and are often hydromorphic in valley bottoms. The elevation varies from 33-297 m  
 157 above sea level. The centre of the catchment is the flattest area (most slopes <5%), with long

158 and regular hillslopes. In the north and south, the relief is more rugged, with shorter and  
159 steeper slopes. The southeast portion of the watershed of the catchment is forested and has  
160 the steepest slopes (5-15%).

## 161 2.2. Subcatchment monitoring

162 The monitoring strategy consisted of repeated synoptic sampling of 19 subcatchments (Fig.  
163 1). The 19 subcatchments were selected based on Strahler order (1-2), size (0.8-12.6 km<sup>2</sup>,  
164 mean=5.1 km<sup>2</sup>), absence of a wastewater treatment plant, and accessibility. Their  
165 percentage of agricultural land-use ranged from 17-94% (mean=74%), with mean slopes  
166 ranging from 2.7-6.6% (mean=4.9%). Together, these 19 subcatchments covered 28% of the  
167 Yvel catchment's area. All 19 monitoring points were sampled approximately every two  
168 weeks from April 2018 to July 2019 (30 dates in total). Samples were filtered *in situ*  
169 immediately after sampling with cellulose acetate filters of 0.20 µm pore size for nitrate (NO<sub>3</sub><sup>-</sup>)  
170 analysis. All filters were rinsed in the laboratory with 20 ml of deionised water before use. An  
171 unfiltered water sample was also collected to analyse TP. The samples were transported to  
172 the laboratory in a cool box and then refrigerated at 4°C until analysis. TP was analysed  
173 within 48 h of sampling, while NO<sub>3</sub><sup>-</sup> was analysed within one week. TP was determined  
174 colorimetrically via reaction with ammonium molybdate (Murphy and Riley 1962) after  
175 digesting the samples in acidic potassium persulfate. The precision of TP measurements  
176 was ±13 µg.L<sup>-1</sup>, while that of NO<sub>3</sub><sup>-</sup> concentrations, analysed by ionic chromatography  
177 (Dionex, DX120), was ±4%. Hydrochemical data are available at  
178 <https://www.hydroshare.org/resource/7c7d7f6dd1f14450883ae1c243c3c28f/> (Dupas 2020).


## 179 2.3. Landscape configuration index

180 We developed the landscape configuration index (LCI) (Eq. 1) to test the hypothesis that the  
181 hydrological proximity of agricultural areas to watercourses and their overlap with flow  
182 accumulation areas influence nutrient exports at the headwater catchment scale. For each  
183 monitored subcatchment, the index was calculated as follows:

$$184 \quad LCI(a,b) = \frac{\frac{1}{n} \times \sum_{i=1}^n LU_i \times FAcc_i^a \times FLS_i^{-b}}{\frac{1}{N} \times \sum_{i=1}^N LU_i \times FAcc_i^a \times FLS_i^{-b}} \quad (Eq. 1)$$

185 where a and b are calibrated parameters, i=1..n are the pixels of a subcatchment; i=1..N are  
186 the pixels of the entire catchment, FAcc<sub>i</sub> is the flow accumulation on pixel i (m<sup>2</sup>), FLS<sub>i</sub> (m) is  
187 the distance along the surface flow line from pixel i to the stream/ditch, and LU<sub>i</sub> equals 1 if  
188 pixel i is a source of nutrients (i.e. an agricultural area), otherwise 0.

189 Fig. 2 details the calculations for two subcatchments with similar land-use composition, but  
 190 whose LCI numerator varies by a factor of 2 when parameters (a, b) = (1.5, 1).



191  
 192 **Fig. 2** Steps used to calculate the index given a=1.5 and b=1.0 for 2 of the 19 monitored  
 193 subcatchments. Note the logarithmic colour scale

194 The denominator is a normalisation factor that corresponds to the mean value of the  
 195 numerator for the entire Yvel catchment. When LCI > 1, a subcatchment's nutrient sources  
 196 are located predominantly in flow accumulation zones and/or near streams. For large values  
 197 of a or b, the LCI assigns high weights to a small percentage of the area. We restricted the  
 198 ranges of parameters a and b so that a few pixels with the highest Facc and lowest FLS  
 199 would not control the values of LCI. We explored the parameter space for pairs of (a, b) for  
 200 which the highest 5% of cumulative LCI in the landscape did not exceed 95% of the overall  
 201 cumulative LCI. When b=0, we varied a within the range 0-2, whereas when a=0, we varied b  
 202 within the range 0-4.

203 By construction, when a and b equal 0 for a subcatchment, its LCI equals its percentage of  
204 agricultural land-use (i.e. landscape composition) divided by the percentage of agricultural  
205 land-use in the entire Yvel catchment. For other values of a and b, the LCI indicates the  
206 landscape configuration. High values of parameter a increase the weight of pixels in flow  
207 accumulation zones in the LCI, while high values of parameter b increase the weight of pixels  
208 in near-stream zones.

## 209 **2.4. Spatial data sources and pre-processing**

### 210 **2.4.1. Hydrographic network**

211 The hydrographic network consisted of both the “natural” stream network and the ditch  
212 network because 1) it is often difficult to distinguish a ditch from a rectified stream and 2) our  
213 synoptic sampling verified that most of the ditches were deep (> 1 m) and flowed in winter  
214 (high-flow season). Ditch networks are a shortcut between agricultural areas and the  
215 "natural" river network (Ahiablame et al. 2011; Buchanan et al. 2013a). Moreover, evidence  
216 suggests that ditches act as 1<sup>st</sup> order streams when considering nutrient spiralling and can  
217 retain and remobilise N and P (Dunne et al. 2007; Hill and Robinson 2012; Smith 2009).  
218 Thus, we considered them part of the hydrographic network. We used the road network as a  
219 proxy for the ditch network, as we observed that ditches bordered all roads in the study area.  
220 Streams (permanent and intermittent) and roads were extracted from vector data (accuracy  
221 of ca. 1.5 m) provided by the *Institut National de l'Information Géographique et Forestière*  
222 (IGN) at 1:25,000 scale. We converted these vector data to raster format and aligned them  
223 with the Digital Elevation Model (DEM) for later analysis.

### 224 **2.4.2. Digital Elevation Model**

225 The DEM, with a native resolution of 5 m (IGN 2018), was resampled to 10 m using cubic  
226 splines in ArcGIS 10.6. Filling was used to remove the depressions on hillslopes (Planchon  
227 and Darboux 2002), and the value "NA" was assigned to the pixels corresponding to roads  
228 and streams. We calculated flow accumulation and the hydrological distances to streams (i.e.  
229 following the surface flow paths) using the multiple flow direction algorithm (Qin et al. 2007).  
230 We chose this algorithm for its ability to generate realistic flow accumulation maps, unlike the  
231 D8 algorithm (O'Callaghan and Mark 1984).

### 232 **2.4.3. Nutrient sources**

233 We extracted the agricultural area from the national Land Parcel Identification System  
234 (Levavasseur et al. 2016). The data are provided as a vector dataset at the 1:5000 scale for  
235 each year since 2010 and contain field boundaries and a code that identifies the crop type.  
236 We used the 2018 dataset and verified the spatial accuracy of the agricultural area  
237 boundaries based on 50 cm orthophotos and the hydrography. We then rasterized this

238 dataset aligned with the DEM. We assigned the value 1 to agricultural areas and 0 to non-  
239 agricultural areas. We included riparian buffer strips in agricultural areas because they are  
240 recent and have been fertilised for years, and are a well-documented legacy source of  
241 nutrients in headwater catchments (Gu et al. 2018).

## 242 **2.5. Optimisation of (a, b) parameters and interpretation**

243 For each pair (a, b) within the previously defined domain of exploration, we calculated  
244 Spearman's rank correlation ( $R_s$ ) between the LCI and median  $\text{NO}_3^-$  and TP concentrations  
245 of each subcatchment (Figure 3). Since several of the subcatchments were intermittent, we  
246 calculated the median concentrations for the 22 dates (out of 30) when at least 17 of the 19  
247 subcatchments were flowing, so as not to bias calculation of the median. We focused on  
248 ranks rather than concentrations because several studies have shown that concentration  
249 estimates had high uncertainty when calculated with low-frequency data (e.g. Cassidy and  
250 Jordan 2011; Moatar et al. 2020), while the ranks of subcatchments could be predicted with  
251 high degree of confidence, as they are stable across flow conditions (Abbott et al. 2018;  
252 Dupas et al. 2019, Gu et al., 2020 - in review). The result interpretation was twofold:

- 253 - The spatial configuration of sources was considered to have an effect if  $R_s$  for at least  
254 one pair (a, b)  $\neq$  (0, 0) was significantly higher than that for (a, b) = (0, 0).
- 255 - Optimal values of a and b were examined to assess the relative importance of  
256 hydrological distance to streams and flow accumulation on hillslopes.


257 We then calculated the ratio of LCI with optimal (a, b) to LCI with (a, b) = (0, 0) for 500  
258 randomly selected subcatchments in the Yvel catchment, and analysed how the relationship  
259 between these two metrics evolved as catchment size increased.

260

## 261 **3. Results**

### 262 **3.1. Comparison of land-use composition and configuration metrics as predictors of** 263 **$\text{NO}_3^-$ and TP concentrations in headwater catchments**

264 Landscape composition (i.e. LCI with (a, b) = (0, 0)) predicted median  $\text{NO}_3^-$  concentrations  
265 well ( $R_s=0.84$ ,  $p<0.001$ ) but not those of TP ( $R_s=0.33$ ,  $p=0.18$ ). Varying (a, b) did not  
266 substantially improved the prediction of median  $\text{NO}_3^-$  concentration rank; thus, landscape  
267 composition predicted  $\text{NO}_3^-$  exports well at the headwater catchment scale, and considering  
268 landscape configuration did not improve the prediction (Fig. 3A). For  $\text{NO}_3^-$ , this result refutes  
269 our first hypothesis that the spatial configuration of nutrient sources influences nutrient  
270 exports.




272

273 **Fig. 3** A) Optimisation plan for parameters a and b used to predict median concentration of  
 274 nitrate ( $NO_3^-$ ) and total phosphorus (TP) in 19 headwater catchments. Rank correlations ( $R_s$ )  
 275 not shown (blank) indicate (a, b) outside of the exploration domain. For each optimisation  
 276 plan, black dots indicate 10% highest  $R_s$ , red square the best correlation B)  $R_s$  between  
 277 median  $[TP]$  and 1) landscape composition and 2) optimised landscape configuration index  
 278 (LCI). Annotations show LCI:median  $[TP]$

279 The correlation between the percentage of agricultural land and median TP concentrations  
280 ((a, b) = (0, 0)) was 0.35 ( $p=0.145$ ) (Fig. 3B). This correlation improved as weights increased  
281 for sources near streams ( $R_s$  increased as  $b$  increased) and for sources that overlapped  
282 surface flow accumulation ( $R_s$  increased as  $a$  increased) (Fig. 3A). This confirms our first  
283 hypothesis that landscape configuration influences P exports, and agricultural areas near  
284 watercourses and that overlap surface flow accumulation areas result in larger P exports.  
285 The optimum correlation was obtained for  $a=1.4$  and  $b=2.2$  ( $R_s=0.81$ ,  $p<0.001$ ), at the limit of  
286 our field of exploration, which means that 5% of the area determined nearly 95% of the  
287 index.

288 Using the same parameter exploration scheme, we optimised (a, b) for TP concentrations on  
289 each sampling date (Fig. 4C). The LCI with (a, b)  $\neq$  (0, 0) always predicted TP concentrations  
290 better than landscape composition (Fig. 4A). The correlation between the optimised LCI and  
291 TP concentrations differed significantly ( $p<0.05$ ) from 0 for all but one sampling date, except  
292 during the low-flow periods (Fig. 4B).



293


294 **Fig. 4** Comparison of landscape composition and optimised landscape configuration index  
 295 (LCI) as predictors of TP concentrations, for all sampling dates. (a) Spearman's rank  
 296 correlation (Rs), (b)  $-\log_{10}$ -transformed p-values (dashed line indicates  $p=0.05$ ), and (c) for  
 297 optimised LCI, parameters a and b applied to flow accumulation and inverse distance to the  
 298 stream/ditch (d). Relative weight of the top 5% of weighted pixels. Grey areas represent low-  
 299 flow periods, when two or more sampled streams were dry

300 During the low-flow season (Aug-Dec 2018) and beginning of the rewetting season (Jul  
 301 2019), TP concentrations in the headwaters were not correlate with the optimised LCI  
 302 ( $p \gg 0.05$ ) (Fig. 4B). Note that 13 of the 19 streams were dry at the peak of the low-flow  
 303 season (Fig. 1S). Outside the low-flow season, optimal values of parameters a and b were  
 304 stable, varying little around a median value of 1.4 and 2.2, respectively. Therefore, the total  
 305 weight of the top 5% of weighted area remained close to 95% (Fig. 4D), except during the  
 306 low-flow season. The optimised LCI based on median TP concentrations, hereafter referred  
 307 to as LCI-TP, appears to be a robust sensitivity index to determine TP CSAs across flow  
 308 conditions.

### 309 **3.2. Spatial aggregations of the LCI-TP**

310 The data-driven optimization of the LCI parameters to predict the ranking of exports at the  
311 catchment scale yielded an index that assigned nearly all of the weights (95%) to small areas  
312 (5%). Most of these areas (Fig. 5A) were located near streams and ditches, but their width  
313 varied. The CSAs often extended ca. 100 m or more into agricultural areas, especially on  
314 long and convex hillslopes. Using LCI-TP values at the pixel level, we calculated  $\text{LCI-TP}_{\text{field}}$   
315 as the mean LCI-TP of each agricultural field. The  $\text{LCI-TP}_{\text{field}}$  (Fig. 5B) revealed that most  
316 fields were weak sources of P (i.e.  $\text{LCI-TP}_{\text{field}} \ll 1$ ), whereas a few fields were CSAs, as their  
317  $\text{LCI-TP}_{\text{field}}$  exceeded 10 and even 40. The associated histogram (Fig. 5C) followed a  
318 lognormal distribution, which confirms the high variability at the field scale ( $Q5-Q95=0.001-$   
319  $14.087$ ,  $sd=39.05$ ). When the LCI-TP was aggregated into  $1 \text{ km}^2$  subcatchments ( $\text{LCI-TP}_{1\text{km}^2}$ )  
320 (Fig. 5D), its variability decreased drastically (Fig. 5F,  $\text{min-max}=0.17-5.38$ ,  $sd=0.49$ ). The  
321 number of subcatchments with  $\text{LCI-TP}_{1\text{km}^2} > 1$  was approximately the same as the number of  
322 subcatchments with  $\text{LCI-TP}_{1\text{km}^2} < 1$  (Fig. 5F,  $\text{median}=0.92$ ), but the distribution remained  
323 lognormal. Following the same pattern, the LCI-TP aggregated into  $25 \text{ km}^2$  subcatchments  
324 ( $\text{LCI-TP}_{25\text{km}^2}$ ) had even lower variability ( $\text{min-max}=0.71-1.36$ ): at this scale the distribution  
325 was symmetrical (apparently normal), and no subcatchment could be considered a CSA. The  
326 information at the subfield scale (Fig. 5A) and field scale (Fig. 5B), which is relevant for  
327 farmers and catchment managers, was generated for the entire study area ([https://antoine-](https://antoine-csgn.github.io/YV1.html)  
328 [csgn.github.io/YV1.html](https://antoine-csgn.github.io/YV1.html)).

329




330

331 **Fig. 5** A) Excerpt of the study area, ca. 6 km × 2 km: top 5% (yellow) and 1% (red) of  
 332 weighted pixels according to LCI-TP, B) Mean LCI-TP aggregated at the field scale (LCI-TP  
 333 *field*), and C) histogram of associated values (note the log scale on the x-axis). D) LCI-TP for 1  
 334 km<sup>2</sup> subcatchments (LCI-TP<sub>1km<sup>2</sup></sub>) and E) 25 km<sup>2</sup> subcatchments (LCI-TP<sub>25km<sup>2</sup></sub>) and associated  
 335 histograms (F and G, respectively)

336 **3.3. Homogenisation of the LCI-TP with increasing catchment size**

337 We delineated 500 subcatchments within the study area based on 500 points randomly  
338 generated over the hydrographic network. For each subcatchment, we calculated the LCI-TP  
339 (i.e. the LCI for  $a=1.4$  and  $b=2.2$ ), the landscape composition index (i.e. the LCI for  $a=b=0$ ),  
340 and their ratio, which we examined as a function of subcatchment area (Fig. 6).



341  
342 **Fig. 6** Stabilisation of the ratio of the LCI-TP (landscape configuration index (LCI) for  $a=1.4$   
343 and  $b=2.2$ ) to landscape composition index (LCI for  $a=b=0$ ) as a function of area (square-  
344 root-transformed x-axis) of subcatchments in the study area (points)

345 This ratio varied greatly (0.02-3.37) for subcatchments smaller than 10 km<sup>2</sup>, the typical size  
346 of 1<sup>st</sup> or 2<sup>nd</sup> order stream catchments but varied less (0.55-1.53) for subcatchments of 10-50  
347 km<sup>2</sup>. The high variability in this ratio, even for heavily farmed subcatchments, indicates that  
348 the landscape can still be optimised at this scale. For catchments larger than 50 km<sup>2</sup>, which  
349 corresponds to 4<sup>th</sup> order rivers in the study area, the ratio converged to 1. The main  
350 implication is that for headwater catchments (<50 km<sup>2</sup>) the correlation between landscape  
351 composition and configuration (as defined by the LCI-TP) is weak and non-significant (Fig.  
352 2S), while for larger catchments the correlation is strong ( $R^2$  near 1) (Fig. 2S).

353  
354 **4. Discussion**

355 **4.1. Spatial configuration of nutrient sources influences P but not N exports**

356 Landscape composition is a strong predictor of median N concentrations, while landscape  
357 configuration as defined here, seems to have no influence (Fig. 2A). This result agrees with  
358 previous research that indicates a strong correlation between metrics of agricultural N  
359 pressure intensity, such as N surplus or N concentrations in streams (Dupas et al. 2015;

360 Kronvang et al. 2005), leaving little space for other controlling factors, such as landscape  
361 configuration. However, distributed process-based modelling of N fate in agricultural  
362 catchments has shown that landscape configuration can have a of second-order influence on  
363 N concentrations (Beaujouan et al. 2002; Casal et al. 2019; McDowell et al. 2014), which  
364 may not have been captured in our study. P concentrations are not related to landscape  
365 composition (Bol et al. 2018) or even P inputs (Dupas et al. 2015; Frei et al. 2020) in  
366 headwater catchments. Multiple factors influence P transfers to streams: soil type and P  
367 content (Djodjic et al. 2004), tile-drainage (King et al. 2015a; King et al. 2015b), small ponds  
368 and hillside storage reservoirs (Schmadel et al. 2019), in-hillslope depressions (Smith and  
369 Livingston 2013), leaks from septic tanks (Withers et al. 2011) and livestock buildings, bank  
370 erosion (Kronvang et al. 2012), and ditch-dredging management (Smith and Pappas 2007).  
371 While hedgerows have little influence on N transfers at the catchment scale (Benhamou et al.  
372 2013; Thomas and Abbott 2018), it is difficult to quantify their combined effects on P  
373 transfers. Without considering these factors, our optimised LCI ranked the headwater  
374 catchments reasonably well according to their P exports (Fig. 2B), although the residuals  
375 may have been due to these other factors. We demonstrate that the spatial configuration of P  
376 sources in the landscape is critical to understand P transfer from land to streams in  
377 headwater catchments. The influence of the spatial configuration of nutrient sources on P but  
378 not N concentrations in agricultural headwater streams results in high variability in N:P ratios  
379 in these ecosystems and has many consequences on stream algae communities (Pringle  
380 1990; Stelzer and Lamberti 2001).

381 The observation that landscape spatial organisation influences N and P transfers is not new.  
382 Nonetheless, our approach differs from previous studies that used single or multiple  
383 regression of several landscape metrics compiled in FRAGSTATS (McGarigal and Marks  
384 1995) to predict nutrient exports at the catchment scale. Despite the large body of studies  
385 that use these landscape metrics (Bu et al. 2014; Lee et al. 2009; Ouyang et al. 2014; Zhang  
386 et al. 2019), no robust cause-effect relationships emerge, and many of them give conflicting  
387 results (Wang et al. 2020). Three reasons may explain why these studies, taken together,  
388 are not conclusive: i) the many pairwise or multiple regressions between water quality  
389 parameters and landscape metrics generate spurious correlations that are not generally  
390 discussed, ii) landscape metrics depend greatly on the resolution of GIS input data, and iii)  
391 the studies do not consider the topological dimension, which is fundamental for explaining  
392 hillslope-to-stream transfers and landscape configuration (Thomas et al. 2016b). Several  
393 approaches have integrated this topological control with the hypothesis that nutrient sources  
394 near streams (Sliva and Dudley Williams 2001; Yates et al. 2014) and/or that overlap flow  
395 accumulation areas (Peterson et al. 2011; Staponites et al. 2019) have a disproportionately

396 higher influence on nutrient exports than other areas. While these approaches still depend on  
397 the spatial resolution of GIS data, they are “hypothesis-driven” rather than purely “data-  
398 driven”, which decreases the risk of spurious correlations. A longstanding weakness of these  
399 approaches, however, is their rigidity due to the lack of calibrated parameters. For example,  
400 the HAiFLS index (Peterson et al. 2011) and Flow-A index (Staponites et al. 2019) included  
401 both the distance to streams and flow accumulation, but the latter dominates the index value  
402 by construction, because it varies more than the former, and neither index allows both factors  
403 to be weighted by calibrated coefficients. The stochastic approach in the LCI developed here  
404 is more flexible, and we show that both hydrological distance and flow accumulation  
405 influence P transfer.

#### 406 **4.2. Spatial variability and temporal stability of critical source areas**

407 The strongest correlation between median  $\text{NO}_3^-$  concentrations and LCI was for  $(a, b) = (0,$   
408  $0)$ , which shows that each source (agricultural area) contributed the same, regardless of its  
409 distance to streams or overlap with flow accumulation areas. This confirms the need to  
410 consider the entire catchment to reduce N loads in agricultural catchments. For median P  
411 concentrations, the optimal was found for  $(a, b) = (1.4, 2.2)$ , a value for which 5% of the  
412 agricultural area concentrated 95% of the weight assigned by our index at the 10 m-pixel  
413 resolution. These values are similar to those found by Thomas et al. (2016a)), who classified  
414 1.6-3.4% of the catchment area (during median storm events) and 2.9-8.5% (during upper-  
415 quartile events) as prone to P transfer, based on a CSA model that also considered land use  
416 and topography as input variables. Summing the LCI-TP at the field scale indicates that the  
417 20% of fields at highest risk represent 85% of the total weights, which are the fields on which  
418 mitigation measures should be prioritised. The distribution is asymmetrical, with 69% of fields  
419 having a mean  $\text{LCI-TP}_{\text{field}}$  less than 1. This information at multiple scales can be a tool to  
420 maximise ecosystem services at the catchment scale, by reorganising landscapes to  
421 decrease P transfer without increasing the percentage of set-aside areas (Doody et al.  
422 2016).

423 The shape and location of the sub-field CSAs overlap both the mandatory riparian buffer  
424 strips (5 m according to local application of the European Union Nitrate Directive (DREAL  
425 2018)) and in-hillslope CSAs based on the Topographical Wetness Index. Because most  
426 buffer strips in the study area were installed recently, we included them in P sources as they  
427 were enriched in P before conversion (Dodd and Sharpley 2016; Gu et al. 2017; Gu et al.  
428 2018; Jarvie et al. 2017; Roberts et al. 2012). The location and shape of the CSAs indicate  
429 the need for new shapes of buffer zones, with variable widths and locations along ditches as  
430 well. Because buffer strips are critical sources of nutrients at the headwater catchment scale,  
431 they require new management practices. Potential solutions include sowing species that can

432 capture more P (William M. Roberts 2019), mowing and exporting the residues each year  
433 (Fiorellino et al. 2017), and applying amendments that have high P-sorbing ability (Borno et  
434 al. 2018).

435 The temporal stability of the optimal parameters (a, b) for TP during the flow period is  
436 consistent with the spatial stability concept (Abbott et al. 2018), but is contrary to the concept  
437 of variable source areas (Collick et al. 2015; Dahlke et al. 2012). Our interpretation of this  
438 temporal stability is that even though the soil-to-stream connectivity varied temporally during  
439 the sampling period, the TP concentrations observed may reflect the remobilisation of  
440 sediments transferred during rare erosion events. Observing storm events that result in  
441 surface transfers requires frequent observations (Cassidy and Jordan 2011). During the low-  
442 flow dates, the correlations with LCI-TP were not significant: different sources and sinks likely  
443 dominate the influence of agricultural areas and their configuration, which predominates  
444 during the flow period. Leaks from septic tanks or animal buildings and desorption from  
445 sediments are the most likely sources of P during this ecologically sensitive period, while  
446 hydrological disconnection, uptake, and sedimentation can be sinks of P (Lannergård et al.  
447 2020; Sandstrom et al. 2020).

#### 448 **4.3. Landscape homogenisation with increasing catchment size: consequences for** 449 **management, monitoring, and modelling**

450 When means were aggregated into 1 km<sup>2</sup> subcatchments, which is the typical size of a 1<sup>st</sup>  
451 order catchment in the study area, the LCI-TP showed high spatial variability (Fig. 5D, 5F).  
452 As mentioned, this variability decreased as the aggregation size increased (Fig. 5E, 5G), and  
453 landscape composition was weakly correlated with landscape configuration for catchments  
454 smaller than 50 km<sup>2</sup> but strongly correlated for those larger than 50 km<sup>2</sup> (Fig. 6, Fig. 2S).  
455 These results explain, at least partially, why relationships between water quality parameters  
456 and landscape composition metrics break down below a certain catchment size (Bol et al.  
457 2018). They also shed light on the long-standing difficulty in scaling from nutrient export  
458 models at the field scale to those at the catchment scale: the spatial configuration of the  
459 nutrient sources can be critical.

460 The variability in the LCI-TP in small subcatchments and its homogenisation in larger  
461 catchments is the expression of a degree of unstructured heterogeneity (i.e. randomness) of  
462 the spatial configuration of agricultural areas (Musolff et al. 2017). Some almost entirely  
463 agricultural subcatchments have a TP sensitivity index less than 1, while some mixed-land-  
464 use catchments have a TP sensitivity index greater than 1. The latter provides the  
465 opportunity to introduce structured heterogeneity, i.e. to reorganise agricultural activities  
466 spatially to reduce P transfers to streams (Musolff et al. 2017).

467 These results have implications for both modelling and monitoring. Semi-distributed models  
468 that simulate P exports from an entire catchment with simulation units smaller than 50 km<sup>2</sup>  
469 should include a coefficient to represent the spatial configuration of the agricultural areas. A  
470 distributed model should consider ditches (as they are the entry point of P into the  
471 hydrographic network) and, especially, the spatial variability in P sources. When monitoring  
472 subcatchments that are smaller than the homogenisation threshold and have similar  
473 agricultural land-use composition, the observed differences in P loads cannot be related  
474 directly to the agricultural practices or soil properties. The spatial configuration of agricultural  
475 areas appears to exert a major control on median P concentrations. This is particularly  
476 important for targeting measures to the most cost-effective fields (Doody et al. 2016) and  
477 could increase in importance due to the recent development of innovative financial tools to  
478 improve water quality, such as payment for ecosystem services schemes, whose obligation  
479 to achieve results is increasing (Hejnowicz et al. 2014). Our research provides a data-driven  
480 method to identify CSAs and thus the most cost-effective fields on which to implement  
481 mitigation measures.

482 The 50 km<sup>2</sup> landscape homogenisation threshold found in this study is similar to the 18-68  
483 km<sup>2</sup> stream-concentration thresholds found by Abbott et al. (2018) for the Rance and Haut-  
484 Couesnon catchments, also located in western France. We assume that the optimised  
485 coefficients and homogenisation threshold would vary with the topo-climatic conditions and  
486 agricultural landscape characteristics. More research is needed to confirm this connection  
487 between landscape configuration and P loads in different environmental settings. Given the  
488 high heterogeneity of landscape configuration, this research calls for a spatially dense  
489 monitoring network of headwater catchments. We also recommend using the LCI to  
490 investigate the influence of landscape configuration on other water contaminants that are  
491 transferred mainly by surface flow paths, such as pesticides and faecal bacteria.

#### 492 **Declarations**

#### 493 **Funding**

494 The authors were supported by the Interreg project Channel Payments for Ecosystem  
495 Services, funded through the European Regional Development Fund (ERDF).

#### 496 **Conflicts of interest/Competing interests (include appropriate disclosures)**

497 The authors declare that they have no known competing financial interests or personal  
498 relationships that could have appeared to influence the work reported in this paper.

#### 499 **Ethics approval (include appropriate approvals or waivers)**

500 Not applicable

501 **Consent to participate (include appropriate statements)**

502 Not applicable

503 **Consent for publication (include appropriate statements)**

504 All co-authors – A. Casquin, R. Dupas, S. Gu, E. Couic, G. Gruau and P. Durand – have  
505 approved the manuscript and agree with its submission to Landscape Ecology

506 **Availability of data and material (data transparency)**

507 This study is based on a recently published dataset entitled *Repeated synoptic sampling for*  
508 *water chemistry monitoring in the Yvel catchment, northwestern France* available at  
509 <https://www.hydroshare.org/resource/7c7d7f6dd1f14450883ae1c243c3c28f/> (Dupas *et al.*  
510 2020). It contains the hydrochemical data and the geographic information of the study site.

511 **Code availability (software application or custom code)**

512 The code (R script) will be made available along with the article upon acceptance on a github  
513 repository.

514 **Authors' contributions**

515 AC: Conceptualization, Methodology, Formal analysis, Visualization, Writing - original draft.  
516 RD: Conceptualization, Supervision, Funding acquisition. SG: Sample collection and  
517 analysis. EC: Sample collection and analysis. GG: Project administration, Funding  
518 acquisition. PD: Supervision. All authors contributed substantially to revisions.

519

520 **References**

- 521 Abbott BW et al. (2018) Unexpected spatial stability of water chemistry in headwater stream  
522 networks *Ecol Lett* 21:296-308 doi:10.1111/ele.12897
- 523 Ahiablame LM, Chaubey I, Smith DR, Engel BA (2011) Effect of tile effluent on nutrient  
524 concentration and retention efficiency in agricultural drainage ditches *Agricultural*  
525 *Water Management* 98:1271-1279 doi:10.1016/j.agwat.2011.03.002
- 526 Beaujouan V, Durand P, Ruiz L, Arousseau P, Cotteret G (2002) A hydrological model  
527 dedicated to topography-based simulation of nitrogen transfer and transformation:  
528 rationale and application to the geomorphology- denitrification relationship  
529 *Hydrological Processes* 16:493-507 doi:10.1002/hyp.327
- 530 Benhamou C, Salmon-Monviola J, Durand P, Grimaldi C, Merot P (2013) Modeling the  
531 interaction between fields and a surrounding hedgerow network and its impact on  
532 water and nitrogen flows of a small watershed *Agricultural Water Management*  
533 121:62-72 doi:10.1016/j.agwat.2013.01.004
- 534 Bishop K, Buffam I, Erlandsson M, Fölster J, Laudon H, Seibert J, Temnerud J (2008) *Aqua*  
535 *Incognita: the unknown headwaters* *Hydrological Processes* 22:1239-1242  
536 doi:10.1002/hyp.7049
- 537 Bol R et al. (2018) Challenges of Reducing Phosphorus Based Water Eutrophication in the  
538 Agricultural Landscapes of Northwest Europe *Frontiers in Marine Science* 5  
539 doi:10.3389/fmars.2018.00276

540 Borno ML, Muller-Stover DS, Liu F (2018) Contrasting effects of biochar on phosphorus  
541 dynamics and bioavailability in different soil types *Sci Total Environ* 627:963-974  
542 doi:10.1016/j.scitotenv.2018.01.283

543 Bu H, Meng W, Zhang Y, Wan J (2014) Relationships between land use patterns and water  
544 quality in the Taizi River basin, China *Ecological Indicators* 41:187-197  
545 doi:10.1016/j.ecolind.2014.02.003

546 Buchanan B, Easton ZM, Schneider RL, Walter MT (2013a) Modeling the hydrologic effects  
547 of roadside ditch networks on receiving waters *Journal of Hydrology* 486:293-305  
548 doi:10.1016/j.jhydrol.2013.01.040

549 Buchanan BP, Archibald JA, Easton ZM, Shaw SB, Schneider RL, Walter MT (2013b) A  
550 phosphorus index that combines critical source areas and transport pathways using a  
551 travel time approach *Journal of Hydrology* 486:123-135  
552 doi:10.1016/j.jhydrol.2013.01.018

553 Burt TP, Pinay G (2005) Linking hydrology and biogeochemistry in complex landscapes  
554 *Progress in Physical Geography: Earth and Environment* 29:297-316  
555 doi:10.1191/0309133305pp450ra

556 Casal L, Durand P, Akkal-Corfini N, Benhamou C, Laurent F, Salmon-Monviola J, Vertès F  
557 (2019) Optimal location of set-aside areas to reduce nitrogen pollution: a modelling  
558 study *The Journal of Agricultural Science* 156:1090-1102  
559 doi:10.1017/s0021859618001144

560 Casquin A, Gu S, Dupas R, Petitjean P, Gruau G, Durand P (2020) River network alteration  
561 of C-N-P dynamics in a mesoscale agricultural catchment *Sci Total Environ*  
562 749:141551 doi:10.1016/j.scitotenv.2020.141551

563 Cassidy R, Jordan P (2011) Limitations of instantaneous water quality sampling in surface-  
564 water catchments: Comparison with near-continuous phosphorus time-series data  
565 *Journal of Hydrology* 405:182-193 doi:10.1016/j.jhydrol.2011.05.020

566 Cole LJ, Stockan J, Helliwell R (2020) Managing riparian buffer strips to optimise ecosystem  
567 services: A review *Agriculture Ecosystems & Environment* 296  
568 doi:10.1016/j.agee.2020.106891

569 Collick AS et al. (2015) Predicting phosphorus dynamics in complex terrains using a variable  
570 source area hydrology model *Hydrological Processes* 29:588-601  
571 doi:10.1002/hyp.10178

572 Collins AL, Zhang Y, McChesney D, Walling DE, Haley SM, Smith P (2012) Sediment source  
573 tracing in a lowland agricultural catchment in southern England using a modified  
574 procedure combining statistical analysis and numerical modelling *Sci Total Environ*  
575 414:301-317 doi:10.1016/j.scitotenv.2011.10.062

576 Dahlke HE, Easton ZM, Lyon SW, Todd Walter M, Destouni G, Steenhuis TS (2012)  
577 Dissecting the variable source area concept – Subsurface flow pathways and water  
578 mixing processes in a hillslope *Journal of Hydrology* 420-421:125-141  
579 doi:10.1016/j.jhydrol.2011.11.052

580 Djodjic F, Borling K, Bergstrom L (2004) Phosphorus leaching in relation to soil type and soil  
581 phosphorus content *J Environ Qual* 33:678-684 doi:10.2134/jeq2004.6780

582 Dodd RJ, Sharpley AN (2016) Conservation practice effectiveness and adoption: unintended  
583 consequences and implications for sustainable phosphorus management *Nutrient  
584 Cycling in Agroecosystems* 104:373-392 doi:10.1007/s10705-015-9748-8

585 Dodds W, Smith V (2016) Nitrogen, phosphorus, and eutrophication in streams *Inland  
586 Waters* 6:155-164 doi:10.5268/iw-6.2.909

587 Dodds WK, Oakes RM (2008) Headwater influences on downstream water quality *Environ  
588 Manage* 41:367-377 doi:10.1007/s00267-007-9033-y

589 Doody DG, Archbold M, Foy RH, Flynn R (2012) Approaches to the implementation of the  
590 Water Framework Directive: targeting mitigation measures at critical source areas of  
591 diffuse phosphorus in Irish catchments *J Environ Manage* 93:225-234  
592 doi:10.1016/j.jenvman.2011.09.002

593 Doody DG et al. (2016) Optimizing land use for the delivery of catchment ecosystem services  
594 *Frontiers in Ecology and the Environment* 14:325-332 doi:10.1002/fee.1296

595 Downing J (2012) Global abundance and size distribution of streams and rivers *Inland*  
596 *Waters* 2:229-236 doi:10.5268/iw-2.4.502

597 DREAL B (2018) Directive nitrates 5ème programme d'actions en Bretagne.  
598 <http://www.bretagne.developpement-durable.gouv.fr/cinquieme-programme-d->  
599 [actions-regional-directive-r837.html](http://www.bretagne.developpement-durable.gouv.fr/cinquieme-programme-d-)[Online]. Accessed 11 January 2021

600 Dunne EJ, Mckee KA, Clark MW, Grunwald S, Reddy KR (2007) Phosphorus in agricultural  
601 ditch soil and potential implications for water quality *Journal of Soil and Water*  
602 *Conservation* 62:244-252

603 Dupas R, Delmas M, Dorioz J-M, Garnier J, Moatar F, Gascuel-Oudoux C (2015) Assessing  
604 the impact of agricultural pressures on N and P loads and eutrophication risk  
605 *Ecological Indicators* 48:396-407 doi:10.1016/j.ecolind.2014.08.007

606 Dupas R, Minaudo C, Abbott BW (2019) Stability of spatial patterns in water chemistry  
607 across temperate ecoregions *Environmental Research Letters* 14:074015  
608 doi:10.1088/1748-9326/ab24f4

609 Dupas R, S. Gu, A. Casquin, G. Gruau, P. Petitjean, P. Durand (2020) Repeated  
610 synoptic sampling for water chemistry monitoring in the Yvel catchment, northwestern  
611 France *Hydroshare*  
612 [url:http://www.hydroshare.org/resource/7c7d7f6dd1f14450883ae1c243c3c28f](http://www.hydroshare.org/resource/7c7d7f6dd1f14450883ae1c243c3c28f)

613 Fiorellino N, Kratochvil R, Coale F (2017) Long-Term Agronomic Drawdown of Soil  
614 Phosphorus in Mid-Atlantic Coastal Plain Soils *Agronomy Journal* 109:455-461  
615 doi:10.2134/agronj2016.07.0409

616 Frei RJ et al. (2020) Predicting Nutrient Incontinence in the Anthropocene at Watershed  
617 Scales *Frontiers in Environmental Science* 7 doi:10.3389/fenvs.2019.00200

618 Gburek WJ, Sharpley AN (1998) Hydrologic Controls on Phosphorus Loss from Upland  
619 Agricultural Watersheds *Journal of Environmental Quality* 27:267-277  
620 doi:10.2134/jeq1998.00472425002700020005x

621 Giri S, Qiu Z, Zhang Z (2018) Assessing the impacts of land use on downstream water  
622 quality using a hydrologically sensitive area concept *Journal of Environmental*  
623 *Management* 213:309-319 doi:10.1016/j.jenvman.2018.02.075

624 Goyette J-O, Bennett EM, Maranger R (2018) Differential influence of landscape features  
625 and climate on nitrogen and phosphorus transport throughout the watershed  
626 *Biogeochemistry* 142:155-174 doi:10.1007/s10533-018-0526-y

627 Gu S et al. (2017) Release of dissolved phosphorus from riparian wetlands: Evidence for  
628 complex interactions among hydroclimate variability, topography and soil properties  
629 *Sci Total Environ* 598:421-431 doi:10.1016/j.scitotenv.2017.04.028

630 Gu S, Gruau G, Malique F, Dupas R, Petitjean P, Gascuel-Oudoux C (2018) Drying/rewetting  
631 cycles stimulate release of colloidal-bound phosphorus in riparian soils *Geoderma*  
632 321:32-41 doi:10.1016/j.geoderma.2018.01.015

633 Hashemi F, Olesen JE, Hansen AL, Borgesen CD, Dalgaard T (2018) Spatially differentiated  
634 strategies for reducing nitrate loads from agriculture in two Danish catchments *J*  
635 *Environ Manage* 208:77-91 doi:10.1016/j.jenvman.2017.12.001

636 Hejnowicz AP, Raffaelli DG, Rudd MA, White PCL (2014) Evaluating the outcomes of  
637 payments for ecosystem services programmes using a capital asset framework  
638 *Ecosystem Services* 9:83-97 doi:10.1016/j.ecoser.2014.05.001

639 Hill CR, Robinson JS (2012) Phosphorus flux from wetland ditch sediments *Sci Total Environ*  
640 437:315-322 doi:10.1016/j.scitotenv.2012.06.109

641 IGN (2018) RGE ALTI Version 2.0 Les Modèles numériques 3D [url:](https://geoservices.ign.fr/ressources_documentaires/Espace_documentaire/MODELE_S_3D/RGE_ALTI/DL_RGEALTI_2-0.pdf)  
642 [https://geoservices.ign.fr/ressources\\_documentaires/Espace\\_documentaire/MODELE](https://geoservices.ign.fr/ressources_documentaires/Espace_documentaire/MODELE_S_3D/RGE_ALTI/DL_RGEALTI_2-0.pdf)  
643 [S\\_3D/RGE\\_ALTI/DL\\_RGEALTI\\_2-0.pdf](https://geoservices.ign.fr/ressources_documentaires/Espace_documentaire/MODELE_S_3D/RGE_ALTI/DL_RGEALTI_2-0.pdf). Accessed 11 January 2021

644 Jarvie HP, Johnson LT, Sharpley AN, Smith DR, Baker DB, Bruulsema TW, Confesor R  
645 (2017) Increased Soluble Phosphorus Loads to Lake Erie: Unintended  
646 Consequences of Conservation Practices? *J Environ Qual* 46:123-132  
647 doi:10.2134/jeq2016.07.0248

648 King KW, Williams MR, Fausey NR (2015a) Contributions of systematic tile drainage to  
649 watershed-scale phosphorus transport *J Environ Qual* 44:486-494  
650 doi:10.2134/jeq2014.04.0149

651 King KW et al. (2015b) Phosphorus transport in agricultural subsurface drainage: a review *J*  
652 *Environ Qual* 44:467-485 doi:10.2134/jeq2014.04.0163

653 Kronvang B, Audet J, Baattrup-Pedersen A, Jensen HS, Larsen SE (2012) Phosphorus load  
654 to surface water from bank erosion in a Danish lowland river basin *J Environ Qual*  
655 41:304-313 doi:10.2134/jeq2010.0434

656 Kronvang B, Jeppesen E, Conley DJ, Søndergaard M, Larsen SE, Ovesen NB, Carstensen J  
657 (2005) Nutrient pressures and ecological responses to nutrient loading reductions in  
658 Danish streams, lakes and coastal waters *Journal of Hydrology* 304:274-288  
659 doi:10.1016/j.jhydrol.2004.07.035

660 Lannergård EE, Agstam-Norlin O, Huser BJ, Sandström S, Rakovic J, Futter MN (2020) New  
661 Insights Into Legacy Phosphorus From Fractionation of Streambed Sediment *Journal*  
662 *of Geophysical Research: Biogeosciences* 125 doi:10.1029/2020jg005763

663 Le Moal M et al. (2018) Eutrophication: A new wine in an old bottle? *Sci Total Environ* 651:1-  
664 11 doi:10.1016/j.scitotenv.2018.09.139

665 Lee S-W, Hwang S-J, Lee S-B, Hwang H-S, Sung H-C (2009) Landscape ecological  
666 approach to the relationships of land use patterns in watersheds to water quality  
667 characteristics *Landscape and Urban Planning* 92:80-89  
668 doi:10.1016/j.landurbplan.2009.02.008

669 Levvasseur F et al. (2016) RPG Explorer: A new tool to ease the analysis of agricultural  
670 landscape dynamics with the Land Parcel Identification System *Computers and*  
671 *Electronics in Agriculture* 127:541-552 doi:10.1016/j.compag.2016.07.015

672 Lintern A et al. (2018) Key factors influencing differences in stream water quality across  
673 space *Wiley Interdisciplinary Reviews: Water* 5:e1260 doi:10.1002/wat2.1260

674 Liu J, Liu X, Wang Y, Li Y, Jiang Y, Fu Y, Wu J (2020) Landscape composition or  
675 configuration: which contributes more to catchment hydrological flows and variations?  
676 *Landsc Ecol* 35:1531-1551 doi:10.1007/s10980-020-01035-3

677 McDowell RW, Moreau P, Salmon-Monviola J, Durand P, Leterme P, Merot P (2014)  
678 Contrasting the spatial management of nitrogen and phosphorus for improved water  
679 quality: Modelling studies in New Zealand and France *European Journal of Agronomy*  
680 57:52-61 doi:10.1016/j.eja.2013.09.011

681 McDowell RW, Srinivasan MS (2009) Identifying critical source areas for water quality: 2.  
682 Validating the approach for phosphorus and sediment losses in grazed headwater  
683 catchments *Journal of Hydrology* 379:68-80 doi:10.1016/j.jhydrol.2009.09.045

684 McGarigal K, Marks BJ (1995) FRAGSTATS: spatial pattern analysis program for quantifying  
685 landscape structure vol 351. doi:10.2737/PNW-GTR-351

686 Moatar F et al. (2020) Stream Solutes and Particulates Export Regimes: A New Framework  
687 to Optimize Their Monitoring *Frontiers in Ecology and Evolution* 7  
688 doi:10.3389/fevo.2019.00516

689 Murphy J, Riley JP (1962) A modified single solution method for the determination of  
690 phosphate in natural waters *Analytica Chimica Acta* 27:31-36 doi:10.1016/s0003-  
691 2670(00)88444-5

692 Musolff A, Fleckenstein JH, Rao PSC, Jawitz JW (2017) Emergent archetype patterns of  
693 coupled hydrologic and biogeochemical responses in catchments *Geophysical*  
694 *Research Letters* 44:4143-4151 doi:10.1002/2017gl072630

695 O'Callaghan JF, Mark DM (1984) The extraction of drainage networks from digital elevation  
696 data *Computer Vision, Graphics, and Image Processing* 28:323-344  
697 doi:10.1016/s0734-189x(84)80011-0

698 ODEM (2012) Apports de phosphore et proliférations de cyanobactéries dans le Lac au Duc  
699 (Morbihan).

700 Ouyang W, Song K, Wang X, Hao F (2014) Non-point source pollution dynamics under long-  
701 term agricultural development and relationship with landscape dynamics *Ecological*  
702 *Indicators* 45:579-589 doi:10.1016/j.ecolind.2014.05.025

703 Page T et al. (2005) Spatial variability of soil phosphorus in relation to the topographic index  
704 and critical source areas: sampling for assessing risk to water quality *J Environ Qual*  
705 34:2263-2277 doi:10.2134/jeq2004.0398

706 Peterson EE, Sheldon F, Darnell R, Bunn SE, Harch BD (2011) A comparison of spatially  
707 explicit landscape representation methods and their relationship to stream condition  
708 *Freshwater Biology* 56:590-610 doi:10.1111/j.1365-2427.2010.02507.x

709 Pionke HB, Gburek WJ, Sharpley AN (2000) Critical source area controls on water quality in  
710 an agricultural watershed located in the Chesapeake Basin *Ecological Engineering*  
711 14:325-335 doi:10.1016/s0925-8574(99)00059-2

712 Planchon O, Darboux F (2002) A fast, simple and versatile algorithm to fill the depressions of  
713 digital elevation models *Catena* 46:159-176

714 Pringle CM (1990) Nutrient Spatial Heterogeneity: Effects on Community Structure,  
715 Physiognomy, and Diversity of Stream Algae *Ecology* 71:905-920  
716 doi:10.2307/1937362

717 Qin C, Zhu AX, Pei T, Li B, Zhou C, Yang L (2007) An adaptive approach to selecting a flow-  
718 partition exponent for a multiple-flow-direction algorithm *International Journal of*  
719 *Geographical Information Science* 21:443-458 doi:10.1080/13658810601073240

720 Reaney SM, Mackay EB, Haygarth PM, Fisher M, Molineux A, Potts M, Benskin CMH (2019)  
721 Identifying critical source areas using multiple methods for effective diffuse pollution  
722 mitigation *J Environ Manage* 250:109366 doi:10.1016/j.jenvman.2019.109366

723 Roberts WM, Stutter MI, Haygarth PM (2012) Phosphorus retention and remobilization in  
724 vegetated buffer strips: a review *J Environ Qual* 41:389-399  
725 doi:10.2134/jeq2010.0543

726 Sandstrom S, Futter MN, Kyllmar K, Bishop K, O'Connell DW, Djodjic F (2020) Particulate  
727 phosphorus and suspended solids losses from small agricultural catchments: Links to  
728 stream and catchment characteristics *Sci Total Environ* 711:134616  
729 doi:10.1016/j.scitotenv.2019.134616

730 Schmadel NM, Harvey JW, Schwarz GE, Alexander RB, Gomez-Velez JD, Scott D, Ator SW  
731 (2019) Small Ponds in Headwater Catchments Are a Dominant Influence on Regional  
732 Nutrient and Sediment Budgets *Geophysical Research Letters* 46:9669-9677  
733 doi:10.1029/2019gl083937

734 Sharpley AN, Kleinman PJA, Flaten DN, Buda AR (2011) Critical source area management  
735 of agricultural phosphorus: experiences, challenges and opportunities *Water Science*  
736 *and Technology* 64:945-952 doi:10.2166/wst.2011.712

737 Shatwell T, Köhler J (2018) Decreased nitrogen loading controls summer cyanobacterial  
738 blooms without promoting nitrogen-fixing taxa: Long-term response of a shallow lake  
739 *Limnol Oceanogr* 64 doi:10.1002/lno.11002

740 Shi ZH, Ai L, Li X, Huang XD, Wu GL, Liao W (2013) Partial least-squares regression for  
741 linking land-cover patterns to soil erosion and sediment yield in watersheds *Journal of*  
742 *Hydrology* 498:165-176 doi:10.1016/j.jhydrol.2013.06.031

743 Shore M, Jordan P, Mellander PE, Kelly-Quinn M, Wall DP, Murphy PN, Melland AR (2014)  
744 Evaluating the critical source area concept of phosphorus loss from soils to water-  
745 bodies in agricultural catchments *Sci Total Environ* 490:405-415  
746 doi:10.1016/j.scitotenv.2014.04.122


747 Sliva L, Dudley Williams D (2001) Buffer Zone versus Whole Catchment Approaches to  
748 Studying Land Use Impact on River Water Quality *Water Research* 35:3462-3472  
749 doi:10.1016/s0043-1354(01)00062-8

750 Smith DR (2009) Assessment of in-stream phosphorus dynamics in agricultural drainage  
751 ditches *Sci Total Environ* 407:3883-3889 doi:10.1016/j.scitotenv.2009.02.038

752 Smith DR, Livingston SJ (2013) Managing farmed closed depressional areas using blind  
753 inlets to minimize phosphorus and nitrogen losses *Soil Use and Management* 29:94-  
754 102 doi:10.1111/j.1475-2743.2012.00441.x


755 Smith DR, Pappas EA (2007) Effect of ditch dredging on the fate of nutrients in deep  
756 drainage ditches of the Midwestern United States *Journal of Soil and Water*  
757 *Conservation* 62:252-261

- 758 Srinivasan MS, McDowell RW (2009) Identifying critical source areas for water quality: 1.  
759 Mapping and validating transport areas in three headwater catchments in Otago, New  
760 Zealand *Journal of Hydrology* 379:54-67 doi:10.1016/j.jhydrol.2009.09.044
- 761 Staponites LR, Bartak V, Bily M, Simon OP (2019) Performance of landscape composition  
762 metrics for predicting water quality in headwater catchments *Sci Rep* 9:14405  
763 doi:10.1038/s41598-019-50895-6
- 764 Steffen W et al. (2015) Sustainability. Planetary boundaries: guiding human development on  
765 a changing planet *Science* 347:1259855 doi:10.1126/science.1259855
- 766 Stelzer RS, Lamberti GA (2001) Effects of N: P ratio and total nutrient concentration on  
767 stream periphyton community structure, biomass, and elemental composition *Limnol*  
768 *Oceanogr* 46:356-367 doi:10.4319/lo.2001.46.2.0356
- 769 Strohmenger L et al. (2020) Multitemporal Relationships Between the Hydroclimate and  
770 Exports of Carbon, Nitrogen, and Phosphorus in a Small Agricultural Watershed  
771 *Water Resources Research* 56 doi:10.1029/2019wr026323
- 772 Temnerud J, Bishop K (2005) Spatial variation of streamwater chemistry in two Swedish  
773 boreal catchments: implications for environmental assessment *Environ Sci Technol*  
774 39:1463-1469 doi:10.1021/es040045q
- 775 Thomas IA et al. (2016a) A sub-field scale critical source area index for legacy phosphorus  
776 management using high resolution data *Agriculture, Ecosystems & Environment*  
777 233:238-252 doi:10.1016/j.agee.2016.09.012
- 778 Thomas Z, Abbott BW (2018) Hedgerows reduce nitrate flux at hillslope and catchment  
779 scales via root uptake and secondary effects *J Contam Hydrol* 215:51-61  
780 doi:10.1016/j.jconhyd.2018.07.002
- 781 Thomas Z, Abbott BW, Troccaz O, Baudry J, Pinay G (2016b) Proximate and ultimate  
782 controls on carbon and nutrient dynamics of small agricultural catchments  
783 *Biogeosciences* 13:1863-1875 doi:10.5194/bg-13-1863-2016
- 784 Uuema E, Roosaare J, Mander Ü (2007) Landscape metrics as indicators of river water  
785 quality at catchment scale *Hydrology Research* 38:125-138 doi:10.2166/nh.2007.002
- 786 Wang M, Wang Y, Li Y, Liu X, Liu J, Wu J (2020) Natural and anthropogenic determinants of  
787 riverine phosphorus concentration and loading variability in subtropical agricultural  
788 catchments *Agriculture Ecosystems & Environment* 287  
789 doi:10.1016/j.agee.2019.106713
- 790 William M. Roberts TSG, Marc I. Stutter, Aránzazu Louro, Mustafa Ali, Philip M. Haygarth  
791 (2019) Phosphorus leaching from riparian soils with differing management histories  
792 under three grass species doi:10.2134/jeq2019.07.0252
- 793 Withers P, Neal C, Jarvie H, Doody D (2014) Agriculture and Eutrophication: Where Do We  
794 Go from Here? *Sustainability* 6:5853-5875 doi:10.3390/su6095853
- 795 Withers PJ, Jarvie HP, Stoate C (2011) Quantifying the impact of septic tank systems on  
796 eutrophication risk in rural headwaters *Environ Int* 37:644-653  
797 doi:10.1016/j.envint.2011.01.002
- 798 Xiao R, Wang G, Zhang Q, Zhang Z (2016) Multi-scale analysis of relationship between  
799 landscape pattern and urban river water quality in different seasons *Sci Rep* 6:25250  
800 doi:10.1038/srep25250
- 801 Yates AG, Brua RB, Corriveau J, Culp JM, Chambers PA (2014) Seasonally Driven Variation  
802 in Spatial Relationships between Agricultural Land Use and in-Stream Nutrient  
803 Concentrations *River Research and Applications* 30:476-493 doi:10.1002/rra.2646
- 804 Zhang W, Pueppke SG, Li H, Geng J, Diao Y, Hyndman DW (2019) Modeling phosphorus  
805 sources and transport in a headwater catchment with rapid agricultural expansion  
806 *Environ Pollut* 255 doi:10.1016/j.envpol.2019.113273



- Monitored subcatchment outlet
- ▨ Monitored subcatchment
- ~ Main river
- ~ Hydrography
- ~ Road/Ditch
- Urban
- Wheat/Barley
- Maize
- Grassland
- Forest
- Water

### Denominator calculation for $a = 1.5$ and $b = 1$



### Numerator and LCI calculation for $a = 1.5$ and $b = 1$

Example of agricultural subcatchments with similar landscape composition but contrasting LCI values





Fig3

Click here to

[access/download/Figure3a.ppt](#)


Rank correlations (Rs) between LCI(0,0) and TP (a) and LCI(1.4,2.2) within the exploration domain

Coefficient (a) applied to flow accumulation



B

Rs = 0.35, p-val = 0.145



C

Rs = 0.81, p-val = 2.27e-05










Fig4

[Click here to access/download;Figure;fig4.pdf](#)






## Editor du Jour

### Certificate of English-Language Proofreading

This certificate confirms that the manuscript entitled **“The influence of landscape spatial configuration on nitrogen and phosphorus exports in agricultural catchments”** by Antoine Casquin et al. was edited and proofread for correct English-language content by two native speakers of American English.



Michelle Corson

29 January 2021

Certificate no. 2021-04

Editor du Jour  
33 rue Émile Bernard  
35700 Rennes, FRANCE  
E-mail: [editordujour@yahoo.fr](mailto:editordujour@yahoo.fr)  
URL: <http://editordujour.corsondna.com/>

[Click here to view linked References](#)



Antoine Casquin  
INRAE - UMR SAS / Institut Agro  
65 rue de St Brieuc  
CS 84215  
35042 Rennes Cedex  
France  
Tel: +33223485142  
E-mail: antoine.casquin@inrae.fr

**Subject: Submission of a manuscript for evaluation and publication in *Landscape Ecology***

Dear Editors,

Enclosed is our manuscript, “The influence of landscape spatial configuration on nitrogen and phosphorus exports in agricultural catchments” by Antoine Casquin et al., which we would like to submit for publication as a Research paper in *Landscape Ecology*.

This manuscript has not been previously published, in whole or in part, and is not under consideration by another journal. All co-authors – A. Casquin, R. Dupas, S. Gu, E. Couic, G. Gruau and P. Durand – have approved the manuscript and agree with its submission to *Landscape Ecology*. We had the English proofread by a professional proofreader (“Editor du Jour”; see enclosed English editing certificate).

The spatial variability in nutrient transfer from land to stream in seemingly similar headwater catchments is currently not fully understood, and upscaling results from the plot scale to the catchment scale remains difficult. Our study indicates that including the influence of the spatial configuration of nutrient sources in relation to the hydrographic network is crucial to explore this issue.


We assessed the influence of landscape spatial arrangement on nitrate ( $\text{NO}_3^-$ ) and total phosphorus (TP) exports at the catchment scale. We developed a novel landscape configuration index that we optimised based on  $\text{NO}_3^-$  and TP concentrations in 19 headwater streams. Results show that the spatial configuration of nutrient sources and sinks in headwaters has a large influence on TP transfers but not  $\text{NO}_3^-$  transfers. We also reveal that the spatial configuration of sources is heterogeneous at small scales ( $<1 \text{ km}^2$ ) but homogeneous at larger scales ( $> 50 \text{ km}^2$ ), and we discuss the implications of these findings.

This study is based on a recently published dataset entitled *Repeated synoptic sampling for water chemistry monitoring in the Yvel catchment, northwestern France*. We plan to make the code (R script) available along with the article upon acceptance.

Thank you for considering our manuscript.

Sincerely,

Antoine Casquin



Click here to access/download  
**Supplementary material**  
SupMat.docx

