



HAL
open science

Soil and Aboveground Carbon Stocks in a Planted Tropical Mangrove Forest (Can Gio, Vietnam)

Truong van Vinh, Cyril Marchand, Khanh Tran Vu Linh, Adrien Jacotot, Thanh Nho Nguyen, Michel Allenbach

► **To cite this version:**

Truong van Vinh, Cyril Marchand, Khanh Tran Vu Linh, Adrien Jacotot, Thanh Nho Nguyen, et al.. Soil and Aboveground Carbon Stocks in a Planted Tropical Mangrove Forest (Can Gio, Vietnam). Wetland Carbon and Environmental Management, 2021. insu-03388141

HAL Id: insu-03388141

<https://insu.hal.science/insu-03388141v1>

Submitted on 21 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Soil and above-ground carbon stocks in a planted tropical mangrove forest (Can Gio,**
2 **Vietnam)**

3
4 Truong Van VINH^{a,b,c}, Cyril MARCHAND^{b,c,*}, Tran Vu Khanh LINH^a, Adrien JACOTOT^{b,c,d},
5 Nguyen Thanh NHO^e, Michel ALLENBACH^c

6
7
8 ^a Department of Forest Resources Management, Faculty of Forestry, Nong Lam University
9 HCMC, Ho Chi Minh City, Vietnam

10 ^b IMPMC, Institut de Recherche pour le Développement (IRD), Sorbonne Université, CNRS,
11 MNHN, Noumea, New Caledonia, France

12 ^c Université de la Nouvelle-Calédonie (UNC), *ISEA, EA 7484*, Noumea, New Caledonia, France

13 ^d Université d'Orléans, ISTO, UMR 7327, 45071, Orleans, France

14 ^e Faculty of Environmental and Food Engineering, Nguyen Tat Thanh University, Vietnam

15
16 * Corresponding author: cyril.marchand@unc.nc

17

18

19

20

21

22

23

24 **Abstract**

25

26 Can Gio mangrove is the largest in Vietnam, developing on approximately 35000
27 hectares. This forest was partially destroyed during the Vietnamese war. A restoration
28 program was developed between the late 70s and the early 90s, using *Rhizophora apiculata*
29 Blume propagules. Currently, the Can Gio mangrove forest regenerates naturally and presents
30 a specific species zonation along the intertidal elevation gradient. *Rhizophora* dominates the
31 inner forest at the highest elevation, while at an intermediate location, *Rhizophora* and
32 *Avicennia* cohabit with other scattered species. The lowest position is colonized by *Avicennia*.
33 Within this context, the main objectives of this study were to determine the soil
34 physicochemical characteristics, as well as the quality (C/N ratios and $\delta^{13}\text{C}$) and the quantity
35 (carbon content and stocks) of the organic matter stored beneath each mangrove stand. In
36 addition, we were interested in determining the above-ground biomass and the total carbon
37 stocks of the ecosystem (without considering the below-ground biomass). Carbon stocks of
38 the Can Gio mangrove forest ranged from 150 to 479 Mg C ha⁻¹, with up to 86 % of the C
39 stored in the upper meter of the soil. The inner forest has the highest stock, followed by the
40 transitional forest, and the fringe forest. The depth extension of the root system of the current
41 forest was estimated, and its contribution to the soil carbon stock was calculated, using the
42 adjacent mudflat as a proxy for the antecedent stocks. Our results show that, for the last 40
43 years, the current mature planted *Rhizophora* forest stored 25.26 Mg C ha⁻¹. Consequently,
44 mangrove plantation and restoration after the war was a success in terms of carbon storing.
45 We suggest that the destruction of the Can Gio mangrove forests for urban development
46 would induce the loss of an efficient CO₂ sink.

47

48 **1. Introduction**

49 Mangroves forests cover around 137760 km² between 30° N and 40° S, with the largest
50 percentage of their surface observed between 20° N and 20° S (Giri et al., 2011). Mangroves
51 provide significant ecosystem services such as habitat for various terrestrial and marine animals
52 that are critical for the coastal biodiversity in the tropics (Alongi, 2008; Mumby et al., 2004;
53 Nagelkerken et al., 2008; Saenger, 2002), reef against erosion and natural disasters (Alongi,
54 2008; Barbier, 2006), or even as a trap for suspended and contaminant materials (Kathiresan,
55 2004; Rivera-Monroy et al., 1999). In addition, mangroves strongly contribute to the economic
56 growth of emerging countries (Mukherjee et al., 2014), providing charcoal, firewood, and
57 construction materials for local communities.

58 Due to their high productivity (average of 218 ± 72 Tg C yr⁻¹) (Bouillon et al., 2008),
59 the anoxic character of their soils that limits organic matter degradation (Kristensen et al.,
60 2008), their high storage capacity (Breithaupt et al., 2012) and their global distribution,
61 mangroves play a key role in carbon cycling in the coastal ocean (Kauffman et al., 2011).
62 Mangroves can store carbon both in their biomass and soils; however, this ability depends on
63 many parameters and may be highly variable. Their biomass varies notably according to latitude
64 with higher biomass under the tropics (Saenger and Snedaker, 1993), climate with higher
65 biomass under the wet than under the dry tropics (Adame et al., 2013), and even the age of the
66 forest (Fromard et al., 1998). In addition, a large part of the carbon content may be stored in
67 their soil, with up to 98 % and up to 90 % for estuarine and coastal mangroves, respectively
68 (Donato et al., 2011). Their potential soil carbon storage was recently estimated at 1023 Mg C
69 ha⁻¹ (Donato et al., 2011), much higher than other highly productive ecosystems, such as rain
70 forests (218.6 Mg C ha⁻¹), peat swamps (370.1 Mg C ha⁻¹), or salt marshes (537.8 Mg C ha⁻¹)
71 (Alongi, 2014). However, mangrove soil carbon stocks depend also on various parameters, such
72 as the latitudinal position of the mangrove, the tree species developing at the surface, soil

73 salinity, and even nutrient availability (Adame et al., 2013; Alongi, 2002; Jacotot et al. 2018;
74 Kauffman et al., 2011; Rahman et al., 2015; Sanders et al., 2010; Wang et al., 2013).
75 Particularly, their position within the tidal zone or along an estuary appears to be critical to their
76 potential soil carbon storage capacities. For example, in an estuarine mangrove forest
77 developing in Mexico, Adame et al. (2015) reported soil carbon stocks ranging from 744 to 912
78 Mg C ha⁻¹ for the upper estuary and from 537 to 1115 Mg C ha⁻¹ for the lower estuary.

79 Recently, Vietnam became the first Asian country to implement a national program of
80 payment for forest environmental services (PFES). In Vietnam, mangroves cover around
81 270000 ha (FAO, 2015) with more than 80 % distributed in the southern part of the country,
82 below 10° N latitude (Hawkins et al., 2010). The Can Gio mangrove, located between Ho Chi
83 Minh City (HCMC) and the South China Sea (Bien Dong in Vietnamese), is actually the largest
84 contiguous mangrove in Vietnam with a surface of around 35000 ha. During the Vietnamese
85 war (1964-1971), approximately 57% of this mangrove was destroyed by the spraying of a
86 mixture of herbicides and defoliants (Ross, 1975). After the war, in the late 70s and early 90s,
87 a vast reforestation program was implemented by the HCMC Forest Department, and most of
88 the Can Gio area was replanted using *Rhizophora apiculata* Blume propagules collected in the
89 Mekong Delta. In January 2000, the mangrove forest of Can Gio was registered in the UNESCO
90 World Network of Biosphere Reserves list. In addition, the environmental value of the Can Gio
91 mangrove forest was taken into account in the National Strategy on Climate Change and Sea
92 Level Rise by the Vietnamese government. Currently, the Can Gio mangrove forest is highly
93 diversified, with a total of 77 mangrove species (35 true mangroves and 42 associates) (Tri et
94 al., 2000), whose main species present a specific zonation along the intertidal elevation
95 gradient. This zonation includes, from the lowest to the highest position: i) a fringe forest that
96 is dominated by *Avicennia alba*; ii) a transitional forest, composed of a mixture of *R. apiculata*
97 *A. alba*, *A. officinalis* and sparse *Excoecaria agallocha* and *Sonneratia alba*; and iii) an inner

98 forest, which is mainly composed of mature *R. apiculata*. It was recently demonstrated that the
99 mineralization of mangrove-derived organic matter has a key role in sustaining coastal food
100 webs in the Can Gio estuary (David et al. 2018a, David et al. 2019). Recent studies also showed
101 that the total carbon stocks in the Can Gio mangrove forest could reach 1000 Mg C ha⁻¹ (Dung
102 et al., 2016; Nam et al., 2016) and that the below-ground carbon accumulation rates may reach
103 3.24 Mg C ha⁻¹ yr⁻¹ (MacKenzie et al., 2016). However, these studies were interested neither in
104 soil organic carbon (SOC) quality nor in the influence of soil elevation on carbon stocks of the
105 different stands.

106 Therefore, the main objectives of this study were: i) to evaluate the soil physicochemical
107 parameters, the organic matter quality, and the vegetation characteristics of three different
108 mangrove stands and of the adjacent mudflat developing along the elevation gradient, ii) to
109 determine the carbon stocks in the biomass and in the first meter of soil, and iii) to calculate the
110 soil carbon stocks related to forest plantation. We hypothesized that carbon stocks in the soil
111 and in the biomass depend on the mangrove species and, therefore, to the position of the stand
112 along the intertidal elevation gradient. To reach our goals, core samples were taken within each
113 of the four stands. Supplementary cores were also collected in the adjacent mudflat to serve as
114 a soil reference (i.e., before the colonization of the soil by the mangrove). Then,
115 physicochemical parameters (pH, redox, and salinity) and the distribution of C/N ratios and
116 $\delta^{13}\text{C}$ stable isotopes with depth in each stand were measured. Carbon stocks were determined
117 by combining bulk density and total organic carbon content for the soil and by using biomass
118 measurements with specific (Vinh et al., 2019) and generic allometric equations for the above-
119 ground stocks.

120 **2. Methods**

121 **2.1. Study site**

122 The present study was conducted in the Can Gio mangrove forest in Southern Vietnam
123 (Fig. 1a). This mangrove forest of around 35000 ha is the largest contiguous mangrove in
124 Vietnam. Located in the district of Can Gio, one of the 24 districts of Ho Chi Minh City, the
125 Can Gio mangrove forest is at the deltaic confluence of the rivers Sai Gon, Dong Nai, and Vam
126 Co, flowing into the East Sea (Bien Dong in Vietnamese). The Can Gio mangrove forest is
127 composed of several species that follow a specific zonation determined by soil elevation. From
128 the lowest to the highest intertidal zone, the forest is composed of a mudflat that is totally
129 denuded of vegetation; a fringe forest, dominated by *Avicennia* spp. (mainly *A. alba*); a
130 transitional forest, mainly composed of *R. apiculata*, *A. alba*, *A. officinalis*, as well as sparse
131 *Excoecaria agallocha* and *Sonneratia alba*; and an inner forest dominated by mature *R.*
132 *apiculata*.

133 The climate in this region is typically monsoonal (type Am in Köppen-Geiger
134 classification) with two main seasons: a wet season from May to November and a dry season
135 from December to April. The annual average rainfall is 1816 mm, spread over 154 rainy days,
136 with approximately 80 % of the rainfall occurring during the wet season. As a result, the flow
137 of sediments transported by the river crossing the mangrove forest during the wet season is
138 elevated (~160 million tons (Milliman and Meade, 1983)), leading to the accretion of the deltaic
139 plain downstream. Conversely, during the dry season, the fresh water flow is strongly reduced,
140 and the mangrove forest is severely affected by saline intrusions. The Can Gio estuary is
141 characterized by irregular semi-diurnal tides with a tidal range of 2 to 4 m.

142 **2.2. Field measurements and carbon stocks**

143 2.2.1 Elevation

144 The elevation of each stand relative to mean sea level (MSL) was measured with a dye-
145 type tide gauge (Clough, 2014; English et al., 1997; Schmitt and Duke, 2014). First, the cotton

146 tape was soaked in a water soluble food dye and attached to a wooden stake of about 2.5 m in
147 length. At low tide, the wooden stakes were inserted deeply into the soil starting from the river
148 edge. The distance between the two columns was 2 m in the mudflat, 5 m in the fringe forest,
149 and then every 10 m until the inner forest. After high tide, we measured the height of the
150 washout line above the ground (Albers and Schmitt, 2015; Clough, 2014). In this study, data
151 from the Vung Tau tide station, Viet Nam (10.3333° N, 107.0667° E) were used to report our
152 data relative to MSL in this zone. The MSL at Vung Tau tide station in 2016 was 2.16 m. To
153 determine MSL at the study site, the following equation was used: $MSL = \text{highest tide value} -$
154 $MSL \text{ at Vung Tau}$. Monthly and annual mean sea levels and the daily tide series are collected
155 and published by the Permanent Service for Mean Sea Level (Pissierssens, 2002).

156 2.2.2. Cores collection and soil physicochemical parameters

157 Soil cores were collected in triplicate with a gouge auger (1 m long, 3 cm wide) attached
158 to a cross handle. Cores were collected during the dry season of 2016 and the wet season of
159 2017 from the surface to a depth of 100 cm in each mangrove stand as S1 in the mudflat, S2 in
160 the fringe forest, S3 in the transitional forest, and S4 in the inner forest (Fig. 1c). The auger was
161 carefully inserted into the soil to minimize disturbance of the core surface, twisted, and finally
162 removed, following the method used by MacKenzie et al. (2016). Each core was then separated
163 into different depth intervals: 0 – 2.5 cm, 2.5 – 5 cm, 5 – 7.5 cm, 7.5 – 10 cm, 10 – 15 cm, 15 –
164 20 cm, 20 – 25 cm, 25 – 30 cm, and then every 10 cm from 30 to 100 cm. For each interval,
165 one subsample of a known volume was collected and placed in a zippered bag that was
166 immediately sealed in aluminum foil to minimize gas exchange and stored in a frozen box until
167 it reached the laboratory. All samples were, then, dried by freeze-drying at -52 °C until a
168 constant weight was achieved. Finally, the dry bulk density (DBD) of each sample was
169 determined by dividing its dry mass by its fresh volume.

170 Additional cores were collected to measure pore-water salinity, redox potential (Eh),
171 and pH. Pore-water salinity was measured with a hand-held refractometer. Redox potential and

172 pH were measured using, respectively, a combined Pt-Ag/Ag-Cl electrode and a glass electrode,
173 both connected to a pH/mV/T meter (Marchand et al., 2004). The pH electrode was calibrated
174 prior to sampling using three standard solutions of pH 4, 7, and 10 at 25 °C (National Institute
175 of Standards and Technology, USA), and the redox electrode was checked prior to utilization
176 with a 0.43 V standard solution and demineralized water (Marchand et al., 2004; Thanh-Nho et
177 al., 2017).

178 2.2.3. Soil TOC, TN, $\delta^{13}\text{C}$, and soil carbon stocks

179 Soil total organic carbon (TOC), total nitrogen (TN), and $\delta^{13}\text{C}$ values were determined
180 using an elemental analyzer coupled to an isotope ratio mass spectrometer (Integra2, Sercon,
181 UK). The analytical precisions of the analyzer were checked using the IAEA-600 caffeine
182 standard (IAEA Nucleus) and were less than 1 % for TOC, 0.15 % for N, and 0.3 % for $\delta^{13}\text{C}$.
183 All the analyses were performed at the French Institute for the Sustainable Development (IRD)
184 of Noumea, New Caledonia, France. The $\delta^{13}\text{C}$ values are reported permil (‰) deviations from
185 a Pee-Dee Belemnite (PDB) limestone carbonate as the standard using the following equation:

$$186 \quad \delta^{13}\text{C} (\text{‰}) = \left(\frac{\frac{^{13}\text{C}}{^{12}\text{C}}_{\text{sample}}}{\frac{^{13}\text{C}}{^{12}\text{C}}_{\text{standard}}} - 1 \right) \times 1000 \quad (\text{Eq. 1})$$

187 The soil carbon stocks of each mangrove stand (Mg C ha^{-1}) were determined using the
188 following equation:

$$189 \quad \text{Soil carbon stock (Mg C ha}^{-1}\text{)} = \text{TOC (\%)} * \text{DBD (g cm}^{-3}\text{)} * \text{depth interval (cm)} \quad (\text{Eq. 2})$$

190 2.2.4. Above-ground carbon stocks

191 In November 2016, three transects (150 m long and 20 m wide) were established
192 following an elevation gradient in the mangrove forest (Fig. 1c). All transects encompassed the
193 four different stands of the mangrove (i.e., the mudflat, the fringe, the transitional, and the inner
194 forests). All trees within the transects were counted and their DBH at 1.3 m above the soil were
195 measured, except for *R. apiculata*, whose diameters were measured just above the highest prop

196 root.

197 For *R. apiculata*, the above-ground biomass was calculated using the allometric
198 equation specifically developed for this species in Southern Vietnam (Vinh et al., 2019):

$$199 \quad WR_{\text{Total}} = 0.38363 * DBH^{2.2348} \quad (\text{Eq. 3})$$

200 For the other species, the above-ground biomass was determined using the common allometric
201 equations developed by Komiyama et al. (2005).

$$202 \quad W_{\text{Total}} = 0.251 * DBH^{2.46} \quad (\text{Eq. 4})$$

203 where, in Eq. 3, WR_{Total} is the total above-ground biomass of *R. apiculata* (kg), and D is the
204 diameter above the highest prop root of *R. apiculata* (cm); and in Eq. 4. W_{Total} is the total above-
205 ground biomass biomass of the other species (kg), and DBH is the diameter at breast height of
206 the other species (cm).

207 Above-ground carbon stocks were then estimated by multiplying the above-ground
208 biomass by a carbon conversion factor. Two conversion factors were used in this study: 0.4409
209 for *R.apiculata*, which is the specific carbon conversion factor for this species (Vinh et al.,
210 2019), and 0.451 for the other species (Hiraishi et al., 2014). Eventually, the total above-ground
211 carbon stocks per area (Mg C ha^{-1}) were calculated by summing the above-ground carbon stock
212 of *R. apiculata* and the one for the other trees using the tree density per hectare.

213 **2.3. Statistical analyses**

214 A parametric two-way analysis of variance (ANOVA) was used to test the significant
215 effects ($p < 0.05$) of seasons and sites on Eh, pH, and pore-water salinity. A two-way ANOVA
216 was also applied to assess the significant differences ($p < 0.05$). A Student's t-test was used to
217 test significant differences ($p < 0.05$) between $\delta^{13}\text{C}$ values, C/N ratios, and C concentrations of
218 an upper layer from 0 to 40 cm and a lower layer from 40 to 100 cm, and for Eh and pH for the
219 transitional forest and inner forest. All statistical analyses were performed using XLSTAT
220 software version 2017.4 for Mac OS10.13.4.

221 **3. Results**

222 **3.1. Can Gio mangrove distribution**

223 The relative elevations of the soil surface at the four sampling stations increased from the
224 river toward the inner forest. The *A. alba* forest developed between +20 and +30 cm above
225 MSL, the transitional forest between +30 and + 40 cm, and the *Rhizophora* stand between +40
226 and + 75 cm above MSWL (Fig. 2).

227 The fringe forest had a mean DBH value of 10.7 cm, a tree density of 1327 ind ha⁻¹, and
228 a basal area of 11.9 m² ha⁻¹. The latter parameter increased to the inner forest with a value
229 reaching 28.6 cm; the inner forest was also characterized by the highest mean DBH, 18.6 cm.
230 The transitional forest was characterized by the highest tree density, 3727 individuals per
231 hectare (ind ha⁻¹), and the lowest DBH, 8.5 cm (Table 1).

232 **3.2. Soil physicochemical parameters (pore-water salinity, Eh, and pH)**

233 Pore-water salinity remained relatively stable along the core profiles during both seasons
234 in the mudflat as well as in the fringe and the transitional forests, with values ranging from 10.0
235 to 23.3, 18.0 to 28.3, and 12.0 to 30.0, respectively (Fig. 3 a, d, g). However, in the inner forest,
236 pore-water salinity increased from 10.0 and 27.3 at the soil surface to 18.0 and 31.7 at 20 cm
237 depth, during the dry and the wet season, respectively. Below 20 cm, pore-water salinity
238 remained stable until 100 cm of depth (Fig. 3 l). When integrating the entire sampled profile,
239 the season had a significant effect ($p < 0.001$) on pore-water salinity, with higher values during
240 the dry season in all stands, with the exception of the transitional forest where no significant
241 differences ($p > 0.05$) in mean pore-water salinity were observed. In addition, pore-water
242 salinity was significantly different between the four zones in both seasons ($p < 0.001$). Mean
243 values were 20.07 ± 1.78 and 12.47 ± 3.48 in the mudflat, 24.8 ± 1.8 and 19.8 ± 1.3 in the fringe
244 forest, 19.0 ± 1.8 and 19.5 ± 3.5 in the transitional forest, and 29.5 ± 2.8 and 16.3 ± 4.5 in the
245 inner forest, for the dry and the wet season, respectively. Consequently, pore-water salinity

246 increased along the intertidal elevation gradient with lower values at the lowest position and
247 the higher values in the higher position.

248 In the four zones, Eh values decreased with depth, from 150 to -50 mV in the mudflat,
249 from -50 to -250 mV in the fringe and the transitional forests and from -50 to -400 mV in the
250 inner forest (Fig. 3 b, e, h, m). When integrating the whole cores, from the top to the bottom, at
251 100 cm depth, Eh significantly ($p < 0.001$) decreased from the mudflat to the inner forest. Mean
252 values were -10.7 ± 104.9 and 46.6 ± 90.9 mV in the mudflat, -52.2 ± 70.0 and -107.9 ± 84.1
253 mV in the fringe forest, 44.7 ± 104.5 and -118.9 ± 93.6 mV in the transitional forest, and -268.9
254 ± 185.9 and -117.4 ± 57.2 mV in the inner forest, for the dry and the wet season, respectively.
255 However, although Eh was higher during the dry season than during the wet season, these
256 differences were not significant in any of the stands ($p > 0.05$, Table 2). Finally, Eh values in
257 the upper layer (0 – 40 cm) were significantly different ($p < 0.05$) from the lower layer (40 –
258 100 cm) for both the transitional forest and inner forests.

259 Concerning pH, values did not vary significantly during both seasons in the mudflat,
260 with values ranging from 6.9 to 7.5 (Fig. 3c, f, k and n). For the other stands, pH values varied
261 with depth and with season, from 6.2 to 7.6. When integrating the whole core (i.e., from the soil
262 surface to 100 cm depth), pH values were significantly different between stands ($p < 0.001$),
263 with higher mean values in the mudflat, followed by the fringe and the transitional forests, and
264 with the lowest mean value in the inner forest. In addition, there were significant differences (p
265 < 0.05) between the upper layer (0 – 40 cm) and the lower layer (40 – 100 cm) for both the
266 transitional and inner forests.

267 **3.3. Soil organic matter characteristics (DBD, TOC, TN, $\delta^{13}\text{C}$, and C/N ratios)**

268 For all stands, DBD values remained stable along the soil profile, from the surface to
269 100 cm depth. However, mean DBD values for the first meter of soil were statistically different
270 ($p < 0.001$) between each stand. The mean DBD values were 0.52 ± 0.06 , 0.62 ± 0.02 , $0.63 \pm$

271 0.03, and $0.64 \pm 0.08 \text{ g cm}^{-3}$ in the mudflat, the fringe, the transitional, and the inner forests,
272 respectively.

273 In the mudflat and in the fringe forest, TOC values were stable from the soil surface to
274 the bottom of the core, with mean values of 2.42 ± 0.34 and $3.2 \pm 0.27 \%$, respectively (Fig. 4
275 a). In the transitional and the inner forests, TOC values between the upper (0 – 40 cm) and the
276 deep layers (40 – 100 cm) were significantly different ($p < 0.05$). Mean TOC values were 4.14
277 ± 1.05 and $7.03 \pm 3.10 \%$ for the transitional forest and 4.36 ± 1.31 and $6.75 \pm 2.80 \%$ for the
278 inner forest for the upper and the deep layers, respectively (Fig. 4 a). In addition, the average
279 TOC values for the first meter of soil were significantly different ($p < 0.001$) between the four
280 zones.

281 C/N ratios were relatively stable throughout the entire sampled soil profile in the mudflat
282 and in the fringe forest with values ranging from 11.2 to 12.1 and from 11.2 to 15.7, respectively
283 (Fig. 4 b). However, similar to TOC, C/N ratios between the upper 40 cm of soil in the
284 transitional and inner forests were statistically different ($p < 0.05$) when compared to the lower
285 60 cm. The mean values of C/N ratios were 12.2 and 18.2 for the transitional forest and 17.5
286 and 25.7 for the inner forest for the upper and the lower layers, respectively (Fig. 4 b). In
287 addition, when considering the complete sampled profile, C/N ratios were significantly
288 different ($p < 0.001$) between the four stands and increased landward. The mean values were
289 11.61 ± 0.22 , 12.08 ± 1.09 , 14.44 ± 4.53 , and 22.78 ± 5.49 for the mudflat, the fringe forest, the
290 transitional forest, and the inner forest, respectively.

291 Concerning $\delta^{13}\text{C}$, in the mudflat and in the inner forest, the values slightly increased
292 from -30 ‰ and -32 ‰ at the soil surface to -27 ‰ and -31 ‰ at 20 cm depth and then remained
293 relatively stable until the bottom of the cores (Fig. 4 c). In the fringe forest, $\delta^{13}\text{C}$ values rapidly
294 dropped from -27 ‰ at the soil surface to -32 ‰ at 10 cm depth, and then gradually increased
295 to a mean value of -29 ‰ from 20 to 80 cm depth. However, a second rapid drop to -34 ‰ was

296 observed in the profile at 90 cm depth (Fig. 4 c). Finally, in the transitional forest, $\delta^{13}\text{C}$ values
297 decreased from -29 ‰ at the soil surface to -30 ‰ at 25 cm depth and then remained stable
298 until 100 cm depth (Fig. 4 c). When integrating the entire core, the mean $\delta^{13}\text{C}$ were significantly
299 different between the four zones ($p < 0.001$), with mean values of -28 ± 0.8 ‰, -30 ± 2.3 ‰, $-$
300 29 ± 1.4 ‰, and -31 ± 0.3 ‰ in the mudflat, the fringe, the transitional forest, and the inner
301 forests, respectively.

302 **3.4. Carbon stocks in the above-ground biomass and in the soils of the different stands**

303 Above-ground carbon stocks were estimated at 24.3 ± 5.1 Mg C ha⁻¹ for the fringe
304 forest, 91.7 ± 29.4 Mg C ha⁻¹ for the transitional forest, and 118.8 ± 9.5 Mg C ha⁻¹ for the inner
305 forest. In addition, above-ground carbon stocks were significantly different between the
306 different stands ($p < 0.001$). No above-ground biomass was measured in the mudflat due to the
307 absence of vegetation in this zone. Soil carbon stocks increased with the elevation gradient,
308 with higher values in the inner forest (360.4 Mg C ha⁻¹), followed by the transitional forest
309 (219.8 Mg C ha⁻¹), the fringe forest (157.6 Mg C ha⁻¹), and the mudflat (150.2 Mg C ha⁻¹) (Fig.
310 5). In addition, soil carbon stocks were significantly different between the different stands ($p <$
311 0.001). Eventually, the total carbon stocks (without the below-ground biomass) were $150.2 \pm$
312 19.6 , 181.9 ± 24.9 , 311.5 ± 28.1 , and 479.2 ± 32.6 Mg C ha⁻¹, for the mudflat, the fringe, the
313 transitional forest, and the inner forest, respectively (Fig. 5).

314 **4. Discussion**

315 **4.1. Mangrove zonation in Can Gio**

316 Between 1978 and 1994, a vast mangrove reforestation program was undertaken with
317 *R. apiculata* as a primary species (Hong and San, 1993). When *Rhizophora* stands were
318 established, the mangrove area extended through natural regeneration and rapid colonization of
319 mudflats along riverbanks notably by *A. alba*, which can be considered as a pioneer species

320 (see for example Balke et al. (2011); Brunt and Davies (2012); Naidoo and Naidoo (2017);
321 Proisy et al. (2009)) (Fig. 2 a, b). This fringe forest developed at lower elevations as compared
322 to the *R. apiculata* forest and was separated from the latter by a transitional forest composed of
323 several species. Mangrove zonation often manifests itself as a mosaic that varies according to
324 physical, biological, and chemical interactions established between plant and substrate in a
325 given area. Pore-water salinity was often considered as the main driver of the zonation
326 (Banerjee et al., 2013; Marchand et al., 2012; 2011), because mangrove plants have different
327 abilities to cope with this factor (Ellison, 1998; Mckee, 1993; Walsh, 1974). For instance, under
328 semi-arid climate, pore-water salinity increased landward, where evaporation processes were
329 intense due to the rare periods of immersions and low precipitation and could reach to a value
330 greater than 50. As a consequence, *Avicennia* trees, which can cope with high salinity (Kendall
331 and Skipwith, 1969; Khan and Aziz, 2001; Marchand et al., 2004; Ukpong, 1997), developed
332 at higher tidal position than the *Rhizophora* trees, which colonized the seaward zone of the
333 mangrove forest. In Can Gio, the zonation was the opposite because pore-water salinity never
334 reached such high values. In fact, pore-water salinity increased from the fringe forest to the
335 interior forest due to its higher elevation that induced more evaporation, but it was never higher
336 than 30. The Can Gio mangrove forest is an estuarine mangrove with freshwater inputs from
337 the Sai Gon and the Dong Nai Rivers. Along the estuary, salinity values ranged from 2 to 26
338 during the year; and even at the mouth of the estuary, salinity never reached the value of
339 seawater, possibly due to the high freshwater inputs from these two rivers and also from the
340 Mekong delta, which is further south of the study site (David et al., 2018b; Thanh-Nho et al.,
341 2018). Furthermore, during the rainy season, the intense rainfall brought additional freshwaters
342 that induce a dilution of pore-water salinity, with mean values for the mature *Rhizophora* forest
343 decreasing from 29 to 16. In the latter stand, salinity also increased with depth, notably because
344 dilution with rainwater occurred in the upper sediment during the rainy season, but also possibly

345 because dissolved salts could migrate at depth through convection processes and accumulate
346 there (Marchand et al., 2004).

347 ***4.2. Influence of mangrove development on soil properties***

348 Organic carbon (OC) content in mangrove sediments usually ranges from 0.5 % to 15 %,
349 with a median value of 2.2 % (Kristensen et al., 2008). In the Can Gio forest, soil OC increased
350 with increasing elevation landward, ranging from 2.42 % in the low intertidal zone to 5.82 %
351 in the high intertidal zone, indicating that the interior forest had accumulated a larger organic
352 carbon stock. The interior forest studied here is composed of mature *Rhizophora* trees, which
353 were planted in 1978. Therefore, mangrove-derived organic matter had accumulated in its soil
354 for almost 40 years at the date of the coring. *Rhizophora* trees under tropical climates as in
355 southern Vietnam are highly productive (Alongi, 2014), which can positively increase soil
356 carbon stocks. In addition, this stand was the furthest from the tidal creek; and as a consequence,
357 tidal flushing of leaf litter was reduced. Leaf litter can accumulate and increase the soil carbon
358 stocks. Conversely, due to their more recent development, lower productivity, and proximity to
359 tidal creeks, the soils of the fringe stand composed of *A. alba* contained less organic carbon
360 than the mature inland *Rhizophora* forest.

361 These gradients of elevation, length of tidal immersion, and carbon content strongly
362 influence redox conditions and pH of the soil. Redox potential (Eh) decreased with increasing
363 elevation from the mudflat to the mature *Rhizophora* stand. Additionally, within the *Rhizophora*
364 stand, the redox condition became rapidly anoxic with depth (Fig. 3). We suggest that the higher
365 organic content beneath this stand induced a higher electron acceptor demand, which were less
366 renewed by the tides due to its high position in the tidal zone. The higher redox values measured
367 beneath the *Avicennia* stands may also be related to the ability of this mangrove species to aerate
368 the sediment through its root system as described by Scholander et al. (1955). Hesse (1961)
369 also observed that *Rhizophora* soils were anoxic most of the time and sulfidic. This difference

370 between the two species was later confirmed in different countries and was suggested to be
371 related to different organic enrichment of the soil, specific abilities of the root system, and
372 different positions in the tidal zone (Marchand et al., 2004; Marchand et al., 2011; Mckee,
373 1993). pH values also varied along the elevation gradient of the studied tidal zone, decreasing
374 from 6.8 in the mudflat to 6.2 in the mature *Rhizophora* stand. We suggest that the increased
375 organic content and its decay processes were responsible for this soil acidification. In addition,
376 in anoxic mangrove soils, sulfides minerals could precipitate (Balk et al., 2016), and slight
377 modifications of the redox conditions could induce their oxidation, which could result in soil
378 acidification (Noel et al., 2017; 2014).

379 ***4.3. Characterization of soil organic matter with depth and along the intertidal elevation*** 380 ***gradient***

381 In mangrove soils, organic matter is usually a mixture between autochthonous organic
382 matter (leaf litter, roots debris, and microphytobenthos) and allochthonous organic matter
383 derived from marine and/or terrestrial origins (Kristensen et al., 2008). However, in a distinct
384 area, the respective contribution of each source in the carbon pool depends on several
385 parameters, including trees productivity, mangrove position, tidal range, freshwater inputs,
386 position of in the intertidal zone, etc. In the Can Gio mangrove soils, $\delta^{13}\text{C}$ and C/N ratios ranged
387 between -28 ‰ to -31 ‰ and between 12 to 22, respectively, and were consistent with those
388 previously reported (Bouillon et al., 2003; Prasad et al., 2017, Jacotot et al., 2018). However,
389 OM quality differed along the intertidal elevation gradient, with higher C/N ratios and depleted
390 $\delta^{13}\text{C}$ values as the elevation increased. These gradients suggested an increased contribution in
391 the upper intertidal zone of mangrove-derived organic matter, characterized by elevated C/N
392 ratios and depleted $\delta^{13}\text{C}$ values (Bosire et al., 2005; Jennerjahn and Ittekkot, 1997; Kristensen
393 et al., 2008; Marchand et al., 2005). These results were consistent with our previous hypothesis
394 of a higher enrichment of the soil resulting from mangrove development and a greater

395 contribution of *Rhizophora* leaf litter in the inner forest due to the higher productivity of the
396 stand, the limited tidal export, and the anoxic character of the soils that limits OM decay process
397 and favors its accumulation. Conversely, towards the tidal creek side of the mangrove, the
398 enriched $\delta^{13}\text{C}$ and lower C/N ratios, close to 12, suggested a greater contribution of
399 phytoplankton or phytobenthos and/or more degraded higher plant debris. In a recent study
400 (Vinh et al., 2020), we showed that leaf litter in the *Avicennia* stand was more rapidly
401 decomposed, and that decay rates were even enhanced during the monsoon. Consequently, we
402 suggest that the position of the stand along the intertidal elevation gradient and its species
403 composition influenced the organic matter characteristics of the soil.

404 Surprisingly, an organic-rich layer was observed at depth beneath the transitional and
405 the inner forests from 40 to 100 cm depth. In the mangroves of southern Vietnam, MacKenzie
406 et al. (2016) determined an average vertical accretion rate of around $0.99 \pm 0.09 \text{ cm yr}^{-1}$. This
407 vertical accretion rate was relatively elevated and reflected notably the high sedimentation rate
408 characterizing most Asian estuaries. During the wet season, strong rainfalls induce high erosion
409 rates in the upper watersheds. As a result, high quantity of sediments are transported by the
410 rivers and deposited along the coastlines, notably in mangrove forests. Following the accretion
411 rate determined by MacKenzie et al. (2016), the upper layer observed in this study (from 0 to
412 40 cm depth) would have started to be deposited 40 years ago, which almost corresponds to the
413 beginning of the reforestation program started in 1978 (Hong and San, 1993). Consequently,
414 we suggest that the upper layer corresponds to the development of the current forest (0 – 40
415 cm), and the lower layer (40 – 100 cm) accumulated before mangrove destruction during the
416 Vietnamese war. Interestingly, this buried layer was at least 50 % enriched in carbon compared
417 to the upper layer for the same sediment thickness and was also characterized by higher C/N
418 ratios (9.7 vs. 22.3 and 14.0 vs. 37.1, for the upper and the lower layers, respectively) and

419 depleted $\delta^{13}\text{C}$ values (-30.7‰), which suggested that the former forest was more productive
420 and/or accumulated organic carbon during a long period.

421 ***4.4. Influence of mangrove development on carbon stocks***

422 Carbon stocks in the above-ground biomass were $118.8 \pm 9.5 \text{ Mg C ha}^{-1}$ for the inner
423 forest, $91.7 \pm 29.4 \text{ Mg C ha}^{-1}$ for the transitional forest, and $23.3 \pm 5.1 \text{ Mg C ha}^{-1}$ in the fringe
424 forest. Differences between stands may be related to the age of the forests, as the inner forest
425 was planted 40 years ago, while the other stands naturally regenerated recently. In addition,
426 *Rhizophora* trees that dominates the inner forest and extends into the transitional forest are
427 generally more productive than *Avicennia* ones (Komiyama et al., 2008) that colonize the fringe
428 forest. Nevertheless, these results were in the range of those previously observed from other
429 studies in Southern Vietnam, with values ranging from 13.4 to 210.7 Mg C ha^{-1} (Dung et al.,
430 2016; Tue et al., 2014). However, the above-ground carbon stocks in Can Gio were much lower
431 than those in other mangrove forests. For example, in Malaysia, the above-ground carbon stocks
432 reached 202.9 Mg C ha^{-1} (Putz and Chan, 1986). This difference may be explained by the age
433 of the forest, as the forest in Malaysia has developed for 80 years—twice the age of the one in
434 Can Gio, and by silvicultural activities (Vinh et al., 2019).

435 In Can Gio, soil carbon stocks ranged from 150 to 360 Mg C ha^{-1} , which was consistent
436 with and even higher than the reported values for other mangrove forests in Vietnam, ranging
437 from 144 to 233 Mg C ha^{-1} (Dung et al., 2016; Tue et al., 2014). However, these values were
438 lower than the ones of tropical mangroves that ranged from 337 to 640 Mg C ha^{-1} (Adame et
439 al., 2013; Castillo et al., 2017; DeVecchia et al., 2014; Hossain, 2014; Kauffman et al., 2011).
440 We suggest that partial destruction of the mangrove during the Vietnamese War may have
441 prevented organic matter to accumulate for a while and probably allowed the existing material
442 to be eroded by tides and freshwater circulation, explaining these low values for soil carbon
443 stocks. Soil carbon stocks in the mangrove of Can Gio represented between 70 to 86 % of its

444 total carbon stocks (without the below-ground biomass) and was consistent with other Indo-
445 Pacific mangrove forests (Donato et al., 2011; Kauffman et al., 2011; Liu et al., 2014).

446 Recently, some authors suggested that integration depth is of major concern when
447 determining carbon stocks in mangrove forests (Lunstrum and Chen, 2014; Marchand, 2017;
448 Jacotot et al., 2018). In Can Gio, integrating the soil carbon stocks down to one meter takes into
449 account the stocks linked to the development of the current forest, replanted after the
450 Vietnamese war as discussed above, and a part of the stocks that accumulated before the
451 destruction of the forest. When considering only the development of the current forest (i.e., 0 –
452 40 cm, assuming a sedimentation rate of 0.99 ± 0.09 cm yr⁻¹ as discussed above), soil carbon
453 stocks were much at 85.7 ± 33.92 Mg C ha⁻¹ in the inner forest. We did not calculate this stock
454 for the other stands considering that it was a natural regeneration that occurred later than the
455 planting and had an OM quality depth profile different from the mature *Rhizophora* forest. In
456 addition, knowing the age of the forest and the amount of carbon stored in its soil since its
457 development allowed the precise contribution of this forest to the enrichment in soil organic
458 matter to be determined. Doing this, the antecedent carbon stocks (i.e., the stocks that were
459 present before the apparition of the forest) must be determined (Lal, 2005). In our study, the
460 stocks in the mudflat were chosen as a proxy for the antecedent carbon stocks. Therefore, for
461 the last 40 years, the actual forest contributed to an enrichment of 25.26 Mg C ha⁻¹ in the inner
462 forest. As a result, the carbon burial rate in the mature *Rhizophora* was ~ 0.6 Mg C ha⁻¹ yr⁻¹,
463 which is lower than the global value of 1.35 Mg C ha⁻¹ yr⁻¹ reported by Bouillon et al. (2008).
464 However, our result may be underestimated, considering that the mudflat was probably enriched
465 by mangrove-derived organic matter, as suggested by the depleted $\delta^{13}\text{C}$ values and the high
466 TOC content. Nevertheless, this study demonstrated that increasing mangrove areas by either
467 restoration or expansion is an important way of increasing carbon storage in the coastal ocean,
468 and mangroves must be considered in future climate change mitigation programs.

469 5. Conclusions

470 Degraded by the spraying of defoliants during the Vietnam War, mangrove forests in
471 Can Gio Estuary successfully recovered through replantation and natural regeneration. They
472 now store a high amount of carbon both in their biomass and in their soils. Their destruction for
473 infrastructure development along the coastline would result in the loss of an efficient CO₂ sink.

474 The main conclusions of this study can be summarized as follow:

475 1. C stocks in the above-ground biomass were significantly different between stands,
476 increasing landward, reaching up to 118.8 ± 9.5 Mg C ha⁻¹ for the mature *Rhizophora* stand.
477 Differences in carbon stocks in the above-ground biomass between stands were related to
478 different forest ages, mangrove species, and tree densities, the latter being managed by thinning.

479 2. The specific zonation, with planted *Rhizophora* trees at the highest elevation in the
480 tidal zone (with a limited pore-water salinity value due to a monsoon-dominated climate), and
481 natural colonization of the river banks by *Avicennia* trees resulted in gradients in the soil
482 physicochemical properties from the mudflat to the inner forest. Due to their more recent
483 development, lower productivity, and proximity to tidal creeks, the soils of the fringe stand
484 composed of *A. alba* contained less organic carbon than the mature inland *Rhizophora* forest.
485 These gradients of elevation and of carbon content strongly influenced the redox conditions and
486 pH in the soil; with both decreasing from the mudflat to the mature *Rhizophora* stand.

487 3. Regarding soil organic matter quality, the $\delta^{13}\text{C}$ and C/N ratio values suggested a
488 higher contribution of mangrove-derived organic matter in the inner forest, most probably
489 because of its age, the high productivity of the stand, and the distance from the tidal creek that
490 limits leaf litter flushing. At depth, beneath the mature *Rhizophora* stand and the transitional
491 forest, an increased organic content combined with depleted $\delta^{13}\text{C}$ values and C/N ratio increase
492 suggested an elevated contribution of vascular plant debris to the soil organic matter pool. We

493 suggest that this enrichment reflected the past mangrove forest before its destruction during the
494 war.

495 4. Soil carbon stocks in the mature *Rhizophora* forest, down to one meter, represented
496 almost three times the stock in the above-ground biomass. However, when considering only the
497 upper soil, which was related to current forest development as evidenced by $\delta^{13}\text{C}$ values and
498 C/N ratios, stocks in the soil and in the above-ground biomass were similar, and the carbon
499 burial rate was lower than $1 \text{ Mg C ha}^{-1} \text{ yr}^{-1}$.

500 In a future research effort, the net ecosystem productivity of the Can Gio mangrove
501 forest, Southern Vietnam's largest, should be studied, possibly using the eddy-covariance
502 technique. The influence of the monsoon on its productivity should also be considered.

503

504 Acknowledgements

505 This study was supported by an ARTS grant from IRD, and by the Air Liquide
506 Foundation. The authors would like to thank Mr. Tran Nhat Toan, Mr. Le Anh Tuan, and Mr.
507 Truong Van Hoang for their valuable assistance during fieldwork. We also thank Leocadie
508 Jamet and Jean-Louis Duprey from the LAMA laboratory for their assistance in the analyses.
509 We also would like to express our gratitude to the authorities of Can Gio District People
510 Committee, CGMBR, Vam Sat Ecological Tourist Centre – Phu Tho Tourist Company who
511 facilitated the collection of field data.

512

513 References

514 Adame, M.F., Kauffman, J.B., Medina, I., Gamboa, J.N., Torres, O., Caamal, J.P., Reza, M.,
515 Herrera-Silveira, J.A., 2013. Carbon Stocks of Tropical Coastal Wetlands within the

516 Karstic Landscape of the Mexican Caribbean. PLoS ONE 8, e56569.
517 <https://doi.org/10.1371/journal.pone.0056569>

518 Adame, M.F., Santini, N.S., Tovilla, C., Vázquez-Lule, A., Castro, L., Guevara, M., 2015.
519 Carbon stocks and soil sequestration rates of tropical riverine wetlands. Biogeosciences
520 12, 3805–3818. <https://doi.org/10.5194/bg-12-3805-2015>

521 Albers, T., Schmitt, K., 2015. Dyke design, floodplain restoration and mangrove co-
522 management as parts of an area coastal protection strategy for the mud coasts of the
523 Mekong Delta, Vietnam. Wetlands ecology and management 23, 991–1004.
524 <https://doi.org/10.1007/s11273-015-9441-3>

525 Alongi, D.M., 2014. Carbon Cycling and Storage in Mangrove Forests. Annual Review of
526 Marine Science 6, 195–219. <https://doi.org/10.1146/annurev-marine-010213-135020>

527 Alongi, D.M., 2012. Carbon sequestration in mangrove forests. Carbon Management 3, 313–
528 322. <https://doi.org/10.4155/cmt.12.20>

529 Alongi, D.M., 2008. Mangrove forests: Resilience, protection from tsunamis, and responses to
530 global climate change. Estuarine, Coastal and Shelf Science 76, 1–13.
531 <https://doi.org/10.1016/j.ecss.2007.08.024>

532 Alongi, D.M., Tirendi, F., Clough, B.F., 2000. Below-ground decomposition of organic matter
533 in forests of the mangroves *Rhizophora stylosa* and *Avicennia marina* along the arid coast
534 of Western Australia. Aquatic Botany 68, 97–122. [https://doi.org/10.1016/S0304-
535 3770\(00\)00110-8](https://doi.org/10.1016/S0304-3770(00)00110-8)

536 Andersen, F., Kristensen, E., 1988. Oxygen microgradients in the rhizosphere of the mangrove
537 *Avicennia marina*. Marine Ecology Progress Series 44, 201–204.
538 <https://doi.org/10.3354/meps044201>

539 Balk, M., Keuskamp, J.A., Laanbroek, H.J., 2016. Potential for Sulfate Reduction in Mangrove
540 Forest Soils: Comparison between Two Dominant Species of the Americas. *Front.*
541 *Microbiol.* 7. <https://doi.org/10.3389/fmicb.2016.01855>

542 Balke, T., Bouma, T.J., Horstman, E.M., Webb, E.L., Erfteimeijer, P.L., Herman, P.M., 2011.
543 Windows of opportunity: thresholds to mangrove seedling establishment on tidal flats.
544 *Marine Ecology Progress Series* 440, 1–9.

545 Banerjee, L.K., 1993. Influence of salinity on mangrove zonation, in: *Towards the Rational Use*
546 *of High Salinity Tolerant Plants, Tasks for Vegetation Science.* Springer, Dordrecht, pp.
547 181–186. https://doi.org/10.1007/978-94-011-1858-3_19

548 Barbier, E.B., 2006. Natural barriers to natural disasters: replanting mangroves after the
549 tsunami. *Frontiers in Ecology and the Environment* 4, 124–131.
550 [https://doi.org/10.1890/1540-9295\(2006\)004\[0124:NBTNDR\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2006)004[0124:NBTNDR]2.0.CO;2)

551 Bosire, J.O., Dahdouh-Guebas, F., Kairo, J.G., Kazungu, J., Dehairs, F., Koedam, N., 2005.
552 Litter degradation and CN dynamics in reforested mangrove plantations at Gazi Bay,
553 Kenya. *Biological Conservation* 126, 287–295.
554 <https://doi.org/10.1016/j.biocon.2005.06.007>

555 Bouillon, S., Borges, A.V., Castañeda-Moya, E., Diele, K., Dittmar, T., Duke, N.C., Kristensen,
556 E., Lee, S.Y., Marchand, C., Middelburg, J.J., others, 2008. Mangrove production and
557 carbon sinks: a revision of global budget estimates. *Global Biogeochemical Cycles* 22.
558 <https://doi.org/10.1029/2007GB003052>

559 Bouillon, S., Dahdouh-Guebas, F., Rao, A., Koedam, N., Dehairs, F., 2003. Sources of organic
560 carbon in mangrove sediments: variability and possible ecological implications.
561 *Hydrobiologia* 495, 33–39. <https://doi.org/10.1023/A:1025411506526>

562 Breithaupt, J.L., Smoak, J.M., Smith, T.J., Sanders, C.J., Hoare, A., 2012. Organic carbon burial
563 rates in mangrove sediments: Strengthening the global budget: Mangrove organic carbon
564 burial rates. *Global Biogeochemical Cycles* 26. <https://doi.org/10.1029/2012GB004375>

565 Brunt, M.A., Davies, J.E. (Eds.), 1994. *The Cayman Islands: Natural History and*
566 *Biogeography, Monographiae Biologicae*. Springer Netherlands.

567 Castillo, J.A.A., Apan, A.A., Maraseni, T.N., Salmo, S.G., 2017. Soil C quantities of mangrove
568 forests, their competing land uses, and their spatial distribution in the coast of Honda Bay,
569 Philippines. *Geoderma* 293, 82–90. <https://doi.org/10.1016/j.geoderma.2017.01.025>

570 Clough, B., 2014. *Site Assessment Guidelines for Mangrove Rehabilitation in Bac Lieu*
571 *Province, Vietnam* 41.

572 Curran, M., Cole, M., Allaway, W.G., 1986. Root Aeration and Respiration in Young Mangrove
573 Plants (*Avicennia marina* (Forsk.) Vierh.). *J Exp Bot* 37, 1225–1233.
574 <https://doi.org/10.1093/jxb/37.8.1225>

575 David, F., C. Marchand, P. Taillardat, N. Thanh Nho, and T. Meziane. (2018a). Nutritional
576 composition of suspended particulate matter in a tropical mangrove creek during a tidal
577 cycle (Can Gio, Vietnam). *Estuarine Coastal and Shelf Sciences* 200, 126-130.
578 <https://doi.org/10.1016/j.ecss.2017.10.017>

579 David, F., Meziane, T., Tran-Thi, N.-T., Truong Van, V., Thanh-Nho, N., Taillardat, P.,
580 Marchand, C., (2018b). Carbon biogeochemistry and CO₂ emissions in a human impacted
581 and mangrove dominated tropical estuary (Can Gio, Vietnam). *Biogeochemistry* 1–15.
582 <https://doi.org/10.1007/s10533-018-0444-z>

583 David, F., C. Marchand, N. Thanh-Nho, P. Taillardat, and T. Meziane. (2019). Trophic
584 relationships and basal resources utilization in the Can Gio Mangrove Biosphere Reserve
585 (Southern Vietnam). *Journal of Sea Research* 145, 35–43.
586 <https://doi.org/10.1016/j.seares.2018.12.006>

587 DelVecchia, A.G., Bruno, J.F., Benninger, L., Alperin, M., Banerjee, O., Morales, J. de D.,
588 2014. Organic carbon inventories in natural and restored Ecuadorian mangrove forests.
589 PeerJ 2, e388. <https://doi.org/10.7717/peerj.388>

590 Donato, D.C., Kauffman, J.B., Murdiyarso, D., Kurnianto, S., Stidham, M., Kanninen, M.,
591 2011. Mangroves among the most carbon-rich forests in the tropics. *Nature Geoscience*
592 4, 293–297. <https://doi.org/10.1038/ngeo1123>

593 Dung, L.V., Tue, N.T., Nhuan, M.T., Omori, K., 2016. Carbon storage in a restored mangrove
594 forest in Can Gio Mangrove Forest Park, Mekong Delta, Vietnam. *Forest Ecology and*
595 *Management* 380, 31–40. <https://doi.org/10.1016/j.foreco.2016.08.032>

596 Ellison, J.C., 1999. Impacts of sediment burial on mangroves. *Marine Pollution Bulletin* 37,
597 420–426. [https://doi.org/10.1016/S0025-326X\(98\)00122-2](https://doi.org/10.1016/S0025-326X(98)00122-2)

598 English, S.S., Wilkinson, C.C., Baker, V.V., 1997. Survey manual for tropical marine resources.
599 Australian Institute of Marine Science.

600 Fromard, F., Puig, H., Mougin, E., Marty, G., Betoulle, J.L., Cadamuro, L., 1998. Structure,
601 above-ground biomass and dynamics of mangrove ecosystems: new data from French
602 Guiana. *Oecologia* 115, 39–53. <https://doi.org/10.1007/s004420050489>

603 Giri, C., Ochieng, E., Tieszen, L.L., Zhu, Z., Singh, A., Loveland, T., Masek, J., Duke, N.,
604 2011. Status and distribution of mangrove forests of the world using earth observation
605 satellite data. *Global Ecology and Biogeography* 20, 154–159.
606 <https://doi.org/10.1111/j.1466-8238.2010.00584.x>

607 Hawkins, S., Robertson, S., Thu Thuy, P., Xuan To, P., McNally, R., Van Cuong, C., Dart, P.,
608 Xuan Phuong, P., Brown, S., Vu, N., 2010. Roots in the water: legal frameworks for
609 mangrove PES in Vietnam.

610 Hesse, P.R., 1961. Some differences between the soils of *Rhizophora* and *Avicennia* mangrove
611 swamps in Sierra Leone. *Plant and Soil* 14, 335–346.
612 <https://doi.org/10.1007/BF01666292>

613 Hiraishi, T., Krug, T., Tanabe, K., Srivastava, N., Baasansuren, J., Fukuda, M., Troxler, T.G.,
614 2014. 2013 supplement to the 2006 IPCC guidelines for national greenhouse gas
615 inventories: Wetlands. IPCC, Switzerland.

616 Hong, P.N., San, H.T., 1993. Mangroves of Vietnam. *Iucn*.

617 Hossain, M., 2014. Carbon pools and fluxes in *Bruguiera parviflora* dominated naturally
618 growing mangrove forest of Peninsular Malaysia. *Wetlands Ecol Manage* 22, 15–23.
619 <https://doi.org/10.1007/s11273-013-9318-2>.

620 Jacotot, A., Marchand, C., Rosenheim, B. E., Domack E., Allenbach, M., 2018. Mangrove
621 soil carbon stocks along an elevation gradient: influence of the late Holocene marine
622 regression (New Caledonia). *Marine Geology* 404, 60670.

623 Jennerjahn, T.C., Ittekkot, V., 1997. Organic matter in sediments in the mangrove areas and
624 adjacent continental margins of Brazil. 1. Amino acids and hexosamines. *Oceanologica Acta*
625 20, 359–369.

626 Kathiresan, K., 2003. How do mangrove forests induce sedimentation? *Revista de Biología*
627 *Tropical* 51, 355–360.

628 Kauffman, J.B., Donato, D., 2012. Protocols for the measurement, monitoring and reporting of
629 structure, biomass and carbon stocks in mangrove forests. Center for International
630 Forestry Research (CIFOR), Bogor, Indonesia.

631 Kauffman, J.B., Heider, C., Cole, T.G., Dwire, K.A., Donato, D.C., 2011. Ecosystem Carbon
632 Stocks of Micronesian Mangrove Forests. *Wetlands* 31, 343–352.
633 <https://doi.org/10.1007/s13157-011-0148-9>

634 Kendall, C.G.S.C., Skipwith, P.A. d. E., 1969. Geomorphology of a Recent Shallow-Water
635 Carbonate Province: Khor Al Bazam, Trucial Coast, Southwest Persian Gulf. GSA
636 Bulletin 80, 865–892. [https://doi.org/10.1130/0016-](https://doi.org/10.1130/0016-7606(1969)80[865:GOARSC]2.0.CO;2)
637 [7606\(1969\)80\[865:GOARSC\]2.0.CO;2](https://doi.org/10.1130/0016-7606(1969)80[865:GOARSC]2.0.CO;2)

638 Khan, M.A., Aziz, I., 2001. Salinity tolerance in some mangrove species from Pakistan.
639 Wetlands Ecology and Management 9, 229–233.
640 <https://doi.org/10.1023/A:1011112908069>

641 Komiyama, A., Ong, J.E., Pongparn, S., 2008. Allometry, biomass, and productivity of
642 mangrove forests: A review. Aquatic Botany 89, 128–137.
643 <https://doi.org/10.1016/j.aquabot.2007.12.006>

644 Komiyama, A., Pongparn, S., Kato, S., 2005. Common allometric equations for estimating the
645 tree weight of mangroves. Journal of Tropical Ecology 21, 471–477.
646 <https://doi.org/10.1017/S0266467405002476>

647 Kristensen, E., Bouillon, S., Dittmar, T., Marchand, C., 2008. Organic carbon dynamics in
648 mangrove ecosystems: A review. Aquatic Botany, Mangrove Ecology – Applications in
649 Forestry and Coastal Zone Management 89, 201–219.
650 <https://doi.org/10.1016/j.aquabot.2007.12.005>

651 Lacerda, L.D., Ittekkot, V., Patchineelam, S.R., 1995. Biogeochemistry of Mangrove Soil
652 Organic Matter: a Comparison Between Rhizophora and Avicennia Soils in South-eastern
653 Brazil. Estuarine, Coastal and Shelf Science 40, 713–720.
654 <https://doi.org/10.1006/ecss.1995.0048>

655 Lal, R., 2005. Forest soils and carbon sequestration. Forest Ecology and Management, Forest
656 Soils Research: Theory, Reality and its Role in Technology 220, 242–258.
657 <https://doi.org/10.1016/j.foreco.2005.08.015>

658 Lamb, A.L., Wilson, G.P., Leng, M.J., 2006. A review of coastal palaeoclimate and relative
659 sea-level reconstructions using $\delta^{13}\text{C}$ and C/N ratios in organic material. *Earth-Science*
660 *Reviews* 75, 29–57. <https://doi.org/10.1016/j.earscirev.2005.10.003>

661 Leopold, A., Marchand, C., Deborde, J., Chaduteau, C., Allenbach, M., 2013. Influence of
662 mangrove zonation on CO₂ fluxes at the sediment–air interface (New Caledonia).
663 *Geoderma* 202, 62–70. <https://doi.org/10.1016/j.geoderma.2013.03.008>

664 Liu, H., Ren, H., Hui, D., Wang, W., Liao, B., Cao, Q., 2014. Carbon stocks and potential
665 carbon storage in the mangrove forests of China. *Journal of Environmental Management*
666 133, 86–93. <https://doi.org/10.1016/j.jenvman.2013.11.037>

667 Lunstrum, A., Chen, L., 2014. Soil carbon stocks and accumulation in young mangrove forests.
668 *Soil Biology and Biochemistry* 75, 223–232.
669 <https://doi.org/10.1016/j.soilbio.2014.04.008>

670 Luong, N.V., Tateishi, R., Hoan, N.T., 2015. Analysis of an impact of succession in mangrove
671 forest association using remote sensing and GIS technology. *Journal of Geography and*
672 *Geology* 7, 106. <https://doi.org/10.5539/jgg.v7n1p106>

673 MacKenzie, R.A., Foulk, P.B., Klump, J.V., Weckerly, K., Purbospito, J., Murdiyarso, D.,
674 Donato, D.C., Nam, V.N., 2016. Sedimentation and belowground carbon accumulation
675 rates in mangrove forests that differ in diversity and land use: a tale of two mangroves.
676 *Wetlands ecology and management* 24, 245–261. [https://doi.org/10.1007/s11273-016-](https://doi.org/10.1007/s11273-016-9481-3)
677 [9481-3](https://doi.org/10.1007/s11273-016-9481-3)

678 Marchand, C., 2017. Soil carbon stocks and burial rates along a mangrove forest
679 chronosequence (French Guiana). *Forest Ecology and Management* 384, 92–99.
680 <https://doi.org/10.1016/j.foreco.2016.10.030>

681 Marchand, C., Allenbach, M., Lallier-Vergès, E., 2011. Relationships between heavy metals
682 distribution and organic matter cycling in mangrove sediments (Conception Bay, New
683 Caledonia). *Geoderma* 160, 444–456. <https://doi.org/10.1016/j.geoderma.2010.10.015>

684 Marchand, C., Baltzer, F., Lallier-Vergès, E., Albéric, P., 2004. Pore-water chemistry in
685 mangrove sediments: relationship with species composition and developmental stages
686 (French Guiana). *Marine Geology* 208, 361–381.
687 <https://doi.org/10.1016/j.margeo.2004.04.015>

688 Marchand, C., Disnar, J.-R., Lallier-Vergès, E., Lottier, N., 2005. Early diagenesis of
689 carbohydrates and lignin in mangrove sediments subject to variable redox conditions
690 (French Guiana). *Geochimica et Cosmochimica Acta* 69, 131–142.
691 <https://doi.org/10.1016/j.gca.2004.06.016>

692 Marchand, C., Fernandez, J.-M., Moreton, B., Landi, L., Lallier-Vergès, E., Baltzer, F., 2012.
693 The partitioning of transitional metals (Fe, Mn, Ni, Cr) in mangrove sediments
694 downstream of a ferralitized ultramafic watershed (New Caledonia). *Chemical Geology*
695 300–301, 70–80. <https://doi.org/10.1016/j.chemgeo.2012.01.018>

696 McKee, K.L., 1993. Soil physicochemical patterns and mangrove species distribution–
697 reciprocal effects. *Journal of ecology* 477–487. <http://dx.doi.org/10.2307/2261526>

698 McKee, K.L., Mendelsohn, I.A., 1987. Root metabolism in the black mangrove (*Avicennia*
699 *germinans* (L.) L): Response to hypoxia. *Environmental and Experimental Botany* 27,
700 147–156. [https://doi.org/10.1016/0098-8472\(87\)90065-7](https://doi.org/10.1016/0098-8472(87)90065-7)

701 Mizanur Rahman, M., Nabiul Islam Khan, M., Fazlul Hoque, A.K., Ahmed, I., 2015. Carbon
702 stock in the Sundarbans mangrove forest: spatial variations in vegetation types and
703 salinity zones. *Wetlands Ecology and Management* 23, 269–283.
704 <https://doi.org/10.1007/s11273-014-9379-x>

705 Mukherjee, N., Sutherland, W.J., Dicks, L., Hugé, J., Koedam, N., Dahdouh-Guebas, F., 2014.
706 Ecosystem Service Valuations of Mangrove Ecosystems to Inform Decision Making and
707 Future Valuation Exercises. *PLoS One* 9. <https://doi.org/10.1371/journal.pone.0107706>

708 Mumby, P.J., Edwards, A.J., Arias-González, J.E., Lindeman, K.C., Blackwell, P.G., Gall, A.,
709 Gorczynska, M.I., Harborne, A.R., Pescod, C.L., Renken, H., 2004. Mangroves enhance
710 the biomass of coral reef fish communities in the Caribbean. *Nature* 427, 533–536.
711 <https://doi.org/10.1038/nature02286>

712 Nagelkerken, I., Blaber, S.J.M., Bouillon, S., Green, P., Haywood, M., Kirton, L.G., Meynecke,
713 J.-O., Pawlik, J., Penrose, H.M., Sasekumar, A., Somerfield, P.J., 2008. The habitat
714 function of mangroves for terrestrial and marine fauna: A review. *Aquatic Botany,*
715 *Mangrove Ecology – Applications in Forestry and Coastal Zone Management* 89, 155–
716 185. <https://doi.org/10.1016/j.aquabot.2007.12.007>

717 Naidoo, G., Naidoo, K., 2017. Are pioneer mangroves more vulnerable to oil pollution than
718 later successional species? *Mar. Pollut. Bull.* 121, 135–142.
719 <https://doi.org/10.1016/j.marpolbul.2017.05.067>

720 Nam, V.N., Sasmito, S.D., Murdiyarso, D., Purbopuspito, J., MacKenzie, R.A., 2016. Carbon
721 stocks in artificially and naturally regenerated mangrove ecosystems in the Mekong
722 Delta. *Wetlands ecology and management* 24, 231–244. [https://doi.org/10.1007/s11273-](https://doi.org/10.1007/s11273-015-9479-2)
723 [015-9479-2](https://doi.org/10.1007/s11273-015-9479-2)

724 Noël, V., Juillot, F., Morin, G., Marchand, C., Ona-Nguema, G., Viollier, E., Prévot, F., Dublet,
725 G., Maillot, F., Delbes, L., Marakovic, G., Bargar, J.R., Brown, G.E., 2017. Oxidation of
726 Ni-Rich Mangrove Sediments after Isolation from the Sea (Dumbea Bay, New
727 Caledonia): Fe and Ni Behavior and Environmental Implications. *ACS Earth and Space*
728 *Chemistry* 1, 455–464. <https://doi.org/10.1021/acsearthspacechem.7b00005>

729 Noël, V., Marchand, C., Juillot, F., Ona-Nguema, G., Viollier, E., Marakovic, G., Olivi, L.,
730 Delbes, L., Gelebart, F., Morin, G., 2014. EXAFS analysis of iron cycling in mangrove
731 sediments downstream a lateritized ultramafic watershed (Vavouto Bay, New Caledonia).
732 *Geochimica et Cosmochimica Acta* 136, 211–228.
733 <https://doi.org/10.1016/j.gca.2014.03.019>

734 Pissierssens, P., 2002. *Manual on Sea Level Measurement and Interpretation*. UNESCO.

735 Prasad, M.B.K., Kumar, A., Ramanathan, A.L., Datta, D.K., 2017. Sources and dynamics of
736 sedimentary organic matter in Sundarban mangrove estuary from Indo-Gangetic delta.
737 *Ecological Processes* 6. <https://doi.org/10.1186/s13717-017-0076-6>

738 Proisy, C., Gratiot, N., Anthony, E.J., Gardel, A., Fromard, F., Heuret, P., 2009. Mud bank
739 colonization by opportunistic mangroves: A case study from French Guiana using lidar
740 data. *Continental Shelf Research, On the dynamics of mud deposits in coastal areas* 29,
741 632–641. <https://doi.org/10.1016/j.csr.2008.09.017>

742 Putz, F.E., Chan, H.T., 1986. Tree growth, dynamics, and productivity in a mature mangrove
743 forest in Malaysia. *Forest ecology and management* 17, 211–230.
744 [https://doi.org/10.1016/0378-1127\(86\)90113-1](https://doi.org/10.1016/0378-1127(86)90113-1)

745 Rivera-Monroy, V.H., Torres, L.A., Bahamon, N., Newmark, F., Twilley, R.R., 1999. The
746 Potential Use of Mangrove Forests as Nitrogen Sinks of Shrimp Aquaculture Pond
747 Effluents: The Role of Denitrification. *Journal of the World Aquaculture Society* 30, 12–
748 25. <https://doi.org/10.1111/j.1749-7345.1999.tb00313.x>

749 Ross, P., 1975. The mangroves of South Vietnam: the impact of military use of herbicides, in:
750 *Proceedings of the International Symposium on Biology and Management of Mangroves*.
751 Gainesville, Florida: Institute of Food and Agricultural Sciences, University of Florida.
752 pp. 695–709.

753 Saenger, P., 2002. Introduction: The Mangrove Environment, in: Mangrove Ecology,
754 Silviculture and Conservation. Springer, pp. 1–10.

755 Saenger, P., Snedaker, S.C., 1993. Pantropical trends in mangrove above-ground biomass and
756 annual litterfall. *Oecologia* 96, 293–299. <https://doi.org/10.1007/BF00317496>

757 Sanders, C.J., Smoak, J.M., Naidu, A.S., Sanders, L.M., Patchineelam, S.R., 2010. Organic
758 carbon burial in a mangrove forest, margin and intertidal mudflat. *Estuarine, Coastal and*
759 *Shelf Science* 90, 168–172. <https://doi.org/10.1016/j.ecss.2010.08.013>

760 Schmitt, K., Duke, N.C., 2016. Mangrove Management, Assessment, and Monitoring, in:
761 Tropical Forestry Handbook. Springer, pp. 1725–1759.

762 Thanh-Nho, N., Strady, E., Nhu-Trang, T.-T., David, F., Marchand, C., 2018. Trace metals
763 partitioning between particulate and dissolved phases along a tropical mangrove estuary
764 (Can Gio, Vietnam). *Chemosphere* 196, 311–322.
765 <https://doi.org/10.1016/j.chemosphere.2017.12.189>

766 Thibodeau, F.R., Nickerson, N.H., 1986. Differential Oxidation of Mangrove Substrate by
767 *Avicennia germinans* and *Rhizophora mangle*. *American Journal of Botany* 73, 512–516.
768 <https://doi.org/10.2307/2444255>

769 Tue, N.T., Dung, L.V., Nhuan, M.T., Omori, K., 2014. Carbon storage of a tropical mangrove
770 forest in Mui Ca Mau National Park, Vietnam. *Catena* 121, 119–126.
771 <https://doi.org/10.1016/j.catena.2014.05.008>

772 Ukpong, I.E., 1997. Vegetation and its relation to soil nutrient and salinity in the Calabar
773 mangrove swamp, Nigeria. *Mangroves and Salt Marshes* 1, 211–218.
774 <https://doi.org/10.1023/A:1009952700317>

775 Ukpong, L.E., 1995. Vegetation and soil acidity of a mangrove swamp in southeastern Nigeria.
776 *Soil use and management* 11, 141–144. [https://doi.org/10.1111/j.1475-](https://doi.org/10.1111/j.1475-2743.1995.tb00512.x)
777 [2743.1995.tb00512.x](https://doi.org/10.1111/j.1475-2743.1995.tb00512.x)

778 Vafaei, S., Soosani, J., Adeli, K., Fadaei, H., Naghavi, H., Pham, T.D., Tien Bui, D., 2018.
779 Improving Accuracy Estimation of Forest Aboveground Biomass Based on Incorporation
780 of ALOS-2 PALSAR-2 and Sentinel-2A Imagery and Machine Learning: A Case Study
781 of the Hyrcanian Forest Area (Iran). *Remote Sensing* 10, 172.
782 <https://doi.org/10.3390/rs10020172>

783 Vinh, T.V., Marchand C., Linh, T.V.K., Vinh, D.D., Allenbach, M. 2019. Allometric models
784 to estimate above-ground biomass and carbon stocks in a *Rhizophora apiculata* tropical
785 mangrove forest (Southern Vietnam). *Forest Ecology and Management*.
786 <https://doi.org/10.1016/j.foreco.2018.12.017>

787 Vinh, T.V., Allenbach, M., Linh, T.V.K., Marchand C., 2020. Changes in leaf litter quality
788 during its decomposition in a tropical planted mangrove forest (Can Gio, Vietnam).
789 *Frontiers in Environmental Science. Biogeochemical Dynamics*. [https://doi:](https://doi.org/10.3389/fenvs.2020.00010)
790 [10.3389/fenvs.2020.00010](https://doi.org/10.3389/fenvs.2020.00010)

791 Vo, Q.T., Oppelt, N., Leinenkugel, P., Kuenzer, C., 2013. Remote sensing in mapping
792 mangrove ecosystems—An object-based approach. *Remote Sensing* 5, 183–201.
793 [https://doi:10.3390/rs501018](https://doi.org/10.3390/rs501018)

794 Wachid, M.N., Hapsara, R.P., Cahyo, R.D., Wahyu, G.N., Syarif, A.M., Umarhadi, D.A.,
795 Fitriani, A.N., Ramadhanningrum, D.P., Widyatmanti, W., 2017. Mangrove canopy
796 density analysis using Sentinel-2A imagery satellite data, in: *IOP Conference Series:*
797 *Earth and Environmental Science*. IOP Publishing, p. 012020.

798 Walsh, G.E., 1974. Mangroves: a review, in: *Ecology of Halophytes*. Elsevier, pp. 51–174.
799 <https://doi.org/10.1016/B978-0-12-586450-3.X5001-X>

800 Wang, G., Guan, D., Peart, M.R., Chen, Y., Peng, Y., 2013. Ecosystem carbon stocks of
801 mangrove forest in Yingluo Bay, Guangdong Province of South China. *Forest Ecology*
802 *and Management* 310, 539–546. <https://doi.org/10.1016/j.foreco.2013.08.045>

803 Table 1. Vegetation structure in the different zones. MF: Mudflat; FF: Fringe forest; TF:
 804 Transition forest; IF: Interior forest; Aa: *Avicennia alba*; Ao: *Avicenia officinalis*; Ct: *Ceriops*
 805 *tagal*; Aa: *Sonneratia alba*; Ra: *Rhizophora apiculata*, AGC above-ground C stock, SC soil C
 806 stock.

Sites	Species	DBH (cm)	Tree density ha ⁻¹	Basal area (m ² ha ⁻¹)
MF	-	-	-	-
FF	<i>Aa, Ao</i>	10.7	1327	11.9
TF	<i>Aa, Ao, Ct, Sa, Ra</i>	8.5	3727	21.2
IF	<i>Ra</i>	18.6	1127	28.4

807

808 Table 2: Two-way ANOVA tests for pH, Eh and salinity values showing the effect of sampling
 809 site and seasons on soil parameters. * and *** indicate statistically significant effects for p-
 810 Values < 0.05 and < 0.001, respectively .ns means no statistically significant difference (p >
 811 0.05).

Parameters	n	Source of variation		
		Sites	Seasons	Interaction
pH	150	48.9***	5.9*	22.8***
Eh	150	19.1***	1.4 ns	11.9***
Pore-water salinity	150	1137.7***	203.4***	23.7***

812

813

814 **Figures captions**

815

816 Figure 1: Study sites location. (A) Vietnam map, (B) Can Gio Estuary and (C) Site study with
817 three transects from the mudflat to the inner forest.

818

819 Figure 2: Mangrove distribution along the transect at Can Gio mangrove, (a) and (b) distribution
820 of mangrove species from river towards the land, (c) elevation of the intertidal zone and
821 mangrove zonation.

822

823 Figure 3: Mean salinity, Eh and pH values beneath the different mangrove zones studied:
824 Mudflat (a, b, c), fringe forest (d, e, f), transitional forest (g, h, k), and inner forest (l, m, n).
825 Orange lines present the values measured during the dry season and blue lines present the values
826 measured during the rainy season.

827

828 Figure 4: C contents (a), C/N ratios (b) and $\delta^{13}\text{C}$ values (c) profiles. Black, light brown, orange,
829 blue, green lines represent mudflat, fringe forest, transition forest, and mature *Rhizophora*, (d)
830 picture of buried dead trunk of *Rhizophora* below the actual root system.

831

832 Figure 5: Ecosystem C stocks in the different sites along the elevation gradient. Green bars
833 represent above-ground C stock, grey bars represent soil C stocks from 0 to 40 cm depth, orange
834 bars represent below-ground C stocks from 40 to 100 cm depth, blue bars represent below-
835 ground C stock from 0 to 100 cm depth. Solid vertical black lines present the standard deviation
836 (SD).

837

838