



HAL
open science

Cenozoic exhumation patterns in the northern Andes: Constraints from the southern Bucaramanga Fault, Eastern Cordillera, Colombia

Francisco Velandia, Mauricio Bermúdez, Barry Kohn, Matthias Bernet, Carlos Zuluaga, Stephanie Brichau

► **To cite this version:**

Francisco Velandia, Mauricio Bermúdez, Barry Kohn, Matthias Bernet, Carlos Zuluaga, et al.. Cenozoic exhumation patterns in the northern Andes: Constraints from the southern Bucaramanga Fault, Eastern Cordillera, Colombia. *Journal of South American Earth Sciences*, 2021, 111, pp.103473. 10.1016/j.jsames.2021.103473 . insu-03466312

HAL Id: insu-03466312

<https://insu.hal.science/insu-03466312>

Submitted on 2 Mar 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Author statement

Velandia: methodology, conceptualization, funding acquisition, software, formal analysis, writing, reviewing and editing. **Bermúdez:** methodology, conceptualization, software, validation, formal analysis, writing, reviewing and editing. **Kohn:** methodology, conceptualization, software, validation, formal analysis, writing, reviewing and editing. **Bernet:** methodology, conceptualization, writing and reviewing. **Zuluaga:** methodology, conceptualization and writing. **Brichau:** methodology, conceptualization and writing.

1 **Cenozoic exhumation patterns in the Northern Andes: constraints from the southern**
 2 **Bucaramanga Fault, Eastern Cordillera, Colombia**

3 Francisco Velandia^{1,2*}, Mauricio A Bermúdez³, Barry Kohn⁴, Matthias Bernet⁵, Carlos A. Zuluaga²,
 4 Stephanie Brichau⁶

5
 6 ¹Escuela de Geología, Universidad Industrial de Santander, Cra. 27 Calle 9, Bucaramanga, Colombia.

7 * Corresponding author, Email: favelanp@uis.edu.co

8 ²Departamento de Geociencias, Universidad Nacional de Colombia, Bogotá, D.C., Colombia.

9 ³Escuela de Ingeniería Geológica, Universidad Pedagógica y Tecnológica de Colombia, Sogamoso,
 10 Colombia

11 ⁴School of Earth Sciences, University of Melbourne, Vic 310, Australia

12 ⁵Institut des Sciences de la Terre, CNRS, Université Grenoble Alpes, Grenoble Cedex 9, France

13 ⁶Géosciences Environnement Toulouse (GET), Université de Toulouse, UPS, CNRS, IRD, CNES, 14 avenue
 14 E. Belin, 31400 Toulouse, France

15

16

17 Francisco Velandia, ORCID-ID: 0000-0001-6263-0903

18 Mauricio A Bermúdez, ORCID-ID: 0000-0003-0584-4790

19 Barry Kohn, ORCID-ID: 0000-0001-5064-5454

20 Matthias Bernet, ORCID-ID: 0000-0001-5046-7520

21 Carlos Zuluaga, ORCID-ID: 0000-0002-2403-2731

22 Stephanie Brichau, ORCID-ID: 0000-0002-8073-7103

23

24 **Abstract**

25 The left lateral strike-slip Bucaramanga Fault exhibits a transpressional southern
 26 termination located towards the axial zone of the Eastern Cordillera of Colombia, where
 27 the Boyacá and Soapaga Faults are also identified as inversion-related structures. To
 28 unravel their exhumation history, we obtained apatite and zircon fission-track and (U-
 29 Th)/He ages from samples collected along different structural domains, along five vertical
 30 profiles. Joint Bayesian inverse modeling of these data reveals at least four different
 31 episodes of cooling. These are approximately: (i) 50 ± 5 Ma, (ii) 20 ± 5 Ma, (iii) 12 ± 3
 32 Ma, and (iv) 5 ± 3 Ma. The earliest pulse is associated with reactivation of the Boyacá and
 33 Soapaga Faults. The second pulse is related to the age of transpressive reactivation along
 34 the southern termination of the Bucaramanga Fault and coincides with a marked increase
 35 in relief. The Miocene-Pliocene pulses are related to Bucaramanga Fault strike-slip
 36 reactivation. Older fission-track ages previously reported from other areas of the Santander
 37 Massif suggest migration of exhumation from north to south. The four cooling episodes
 38 identified in this study can be related, within a broader geodynamic context, to interaction
 39 between the Cocos, Nazca, Caribbean, and South American plates, and the accretion of
 40 large tectonic domains of different affinity (oceanic or continental) against the South
 41 American plate during the Cenozoic. Our results are consistent with previous work

42 reported in the Santander Massif. The ages observed in the in-situ data correspond with the
43 ages found in modern river sediments and support a relief development from the Eocene to
44 the present.

45

46 **Keywords:** Low-temperature thermochronology, thermal history modelling, transpression,
47 strike-slip faults.

48

49 **Highlights**

- 50 - New thermochronologic data reported for the Eastern Cordillera of Colombia
- 51 - An exhumation pulse occurred in early Eocene time
- 52 - Significant cooling at 20 ± 5 Ma defines the onset of fault transpression and a
53 significant increase in relief.
- 54 - Miocene-Pleistocene cooling pulses are related to Bucaramanga Fault reactivation
- 55 - Exhumation along the Bucaramanga Fault migrated from north to south
- 56 - Several structural zones exhibit significant differences in thermal and exhumation
57 histories

58 **1. Introduction**

59 The axial zone of the Eastern Cordillera of Colombia is characterized by regional faults
60 such as the Soapaga, Boyacá and the southern termination of the Bucaramanga master
61 fault, along which crystalline basement outcrops in the Santander and Floresta massifs. The
62 Cordillera comprises a Mesozoic basin, which was inverted tectonically during the
63 Cenozoic (Fabre, 1983; Colletta *et al.*, 1990; Dengo and Covey, 1993; Cooper *et al.*, 1995;
64 Sarmiento, 2001; Toro *et al.*, 2004; Sarmiento-Rojas *et al.*, 2006; Tesón *et al.*, 2013). The
65 boundary between the Santander and Floresta massifs is not clearly defined, so the whole
66 area is generally considered as uplifted Mesozoic basement known as the “Santander
67 High”. Sarmiento (2001) suggested that the extension of this uplifted basement propagated
68 from the Floresta Massif in the south to the Perijá Range in the north (Figure 1).

69

70 To the south, tectonic inversion of the Santander High occurred mainly along the Boyacá
71 and Soapaga faults (Cooper *et al.*, 1995; Sarmiento, 2001; Toro *et al.*, 2004; Tesón *et al.*,
72 2013); whereas to the north, the inversion has been described along the La Salina and
73 Suárez faults (Sarmiento, 2001; Tesón *et al.*, 2013; Caballero *et al.*, 2013). During the

74 Andean orogeny, Cenozoic reactivation of the Bucaramanga, Río Servitá and Baraya
75 faults, located at the center and to the east of the current Santander Massif was significant
76 (Kammer, 1993; Corredor, 2003; Forero-Ortega *et al.*, 2020). The origin of the Eastern
77 Cordillera has been explained from the point of view of transpressional models by several
78 authors (e.g., Kammer, 1999; Taboada *et al.*, 2000; Sarmiento, 2001; Sarmiento-Rojas *et*
79 *al.*, 2006; Velandia *et al.*, 2020). Transpression is especially pronounced at the southern
80 termination of the sinistral Bucaramanga Fault (Velandia and Bermúdez, 2018; Bermúdez
81 *et al.*, 2021), the study area of this work.

82

83 Based on field observations and geo-thermochronological data, different tectonic models
84 have been proposed to explain the evolution of the Santander and Floresta massifs
85 (Shagam *et al.*, 1984; Parra *et al.*, 2009; Villagómez *et al.*, 2011; Tesón *et al.*, 2013; van
86 der Lelij, 2013; van der Lelij *et al.*, 2016a, b; Amaya *et al.*, 2017; Siravo *et al.*, 2019),
87 which record exhumation events from the Paleocene for the Eastern Cordillera, related also
88 with the initial shortening of its central part during the late Oligocene (Villamil, 1999;
89 Mora *et al.*, 2013; Egbue *et al.*, 2014; Horton *et al.*, 2020). Thermochronological studies in
90 the Santander and Floresta massifs have a dispersed regional sampling approach (Shagam
91 *et al.*, 1984; Villagómez *et al.*, 2011; van der Lelij, 2013; van der Lelij *et al.*, 2016a), or are
92 located too far from our study area (Saylor *et al.*, 2012a,b; Parra *et al.*, 2009; Ramírez-
93 Arias *et al.*, 2012; Amaya *et al.*, 2017, 2020). All these studies report cooling episodes,
94 especially the Miocene-Pliocene Andean Orogeny. At the southern Bucaramanga Fault,
95 only three thermochronological ages in the literature are from locations close to our study
96 area: two reported by van der Lelij (2016a) in San Joaquín and Onzaga, and one by Toro
97 (1990). These are insufficient however to understand the exhumation history of this part of
98 the axial zone of the Eastern Cordillera, where besides compressional structures, strike-slip
99 deformation is also reported (Velandia and Bermúdez, 2018).

100

101 Exhumation along strike-slip faults has been studied globally in several regions. Some of
102 the processes that contribute to exhumation in these zones have been described by Cao and
103 Neubauer (2016): (i) along transtensive and transpressive zones, (ii) overlapping of ancient
104 strike-slip faults, (iii) reactivation of normal or reverse faults, and (iv) oblique shear with
105 magma ascent and buoyancy by density contrasts. Exhumation along strike-slip fault
106 terminations, some related to transpression, have been studied by Spotila *et al.* (2007a, b),

107 Cruz *et al.* (2007), Furlong (2007), Umhoefer *et al.* (2007), Bermúdez *et al.* (2010; 2011),
108 and Cochran *et al.* (2017). Our research constitutes another example of exhumation along a
109 transpressive system where tectonics is the main driver for the exhumation (García-
110 Delgado *et al.*, 2021).

111

112 The goal of the present study is to characterize the exhumation and tectonic evolution at
113 the southern termination of the Bucaramanga Fault, where the spatial and temporal pattern
114 of rock exhumation is poorly constrained. For this purpose, we present new thermal history
115 models based on low-temperature thermochronology data from bedrock along five
116 “vertical” elevation profiles in that area. Since Velandia and Bermúdez (2018) described
117 the structural style for the transpressive system, including the relationship with inverted
118 faults, in this study we further elucidate the timing of the exhumation of this deformational
119 system. A corollary of this work will be to understand the mechanism of how oblique
120 motion is partitioned into vertical strain. Further, in the middle part of the Bucaramanga
121 Fault a major cooling event occurring at ~25 Ma has been reported by Amaya *et al.* (2017,
122 2020). We will also evaluate whether such cooling extended southward to the study area,
123 and if so, explain the implications in a regional context.

124

125 In this study we plan to address the following research questions: What is the role of the
126 Santa Marta-Bucaramanga strike-slip fault in the study area?. How does this fault influence
127 the exhumation patterns?. To what degree are other faults (Boyacá and Soapaga) also
128 involved in the exhumation?. Will there be differences in terms of exhumation between the
129 southern termination of the Bucaramanga fault and the Santander Massif or the Floresta
130 Massif?. Is it possible to define tectonic blocks or structural domains that exhume at
131 different rates and times?.

132

133 **2. Geological setting**

134 The core of the northern Eastern Cordillera of Colombia consists of the Floresta Massif,
135 the Santander Massif and the Perijá Range (Figure 1), which together formed the so-called
136 *Santander High* during Mesozoic extension (De Freitas *et al.*, 1997; Sarmiento, 2001;
137 Sarmiento-Rojas *et al.*, 2006). In the area where the Bucaramanga, Boyacá and Soapaga
138 faults converge, crystalline basement comprises Paleozoic to Jurassic metamorphic and
139 igneous rocks (Figure 1). This area is bounded by folded Mesozoic sedimentary rocks,

140 which once covered the Santander High. Subsequently, these sediments were eroded from
141 the central part of the mountain range during Cenozoic exhumation (Horton, *et al.*, 2010,
142 2020; Parra *et al.*, 2010; Amaya *et al.*, 2017), when tectonic activity and transpressive
143 deformation were predominant (Ramírez-Arias *et al.*, 2012).

144

145 The crystalline basement of the Santander and Floresta massifs comprises diverse rock
146 types from high-grade metamorphic rocks with sedimentary and igneous protoliths (Vargas
147 *et al.*, 1976, 1987; Ward *et al.*, 1977 a, b). These include; Precambrian (Ward *et al.*, 1977a,
148 b) or Paleozoic Bucaramanga Gneiss (Zuluaga *et al.*, 2017; Zuluaga and López, 2019),
149 Early Paleozoic migmatites, Paleozoic orthogneiss, Silurian-Ordovician low-grade
150 metamorphic (Silgará Fm.) and Silurian-Devonian metamorphic rocks (Guaca - La Virgen
151 / Floresta Fm.) (Vargas *et al.*, 1976, 1987; Ward *et al.*, 1977 a, b; Mantilla-Figueroa *et al.*,
152 2016; van der Lelij *et al.*, 2016b; Zuluaga *et al.*, 2017). Sequences of siltstones,
153 conglomerates, sandstones and limestones are also present in the massifs (Devonian
154 Floresta and Tibet formations; Carboniferous-Permian Cucho Formation, and the Paleozoic
155 Río Nevado Formation). In addition, Ordovician-Silurian and Jurassic intrusive rocks,
156 Early to Middle Jurassic volcanic and volcanoclastic rocks, as well as Late Jurassic
157 sedimentary rocks (Girón Formation) also crop out in the area (Vargas *et al.*, 1976, 1987;
158 Ward *et al.*, 1977 a, b).

159

160 A Cretaceous marine succession overlies the Girón Group, but there is no consensus about
161 the nature of the contact. The Early to Late Cretaceous marine sequence comprises units of
162 varied lithology: claystones, sandstones, limestones, grouped in the Río Negro, Tibú-
163 Mercedes, Aguardiente, Capacho, La Luna and Colón-Mito Juan formations. A Cenozoic
164 sedimentary succession overlying this sequence includes the Catatumbo basin formations
165 (Barco, Cuervos, Mirador and Carbonera formations) correlated in the Eastern Cordillera
166 with the Socha sandstones, Socha siltstones, Picacho and Concentración formations toward
167 the adjacent El Cocuy and El Tablazo sub-basins (Fabre, 1983; Jiménez *et al.*, 2014)
168 (Figure 2). Quaternary deposits are commonly found in the northeastern block of the
169 Soapaga and Bucaramanga faults, and basins located between the Soapaga and Boyacá
170 faults (Figure 1). These structures played an important role in the regional tectonic
171 framework since the early Mesozoic, which involved their activity in the half-graben
172 systems as the Cretaceous Tablazo and El Cocuy basins (Figure 1) (Fabre, 1983; Mojica

173 and Kammer, 1995; Cooper *et al.*, 1995), and others as the Yariguíes or Arcabuco –
174 Norean for the Jurassic (Velandia, 2017; Osorio-Afanador and Velandia, 2021). These
175 structures probably originated or were reactivated from a combination of rifting and back-
176 arc extensional activity related to the spreading of Pangea and the subduction of the
177 Farallón Plate beneath the South and North American plates (Pindell and Dewey, 1982).
178 The faults bounding the basins, such as the Boyacá, Bucaramanga, Soapaga or Río Servitá
179 (Figure 1), were affected by tectonic inversion during the Cenozoic and this has also been
180 documented in the northern Eastern Cordillera (Fabre, 1983; Colletta *et al.*, 1990; Dengo
181 and Covey, 1993; Cooper *et al.*, 1995; Kammer and Sánchez, 2006; Tesón *et al.*, 2013).
182 This tectonism is related to the eastern advance of the deformation, with orogenic pulses
183 occurring during the Oligocene, Miocene and Plio–Pleistocene (Parra *et al.*, 2009; Mora *et*
184 *al.*, 2013, 2015), although the compressional regime related to tectonic inversion was
185 initiated along the western flank of the cordillera in the late Paleocene – early Eocene
186 (Cortés *et al.*, 2005; Parra *et al.*, 2009; Nie *et al.*, 2012). In some cases, exhumation during
187 this period was interpreted as the product of crystalline basement erosion (Shagam *et al.*
188 1984; Horton *et al.*, 2010; van der Lelij *et al.*, 2016a). Structural models for the Eastern
189 Cordillera of Colombia and Mérida Andes in Venezuela are consistent in their agreement
190 on the role of preexisting discontinuities for controlling partition between structural
191 domains, deformation, and the erosional exhumation of these mountain belts. A large part
192 of these pre-existing discontinuities developed along the former passive margin of the
193 South American plate and played an important role in the evolution of the orogen (Jacques,
194 2004; Mora *et al.*, 2009). The distribution of thermochronological ages and the time-
195 temperature history arising from these data have allowed the definition of tectonic blocks
196 with different ages and cooling rates along the Eastern Cordillera (Shagam *et al.*, 1984;
197 Mora *et al.*, 2008, 2013, 2015; Parra *et al.*, 2009; Amaya *et al.* 2017; Pérez-Consuegra *et*
198 *al.*, 2021) and the Mérida Andes (Kohn *et al.*, 1984; Bermúdez *et al.*, 2010, 2011; van der
199 Lelij *et al.*, 2016a).

200

201 In the Floresta Massif, two N30E oriented reverse faults of possible Jurassic origin, the
202 Boyacá and Soapaga faults, were reactivated during the Andean orogeny. As these faults
203 approach the strike-slip Bucaramanga Fault, trending N15W along the Eastern Cordillera,
204 their orientation changes to the north. Most structural style models show thick and thin-
205 skinned tectonics for the boundaries of the belt (at the southern part of the Floresta Massif),

206 where pure shear compression is the predominant mechanism (Colletta *et al.*, 1990; Toro,
207 1990; Dengo and Covey, 1993; Cooper *et al.*, 1995; Roeder and Chamberlain, 1995; Toro
208 *et al.*, 2004, Saylor, *et al.*, 2012a, Tesón *et al.*, 2013). However, simple shear and sinistral
209 kinematics have also been recognized along the Eastern Cordillera (Kammer, 1996, 1999;
210 Taboada *et al.*, 2000; Sarmiento, 2001; Cediel *et al.*, 2003; Velandia, 2005; Acosta *et al.*,
211 2007, García-Delgado *et al.*, 2020; Velandia *et al.*, 2020; Machuca *et al.*, 2021).

212

213 The southern termination of the Bucaramanga Fault, where the present
214 thermochronological study was carried out, is characterized as a transpressional fault
215 system in a regional restraining bend induced by sinistral kinematics (Velandia, 2017;
216 Velandia and Bermúdez, 2018). The main trace of the strike-slip fault defines a 5 to 8 km
217 wide damage zone up to the Ricaurte site, from where the structure branches into two
218 traces; the Chaguacá and Los Micos faults (Figure 1). These form the main fractures of a
219 positive flower structure, which includes other longitudinal faults such as La Chorrera and
220 Chicamocha, together with some minor NE transverse faults (Figure 2). Taking into
221 account the sinistral kinematics of the major faults and dextral movement of the bounded
222 transverse traces, the transpressional set is defined as a domino structural style. The
223 southward advance of deformation along the transpressional system enabled the capture of
224 the Boyacá and Soapaga faults, which constitute the main inverted structures of the axial
225 zone of the Eastern Cordillera (Velandia and Bermúdez, 2018).

226

227 **3. Methods**

228 A total of 36 samples of crystalline basement (igneous and metamorphic rocks) were
229 collected along the southern termination of the Bucaramanga Fault between the town of
230 Cepitá and the Páramo (Figures 1 and 3). 21 of these samples belong to five age-elevation
231 profiles (Figures 3 and 4), with samples collected as near as vertically as possible from the
232 highest to the lowest part within the blocks. It is important to clarify that locally no faults
233 cross or cut the blocks. These profile are: (i) Ricaurte profile (samples FV-55 to 58, from
234 1026 m to 1734 m elevation), located west of the Bucaramanga Fault; (ii) El Tendido
235 profile (samples FV-43 to 46, from 1038 m to 1613 m elevation), located along the
236 Bucaramanga Fault (Figure 3); (iii) Chaguacá profile (samples FV-31, 37-39, from 2415 m
237 to 3344 m elevation), (iv) the Páramo profile (samples FV-31-35, from 3422 m and 3628
238 m elevation), located in the area delimited by the Chaguacá and Los Micos faults (Figure

239 3), and (v) San Joaquin profile (samples FV-48 to 51, from 2027 m to 2567 m elevation),
240 located between the Bucaramanga and Lebrija faults (Figure 3).

241

242 Detailed sample information is presented in Tables 1, 2 and 3. Apatite and zircon grains
243 were separated using conventional mineral separation techniques in three different
244 laboratories: Servicio Geológico Colombiano (Bogotá, Colombia), Universidad EAFIT
245 (Medellín, Colombia) and at the Université Paul Sabatier (Toulouse, France).

246

247 **3.1. Fission-track analysis**

248 The apatite and zircon fission-track (AFT and ZFT, respectively) method is based on the
249 accumulation of fission tracks or radiation damage trails (Wagner and van den Haute,
250 1992; Gallagher *et al.*, 1998). The tracks result from the spontaneous fission of heavy
251 radionuclides (with an atomic number ≥ 90 and a mass number ≥ 230). Among these
252 elements ^{232}Th , ^{235}U and ^{238}U can be measured, but it is ^{238}U which has the abundance and
253 half-life significant for geology (Tagami and O'Sullivan, 2005). For the AFT and ZFT two
254 accepted age determination are used currently: i) the external detector method (EDM), and
255 ii) the fission-track Laser Ablation Quadrupole Inductively-Coupled-Plasma Mass
256 Spectrometer (LA-Q-ICP-MS) method. For the first method, the age can be calculated
257 from counting spontaneous tracks in crystals (e.g. apatite or zircon) and induced tracks in
258 an external detector (ultra-low uranium mica), held in close contact with the crystal during
259 neutron irradiation (e.g., see Hurford and Green, 1982; Wagner and Van den Haute, 1992,
260 Gallagher *et al.*, 1998, Donelick *et al.*, 2005, and Tagami and O'Sullivan, 2005). The LA-
261 Q-ICP-MS spot analysis has been successfully employed for in-situ measurements in AFT
262 dating (e.g., Hasebe *et al.*, 2004, 2013; Chew and Donelick, 2012; Vermeesch, 2018;
263 Cogné *et al.*, 2020; Ansberque *et al.*, 2021) and ZFT dating (e.g., Hasebe *et al.*, 2013). This
264 approach is rapid as it avoids the need for irradiation and counting of induced fission-
265 tracks, limits the use of hazardous acids and can also yield additional information such as
266 apatite Cl content (Chew *et al.*, 2014).

267

268 Three apatite samples (FV-07, FV-20 and FV-21) and eight zircon samples (FV-07, FV-
269 17, FV-18, FV-20, FV-21, FV-23, FV-28 and FV-29) were prepared and processed in the
270 fission-track laboratory of the ISTERre thermochronology laboratory in Grenoble, following
271 the external detector method and zeta calibration procedure. Apatites were etched in HNO_3

272 5M, for 20 seconds at 21° C, while zircons grains were etched in a NaOH-KOH eutectic
273 solution at 228° C for 10-20 hours. Samples were irradiated at the FRM II research reactor
274 in Garching, Germany. After irradiation, muscovite external detectors were etched for 18
275 min at 20°C in 48% HF to reveal induced tracks. Samples were analyzed using a 1250×
276 dry objective with an Olympus BH2 optical microscope and the FTStage 4.04 system in
277 the ISTerre thermochronology laboratory.

278

279 Four more apatite samples (FV-44, FV-46, FV-49, and FV-57; Table 2) were analysed
280 using the fission-track LA-Q-ICP-MS method at the University of Melbourne following
281 the methodology described by Gleadow *et al.* (2015). Results for all AFT and ZFT samples
282 are presented in Tables 1 and 2. Due to sample problems, it was only possible to obtain
283 confined track lengths for two samples (FV-44 and FV-57; Table 2), these lengths
284 measurements were corrected as a function of their angle to the c-axis. Aliquots of these
285 samples were irradiated on a second mount using a ^{252}Cf source under vacuum for 140 min
286 at the Thermochronology Laboratory at the University of Melbourne to produce more
287 confined tracks for length measurement. Regardless of the fission-tracks method used
288 (EDM or LA-ICPMS), Dpar was measured and used as kinetic parameter only for those
289 samples for which confined fission-track lengths were obtained.

290

291 **3.2. (U-Th)/He Analysis**

292 (U-Th-Sm)/He thermochronology is based upon the accumulation of ^4He in minerals and
293 rocks over geologic time by the α -decay of ^{238}U , ^{235}U , ^{232}Th , and ^{147}Sm (Farley, 2002). For
294 detailed reviews of the apatite (U-Th-Sm)/He (or AHe) and zircon (U-Th)/He (or ZHe)
295 methodology see Farley and Stockli (2002) and Reiners (2005), respectively. The
296 methodology reported here was described by Gleadow *et al.* (2015). Furthermore, Durango
297 apatite (AHe reference age = 31.02 ± 1.01 Ma, McDowell *et al.*, 2005) and Fish Canyon
298 Tuff zircon (ZHe reference age = 28.3 ± 0.4 Ma, Gleadow *et al.*, 2015) were also analyzed
299 as ‘unknowns’ with each batch of apatite or zircons samples run and served as an internal
300 check on analytical accuracy (see Tables 3 and 4, respectively). AHe data for 52 single
301 grains from 16 samples were acquired, additionally ZHe data from 121 individual grains
302 from 32 samples were obtained, and results are presented in Tables 3 and 4. In these tables
303 we use the age columns, since they were corrected for factors such as grain geometry,
304 alpha radiation damage, amongst others. Different factors that can contribute in different

305 degrees to single-grain AHe and ZHe age dispersion include grain size, radiation damage,
306 grain fragmentation, U and/or Th zonation, complex eU intra-grain relationships, He
307 implantation, uncertainties in the alpha ejection factor calculation, mineral inclusions, He-
308 rich fluid inclusions and lengthy residence in the He partial retention zone (e.g., Wildman
309 *et al.*, 2016; Danišik *et al.*, 2017; Hueck *et al.*, 2018). Furthermore, imperfections, such as
310 vacancy damage and crystallographic micro-voids may also affect He diffusion (Gerin *et*
311 *al.*, 2017; Zeitler *et al.*, 2017). From all these factors a significant variation of
312 compositional and crystallographic characteristics could produce an intra-sample
313 dispersion at He ages above 50% (e.g. Wolfe and Stockli, 2010; Wildman *et al.*, 2016). It
314 is possible to reduce a part of this possible source of error by crystal selection (Danišik *et*
315 *al.*, 2017). In this study, we took into account grain size, morphology and accumulated
316 radiation damage. The corrected age was calculated as the measured age/Ft, where Ft is the
317 total fraction of alpha decays (see Farley *et al.*, 1996; Fitzgerald *et al.*, 2006, and Zeitler,
318 2015 for more details). For data interpretation and thermal history modelling, similar to
319 other works (e.g. Boone *et al.*, 2018), we use the effective uranium concentration
320 $eU=U+0.235\times Th$ as a proxy for α -radiation damage.

321

322 **4. Results**

323 **4.1 Fission/track analysis**

324 Regarding the three AFT ages obtained using the external detector method, the late
325 Oligocene-early Miocene age for FV-07 (Table 1) located near Cepitá town (Figure 3), is
326 comparable with an age reported by Amaya *et al.* (2017, 2020) for the same western block
327 of the Bucaramanga Fault. The other two samples, which are separated by the Pueblo Viejo
328 fault and sinistral Chaguacá fault (Figure 3), yield early Miocene (FV-21 to the west) and
329 late Miocene (FV-20 to the east) ages, which are similar to AFT ages of two samples
330 previously reported for the area (10VDL32, and 10VDL28 in van der Lelij *et al.*, 2016a)
331 (Figure 3). Moreover, for AFT ages determined using the LA-ICP-MS method (Table 2,
332 Figure 3) three show an early Miocene age and one a late Miocene age.

333

334 ZFT data show the oldest ages (Lower Cretaceous) in two samples located in the southern
335 part of study area (FV-23 and FV-28), separated by the Boyacá and Chaguacá faults
336 (Figure 3). The two ZFT samples for Chaguacá fault (FV-21 and FV-20) show Paleocene-

337 Eocene ages, while three other ZFT samples yield Eocene-Oligocene ages (FV-17, FV-18,
338 FV-29) (Table 1, Figure 3).

339

340 **4.2 (U-Th)/He analysis**

341 Single grain AHe range from 3.7 ± 0.2 to 42.5 ± 2.6 Ma (Table 3). Table 4 summarizes
342 ZHe data for 118 individual grains from 31 samples. Except for two single grains the
343 morphology for zircons has both terminations (2T). Single-grain ZHe ages range from 3.5
344 ± 0.2 Ma to 227.7 ± 14.1 Ma, and intra-sample dispersion is large in 17 samples. Before
345 estimating weighted (U-Th)/He ages because of issues which may result in age dispersion
346 mentioned above (section 3.2), it is necessary to justify what is considered to be an outlier
347 in an intra-sample data set. However, in some samples ages replicate reasonably well
348 within associated uncertainties (e.g. AHe FV-40, FV-44, FV-45, FV-54, FV-56, FV-61 or
349 ZHe FV-31, FV-35, FV-39, FV-44, FV-45, FV-47, FV-52, FV-59, FV-60 and FV-61).

350

351 We used the following as exclusion criteria: i) Grain morphology, grains without one or
352 both terminations were excluded; ii) seemingly lower or higher [eU] ppm outliers,
353 compared to the remaining grains, although higher [eU] ppm could affect closure
354 temperature and could provide some additional information on the cooling history during
355 numerical inversion of our dataset. We tested the inclusion of these grains but did not find
356 any significant differences in numerical models when including these ages and when
357 considering constraint-free models. This is shown in Figure DR-1; iii) Following Fukuda *et*
358 *al.* (2020), we calculated weighted mean age of samples, where more than three intra-
359 sample analyses were used, employing IsoplotR (Vermeesch, 2018). We verified whether
360 grain ages within the error range of $\pm 3\sigma$ were consistent with a weighted mean age ranging
361 within $\pm 95\%$ confidence interval. When single grain ages did not overlap within the
362 weighted mean age of a sample, the grain ages were rejected as outliers and the weighted
363 mean age without such outliers was recalculated, and iv) based on a “quasi-plateau age” or
364 flatline trend (e.g. Johnson *et al.*, 2017) to $\sim 20 \pm 5$ Ma. We used this “plateau” criterion,
365 which was derived from a first-order observation. In order to understand the different
366 factors that could produce age dispersion, a plot of [eU] versus ZHe ages is shown in
367 Figure 5A which considers all grains including “outliers”. Although no trend is observed, it
368 is possible to discriminate an alignment or “plateau” of significant number of individual
369 grains ages around 20 ± 5 Ma. This plateau is observed for both high and relatively low

370 values of [eU] (Figures 5B and C). It is suggested that this plateau may be associated with
371 a distinct cooling event in the study area, which is the reason why it is taken into account
372 as a reference value when calculating the weighted mean AHe and ZHe ages for the
373 individual grains in a sample. Using IsoplotR (Vermeesch, 2018) we calculated weighted
374 mean ages for each sample (Tables 3 and 4).

375

376 **5. Interpretation and discussion**

377 A common approach to estimating exhumation rates is to collect and date a suite of
378 samples within a tectonic block according to elevation, in certain cases (e.g. when there is
379 not a combination of advection and isotherm compression, the effect of topography on
380 near-surface isotherms, and where samples are collected across the topography or the
381 topographic wavelength) the slope of these age-elevation relationship (AER) can be
382 considered as the exhumation rate (Braun, 2002), where changes or breaks in slope can be
383 related as changes in exhumation rate through time (e.g. Wagner and Reimer, 1972;
384 Wagner *et al.*, 1977; Gleadow and Fitzgerald, 1987; Fitzgerald *et al.*, 1995; Fitzgerald and
385 Malusà, 2019). Thus, the age-elevation approach can be considered as a 1D-interpretative
386 approach considering the effect of the cooling age distribution at the surface, because
387 cooling ages increase with sample elevation in proportion to the exhumation rate. This
388 way, thermal models are frequently used to investigate exhumation rates (Huntington *et*
389 *al.*, 2007).

390

391 During the exhumation/cooling process the concept of closure temperature defined by
392 Dodson (1973) as the temperature of a mineral (e.g., apatite or zircon) at the time given by
393 its radiometric age is very important, because the daughter product is essentially retained in
394 a crystal below the closure temperature range but not above it. This facilitates the
395 interpretation of thermochronological data, but often these are only valid for minerals that
396 experienced decreasing temperatures with time (or monotonic cooling; Dodson, 1973).
397 Closure temperature will vary depending on how quickly a sample is cooled, grain size and
398 mineral composition. Thus, faster cooling results in higher closure temperatures and vice
399 versa (e.g., Reiners and Brandon, 2006). Each thermochronometer has a temperature
400 window over which the daughter product starts to be retained or annealed within the
401 system, since it was related to the He Partial Retention Zone (HePRZ) (Wolf *et al.*, 1998)
402 and Partial Annealing Zone (PAZ) for the fission-track method (Gleadow and Fitzgerald,

403 1987), respectively. In the PAZ fission-tracks lengths are shortened. ^4He can be considered
404 as an almost open system (Wolfe and Stockli, 2010), where some He diffusion information
405 is retained, particularly for apatite the information is partially retained between $\sim 40^\circ\text{C}$ and
406 70°C , and typically not retained above $\sim 80^\circ\text{C}$ (Farley, 2000, 2002). The closure
407 temperature for He in zircon, in contrast, is typically $\sim 170\text{--}190^\circ\text{C}$ (Reiners *et al.*, 2004),
408 and the PRZ $\sim 130\text{--}180^\circ\text{C}$ (Reiners and Brandon, 2006; Wolfe and Stockli, 2010), but these
409 ranges may vary considerably as a function of radiation damage (e.g. Shuster *et al.*, 2006;
410 Guenther *et al.*, 2013). For the fission-track technique, all fission-tracks are annealed and
411 their concentration, and thus age, is zero above $\sim 120^\circ\text{C}$ in apatite (Laslett *et al.*, 1987;
412 Ketcham *et al.*, 1999), and $\sim 240^\circ\text{C}$ in zircon (Zaun and Wagner, 1985). Partial annealing
413 of fission tracks typically occurs between $\sim 60^\circ\text{C}$ and 120°C in apatite, depending on
414 apatite composition (Green *et al.*, 1989) and between $\sim 180^\circ\text{C}$ and 350°C for zircon,
415 depending on the degree of radiation damage (Tagami, 2005). An exhumed PAZ/PRZ may
416 be revealed in an AER when, following a period of uplift/exhumation, the form of the
417 PAZ/PRZ is retained with a 'sharp' convex inflexion (the 'break in slope') marking the base
418 of the exhumed PAZ/PRZ with reset samples structurally beneath (Gleadow and
419 Fitzgerald, 1987; Fitzgerald and Gleadow, 1988, 1990; Gleadow, 1990; Stockli *et al.*,
420 2000; Fitzgerald and Malusà, 2019).

421

422 In order to facilitate discussion and discriminate between some trends in the data, the
423 weighted mean ages (Tables 3 and 4) plotted versus elevation are presented in Figure 6.
424 We show AER for both methods AHe (Figure 6A) and ZHe (Figure 6B). Based on the
425 frequency of the ages and the errors, we discriminate some rapid cooling events, which are
426 drawn as grey bands in this figure. In particular, the age of 20 ± 5 Ma discriminated as a
427 'plateau' age (Figure 4). This age and other three populations (5 ± 3 Ma, 12 ± 3 Ma, and 50
428 ± 5 Ma) are defined based on detrital modern river apatite fission-track ages distributions
429 (Bermúdez *et al.*, 2021) around the Onzaga and Los Micos river (Figure 1), these two
430 basins cover a considerable portion of our study area. We use weighted ages in this figure
431 because it allows us to show more clearly the most important cooling events in the study
432 area, either by the distribution of detrital ages or by important regional events reported by
433 van der Hammen (1961) and summarized by Restrepo-Moreno *et al.* (2019).

434

435 Assuming that the isotherms corresponding to the closure temperature remained essentially
436 parallel, and the relief did not change through exhumation, in Figure 6 the AER are used to
437 infer “apparent” exhumation rates which could be compared with cooling rates derived
438 from the numerical model of thermal histories. Table 5 summarizes the estimated
439 confidence interval for exhumation rates based on AER. These “apparent” exhumation
440 rates were calculated using the Williamson (1968) regression line, which takes into
441 account the errors on ages. For this regression line we assume that the dependent variable
442 (y-axis) is the (U-Th)/He age and the independent variable (x-axis) is the elevation, thus
443 exhumation rates are calculated as the reciprocal of the slope. When the rate of exhumation
444 is very high, or the regression line is near vertical (infinite slope) this method has the
445 advantage of allowing the detection of variations in exhumation rates by considering errors
446 on slopes and intercepts. Basically, in our dataset it is possible to discriminate three
447 different trends. AER relationships for the Páramo and San Joaquín profiles indicate
448 exhumation rates below 0.5 km my^{-1} at $50 \pm 5 \text{ Ma}$. Regardless of our assumptions for
449 calculating apparent exhumation rates from the AER based on ZHe data, the result is
450 negative or very high at $20 \pm 5 \text{ Ma}$. This suggests a significant change in relief for the
451 Chaguacá and Los Micos profiles. In these areas exhumation rates are $> 1 \text{ km my}^{-1}$. This
452 rate is in good agreement with exhumation rates calculated in these zones by Bermúdez *et*
453 *al.* (2021) assuming that the topography is spatially variable using two different methods
454 BIN2E (Braun *et al.*, 2018) and thermo-kinematic 3D modelling (PeCube; Braun, 2003;
455 Braun *et al.*, 2012). Our results suggest that the ZHe ‘plateau’ age of $20 \pm 5 \text{ Ma}$ could be
456 related to an important cooling phase related to significant topographic amplification. We
457 emphasize that not all profiles have a considerable difference in elevation range (>1000
458 meters) and this could explain why some exhumation rates are extremely low. Exhumation
459 rates derived from AHe ages in the 5 to 15 Ma time range for the Los Micos fault profile
460 and Ricaurte profile are $< 0.2 \text{ km my}^{-1}$.

461

462 It should be noted that based on scarce data in Figure 6, we do speculate on the possibility
463 of the existence of two intervals of tectonic quiescence: (i) 68 to 60 Ma (Maastrichtian to
464 early Paleocene) and (ii) 36 to 34 Ma (late Eocene). These intervals of “apparent” tectonic
465 quiescent allow for the separation of three long intervals where data could indicate
466 exhumation phases: a first from 84 to 72 Ma (Santonian to Campanian), a second with a
467 significant increase in data between 60 and 33 Ma (middle Paleocene to late Eocene) and a

468 final phase based on more data between 32 and 5 Ma (Oligocene-Miocene). To avoid over-
469 interpretation of our data, numerical thermal history models have been generated for
470 comparison.

471

472 **5.1. Inverse Thermal History Modeling**

473 For all the models, AHe, ZHe, and AFT ages of representative samples were used,
474 however in those cases where ZFT ages existed, these were incorporated for determining
475 time-temperature paths through inverse Bayesian transdimensional Markov chain Monte
476 Carlo modeling using the QTQt 5.7.3 program (Gallagher *et al.*, 2009; Gallagher, 2012).
477 This software permits thermal histories to be calculated from low temperature
478 thermochronology data using multiple samples. Additionally, QTQt has an option for
479 considering burial and reheating episodes scenarios (Fox and Carter, 2020). This option is
480 useful to infer possible events associated with major fault activity. Although the software
481 produces four different time-temperature statistical models (expected, mode, maximum-
482 Likelihood and posterior models), here we only considered the Expected model, which
483 corresponds to the weighted mean of the posterior distribution and provides more regular
484 curves and variation rates, compared to other statistical outputs.

485

486 For each sample the oldest ZHe weighted age (Tables 3 and 4) and temperature ($100 \pm$
487 100°C) ranges were used for the general prior (model space). The present-day surface
488 temperature was set as $15 \pm 5^\circ\text{C}$. Although it appears that the constraints used in our QTQt
489 philosophy modeling were forced, we considered four different independent constraints
490 based on two aspects: first, the four main reported pulses thermotectonic/orogenic phases
491 in the Eastern Cordillera of Colombia (Restrepo-Moreno *et al.*, 2019). Second, the
492 preferred time-temperature history derived from the best PeCube model for the study area
493 (see Figure 12c in Bermúdez *et al.*, 2021). We are interested in corroborating the timing of
494 exhumation onset in the study area from at least 35 Ma to the present (see Figure 12C in
495 Bermúdez *et al.*, 2021). That PeCube model (Bermúdez *et al.*, 2021) is constraint
496 independent and the range of parameters run was quite wide, plus it was possible to
497 corroborate equilibrium versus time-varying topography conditions. Thus, we used the
498 following four constraints: (i) 50 ± 10 Ma with temperatures of $160^\circ \pm 40^\circ\text{C}$; (ii) 20 ± 5
499 Ma with temperatures of $120^\circ \pm 60^\circ\text{C}$, (iii) 12 ± 3 Ma with temperatures of $100^\circ \pm 40^\circ\text{C}$,
500 and (iv) 5 ± 3 Ma with temperatures of $80^\circ \pm 40^\circ\text{C}$. Because models start at 50 ± 10 Ma

501 and all temperatures overlap for these different time intervals, the constraints used are even
502 wider than those provided by the best PeCube model (Bermúdez *et al.*, 2021), to give some
503 freedom to the thermal histories. QTQt generates time-Temperature paths for two different
504 Markov chains (birth and death), where a chain is a statistical perturbation of the other
505 (Gallagher, 2012). Our modeling strategy was designed to use the QTQt software with a
506 dual purpose: (i) to model individual samples closer to and away from the main faults
507 (Figure 7) and (ii) to model all samples belonging to each vertical profile separately
508 considered in this study (Figure 8).

509

510 For two samples FV-44 (El Tendido) and FV-57 (Ricaurte) AFT age, track lengths, Dpar
511 measurements and individual AHe, and ZHe ages were used to determine time-temperature
512 paths (Figure 9A, B). For AFT data we selected the multi-kinetic annealing model of
513 Ketcham *et al.* (2007b), with c- axis projected track length data (Ketcham *et al.*, 2007a).
514 As there were AFT, AHe, ZHe ages and MTL measurements from these two samples,
515 which belong to distinct blocks, they were modeled individually and compared with
516 models from the different profiles.

517

518 **5.1.1 Cooling histories along the Los Micos and Chaguacá faults**

519 The time-temperature path for the Chaguacá Fault (Figure 7A) suggests an onset of cooling
520 at ~24 Ma for all samples, and a long residence or slight heating pulse from 24 to 8 Ma
521 closer to the apatite PAZ occurred. Between 8 and 5 Ma a cooling pulse is recorded which
522 occurred at much faster rate than the first pulse (24 Ma). Probably in this short period the
523 rocks reached the near-surface. Finally, from 5 Ma to the present, the cooling rate
524 decreased or possibly there was heating due to fault activity. This last possibly could also
525 be an artifact of the modeling to be able to adjust to the present-day temperature. Although
526 our data do not support this assumption, when we analyzed models for the El Tendido
527 profile, in particular, sample FV-44 the data does support the result. In this case there are
528 individual grains (younger ages) that can support the modeled time-Temperature path for
529 this sample. For modeling we mainly used samples where both AHe and ZHe data (FV-31
530 and FV-59) were available, and we also added samples FV-38 and FV-39 that only had
531 ZHe ages (Tables 3 and 4) because they are located parallel to the Chaguacá Fault. Of all
532 the samples used to model this area (FV31, FV-38, FV-59 and FV-60) the northernmost
533 sample FV-59 exhibits a more complex cooling-heating history. This could be explained

534 by its location in a small tectonic block, where the Bucaramanga Fault bifurcates into the
535 Chaguacá and Los Micos faults. A comparison of predicted versus observed ages is shown
536 in the center of Figure 7A, the colors in the figure represent the elevation of the samples;
537 note that the obtained likelihood is -13.48 which means that the differences between the
538 observed and model-predicted ages are very small. At the base of Figure 7A the
539 predictions of the model are shown in the form of AER.

540

541 Expected t-T models derived for each sample located along the Los Micos Fault (Figure
542 7B), suggest prolonged residence within the ZHe PRZ before 25 Ma. Following this time,
543 three successive cooling phases are observed: 1) a fast pulse between ~25 to 16 Ma, 2) a
544 slow pulse compared to the first one between ~16 and 7 Ma, and finally 3) a fast-cooling
545 pulse between ~7 Ma to present. This last episode is consistent with AFT and AHe ages in
546 that area (e.g. FV-20, 10VDL28 and FV-45; Figure 3). Figure 7B (central part of the panel)
547 compares ages predicted by the model and observed ages. Again, for this model, samples
548 FV-40, FV-54 and FV-61 were selected, which had both AHe and ZHe age data (Tables 2,
549 3 and 4). The comparison between observed and predicted ages in the form of AER is
550 shown at the base of Figure 7B. At an elevation between 2000 to 2500 m, a break in the
551 EAR slope is observed. The model allows kinetic parameters to be predicted for the AFT
552 method, such as Dpar.

553

554 **5.1.2 Cooling histories for vertical profiles**

555 All AFT, AHe, and ZHe ages for each profile (Figure 6) were modeled jointly (Figure 8).
556 It was not possible to incorporate ZFT ages here since such data was not available for the
557 profiles. Figure 8 is again split into three sectors; a left panel presenting the time-
558 Temperature paths predicted by QtQT, a central panel presenting comparisons between
559 predicted and observed ages, and a right panel showing the central panel comparisons in
560 the form of age-elevation relationships (AER). In general, in the interval between 60 and
561 ~25 Ma, the different tectonic blocks experience a long residence in the ZHe PRZ (140-
562 180 °C; Ricaurte), AFT PAZ (San Joaquín), or very slow cooling (El Tendido, Chaguacá,
563 and Páramo). From ~25 Ma to present the blocks exhibit contrasting time-Temperature
564 histories. The Ricaurte profile exhibits rapid cooling from 25 Ma to 12 Ma (Oligocene -
565 Middle Miocene), slow cooling from ~12 Ma to 4 Ma, and an increased cooling rate from
566 4 Ma to the present. The El Tendido profile experienced a slow cooling pulse from ~50 Ma

567 to 25 Ma, a slight increase in cooling rates from ~18 Ma to 14 Ma (Oligocene - early
568 Miocene), a slight reduction in cooling rate from ~16 Ma to 5 Ma and finally from ~5 Ma
569 to the present a rapid cooling pulse. In the San Joaquin structural domain a lengthy
570 residence in the PAZ AFT from ~60 Ma to 22 Ma appears to be followed by a relatively
571 fast continuous (4.4 °C/Ma) cooling pulse from ~22 Ma to the present. Although mean
572 fission-tracks lengths measurements would be required to confirm that scenario. For the
573 Chaguacá profile only three samples with ZHe ages were used to model numerically the
574 time-Temperature history. Thus, the time-temperature history below the ZHe PRZ cannot
575 be interpreted. In the Páramo, slow and continuous cooling occurred through the ZHe PRZ
576 from ~50 Ma to 16 Ma, followed by an increase in the cooling rate from ~16 Ma to 6 Ma
577 and a slight reduction in the cooling rate from ~6 Ma to the present.

578

579 Based on the four well defined thermotectonic/orogenic pulses defined by van der
580 Hammen (1961), supported by low temperature thermochronology (Restrepo-Moreno *et*
581 *al.*, 2019) and the discrete tectonic pulses discriminated from detrital dataset and PeCube
582 modelling (Bermúdez *et al.*, 2021) we can confirm several cooling pulses previously
583 identified. The first pulse occurred before ~25 Ma (Oligocene) was discerned in the San
584 Joaquín and Páramo profiles, and in the western side of the Bucaramanga Fault (Figure 1).
585 This is also observed in the San Joaquín and Páramo profiles, with slow cooling rates
586 (<4°C/Ma), which are located furthest away from the main strand of the Bucaramanga
587 Fault. The similarity between both profiles may indicate that the Chaguacá Fault might not
588 have been active at that time. In the models, most samples from the four profiles indicate
589 the onset of a cooling pulse between ~22 to 15 Ma (Oligocene - early Miocene). This
590 coincides with the onset of Bucaramanga Fault activity, while the most recent pulse
591 recorded between ~6 to 2 Ma for samples in the El Tendido, Chaguacá and Ricaurte
592 profiles is associated with subsequent recent activity of the Bucaramanga Fault and its
593 subsidiary branches in that area. The Ricaurte and El Tendido profiles exhibit complex
594 thermal histories, possibly related to reactivation of the Los Micos and Bucaramanga
595 faults.

596

597 For samples FV-44 (El Tendido) and FV-57 (Ricaurte) weighted AHe and ZHe ages, AFT
598 ages, fission-track lengths and Dpar measurements, were used as input for determining
599 time- temperature paths (Figure 9A and B) using QTQt. El Tendido shows a very slow

600 cooling rate ($2^{\circ}\text{C}/\text{Ma}$) from ~ 50 Ma to 8 Ma with temperatures ranging between $\sim 200^{\circ}\text{C}$ to
601 125°C . From 8 Ma to the present this sample exhibited a fast cooling rate ($\sim 13^{\circ}\text{C}/\text{Ma}$). The
602 histogram located at the right of the Figure 9A represent the fission-track length
603 distribution. The values in this figure correspond to observed (O) and predicted (P) data.

604

605 Figure 9B shows the predicted t-T path for the Ricaurte profile, the Expected model
606 suggests lengthy residence in the zircon PRZ, with heating, possibly due to burial episode
607 between ~ 22 Ma to 16 Ma, whereas a faster cooling episode ($\sim 21^{\circ}\text{C}/\text{Ma}$) began from ~ 16
608 Ma to 12 Ma. From ~ 12 Ma to 2.5 Ma the cooling rate decreased to $\sim 0.1^{\circ}\text{C}/\text{Ma}$ and the
609 sample remained within the AHe PRZ. From ~ 2.5 Ma to present the cooling rate appears to
610 have increased, although this should be confirmed through the use of other more
611 strategically located samples.

612

613 **5.2 Relationship between cooling/heating pulses and fault reactivation**

614 In most of the profiles cooling episodes are recorded between ~ 25 to 15 Ma. For the San
615 Joaquin, and Páramo profiles (Figure 8) cooling events prior to ~ 25 Ma could be not
616 associated with the Bucaramanga Fault reactivation, instead, these were probably related to
617 reactivation of the Boyacá and Soapaga faults at two different times, $\sim 50 \pm 5$ Ma and $\sim 36 \pm$
618 3 Ma (Ricaurte and Páramo; Fig. 7C), i.e. in early and late Eocene time, respectively. The
619 late Eocene pulse is comparable with the onset of cooling between ~ 40 and 30 Ma for the
620 Floresta Massif (Figure 1), based on AFT data reported by Parra *et al.* (2009). These pulses
621 are related to displacement of the Soapaga Fault, whose onset in turn extends to ~ 50 Ma,
622 similarly to ages reported in the present work. In contrast, heating is observed in the
623 Chaguacá Fault (Figure 7A) and in sample FV-57 from the Ricaurte profiles (Figure 9B).
624 These apparent heating "pulses" are possibly related to local advection of isotherms
625 resulting from fault activity, or associated phenomena resulting from either shear heating
626 or fluid flow, which could perturb the AHe ages (Reiners *et al.*, 2005).

627

628 Our modeled ages indicate that mountain building mainly occurred during the Cenozoic,
629 with significant exhumation, especially in the southern and western Santander High.
630 Eocene exhumation is a well-defined pulse observed in the detrital dataset (Bermúdez *et*
631 *al.*, 2021) and this was probably controlled by early activity of the Boyacá fault. The low
632 Oligocene exhumation rates are in concordance with the phase of orogen stabilization

633 reported by Mora *et al.* (2010, 2015) for this time in the northeastern Eastern Cordillera.
634 This differs from ages reported in other sectors around the study area, particularly to the
635 east of the Soapaga Fault (Figure 1). We note that this sector, located on the hanging wall
636 (west part) of the fault, exhibits an apparent tectonic quiescence or slow exhumation
637 between ~36 and 22 Ma. The reported range of ages here is similar to those reported by
638 Saylor *et al.* (2012a) suggesting that to the south, in the Floresta Massif, the exhumation of
639 the Soapaga Fault occurred after that related to activity of the Boyacá Fault (between 3~9
640 and 36 Ma), suggesting a migration of deformation to the east, preceding activity of the
641 Bucaramanga Fault.

642

643 A strong regional exhumation pulse during the early Miocene (between 20 ± 5 Ma, and 12
644 ± 3 Ma) suggests an important role of the Bucaramanga Fault. Similar ages are also
645 reported in AFT samples from this work (FV-17 and FV-21, Figure 3) and from Shagam *et*
646 *al.* (1984), van der Lelij *et al.* (2016b) and Amaya *et al.* (2017, 2020) in the Cepitá sector,
647 in addition to Parra *et al.* (2009) in the Massif of Floresta, where Toro (1990) also reports
648 an AFT age of 22.3 ± 4 Ma. Villagómez (2010) using a sparse AFT age distribution on the
649 west side of the Santander Massif suggests that cooling occurred between 23 and 16 Ma
650 (early Oligocene-Miocene). The timing of ~25 to 20 Ma that Amaya *et al.* (2017) report for
651 cooling in the Cepitá zone (Figure 3) may indicate that the pulse had migrated to the east
652 and in that case, the onset of the tectonic activity of the Bucaramanga Fault and associated
653 exhumation may have migrated from north to south. Van der Lelij *et al.* (2016b) suggest
654 that the fault was already active at 40 Ma (37 Ma for Amaya *et al.*, 2017) and that
655 exhumation was variable along it. However, at least to the south (in our study area), the
656 ~35 Ma exhumation pulse (Bermúdez *et al.*, 2021) is more related to the late Eocene pulse
657 that exhumed the western part of the Floresta Massif due to the inversion of the Boyacá
658 Fault, whose prolonged activity to the north could also have had some influence on
659 Santander Massif tectonism.

660

661 The last cooling pulses from ~5 Ma in the El Tendido profile and ~2 Ma in the Ricaurte
662 profile coincide with exhumation events reported by Shagam *et al.* (1984) and Amaya *et*
663 *al.* (2017, 2020) with similar AFT ages of 5 Ma, in samples located closer to secondary
664 faults associated with the Bucaramanga Fault (Figure 3). This pulse is of relevance for
665 some sectors across the Eastern Cordillera and can be related to an increase in tectonic

666 activity between ~6 and 3 Ma as evidenced by data from the Guaicáramo Fault system in
667 the piedmont (Bayona *et al.*, 2008) and sedimentation of a thick late Miocene to Pliocene
668 facies for the Los Llanos basin reported by Parra *et al.* (2010). Miocene-Pliocene tectonic-
669 exhumation events have also been reported from across the entire Maracaibo block
670 boundaries (Shagam *et al.*, 1984; Kohn *et al.*, 1984; Audemard and Audemard, 2002;
671 Audemard, 2003; Bermúdez *et al.*, 2019).

672

673 **5.3 Relief change, exhumation history, sedimentary record and geodynamic context**

674 A decrease in relief leads to a further overestimate of the exhumation rate from the AER
675 (Braun, 2002). Thus, a significant decrease in relief could lead to a negative slope. Such a
676 negative AER is observed in Figure 6B for the El Tendido, Ricaurte, Los Micos Fault and
677 Chaguaca profiles (Figure 3) around the interval $\sim 20 \pm 5$ Ma. Based on the previous
678 information, the calculation of exhumation rates from age-elevation profiles should not be
679 carried out, but rather we did the exercise for the purpose of comparison with exhumation
680 rates derived from detrital thermochronological data for the same study area by Bermúdez
681 *et al.* (2021). Assuming steady-state topography and spatially variable topography over
682 time for the Chaguacá and Onzaga river catchments (Figure 1), Bermúdez *et al.* (2021)
683 using 3D thermos-kinematic modeling (PeCube; Braun, 2003; Braun *et al.*, 2012)
684 reproduced the distributions of detrital apatite and zircon fission-track ages. The best fits
685 between the predicted and observed cumulative density function (CDF) are achieved when
686 the amplification factor for topography or relief variation is assumed to be changing over
687 time. These results suggest that prior to ~40 Ma some 10 to 50% of the present-day
688 topography existed, with the balance being created in the interval between ~40 and 20 Ma.
689 The PeCube modelling of Bermúdez *et al.* (2021) suggests that exhumation rates in the
690 study area evolved from ~ 2 km my^{-1} between ~40 and 30 Ma, decreasing to values < 0.1
691 km my^{-1} from ~30 to 25 Ma, and again increased to 0.5 km my^{-1} from ~20 Ma to the
692 present. In Table 5, numbers in italics show that despite the possible overestimation of
693 AER ratios, the estimated rates are very close to those reported by Bermúdez *et al.* (2021).

694

695 The late Miocene cooling pulse recorded in our study area can be related to higher
696 exhumation rates resulting in coarse-grained geological units in adjacent basins to the
697 study area (El Cocuy and El Tablazo; Fabre, 1983; Jiménez *et al.*, 2014 – Figures 1, and 2).

698 In these basins during the late Miocene pulse there is important stratigraphic evidence for
699 erosion such as the Tilatá Formation, which outcrops in the axial zone of the Cordillera.

700

701 The latest exhumation pulse that commenced between ~5 and 2 Ma (Plio-Pleistocene,
702 Figure 7) can be linked to exhumation observed along the Bucaramanga Fault, that is
703 associated with the El Tendido shutter ridge in the northern part of the transpressive
704 southern termination of the structure (Velandia and Bermúdez, 2018) and the subsequent
705 impoundment of the Chicamocha River, suggested by Galvis (2016) to be associated with
706 the origin of a Pleistocene lake reported by Villarroel *et al.* (2001) in that area. Further,
707 geomorphological evidence for current tectonic activity along the Bucaramanga Fault is
708 provided by Diederix *et al.* (2009), e.g. eight Holocene earthquakes of magnitude Ms 6.6
709 to 7.0 spanning from 8300 to 930 yr BP and morphometric indices pointing to fault activity
710 as a leading contributor to regional hillslope instability (García-Delgado *et al.*, 2019,
711 2020).

712

713 Similar to the Venezuelan Andes (e.g., Kohn *et al.*, 1984; Colletta *et al.*, 1997; Audemard
714 and Audemard, 2002; Bermúdez *et al.*, 2010), the evolution of the Eastern Cordillera of
715 Colombia has been controlled by the reactivation of preexisting faults (Corredor 2003;
716 Cortés *et al.*, 2005; Mora *et al.*, 2009; Parra *et al.*, 2009). As observed in other
717 transpressional mountain belts (e.g., Spotila *et al.*, 2007a), these faults delimit tectonic
718 blocks characterized by contrasting exhumation histories. Unlike the Venezuelan Andes,
719 the signal of early (pre- Miocene) exhumation across the Eastern Cordillera is significant.
720 These tectonic blocks at the southern Bucaramanga Fault termination responded in
721 different ways to tectonic events that occurred in this zone, which were driven by the triple
722 convergence of the South American, Caribbean and Farallon/Nazca plates. In Paramo of
723 Güina and San Joaquín, older weighted mean ZHe ages ranging from ~80 to 50 Ma are
724 similar to AFT and ZFT ages reported by different authors for other sectors of the Eastern
725 and Central Cordillera (Vinasco *et al.*, 2006; Montes *et al.*, 2010; Cardona *et al.*, 2010;
726 Mora *et al.*, 2017; Restrepo-Moreno *et al.*, 2019) and at the Sierra Nevada de Santa Marta
727 (Patiño *et al.*, 2019) suggesting widespread and significant exhumation at that time. This
728 exhumation could be related to accretion of the Caribbean Plate on northwestern South
729 America. A significant phase of mountain building with changes in exhumation rates
730 occurring closer to $\sim 20 \pm 5$ Ma has been discerned in other sectors of the Colombian

731 Andes, such as: the Central Cordillera, Eastern Cordillera, Garzón Massif and Sierra
732 Nevada de Santa Marta (Amaya *et al.*, 2017, 2020; Patiño *et al.*, 2019; Restrepo-Moreno *et*
733 *al.*, 2019), but also from the Caparo and Trujillo blocks of the Venezuelan Andes
734 (Bermúdez *et al.*, 2010). This period of significant relief growth coincides with important
735 sediment deposition in the Maracaibo basin (Lugo and Mann, 1995; Guzmán and Fisher,
736 2006; Erickson *et al.*, 2012; Bermúdez *et al.*, 2017).

737

738 An exhumation phase at $\sim 12 \pm 3$ Ma coincides with the main surface uplift of the
739 Venezuelan Andes (Bermúdez *et al.*, 2011) and events reported for the Perijá (Shagam *et*
740 *al.*, 1984) and Santa Marta (Villagómez *et al.*, 2011). Finally, a more recent episode
741 between ~ 5 to 2 Ma is similar to that reported by Kohn *et al.* (1984) for the El Carmen
742 block in the Venezuelan Andes. This last event is associated with an increase in oblique
743 convergence between the Maracaibo block and South American plate, which resulted in
744 transpression and exhumation of the central part of the Venezuelan Andes (Kohn *et al.*,
745 1984; Audemard and Audemard, 2002; Bermúdez *et al.*, 2010) and other sectors at the
746 circum-Maracaibo block (e.g. Shagam *et al.*, 1984; Kohn *et al.*, 1984; Bermúdez *et al.*,
747 2017; 2019; Restrepo-Moreno *et al.*, 2019). This episode starting at ~ 5 Ma continues to the
748 present day, particularly in our study area where this event is observed in some regions:
749 circum Chaguacá Fault (Figure 7A), El Tendido, Chaguacá and Ricaurte profiles, but not
750 in all sectors. Convergence between the Maracaibo block and South American plate is also
751 consistent with the current stress field in the northern Andes as resolved from fault
752 striations, earthquake focal mechanisms and GPS data (Taboada *et al.*, 2000; Audemard
753 and Audemard, 2002; Colmenares and Zoback, 2003; Audemard *et al.*, 2005; Cortés and
754 Angelier, 2005; Audemard and Castilla, 2016; Pérez *et al.*, 2018; Mora-Páez *et al.*, 2019;
755 Velandia *et al.*, 2020).

756

757 The timing of the four main cooling episodes reported here is in accordance with tectonic
758 events reconstructed from coarse facies (Van der Hammen, 1961). These last four major
759 events are known as the Laramic, Pre-Andina, Proto-Andina, and Eu-Andina orogenic
760 events and occurred during the Late Cretaceous-Early Paleocene, the middle Eocene, the
761 Late Oligocene, and the Miocene-Pleistocene, respectively (Van der Hammen, 1961).
762 These events are the results of the complex interactions between Nazca, Cocos, Caribbean
763 and South American plate, which control the accretion of continental and/or oceanic

764 tectonic realms as Cauca-Patía, Chocó, Gorgona, Baudó and Maracaibo (Cediel *et al.*,
765 2003; Restrepo-Pace and Cediel, 2019).

766

767 **6. Conclusions**

768 Numerical models based on low temperature thermochronology data suggest contrasting
769 thermal histories between tectonic blocks prior to transpressional Bucaramanga Fault
770 reactivation in the Eastern Cordillera of Colombia. Some blocks experienced slow cooling
771 through the PAZ and PRZ, but some heating pulses occurred prior to the fault
772 displacement, after that, short and fast cooling pulses are discerned, possibly as a result of
773 Bucaramanga Fault activity.

774

775 Although Eocene exhumation of the Eastern Cordillera has been recorded in different
776 works, in this study numerical models of elevation profiles show that in the study area the
777 main cooling pulses occurred at the beginning of the Eocene ($\sim 50 \pm 5$ Ma). These episodes
778 are probably related to activity of the Boyacá and Soapaga faults during tectonic inversion
779 of the study area.

780

781 The onset of the most evident exhumation pulse occurred at $\sim 20 \pm 5$ Ma (late Oligocene-
782 early Miocene) and is characterised by high cooling rates associated with the onset of
783 Bucaramanga Fault transpressive activity along its southern termination, including one of
784 its main splays: the Chaguacá Fault. Older ages reported within the region of the Santander
785 Massif may indicate that exhumation across the Bucaramanga Fault has migrated from
786 north to south.

787

788 An exhumation phase closer to $\sim 12 \pm 3$ Ma is identified as being associated with a regional
789 event that coincided with the main surface uplift of the Venezuelan Andes, Perijá Range
790 and Sierra Nevada de Santa Marta. Another important exhumation pulse is identified
791 during the early Plio-Pleistocene (5-2 Ma) also with a high cooling rate, which is
792 considered to be mainly associated with reactivation of the Bucaramanga Fault.

793

794 **Acknowledgments**

795 This work is part of the doctoral thesis of Francisco Velandia at the Universidad Nacional
796 de Colombia, supported by Colciencias-Colfuturo grant (PDBC 6172) and a study

797 commission given by the Universidad Industrial de Santander. We are indebted to María
 798 Isabel Marín and Wilton Echavarría (Universidad EAFIT), Héctor Enciso, Francisco
 799 Santaella, Sergio Amaya and Mary Luz Peña (laboratories of the Servicio Geológico
 800 Colombiano). We are grateful to Sergio Restrepo-Moreno for remarks on the first version
 801 of this work. Thanks to the School of Geology, Universidad Industrial de Santander,
 802 especially Juan Diego Colegial and Giovanny Jiménez for facilitating samples delivery,
 803 and also to Nicolás Villamizar and Milton Galvis for accompanying the first field trip.
 804 Bermúdez acknowledges support from the Institut de Recherche pour le Développement
 805 (IRD) for the BEST Project and the convention between Universidad de Ibagué and IRD
 806 which allowed to stay in GET lab at Toulouse, France, where apatite and zircon were
 807 separated, also to Universidad de Ibagué (Project 15-377-INT) and Universidad
 808 Pedagógica y Tecnológica de Colombia (UPTC), DIN SGI Project 3104. The University of
 809 Melbourne thermochronology laboratory receives support under the AuScope program of
 810 the National Collaborative Research Infrastructure Strategy (NCRIS). Finally, we thanks to
 811 the editor Franck Audemard, Melanie Noury and an anonymous reviewer for their valuable
 812 comments and suggestions.

813 FIGURES CAPTIONS

814 **Figure 1.** Geological map of the axial zone of the Eastern Cordillera of Colombia. Red boxes
 815 indicate the studied area (modified from Gómez *et al.*, 2015).

816
 817 **Figure 2. A.** Chronostratigraphic compilation in basins adjacent to the Santander and Floresta
 818 massifs. Dark grey = mainly coarse clastic units, light grey = fine-grained textures (modified from
 819 Jiménez *et al.*, 2014). **B.** Thermo-tectonic/orogenic phases from based on a compilation of in-situ
 820 (Restrepo-Moreno *et al.*, 2019) and detrital (Bermúdez *et al.*, 2021) low-temperature
 821 thermochronology data.

822
 823 **Figure 3.** Map of the main geological faults and sample location in the studied area. The southern
 824 termination of the Bucaramanga Fault is characterized by a transpressive system, including the
 825 Chaguacá and Los Micos faults. The figures include location of samples analyzed in this work
 826 (AFT, ZFT, AHe and ZHe), as well as those from previous work by Shagam *et al.* (1984), Toro
 827 (1990), van der Lelij *et al.* (2016a), and Amaya (2016). Only those samples enclosed in red ellipses
 828 belonging to vertical profiles were considered for the modeling of thermal histories. Samples in red
 829 triangles were used to model the thermal history of the Los Micos and Chaguacá faults.

830
 831 **Figure 4.** Panoramic views and Google Earth images of sample locations for vertical profiles. A,
 832 Ricaurte. B, El Tendido. C, San Joaquín. D, Páramo. E, Chaguacá. F. Photograph of the valley
 833 controlled by the sinistral, transpressive Chaguacá Fault.

834
 835 **Figure 5.** Comparisons between [eU] ppm versus individual ZHe grain age for (A) all samples. (B)
 836 and (C) correspond to a close-up of panel (A). In these figures all individual grain ages have been
 837 considered without making any exclusions.

838 **Figure 6.** Weighted (U-Th)/He age-elevation relationships: (A) AHe ages and (B) ZHe ages. AFT
 839 data has also been added in panel (B) for comparison (unfilled diamonds whose outline colors
 840 follow the same nomenclature of the different profiles or areas). Grey bands are considered
 841 significant regional geological events derived from detrital thermochronology (Bermúdez *et al.*,
 842 2021) and other Colombian regions (Restrepo-Moreno *et al.*, 2019). By contrast, based on scarce
 843 ages, we relate white bands with period of “apparent” tectonic quiescence. Superscripts 1 and 2
 844 refer to profile or fault, respectively.

845
 846 **Figure 7.** Time-Temperature expected paths predicted from QTQt software (Gallagher, 2012)
 847 along the (A) Los Micos Fault and (B) Chaguacá Fault. Selected samples for the Los Micos Fault
 848 are shown in red, while selected samples for the Chaguacá Fault models are shown in blue. Green
 849 and transparent triangles correspond to samples from other blocks, which were not considered for
 850 modelling.

851
 852 **Figure 8.** (Left panel) Thermal history models for the different profiles across the southern
 853 Bucaramanga Fault samples determined using the QTQt software (Figure 5). The dashed rectangles
 854 correspond to the four time-Temperature constraints: (i) 50 ± 10 Ma, (ii) 20 ± 5 Ma, (iii) 12 ± 3
 855 Ma, and 5 ± 3 Ma with temperatures of $160^\circ \pm 40^\circ\text{C}$, $120^\circ \pm 60^\circ\text{C}$, $100^\circ \pm 40^\circ\text{C}$, and $80^\circ \pm 40^\circ\text{C}$,
 856 respectively. (Central panel) Observed versus predicted thermochronology ages (note that He ages
 857 are displayed with resampling errors). (Right panel) Age-elevations relationships (AER) showing
 858 comparisons between predicted and observed age and/or kinematic parameter (Dpar or Mean
 859 Track-Length). In these models the AFT ages, mean track lengths, weighted ZHe and AHe ages
 860 were integrated. For AFT data the multikinetic annealing model of Ketcham *et al.* (2007b), with c-
 861 axis projected track length data (Ketcham *et al.*, 2007a) was used. For the radiation damage
 862 accumulation and annealing model, we used the Gautheron *et al.* (2009) model for AHe and the
 863 Guenther *et al.* (2013) model for ZHe ages, respectively.

864
 865
 866 **Figure 9.** (A) Individual time-Temperature path for samples FV-44 (El Tendido profile) and (B)
 867 FV-57 (Ricaurte profile). In these models we integrated AFT ages, mean track lengths, weighted
 868 ZHe and AHe ages. For AFT data the multikinetic annealing model of Ketcham *et al.* (2007b), with
 869 c- axis projected track length data (Ketcham *et al.*, 2007a) was used. For the radiation damage
 870 accumulation and annealing model, we used the Gautheron *et al.* (2009) model for AHe and the
 871 Guenther *et al.* (2013) model for ZHe ages, respectively. (O) = observed and (P) = predicted data.

872
 873

874 TABLE CAPTIONS

875

876 **Table 1.** Apatite and zircon fission-track data from the Bucaramanga Fault southern termination
 877 using the External Detector Method (EDM).

878

879 **Table 2.** Apatite fission-track data from the Bucaramanga Fault southern termination using the LA-
 880 ICP-MS method. In this table, n: number of analyzed crystals. Ns: number of spontaneous tracks
 881 counted. RhoS: spontaneous track density. ^{238}U : average uranium concentration. SD: Standard
 882 deviation of the track length distribution. Dpar: long axis of track etch pit. MTL: mean confined
 883 track length and SE: standard error. NL: number of confined track lengths (TINTs) measured.
 884 Single apatite fission track ages were calculated using an absolute calibration based on primary
 885 constants (aggregate $\xi = 2.176 \times 10^{-3}$). Pooled ages were calculated according to Hasebe *et al.*
 886 (2004). Central ages, as well as dispersion, were calculated using the RadialPlotter program
 887 (Vermeesch, 2009).

888

889 **Table 3.** Apatite (U-Th-Sm)/He bedrock data from the Bucaramanga Fault southern termination.

890

891 **Table 4.** Zircon (U-Th)/He bedrock data from the Bucaramanga Fault southern termination.

892

893 **Table 5.** Average exhumation rates and confidence intervals estimated from Age-Elevation
894 Relationships (AER) for the different profiles. Exhumation rates were calculated using the York
895 method (Williamson, 1968) considering the 1σ error on the age. See text for further details.

896

897 **REFERENCES**898 Acosta, J., Velandia, F., Osorio, J., Lonergan, L., Mora, H., 2007. Strike-Slip deformation within
899 the Colombian Andes. In: Ries, A. C., Butler, R. W. H. and Graham, R. H. (eds) 2007.
900 Deformation of the Continental Crust: The Legacy of Mike Coward. Geological Society, London,
901 Special Publications 272, 303-319.902 Amaya, S., 2016. Termocronología y geocronología del basamento metamórfico del Macizo de
903 Santander, Departamento de Santander. Universidad Nacional de Colombia, Bogotá. Tesis de
904 Doctorado. 174p.905 Amaya, S., Zuluaga, C., Bernet, M., 2017. New fission-track age constraints on the exhumation of
906 the central Santander Massif: Implications for the tectonic evolution of the Northern Andes,
907 Colombia. *Lithos* 282-283, 388-402.908 Amaya-Ferreira, S., Zuluaga, C.A., Bernet, M., 2020. Different levels of exhumation across the
909 Bucaramanga Fault in the Cepitá area of the southwestern Santander Massif, Colombia:
910 Implications for the tectonic evolution of the northern Andes in northwestern South America. In:
911 Gómez, J., Mateus-Zabala, D. (editors), *The Geology of Colombia, Volume 3 Paleogene –*
912 *Neogene*. Servicio Geológico Colombiano, Publicaciones Geológicas Especiales 37, 17p. Bogotá.
913 <https://doi.org/10.32685/pub.esp.37.2019.17>914 Ansberque, C., Chew, D. M., Drost, K., 2021. Apatite fission-track dating by LA-Q-ICP-MS
915 imaging. *Chemical Geology*, 560, 119977. doi:<https://doi.org/10.1016/j.chemgeo.2020.119977>916 Audemard, F.E., Audemard, F.A., 2002. Structure of the Mérida Andes, Venezuela: Relations with
917 the South America-Caribbean geodynamic interaction. *Tectonophysics* 345, 299–327.918 Audemard, F.A., 2003. Geomorphic and geologic evidence of ongoing uplift and deformation in
919 the Mérida Andes, Venezuela. *Quaternary International* 101-102: 43-65.920 Audemard, F.A., Romero, G. Rendón, H., Cano, V., 2005. Quaternary fault kinematics and stress
921 tensors along the southern Caribbean from microtectonic data and focal mechanism solutions.
922 *Earth-Science Reviews*, 69 (3-4): 181-233. doi: 10.1016/j.earscirev.2004.08.001.923 Audemard, F.A., Castilla, R., 2016. Present-day stress tensors along the southern Caribbean plate
924 boundary zone from inversion of focal mechanism solutions: A successful trial. *Journal of South*
925 *American Earth Sciences* 71, 309-319.926 Bayona, G., Cortes, M., Jaramillo, C., Ojeda, G., Aristizábal, J., Reyes-Harker, A., 2008. An
927 integrated analysis of an orogen-sedimentary basin pair: Latest Cretaceous-Cenozoic evolution of
928 the linked Eastern Cordillera orogen and the Llanos foreland basin of Colombia. *Geological*
929 *Society of America Bulletin* 120 (9-10), 1171-1197.930 Bermúdez, M.A., Kohn, B.P., van der Beek, P.A., Bernet, M., O’Sullivan, P.B., Shagam, R., 2010.
931 Spatial and temporal patterns of exhumation across the Venezuelan Andes: implications for
932 Cenozoic Caribbean geodynamics. *Tectonics* 29. TC5009, doi: 10.1029/2009TC002635.933 Bermúdez, M.A., van der Beek, P., Bernet, M., 2011. Asynchronous Miocene–Pliocene
934 exhumation of the central Venezuelan Andes: *Geology* 39, 139–142.

- 935 Bermúdez, M. A., Hoorn, C., Bernet, M., Carrillo, E., van der Beek, P. A., Garver, J. I., Mora, J.,
936 Mehrkian, K., 2017. The detrital record of late-Miocene to Pliocene surface uplift and exhumation
937 of the Venezuelan Andes in the Maracaibo and Barinas foreland basins. *Basin Research* 29, 370-
938 395. doi:10.1111/bre.12154.
- 939 Bermúdez, M.A., Bernet, M., Kohn, B., Bricchau, S., 2019. Exhumation-denudation history of the
940 Maracaibo Block, Northwestern South America: Insights from thermochronology, In Cediél, F and
941 Shaw, R.P. (eds) (*Geology and Tectonics of Northwestern South America: The Pacific-Caribbean-
942 Andean Junction*, pp. 879-898 - *Frontiers in Earth Sciences Series*, Springer Verlag.
943 https://doi.org/10.1007/978-3-319-76132-9_13.
- 944 Bermúdez, M. A., Velandia, F., García-Delgado, H., Jiménez, D., Bernet, M., 2021. Exhumation of
945 the southern transpressive Bucaramanga fault, eastern Cordillera of Colombia: Insights from
946 detrital, quantitative thermochronology and geomorphology. *Journal of South American Earth
947 Sciences* 106, 103057. doi:<https://doi.org/10.1016/j.jsames.2020.103057>.
- 948 Boone, S. C., Kohn, B. P., Gleadow, A. J. W., Morley, C. K., Seiler, C., Foster, D. A., Chung, L.,
949 2018. Tectono-thermal evolution of a long-lived segment of the East African Rift System:
950 Thermochronological insights from the North Lokichar Basin, Turkana, Kenya. *Tectonophysics*,
951 744, 23-46. doi:<https://doi.org/10.1016/j.tecto.2018.06.010>.
- 952 Braun, J., 2002. Quantifying the effect of recent relief changes on age–elevation relationships.
953 *Earth and Planetary Science Letters* 200, 331-343. doi:[https://doi.org/10.1016/S0012-
954 821X\(02\)00638-6](https://doi.org/10.1016/S0012-821X(02)00638-6)
- 955 Braun, J., 2003. PECUBE: A new finite-element code to solve the 3D heat transport equation
956 including the effects of a time-varying, finite amplitude surface topography. *Computers and
957 Geosciences* 29(6), 787–794. [https://doi.org/10.1016/S0098-3004\(03\)00052-9](https://doi.org/10.1016/S0098-3004(03)00052-9).
- 958 Braun, J., van der Beek, P., Valla, P., Robert, X., Herman, F., Glotzbach, C., Pedersen, V., Perry,
959 C., Simon-Labric, T., Prigent, C., 2012. Quantifying rates of landscape evolution and tectonic
960 processes by thermochronology and numerical modeling of crustal heat transport using PECUBE.
961 *Tectonophysics* 524–525, 1–28. <https://doi.org/10.1016/j.tecto.2011.12.035>.
- 962 Braun, J., Gemignani, L., van der Beek, P., 2018. Extracting information on the spatial variability
963 in erosion rate stored in detrital cooling age distributions in river sands. *Earth Surf. Dynam.*
964 *Discuss.* <https://doi.org/10.5194/esurf-2017-42>.
- 965 Caballero, V., Mora, A., Quintero, I., Blanco, W., Parra, M., Rojas, L.E., López, C., Sánchez, N.,
966 Horton, B.K., Stockli, D., Duddy, I., 2013. Tectonic controls on sedimentation in an intermontane
967 hinterland basin adjacent to inversion structures: the Nuevo Mundo Syncline, Middle Magdalena
968 Valley, Colombia. In: Némčok, M., Mora, A. and Cosgrove, J. W. (eds) *Thick-Skin-Dominated
969 Orogens: From Initial Inversion to Full Accretion*. Geological Society, London, Special
970 Publications 377, 315-342.
- 971 Cao, S., Neubauer, F., 2016. Deep crustal expressions of exhumated strike-slip fault systems: Shear
972 zone initiation on rheological boundaries. *Earth-Science Reviews* 162, 155-176.
- 973 Cardona, A., Valencia, V., Bustamante, C., García-Casco, A., Ojeda, G., Ruiz, J., Saldarriaga, M.,
974 Weber, M., 2010. Tectonomagmatic setting and provenance of the Santa Marta Schists, northern
975 Colombia: insights on the growth and approach of Cretaceous Caribbean oceanic terranes to the
976 South American continent. *J. S. Am. Earth Sci.* 29, 784–804.
977 <https://doi.org/10.1016/j.jsames.2009.08.012>.
- 978 Cediél, F., Shaw, R., Cáceres, C., 2003. Tectonic assembly of the northern Andean Block, in
979 Bartolini, C., Buffler, R. T., and Blickwede, J., eds., *The Circum-Gulf of Mexico and the
980 Caribbean: Hydrocarbon habitats, basin formation and plate tectonics*, American Association of
981 Petroleum Geologists Memoir 79, 815-848.

- 982 Chew, D.M., Donelick, R.A. 2012. Combined apatite fission track and U-Pb dating by LA-ICP-MS
983 and its application in apatite provenance studies. *Mineral Assoc Canada Short Course* 42:219–247
984 (St Johns, NL).
- 985 Chew D, Donelick, R.A., Donelick, M.B., Kamber, B.S., Stock, M.J., 2014. Apatite chlorine
986 concentration measurements by LA-ICP-MS. *Geostand Geoanal Res* 38:23–35
- 987 Cochran, W.J., Spotila, J.A., Prince, P.S., McAleer, R.J., 2017. Rapid exhumation of Cretaceous
988 arc-rocks along the Blue Mountains restraining bend of the Enriquillo-Plantain Garden fault,
989 Jamaica, using thermochronometry from multiple closure systems. *Tectonophysics* 721, 292-309.
- 990 Colletta, B., Hebrard, F., Letouzey, J., Werner, P., Rudkiewicz, J., 1990. Tectonic style and crustal
991 structure of the Eastern Cordillera (Colombia) from a balanced cross-section. In Letouzey, J., ed.,
992 *Petroleum and tectonics in mobile belts*: Paris, Editions Technip, pp. 81-100.
- 993 Cogné, N., Chew, D.M., Donelick, R.A., Ansberque, C., 2020. LA-ICP-MS apatite fission track
994 dating: a practical zeta-based approach *Chem. Geol.*, 531, 10.1016/j.chemgeo.2019.119302.
- 995 Colmenares, L., Zoback, M. D., 2003. Stress field and seismotectonics of northern South America,
996 *Geology* 31, 721–724, doi:10.1130/G19409.1.
- 997 Corredor, F., 2003. Eastward extent of the late Eocene early Oligocene onset of deformation across
998 the northern Andes: constraints from the northern portion of the eastern Cordillera fold belt,
999 Colombia. *J. S. Am. Earth Sci.* 16 (6), 445–457.
- 1000 Cooper, M., Addison, F., Alvarez, R., Coral, M., Graham, R., Hayward, A., Howe, S., Martínez, J.,
1001 Naar, J., Peñas, R., Pulham, J., Taborda, A., 1995. Basin development and tectonic history of the
1002 Llanos Basin, Eastern Cordillera, and Middle Magdalena Valley, Colombia. *AAPG Bulletin* 79
1003 (10), 1421-1443.
- 1004 Cortés, M., Angelier, J., 2005. Current states of stress in the northern Andes as indicated by focal
1005 mechanisms of earthquakes, *Tectonophysics* 403, 29–58, doi:10.1016/j.tecto.2005.03.020.
- 1006 Cortés, M., Colletta, B., Angelier, J., 2005. Structure and tectonics of the central segment of the
1007 Eastern Cordillera of Colombia, *J. South Am. Earth Sci.*, 21, 437–465,
1008 doi:10.1016/j.jsames.2006.07.004.
- 1009 Cruz, L., Fayon, A., Teyssier, C., Weber, J., 2007. Exhumation and deformation processes in
1010 transpressional orogens: The Venezuelan Paria peninsula, SE Caribbean-South American plate
1011 boundary. *Geological Society of America, Special Paper* 434, 149-165.
- 1012 Danišik, M., McInnes, B. I. A., Kirkland, C. L., McDonald, B. J., Evans, N. J., Becker, T., 2017,
1013 Seeing is believing: Visualization of He distribution in zircon and implications for thermal history
1014 reconstruction on single crystals, *J. Science Advances* 3(2), e1601121.
1015 doi:10.1126/sciadv.1601121.
- 1016 De Freitas, M., Froncolin, J., Cobbold, O., 1997. The structure of the axial zone of the Cordillera
1017 Oriental, Colombia. VI Simposio Bolivariano “Exploración petrolera en las cuencas subandinas”.
1018 Cartagena. *Memorias*, II: 38-41.
- 1019 Dengo, C., Covey, M., 1993. Structure of the Eastern Cordillera of Colombia: implications for trap
1020 styles and regional tectonics. *AAPG Bulletin* 77 (8), 1315-1337.
- 1021 Diederix, H., Hernández, C., Torres, E., Osorio, J., Botero, P., 2009. Resultados preliminares del
1022 primer estudio paleosimológico a lo largo de la Falla de Bucaramanga. XII Congreso Colombiano
1023 de Geología. Paipa. *Memorias* 18pp.
- 1024 Dodson, M. H., 1973. Closure temperature in cooling geochronological and petrological systems:
1025 *Contributions to Mineralogy and Petrology* 40, 259–274.

- 1026 Donelick R. A., O'Sullivan P. B., Ketcham R. A., 2005. Apatite fission-track analysis. *Reviews in*
1027 *Mineralogy and Geochemistry* 58, 49–94.
- 1028 Egbue, O., Kellogg, J., Aguirre, H., Torres, C., 2014. Evolution of the stress and strain
1029 fields in the Eastern Cordillera, Colombia. *J. S. Am. Earth Sci.* 58, 8-21.
- 1030 Erikson, J.P., Kelley, S.A., Osmolovsky, P., Verosub, K.L., 2012. Linked basin sedimentation and
1031 orogenic uplift: the Neogene Barinas basin sediments derived from the Venezuelan Andes. *J. S.*
1032 *Am. Earth Sci.* 39, 138–156.
- 1033 Fabre, A., 1983. La subsidencia de la Cuenca del Cocuy (Cordillera Oriental de Colombia) durante
1034 el Cretáceo y el Terciario. Segunda parte: esquema de evolución tectónica. *Revista Norandina* 8,
1035 21-27.
- 1036 Farley K. A., Wolf R. A., Silver L. T., 1996. The effects of long alpha-stopping distances on (U-
1037 Th)/He ages. *Geochim. Cosmochim. Acta* 60, 4223–4229.
- 1038 Farley, K. A., 2000. Helium diffusion from apatite; general behavior as illustrated by Durango
1039 fluorapatite: *Journal of Geophysical Research* 105, 2903–2914.
- 1040 Farley, K. A., 2002. (U-Th)/He dating: Techniques, calibrations, and applications. *Reviews in*
1041 *Mineralogy and Geochemistry* 47(1), 819–844. <https://doi.org/10.2138/rmg.2002.47.18>.
- 1042 Farley, K. A., Stockli, D. F., 2002. (U-Th)/He dating of phosphates: Apatite, monazite, and
1043 xenotime, *Rev. Mineral. Geochemistry* 48(1), 559–577. <https://doi.org/10.2138/rmg.2002.48.15>.
- 1044 Fitzgerald, P.G., Gleadow, A. J. W., 1988. Fission-track geochronology, tectonics and structure of
1045 the Transantarctic Mountains in northern Victoria Land, Antarctica. *Chem Geol* 73, 169–198.
- 1046 Fitzgerald, P.G., Gleadow, A. J. W., 1990. New approaches in fission track geochronology as a
1047 tectonic tool: examples from the Transantarctic Mountains. *Nucl Tracks Radiat Meas* 17, 351–357.
- 1048 Fitzgerald, P., Sorkhabi, R., Redfield, T., Stump, E., 1995. Uplift and denudation of the central
1049 Alaska Range: a case study in the use of apatite fission track thermochronology to determine
1050 absolute uplift parameters. *Journal of Geophysical Research* 100, 20175–20191.
- 1051 Fitzgerald, P. G., Baldwin, S.L., Webb, L.E., O'Sullivan, P.B., 2006. Interpretation of (U-Th)/He
1052 single grain ages from slowly cooled crustal terranes: A case study from the Transantarctic
1053 Mountains of southern Victoria Land. *Chemical Geology* 225(1-2): 91-120.
- 1054 Fitzgerald, P. G., Malusà, M. G., 2019. Concept of the exhumed partial annealing (retention) zone
1055 and age-elevation profiles in thermochronology. In: Malusà M.G., Fitzgerald, P.G. (Eds.), *Fission-*
1056 *Track Thermochronology and its Application to Geology*, p. 165-189, Springer Textbooks in Earth
1057 Sciences, Geography and Environment- doi: 10.1007/978-3-319-89421-8_9.
- 1058 Forero-Ortega, A.J., Velandia, F., Barragán-Coy, E.K., 2020. Estilos estructurales y tensores de
1059 esfuerzo hacia el suroriente del Macizo de Santander (Colombia). *Bol. Geol.* 42 (2), 129-145, doi:
1060 10.18273/revbol.v42n2-2020007.
- 1061 Fox, M., Carter, A., 2020. Heated Topics in Thermochronology and Paths towards Resolution.
1062 *Geosciences* 10, 375. Furlong, K.P., 2007. Locating the deep extent of the plate boundary along the
1063 Alpine Fault zone, New Zealand: Implications for patterns of exhumation in the Southern Alps.
1064 *Geological Society of America Special Paper*, 434, 1-14.
- 1065 Hueck, M., Dunkl, I., Heller, B., Stipp Basei, M. A., Siegesmund, S., 2018. (U-Th)/He
1066 Thermochronology and Zircon Radiation Damage in the South American Passive Margin: Thermal
1067 Overprint of the Paraná LIP? , 37(10), 4068-4085. doi:10.1029/2018tc005041.

- 1068 Gallagher, K., 2012. Transdimensional inverse thermal history modeling for quantitative
1069 thermochronology. *Journal of Geophysical Research*, vol. 117, B02408,
1070 doi:10.1029/2011JB008825.
- 1071 Gallagher, K., Brown, R., Johnson, C., 1998. Fission track analysis and its applications to
1072 geological problems. *Ann Rev Earth Planet Sci* 26 (1):519–572.
- 1073 Gallagher, K., Charvin, K., Nielsen, S., Sambridge, M., Stephenson, J., 2009. Markov chain Monte
1074 Carlo (MCMC) sampling methods to determine optimal models, model resolution and model
1075 choice for Earth Science problems. *Marine and Petroleum Geology* 26, 525-535.
- 1076 Galvis, M., 2016. Análisis estructural del segmento sur de la Falla de Bucaramanga en el municipio
1077 de Onzagá, Santander. Tesis de grado, Universidad Industrial de Santander. 192p.
- 1078 García-Delgado, H., Villamizar, N., Bernet, M., 2019. Recent tectonic activity along the
1079 Bucaramanga Fault System (Chicamocha River Canyon, Eastern Cordillera of Colombia): a
1080 geomorphological approach. *Zeitschrift für Geomorphologie* 62 (3), 199-215.
- 1081 García-Delgado, H., Machuca, S., Velandia, F., Audemard, F.A., 2020. Along-strike variations in
1082 recent tectonic activity in the Santander Massif: New insights on landscape evolution in the
1083 Northern Andes. *Journal of South American Earth Sciences*, 98.
1084 <https://doi.org/10.1016/j.jsames.2019.102472>
- 1085 García-Delgado, H., Villamizar-Escalante, N., Bermúdez, M. A., Bernet, M., Velandia, F., 2021.
1086 Climate or tectonics? What controls the spatial-temporal variations in erosion rates across the
1087 Eastern Cordillera of Colombia? *Global and Planetary Change*, 203, 103541.
1088 doi:<https://doi.org/10.1016/j.gloplacha.2021.103541>.
- 1089 Gautheron, C., Tassan-Got, L., Barbarand, J., Pagel, M., 2009. Effect of alpha-damage annealing
1090 on apatite (U-Th)/He thermochronology: *Chemical Geology* 266, 157–170, [https://doi.org/10.1016](https://doi.org/10.1016/j.chemgeo.2009.06.001)
1091 [/j.chemgeo.2009.06.001](https://doi.org/10.1016/j.chemgeo.2009.06.001).
- 1092 Gerin, C., Gautheron, C., Oliviero, E., Bachelet, C., Mbongo Djimbi, D., Seydoux-Guillaume,
1093 A.M., Tassan-Got, L., Sarda, P., Roques, J., Garrido, F., 2017. Influence of vacancy damage on He
1094 diffusion in apatite, investigated at atomic to mineralogical scales: *Geochimica et Cosmochimica*
1095 *Acta* 197, 87-103.
- 1096 Gleadow, A. J. W., 1990. Fission track thermochronology—reconstructing the thermal and tectonic
1097 evolution of the crust. In: *Pacific Rim Congress, Gold Coast, Queensland, 1990*. Australasian
1098 Institute of Mining Metallurgy, pp 15–21. Gleadow, A.J.W., Fitzgerald, P.G. 1987. Uplift history
1099 and structure of the Transantarctic Mountains: new evidence from fission track dating of basement
1100 apatites in the Dry Valleys area, southern Victoria Land, *Earth Planet. Sci. Lett.* 82, 1–14.
- 1101 Gleadow, A., Harrison, M., Kohn, B., Lugo-Zazueta, R., Phillips, D., 2015. The Fish Canyon Tuff:
1102 A new look at an old low-temperature thermochronology standard. *Earth and Planetary Science*
1103 *Letters* 424, 95–108.
- 1104 Gómez, J., Montes, N., Nivia, A., Diederix, H. Compiladores., 2015. Mapa Geológico de
1105 Colombia, escala 1:1'000.000. Servicio Geológico Colombiano. Bogotá.
- 1106 Green, P. F., Duddy, I. R., Laslett, G. M., Hegarty, K. A., Gleadow, A. J. W., Lovering, J. F.,
1107 1989. Thermal annealing of fission tracks in apatite: 4. Quantitative modelling techniques and
1108 extension to geological timescales: *Chemical Geology* 79, 155–182.
- 1109 Guenther, W. R., Reiners, P. W., Ketcham, R. A., Nasdala, L., Giester, G., 2013. Helium diffusion
1110 in natural zircon: radiation damage, anisotropy, and the interpretation of zircon (U-TH)/He
1111 thermochronology. *American Journal of Science* 313, 145–198. <https://doi.org/10.2475/03.2013.01>.

- 1112 Guzmán, J.I. and Fisher, W.L., 2006. Early and middle Miocene depositional history of the
1113 Maracaibo basin, western Venezuela. *Am. Assoc. Pet. Geol. Bull.*, 90, 625–655.
- 1114 Hasebe, N., Barbarand, J., Jarvis, K., Carter, A., Hurford, A.J., 2004. Apatite fission-track
1115 chronometry using laser ablation ICP-MS. *Chem. Geol.* 207, 134–145.
- 1116 Hasebe N, Tamura A, Arai, S., 2013. Zeta equivalent fission-track dating using LA-ICP-MS and
1117 examples with simultaneous U-Pb dating. *Island Arc* 22, 280–291.
- 1118 Horton, B.K., Saylor, J.E., Nie, J., Mora, A., Parra, M., Reyes-Harker, A., Stockli, D.F., 2010.
1119 Linking sedimentation in the northern Andes to basement configuration, Mesozoic extension, and
1120 Cenozoic shortening: Evidence from detrital zircon U-Pb ages, Eastern Cordillera, Colombia:
1121 *Geological Society of America Bulletin* 122 (9-10), 1423– 1442.
- 1122 Horton, B.K., Parra, M., Mora, A., 2020. Construction of the Eastern Cordillera of Colombia:
1123 Insights from the sedimentary record. In: Gómez, J., Mateus-Zabala, D. (editors), *The Geology of*
1124 *Colombia, Volume 3 Paleogene – Neogene*. Servicio Geológico Colombiano, Publicaciones
1125 *Geológicas Especiales* 37, 22 p. Bogotá. <https://doi.org/10.32685/pub.esp.37.2019.03>.
- 1126 Hueck, M., Dunkl, I., Heller, B., Stipp Basei, M. A., Siegesmund, S., 2018. (U-Th)/He 1044
1127 Thermochronology and Zircon Radiation Damage in the South American Passive Margin: Thermal
1128 1045 Overprint of the Paraná LIP?, *37, 10*, 4068-4085. doi:10.1029/2018tc005041.
- 1129 Huntington, K. W., Ehlers, T. A., Hodges, K. V., Whipp, D. M., 2007. Topography, exhumation
1130 pathway, age uncertainties, and the interpretation of thermochronometer data, *Tectonics* 26,
1131 TC4012, doi:10.1029/2007TC002108.
- 1132 Hurford, A.J., Green, P.F., 1982. A user's guide to fission track dating calibration. *Earth Planet Sci*
1133 *Lett* 59:343–354.
- 1134 Jacques, J. M., 2004. The influence of intraplate structural accommodation zones on delineating
1135 petroleum provinces of the Sub-Andean foreland basins. *Petroleum Geoscience*, 10(1), 1.
1136 doi:10.1144/1354-079303-582.
- 1137 Jiménez, G., Speranza, F., Faccenna, C., Bayona, G., Mora, A., 2014. Paleomagnetism and
1138 magnetic fabric of the Eastern Cordillera of Colombia: Evidence for oblique convergence and
1139 nonrotational reactivation of a Mesozoic intracontinental rift, *Tectonics* 33,
1140 doi:10.1002/2014TC003532.
- 1141 Johnson, J. E., Flowers, R. M., Baird, G. B., Mahan, K. H., 2017. Inverted” zircon and apatite (U–
1142 Th)/He dates from the Front Range, Colorado: High-damage zircon as a low-temperature (<50 °C)
1143 thermochronometer. *Earth and Planetary Science Letters* 466, 80-90. Doi:
1144 <https://doi.org/10.1016/j.epsl.2017.03.002>.
- 1145 Kammer, A., 1993. Steeply dipping basement faults and associated structures of the Santander
1146 massif, Eastern Cordillera, Colombian Andes. *Geología Colombiana*, 18, 47–62.
- 1147 Kammer, A., 1996. Estructuras y deformaciones del borde oriental del Macizo de Floresta.
1148 *Geología Colombiana* 21, 65-80.
- 1149 Kammer, A., 1999. Observaciones acerca de un origen transpresivo de la Cordillera Oriental.
1150 *Geología Colombiana*, 24, 29-53.
- 1151 Kammer, A., Sánchez, J., 2006. Early Jurassic rift structures associated with the Soapaga and
1152 Boyaca faults of the Eastern Cordillera, Colombia: Sedimentological inferences and regional
1153 implications. *Journal of South American Earth Sciences* 21(4), 412-422. Ketcham, R. A., Donelick,
1154 R. A., Carlson, W. D. 1999. Variability of apatite fission-track annealing kinetics; III.
1155 Extrapolation to geological time scales: *American Mineralogist* 84, 1235–1255

- 1156 Ketcham, R. A., Carter, A., Donelick, R. A., Barbarand, J., Hurford, A. J., 2007a. Improved
1157 measurement of fission-track annealing in apatite using c-axis projection, *Am. Mineral.* 92, 789-
1158 798, doi:10.2138/am.2007.2280.
- 1159 Ketcham, R. A., Carter, A., Donelick, R. A., Barbarand, J., Hurford, A. J., 2007b. Improved
1160 modeling of fission-track annealing in apatite, *Am. Mineral.* 92, 799-810,
1161 doi:10.2138/am.2007.2281.
- 1162 Kohn, B., Shagam, R., Banks, P., Burkley, L., 1984. Mesozoic–Pleistocene fission track ages on
1163 rocks of the Venezuelan Andes and their tectonic implications, *Mem. Geol. Soc. Am.* 162, 365–
1164 384.
- 1165 Laslett, G. M., Green, P. F., Duddy, I. R., Gleadow, A.J.W., 1987. Thermal annealing of fission
1166 tracks in apatite. 2. A quantitative analysis: *Chemical Geology* 65, 1–13.
- 1167 Lugo, J., Mann, P., 1995, Jurassic–Eocene tectonic evolution of Maracaibo basin, Venezuela, in A.
1168 J. Tankard, R. Suárez S., and H. J. Welsink, *Petroleum basins of South America: AAPG Memoir*
1169 62, p. 699–725.
- 1170 Machuca, S., García-Delgado, H., Velandia, F., 2021. Studying active fault-related folding on
1171 tectonically inverted orogens: A case study at the Yariquíes Range in the Colombian Northern
1172 Andes. *Geomorphology* 375, 107515. doi: 10.1016/j.geomorph.2020.107515.
- 1173 Mantilla-Figueroa, L.C., García-Ramírez, C.A., Valencia, V.A., 2016. Propuesta de escisión de la
1174 denominada ‘Formación Silgará’ (Macizo de Santander, Colombia), a partir de edades U-Pb en
1175 circones detríticos. *Boletín de Geología*, 38 (1), 33-50.
- 1176 McDowell, F. W., McIntosh, W. C., Farley, K. A., 2005. A precise ^{40}Ar - ^{39}Ar reference age for the
1177 Durango apatite (U-Th)/He and fission-track dating standard. *Chemical Geology* 214, 249-263.
1178
- 1179 Mojica, J., Kammer, A., 1995. Eventos jurásicos en Colombia. *Geología Colombiana*, 19, 165-172.
1180
- 1181 Montes, C., Guzmán, G., Bayona, G., Cardona, A., Valencia, V., Jaramillo, C., 2010. Clockwise
1182 rotation of the Santa Marta massif and simultaneous Paleogene to Neogene deformation of the
1183 Plato-San Jorge and Cesar-Ranchería basins. *J. S. Am. Earth Sci.* 29, 832–848.
1184 <https://doi.org/10.1016/j.jsames.2009.07.010>.
1185
- 1186 Mora, A., Parra, M., Strecker, M.R., Sobel, E.R., Hooghiemstra, H., Torres, V., Jaramillo, J.V.,
1187 2008. Climatic forcing of asymmetric orogenic evolution in the Eastern Cordillera of Colombia.
1188 *Geological Society of America Bulletin* 120, 930–949.
- 1189 Mora, A., Gaona, T., Kley, J., Montoya, D., Parra, M., Quiroz, L.I., Reyes, G., Strecker, M.R.,
1190 2009.
1191 The role of inherited extensional fault segmentation and linkage in contractional orogenesis: a
1192 reconstruction of Lower Cretaceous inverted rift basins in the Eastern Cordillera of Colombia.
1193 *Basin*
1194 *Research* 21, 111-137.
- 1195 Mora, A., Baby, P., Roddaz, M., Parra, M., Brusset, S., Hermoza, W., Espurt, N., 2010. Tectonic
1196 history of the Andes and Subandean zones: Implications for Neogene Amazonia, in C. Hoorn and
1197 F. Wesseling, eds., *Amazonia: Landscape and species evolution a look into the past: Wiley-*
1198 *Blackwell*, p. 38–61.
- 1199 Mora, A., Reyes-Harker, A., Rodríguez, G., Tesón, E., Ramírez-Arias, J.C., Parra, M., Caballero,
1200 V., Mora, J.P., Quintero, I., Valencia, V., Ibáñez, M., Horton, B.K., Stockli, D.F., 2013. Inversion
1201 tectonics under increasing rates of shortening and sedimentation: Cenozoic example from the
1202 Eastern Cordillera of Colombia. In: Nemečok, M., Mora, A., and Cosgrove, J.W., eds., *Thick-Skin-*

- 1203 Dominated Orogens: From Initial Inversion to Full Accretion: Geological Society of London,
1204 Special Publication 377, 411-442. doi:10.1144/SP377.6
- 1205 Mora, A., Parra, M., Forero, G., Blanco, V., Moreno, N., Caballero, V., Stockli, D., Duddy, I.,
1206 Ghorbal, B., 2015, What drives orogenic asymmetry in the Northern Andes? A case study from the
1207 apex of the Northern Andean Orocline, in C. Bartolini and P. Mann, eds., Petroleum geology and
1208 potential of the Colombian Caribbean Margin: AAPG Memoir 108, 547–586.
- 1209 Mora, J.A., Oncken, O., Le Breton, E., Ibáñez-Mejía, M., Faccenna, C., Veloza, G., Vélez, V.,
1210 Freitas, M., Mesa, A., 2017. Linking Late Cretaceous to Eocene tectonostratigraphy of the San
1211 Jacinto fold belt of NW Colombia with Caribbean Plateau collision and flat subduction. *Tectonics*
1212 36, 2599–2629.
- 1213 Mora-Páez, H., Kellogg, J. N., Freymueller, J. T., Mencin, D., Fernandes, R. M. S., Diederix, H.,
1214 LaFemina, P., Cardona-Piedrahita, L., Lizarazo, S., Peláez-Gaviria, J.R., Díaz-Mila, F., Bohórquez-
1215 Orozco, O., Giraldo-Londoño, L., Corchuelo-Cuervo, Y., 2019. Crustal deformation in the northern
1216 Andes – A new GPS velocity field. *Journal of South American Earth Sciences*.
1217 doi:10.1016/j.jsames.2018.11.002.
- 1218 Nie, J., Horton, B.K., Saylor, J.E., Mora, A., Mange, M., Garzzone, C.N., Basu, A., Moreno, C.J.,
1219 Caballero, V., Parra, M., 2012. Integrated provenance analysis of a convergent retroarc foreland
1220 system: U-Pb ages, heavy minerals, Nd isotopes, and sandstone compositions of the Middle
1221 Magdalena Valley basin, northern Andes, Colombia. *Earth Sci. Rev.* 110, 111-126.
- 1222 Osorio-Afanador, D., Velandia, F., 2021. Late Jurassic syn-extensional sedimentary deposition and
1223 Cenozoic basin inversion as recorded in The Girón Formation, northern Andes of Colombia.
1224 *Andean Geology* 48 (2), 219-236.
- 1225 Parra, M., Mora, A., Sobel, E. R., Strecker, M.R., and González, R., 2009. Episodic orogenic front
1226 migration in the northern Andes: Constraints from low-temperature thermochronology in the
1227 Eastern Cordillera, Colombia. *Tectonics* 28, TC4004, doi:10.1029/2008TC002423.
- 1228 Parra, M., A. Mora, C. Jaramillo, V. Torres, G. Zeilinger, M. R. Strecker., 2010, Tectonic controls
1229 on Cenozoic foreland basin development in the north-eastern Andes, Colombia: *Basin Research* 22
1230 (6), 874–903.
- 1231 Patiño, A. M., Parra, M., Ramírez, J. C., Sobel, E. R., Glodny, J., Almendral, A., Echeverri, S.,
1232 2019. Chapter 5 - Thermochronological constraints on Cenozoic exhumation along the southern
1233 Caribbean: The Santa Marta range, northern Colombia. In B. K. Horton, A. Folguera (Eds.),
1234 *Andean Tectonics*, 103-132.
- 1235 Pérez, O. J., Wesnousky, S. G., De La Rosa, R., Márquez, J., Uzcátegui, R., Quintero, C., Liberal,
1236 L., Mora-Páez, H., Szeliga, W., 2018. On the interaction of the North Andes plate with the
1237 Caribbean and South American plates in northwestern South America from GPS geodesy and
1238 seismic data. *Geophysical Journal International* 214(3), 1986-2001. doi:10.1093/gji/ggy230 %J
1239 *Geophysical Journal International*.
- 1240 Pérez-Consuegra, N., Hoke, G. D., Mora, A., Fitzgerald, P., Sobel, E. R., Sandoval, J. R., Glodny,
1241 J., Valencia, V., Parra, M., Zapata, S., 2021. The case for tectonic control on erosional exhumation
1242 on the tropical Northern Andes based on thermochronology data. *Tectonics*, 40, e2020TC006652.
1243 <https://doi.org/10.1029/2020TC006652>.
- 1244 Pindell, J., Dewey, J. 1982., Permo-Triassic reconstruction of western Pangea and the evolution of
1245 the Gulf of Mexico/Caribbean region. *Tectonics* 1 (2), 179-211.
- 1246 Ramírez-Arias, J. C., Mora, A., Rubiano, J., Duddy, I., Parra, M., Moreno, N., Casallas, W., 2012.
1247 The asymmetric evolution of the Colombian Eastern Cordillera. Tectonic inheritance or climatic
1248 forcing? New evidence from thermochronology and sedimentology. *Journal of South American*

- 1249 Earth Sciences 39, 112-137. Reiners, P.W. 2005. Zircon (U-Th)/He Thermochronometry. Reviews
1250 in Mineralogy and Geochemistry 58, 151-179.
- 1251 Reiners, P.W., Spell, T.L., Nicolescu, S., and Zanetti, K.A., 2004. Zircon (U-Th)/He
1252 thermochronometry: He diffusion and comparisons with $^{40}\text{Ar}/^{39}\text{Ar}$ dating, *Geochim. Cosmochim.*
1253 *Acta* 68 (8), 1857–1887.
- 1254 Reiners, P. W., Brandon, M. T., 2006. Using thermochronology to understand orogenic erosion:
1255 *Annual Review of Earth and Planetary Sciences* 34, 419–466.
- 1256 Restrepo-Moreno, S. A., Foster, D. A., Bernet, M., Min, K., Noriega, S., 2019. Morphotectonic and
1257 Orogenic Development of the Northern Andes of Colombia: A Low-Temperature
1258 Thermochronology Perspective. In F. Cediél, R. P. Shaw (Eds.), *Geology and Tectonics of*
1259 *Northwestern South America: The Pacific-Caribbean-Andean Junction* (pp. 749-832). Cham:
1260 Springer International Publishing.
- 1261 Restrepo-Pace, P.A., Cediél, F., 2019. Proterozoic Basement, Paleozoic Tectonics of NW South
1262 America, and Implications for Paleocontinental Reconstruction of the Americas. In F. Cediél, R. P.
1263 Shaw (Eds.), *Geology and Tectonics of Northwestern South America: The Pacific-Caribbean-*
1264 *Andean Junction* (pp. 97-112). Cham: Springer International Publishing.
- 1265 Roeder, D., Chamberlain, R., 1995. Eastern Cordillera of Colombia: Jurassic-Neogene crustal
1266 evolution. In: Tankard, A., Suarez, R., Welsink, H. (Eds.), *Petroleum Basins of South America.*
1267 *American Association of Petroleum Geologists* 62, 633-645.
- 1268 Sarmiento, L.F., 2001. Mesozoic rifting and Cenozoic basin inversion history of the Eastern
1269 Cordillera, Colombian Andes inferences from tectonic models. Tectonics Department. Vrije
1270 Universiteit, Amsterdam, p. 319.
- 1271 Sarmiento-Rojas, L.F., Van Wess, J.D., Cloetingh, S., 2006. Mesozoic transtensional basin history
1272 of the eastern Cordillera, Colombian Andes: Inferences from tectonic models. *Journal of South*
1273 *American Earth Sciences* 21, 383-411.
- 1274 Saylor, J., Horton, B., Stockli, D., Mora, A., Corredor, J., 2012a. Structural and
1275 thermochronological evidence for Paleogene basement-involved shortening in the axial Eastern
1276 Cordillera, Colombia. *J. S. Am. Earth Sci.* 39, 202–215.
- 1277 Saylor, J.E., Stockli, D.F., Horton, B.K., Nie, J., Mora, A., 2012b. Discriminating rapid exhumation
1278 from syndepositional volcanism using detrital zircon double dating: Implications for the tectonic
1279 history of the Eastern Cordillera, Colombia. *Geological Society of America Bulletin* 124 (5-6),
1280 762–79.
- 1281 Shagam, R., Kohn, B.P., Banks, P.O., Dasch, L.E., Vargas, R., Rodríguez, G.I., Pimentel, N., 1984.
1282 Tectonic implications of Cretaceous-Pliocene fission-track ages from rocks of the circum-
1283 Maracaibo basin región of western Venezuela and eastern Colombia. In: Bonini, W.E., Hargraves,
1284 R., Shagam, R. (Eds.), *The Caribbean-south American Plate Boundary and Regional Tectonics,*
1285 *GSA Memoir*, 162, 385-412.
- 1286 Shuster, D. L., Flowers, R. M., Farley, K. A., 2006. The influence of natural radiation damage on
1287 helium diffusion kinetics in apatite. *Earth and Planetary Science Letters* 249, 148-161.
- 1288 Siravo, G., Fellin, M.G., Faccena, C., Maden, C., 2019. Transpression and build-up of the
1289 cordillera: the example of the Bucaramanga Fault (eastern cordillera, Colombia) *J. Geol. Soc.*, 177,
1290 14-30.
- 1291 Spotila, J.A., House, M.A., Niemi, N.A., Brady, R.C., Oskin, M., Buscher, J.T., 2007a. Patterns of
1292 bedrock uplift along the San Andreas fault and implications for mechanisms of transpression.
1293 *Geological Society of America, Special Paper* 434, 15-33.

- 1294 Spotila, J.A., Niemi, N., Brady, R., House, M., Buscher, J., 2007b. Long-term continental
1295 deformation associated with transpressive plate motion: The San Andreas fault. *Geology* 35 (11),
1296 967-970., doi: 10.1130/G23816A.1.
- 1297 Stockli, D. F., Farley, K. A., Dumitru, T. A., 2000. Calibration of the apatite (U-Th)/He
1298 thermochronometer on an exhumed fault block, White Mountains, California. *Geology* 28(11),
1299 983-986. doi:10.1130/0091-7613(2000)28<983:COTAHT>2.0.CO;2 %J *Geology*
- 1300 Taboada, A., Rivera, L., Fuenzalida, A., Cisternas, A., Philip, H., Bijwaard, H., Olaya, J., Rivera,
1301 C., 2000. Geodynamics of the northern Andes: Subductions and Intracontinental deformation
1302 (Colombia). *Tectonics* 19 (5), 787-813.
- 1303 Tagami, T., 2005. Zircon fission-track thermochronology and applications to fault studies: Reviews
1304 in *Mineralogy and Geochemistry* 58, 95–122.
- 1305 Tagami, T., O’Sullivan, P.B., 2005. Fundamentals of fission-track thermochronology. *Rev Mineral*
1306 *Geochem.*, 58, 19–47.
- 1307 Tesón, E., Mora, A., Silva, A., Namson, J., Teixell, A., Castellanos, J., Casallas, W., Julivert, M.,
1308 Taylor, M., Ibáñez, M., Valencia, V., 2013. Relationship of Mesozoic graben development, stress,
1309 shortening magnitude, and structural style in the Eastern Cordillera of the Colombian Andes. In:
1310 Nemečok, M., Mora, A., and Cosgrove, J.W. (eds) *Thick-Skin-Dominated Orogens: From Initial*
1311 *Inversion to Full Accretion*. Geological Society, London, Special Publications 377 (1), 257-283.
- 1312 Toro, J., 1990. The Termination of the Bucaramanga Fault in the Cordillera Oriental, Colombia.
1313 M.Sc. thesis. Tucson, University of Arizona, 60p.
- 1314 Toro, J., Roure, F., Bordas-Le Floch, N., Le Cornec-Lance, S., Sassi, W., 2004. Thermal and
1315 kinematic evolution of the eastern Cordillera fold-and-thrust-belt, Colombia. In: Swennen, R.,
1316 Roure, F., Granath, J.W. (Eds.), *Deformation, Fluid Flow, and Reservoir Appraisal in Foreland Fold*
1317 *and Thrust Belts*. American Association of Petroleum Geologists, Hedberg Series 1, 79-115.
- 1318 Umhoefer, P. J., Whitney, D. L., Teyssier, C., Fayon, A. K., Casale, G., Heizler, M. T., 2007. Yo-
1319 yo tectonics in a wrench zone, Central Anatolian fault zone, Turkey. *Special Paper of the*
1320 *Geological Society of America* 434, 35-57. doi: 10.1130/2007.2434(03).
- 1321 Van der Hammen, T., 1961. Late Cretaceous and Tertiary stratigraphy and tectogenesis of the
1322 Colombian Andes. *Geologie en Mijnbouw* 40, 181–188.
- 1323 van der Lelij, R., 2013. Reconstructing North-Western Gondwana with Implications for the
1324 Evolution of the Iapetus and Rheic Oceans: A Geochronological, Thermochronological and
1325 Geochemical Study. (Ph.D. Thesis). University of Geneva, 248 p, Geneva.
- 1326 van der Lelij, R., Spikings, R., Mora, A., 2016a. Thermochronology and Tectonics of the Mérida
1327 Andes and the Santander Massif, NW South America. *Lithos* 3803. PII: S0024-4937(16)00032-3,
1328 doi: 10.1016/j.lithos.2016.01.006.
- 1329 van der Lelij, R., Spikings, R., Ulianov, A., Chiaradia, M., Mora, A., 2016b. Palaeozoic to Early
1330 Jurassic history of the northwestern corner of Gondwana, and implications for the evolution of the
1331 Iapetus, Rheic and Pacific Oceans: *Gondwana Research* 31, 271-294.
- 1332 Vargas, R., Arias A., Jaramillo L., Téllez N., 1976. Geología de la Plancha 136 - Málaga, escala
1333 1:100.000. INGEOMINAS.
- 1334 Vargas, R., Arias A., Jaramillo L., Téllez N., 1987. Geología de la Plancha 152 - Soatá, escala
1335 1:100.000. INGEOMINAS.
- 1336 Velandia, F., 2005. Interpretación de transcurrencia de las fallas Soapaga y Boyacá a partir de
1337 Imágenes LANDSAT TM, *Boletín de Geología* 27 (1), 81-94.

- 1338 Velandia, F., 2017. Cinemática de las fallas mayores del Macizo de Santander – énfasis en el
1339 modelo estructural y temporalidad al sur de la Falla de Bucaramanga. PhD Thesis. Universidad
1340 Nacional de Colombia, Bogotá, 222pp.
- 1341 Velandia, F., Bermúdez, M., 2018. The transpressive southern termination of the Bucaramanga
1342 Fault (Colombia): insights from geological mapping, stress tensors, and fractal analysis. *Journal of*
1343 *Structural Geology* 115, 190-207, doi: 10.1016/j.jsg.2018.07.020.
- 1344 Velandia, F., García-Delgado, H., Zuluaga, C.A., López, J.A., Bermúdez, M.A., Audemard, F.A.,
1345 2020. Present-day structural frame of the Santander Massif and Pamplona Wedge: The interaction
1346 of the Northern Andes. *Journal of Structural Geology* 137, 104087, doi: 10.1016/j.jsg.2020.104087.
- 1347 Vermeesch, P., 2009. Radial Plotter: a Java application for fission track, luminescence and other
1348 radial plots. *Radiation Measurements*, 44: 409-410. Software is freely available at
1349 <http://www.ucl.ac.uk/~ucfbpve/radialplotter/>.
- 1350 Vermeesch, P., 2018., IsoplotR: A free and open toolbox for geochronology. *Geoscience Frontiers*,
1351 9(5), 1479-1493. doi:<https://doi.org/10.1016/j.gsf.2018.04.001>.
- 1352 Villagómez, D., 2010. Thermochronology, Geochronology and Geochemistry of the Western and
1353 Central Cordilleras and Sierra Nevada de Santa Marta, Colombia: The Tectonic Evolution of NW
1354 South America. PhD Thesis No4277. Université de Genève. 166p.
- 1355 Villagómez, D., Spikings, R., Magna, T., Kammer, A., Winkler, W., and Beltrán, A., 2011,
1356 Geochronology, geochemistry and tectonic evolution of the Western and Central cordilleras of
1357 Colombia: *Lithos* 125 (3), 875–896.
- 1358 Villamil, T., 1999. Campanian-Miocene tectonostratigraphy, depocenter evolution and basin
1359 development of Colombia and western Venezuela. *Paleogeography, Paleoclimatology,*
1360 *Paleoecology* 153, 239-275.
- 1361 Villarroel, C., Concha, A. Macías, C., 2001. El lago Pleistoceno de Soatá (Boyacá, Colombia):
1362 Consideraciones estratigráficas, paleontológicas y paleoecológicas. *Geología Colombiana*, 26, 79-
1363 93.
- 1364 Vinasco, C.J., Cordani, U.G., González, H., Weber, M., Peláez, C., 2006. Geochronological,
1365 isotopic, and geochemical data from Permo-Triassic granitic gneisses and granitoids of the
1366 Colombian Central Andes. *J. S. Am. Earth Sci.* 21, 355–371.
1367 <https://doi.org/10.1016/j.jsames.2006.07.007>.
- 1368 Wagner, G. A., Reimer, G. M., 1972. Fission track tectonics: The tectonic interpretation of fission
1369 track apatite ages, *Earth Planet. Sci. Lett.* 14, 263 – 268.
- 1370 Wagner, G. A., Reimer, G. M., Jaeger, E., 1977. Cooling ages derived by apatite fission track, mica
1371 Rb-Sr, and K-Ar dating: The uplift and cooling history of the central Alps, *Mem. Ist. Geol. Min.*
1372 *Univ. Padova* 30, 1 – 27.
- 1373 Wagner, G.A., van den Haute, P., 1992. Fission-track dating. Kluwer Academic Publishers,
1374 Dordrecht, *Solid Earth Sciences Library* vol 6, 285p.
- 1375 Ward, D., Goldsmith, R., Cruz, J., Jaramillo, L., Vargas, R., 1977a. Mapa geológico del
1376 cuadrángulo typically H-13 Pamplona, escala 1:100.000. INGEOMINAS.
- 1377 Ward, D., Goldsmith, R., Jimeno, A., Cruz, J., Restrepo, H., Gómez, E., 1977b. Mapa geológico del
1378 cuadrángulo H-12 Bucaramanga, escala 1:100.000. INGEOMINAS.
- 1379 Wildman, M., Brown, R., Beucher, R., Persano, C., Stuart, F., Gallagher, K., Schwanethal, J.,
1380 Carter, A., 2016. The chronology and tectonic style of landscape evolution along the elevated

- 1381 Atlantic continental margin of South Africa resolved by joint apatite fission track and (U-Th-
1382 Sm)/He thermochronology, *Tectonics* 35, 511–545, doi:10.1002/2015TC004042.
- 1383 Williamson, J.H., 1968. Least-squares fitting of a straight line: *Canadian Journal of Physics* 46,
1384 1845–1847, doi:10.1139/p68-523.
- 1385 Wolf, R.A., Farley, K.A., Kass, D.M., 1998. Modeling of the temperature sensitivity of the apatite
1386 (U–Th)/He thermochronometer. *Chemical Geology* 148, 105–114.
- 1387 Wolfe, M. R., Stockli, D. F., 2010. Zircon (U–Th)/He thermochronometry in the KTB drill hole,
1388 Germany, and its implications for bulk He diffusion kinetics in zircon. *Earth and Planetary Science*
1389 *Letters* 295(1), 69-82. doi:<https://doi.org/10.1016/j.epsl.2010.03.025>
- 1390 Zaun, P. E., Wagner, G. A., 1985. Fission-track stability in zircons under geological conditions:
1391 Nuclear tracks and radiation measurements, 10, 303–307.
- 1392 Zeitler, P.K., 2015. U-Th/He Dating in In Rink, W. J., In Thompson, J. W., In Heaman, L. M., In
1393 Jull, A. J. T., & In Paces, J. B. *Encyclopedia of scientific dating methods*. 932-941, DOI
1394 10.1007/978-94-007-6326-5_131-1.
- 1395 Zeitler, P.K., Enkelmann, E., Thomas, J.B., Watson, E.B., Ancuta, L.D., Idleman, B.D., 2017.
1396 Solubility and trapping of helium in apatite: *Geochimica et Cosmochimica Acta* 209, 1-8.
1397
- 1398 Zuluaga, C.A., Amaya, S., Urueña, C., Bernet, M., 2017. Migmatization and low-pressure
1399 overprinting metamorphism as record of two pre Cretaceous tectonic episodes in the Santander
1400 Massif of the Andean basement in northern Colombia (NW South America). *Lithos* 274, 123-146.
1401
- 1402 Zuluaga, C.A., López, J.A., 2019. Ordovician Orogeny and Jurassic Low-Lying Orogen in the
1403 Santander Massif, Northern Andes (Colombia). In: Cediel, F., Shaw, R.P. (eds.), *Geology and*
1404 *Tectonics of Northwestern South America*. *Frontiers in Earth Sciences*. Springer, Chapter 4, 195-
1405 250. https://doi.org/10.1007/978-3-319-76132-9_4.
1406

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Journal Pre-proof