



HAL
open science

Shedding light on tropospheric H₂SO₄ production from Criegee intermediates + SO₂: a comprehensive laboratory chamber study using the highly instrumented HELIOS platform

Yangang Ren, Max McGillen, Alexandre Kukui, Véronique Daële, Wahid Mellouki

► To cite this version:

Yangang Ren, Max McGillen, Alexandre Kukui, Véronique Daële, Wahid Mellouki. Shedding light on tropospheric H₂SO₄ production from Criegee intermediates + SO₂: a comprehensive laboratory chamber study using the highly instrumented HELIOS platform. 23rd EGU General Assembly, held online, Apr 2021, Vienna, Austria. 10.5194/egusphere-egu21-14862 . insu-03520159

HAL Id: insu-03520159

<https://insu.hal.science/insu-03520159>

Submitted on 6 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Shedding light on tropospheric H₂SO₄ production from Criegee intermediates + SO₂: a comprehensive laboratory chamber study using the highly instrumented HELIOS platform

Yangang Ren¹, Max McGillen¹, Alexandre Kukui², Veronique Daële¹, and Abdelwahid Mellouki¹

¹Institut de Combustion, Aérothermique, Réactivité, Environnement (ICARE – CNRS), Orléans, France (yangang.ren@cnrs-orleans.fr)

²Laboratoire de Physique et Chimie de l'Environnement et de l'Espace (LPC2E – CNRS/Université Orléans/CNES), Orléans, France

Although the dominant source of H₂SO₄ in the atmosphere is generally considered to be the reaction of SO₂ with OH, it is probable that the rapid reactions of Criegee intermediates (CIs) with SO₂ can contribute significantly to the tropospheric H₂SO₄ budget under certain conditions. CIs are produced from alkene ozonolysis, and the vast quantities of unsaturated biogenic and anthropogenic volatile organic compounds emitted could provide a large and diverse flux of CIs to the atmosphere. There remain several key uncertainties regarding the global importance of CIs towards SO₂ oxidation, which are principally related to the ambient concentrations of CIs and the competition between CI reaction with SO₂ against the many other bimolecular and unimolecular loss processes. This is especially true of the larger, more complex CIs that are produced from terpene ozonolysis.

We present experimental studies of the ozonolysis of tetramethylethylene, α-pinene and limonene, using the HELIOS chamber. HELIOS is a highly instrumented large-scale outdoor atmospheric simulation chamber and consists of a hemispheric 90 m³ Teflon-foil reactor, which is interfaced to a variety of on-line measurements including FTIR, PTR-ToF-MS, FIGAERO-ToF-CIMS, OH/H₂SO₄-CIMS, Aerolaser HCHO, LOPAP and SMPS, together with several GC-MS/FID and LC-MS instruments and a suite of monitors (NO, NO₂, O₃). Equipped with this range of instrumentation we are able to conduct alkene ozonolysis under near-ambient conditions, whilst we also have a high coverage of key reactive species in the systems of interest.

From our results, we are able to provide new information regarding kinetic and mechanistic behaviour of several atmospherically important CIs and their reactive intermediates, providing new constraints on the role of CIs on the tropospheric H₂SO₄ budget.

Keywords: ozonolysis, Criegee Intermediate, sulfur dioxide, sulfuric acid, kinetics