



HAL
open science

The first fossil Parascleroderma (Hymenoptera: Bethylidae): a new species in mid-miocene zhangpu amber

Manuel Brazidec, Vincent Perrichot

► **To cite this version:**

Manuel Brazidec, Vincent Perrichot. The first fossil Parascleroderma (Hymenoptera: Bethylidae): a new species in mid-miocene zhangpu amber. *Palaeoworld*, 2023, 32 (4), pp.709-715. 10.1016/j.palwor.2022.01.009 . insu-03557051

HAL Id: insu-03557051

<https://insu.hal.science/insu-03557051>

Submitted on 4 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Journal Pre-proofs

The first fossil *Parascleroderma* (Hymenoptera: Bethylidae): a new species in mid-miocene zhangpu amber

Manuel Brazidec, Vincent Perrichot

PII: S1871-174X(22)00009-9

DOI: <https://doi.org/10.1016/j.palwor.2022.01.009>

Reference: PALWOR 673

To appear in: *Palaeoworld*

Received Date: 23 November 2021


Revised Date: 18 January 2022

Accepted Date: 24 January 2022

Please cite this article as: M. Brazidec, V. Perrichot, The first fossil *Parascleroderma* (Hymenoptera: Bethylidae): a new species in mid-miocene zhangpu amber, *Palaeoworld* (2022), doi: <https://doi.org/10.1016/j.palwor.2022.01.009>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2022 Elsevier B.V. and Nanjing Institute of Geology and Palaeontology, CAS.



The first fossil *Parascleroderma* (Hymenoptera: Bethylidae): a new species in mid-Miocene Zhangpu amber

Manuel Brazidec ^{a, b} *, Vincent Perrichot ^a

^a Univ. Rennes, CNRS, Géosciences Rennes, UMR 6118, 35000 Rennes, France

^b State Key Laboratory of Palaeobiology and Stratigraphy, Nanjing Institute of Geology and Palaeontology and Center for Excellence in Life and Palaeoenvironment, Chinese Academy of Sciences, 39 East Beijing Road, Nanjing 210008, China

* Corresponding author.

E-mail addresses: manuel.brazidec@gmail.com (M. Brazidec),

vincent.perrichot@univ-rennes1.fr (V. Perrichot)

Abstract

The recently discovered amber biota from Zhangpu, China, offers a unique window into the tropical ecosystem of an Asian rainforest during the Mid-Miocene Climatic Optimum (MMCO). Here, the first flat wasp (Bethylidae) from this deposit is documented: *Parascleroderma palaeosinica* n. sp. is described and figured from a single male, and is the first fossil *Parascleroderma* known to date. The new extinct species is characterized by its minute size, the pedicel longer than the flagellomeres, the posterior ocelli inserted much posteriad on head, the notauli not impressed on anterior third of anteromesoscutum, and the hypopygium simple, with posterior margin slightly incurved medially. *Parascleroderma* is a cosmopolitan genus within the Pristocerinae and it is therefore not surprising to find its record in the Zhangpu biota, simultaneously with the northern expansion of the southeastern Asian tropical forest during the MMCO.

Keywords: Chrysidoidea; Bethylidae; Pristocerinae; Cenozoic; Miocene; China

1. Introduction

With up to 3000 current species arranged in 96 valid genera (Azevedo et al., 2018), the Bethylidae form the largest family among the Chrysidoidea. This family of wasps displays its greatest diversity in tropical regions but is cosmopolitan in its

global distribution. Bethylid wasps are generally considered as external parasitoids of lepidopteran and coleopteran larvae that they paralyze and drag into cracks (Evans, 1964; Rubink and Evans, 1979).

Within Chrysidoidea, Bethylidae are generally retrieved as the sister-group of Chrysididae in phenotypic studies (Brothers and Carpenter, 1993; Carpenter, 1999; Brothers, 2011; Brothers and Melo, 2021), but sister to Plumariidae in phylogenomic studies (Branstetter et al., 2017; Peters et al., 2017). Grimaldi and Engel (2005) indicate that the narrowly transverse and nearly completely hidden metanotum would be the synapomorphy of the clade (Bethylidae + Chrysididae). However, in the bethylid subfamily Pristocerinae, the metanotum is developed medially and tends to overlap the mesoscutellum (Azevedo et al., 2018). Bethylids are recognizable by the prognathous head, the antennae with the same number of antennomeres in males and females (11 or 10, rarely 8 or 7 flagellomeres), the pronotum with anterior flange, the prosternum small, transverse, and the six or seven exposed metasomal terga (Finnamore and Brothers, 1993). The flattened body, evolved to sneak into their host habitats, is also retained as a characteristic feature of the family, hence their nickname 'Flat wasps'. Nine subfamilies were recently recognized in the Bethylidae (Colombo et al., 2020): the extant Bethylinae, Epyrinae, Mesitiinae, Pristocerinae and Scleroderminae, and the extinct Elektroepyrinae, Holopsenellinae, Lancepyrinae and Protopristocerinae. Since then, however, *Holopsenella* Engel et al., 2016, type genus of the Holopsenellinae, has been removed from Bethylidae and elevated to a family left as *incertae sedis* in the Aculeata (Lepeco and Melo, 2022).

With more than 90 species in seven of the eight subfamilies (no fossil Mesitiinae has been described yet), the fossil record of Bethylidae spans nearly 130 million years from the Lower Cretaceous to the Holocene (see Martynova et al., 2019; and the posterior discoveries of: Engel, 2019; Falières and Nel, 2019a, 2019b, 2019c, 2020; Colombo et al., 2020, 2021a, 2021b; Jouault et al., 2020, 2021; Colombo and Azevedo, 2021; Jouault and Brazidec, 2021; Tribull et al., 2021). Despite this relatively high richness compared to other chrysidoid lineages (e.g., Chrysididae), lots of new species are likely to be described from Cretaceous or Cenozoic deposits, particularly from the recently discovered Miocene amber of Ethiopia and China (Bouju and Perrichot, 2020; Wang et al., 2021). Even from the long-surveyed deposits such as Baltic amber, new taxa may be recovered, as exemplified by the recent discovery of a new pristocerine genus (Tribull et al., 2021).

Here, we report the first Bethylidae from the rich mid-Miocene amber biota of Zhangpu, in Fujian Province, China, and we discuss its taxonomic affinities.

2. Material and methods

Zhangpu amber is found with plant impressions in two layers of sandy mudstone interbedded with coal seams that belong to the Fotan Group, a geological unit that occurs widely in Zhangpu County, Fujian Province, southeastern China (Wang et al., 2021, fig. 1). Under- and overlying basalt layers allow constrained dating of the amber between 14.8 ± 0.6 Ma and 14.7 ± 0.4 Ma, which corresponds to the middle Miocene (Langhian; Zheng et al., 2019).

This study is based on a complete male specimen kindly lent by Prof. Bo Wang (Nanjing, China) to the authors for study, and housed in the Nanjing Institute of Geology and Palaeontology (NIGP), Chinese Academy of Sciences, Nanjing, China. The fragile amber piece was embedded in a block of epoxy resin (Araldite® 2020) for consolidation and posteriorly polished to facilitate the observation of the specimen, using thin silicon carbide sanding papers on a grinder polisher (Buehler EcoMet 30). The examination and photographs were conducted with a Leica DMC4500 camera attached to a Leica M205C stereomicroscope. All images are digitally stacked photomicrographic composites of several focal planes, which were obtained using Helicon Focus 6.7 software. Adobe Illustrator CC2019 and Photoshop CC2019 software were used to compose the figures and ImageJ 1.53 for measurements. The description of the characters follows the nomenclature of Lanes et al. (2020).

This published work and its new nomenclatural act are registered in ZooBank with the following LSID (reference): urn:lsid:zoobank.org:pub:570AA2F6-4E8D-4DD0-B01A-9D824F8CD4F6.

3. Systematic palaeontology

Superfamily Chrysidoidea Latreille, 1802

Family Bethylidae Haliday, 1839

Subfamily Pristocerinae Mocsáry, 1881

Genus *Parascleroderma* Kieffer, 1904

Parascleroderma palaeosinica n. sp.

(Figs. 1, 2)

LSID: urn:lsid:zoobank.org:act:1E0E3362-9053-47AA-A6A2-457880270E5E.

Etymology: The species name combines ‘*palaiós*’, meaning ‘ancient’ in Greek and ‘*sinica*’, in reference to the geographic origin of Zhangpu amber.

Material: Holotype male, NIGP178180.

Type locality: Zhangpu County, Zhangzhou Prefecture, Fujian Province, China.

Horizon: Sedimentary layer II, Fotan Group, middle Miocene: Langhian, 14.7 Ma.

Diagnosis (male): Body minute-sized, 1.67 mm long, pale brown; pedicel longer than each flagellomere (Fig. 1C); posterior ocelli inserted much posteriad on head (Fig. 1D); notauli not impressed on anterior third of anteromesoscutum, well-marked on posterior two-thirds (Fig. 1D); hypopygium simple with posterior margin slightly incurved medially (Figs. 1A, 2).

Description: Male. Body flattened and narrow, not visibly sculptured, pale brown. Head prognathous, longer than high, longer than wide, oval in dorsal view; frons smooth, concave, with faint median sulcus originating from anterior ocellus, frons with sparse long setae; compound eye oval, longer than high, without visible setae, located in anterior part of head; mandibles only slightly overlapping at tip; maxillary palpus with at least 4 palpomeres and long apical setae; 13 antennomeres, all antennomeres bearing numerous long setae at most as long as one third of flagellomere length; scape elongate, 2.5 times longer than wide, slightly curved, pedicel as long as apical flagellomere; all flagellomeres cylindrical, longer than wide, flagellomeres I–X subequal in length, only slightly shorter than flagellomere XI; anterior ocellus located at back of frons, posterior ocelli pushed at extreme posterior of frons and widely separated from each other; occipital carina present, forming ridge extending from occipital foramen to mandibles.

Mesosoma shorter than metasoma; propleuron elongate, slightly exposed dorsally, ‘neck-shaped’; pronotal flange developed, dorsal pronotal area longer than anteromesoscutum, narrowed anteriorly; prosternum reduced between procoxae; mesoscutum with notauli impressed on posterior two thirds, slightly converging posteriorly, absent anteriorly; parapsidal signum not marked; mesoscutellum with blunted apex; metanotum large, overlapping mesoscutellum posteriorly, transversally carinate; metapectal-propodeal complex longer than wide, without posterolateral spines, smooth, without sharp longitudinal carinae. Forewing with only costal [C], radial [R] and cubital [1Cu] cells closed by tubular veins; C, Sc+R, M+Cu, A, Rs, cu-

a, basal segment of M and Cu tubular; basal segments of Rs and M forming contiguous line; cu-a angled at junction with Cu; Cu nebulous posteriorly to junction with 1m-cu then completely fading; 1m-cu nebulous; pterostigma reduced in length, rapidly vanishing after 2r-rs; R1 not apparent after pterostigma; 2r-rs and Rs continuously arched, without marked angles between sections, distal segment of Rs tubular for much of its length, then reaching wing margin as nebulous vein, thus not closing radial cell [2R1]; distal segment of M completely spectral. Hind wing slightly shorter than forewing; C only pigmented vein; Rs originating from C in anterior half of wing, nebulous to spectral; no closed cells. Legs sparsely pubescent; profemur swollen; first tarsomere longer than combined length of following tarsomeres; pro- and mesotibiae with two spurs, metatibial spurs not visible; metafemur enlarged, without spines.

Metasoma sparsely setose; metasomal tergites without modifications; metasomal tergite I as long as wide; metasomal tergite II shorter than I; hypopygium not divided, with outer surface flat and posterior margin weakly incurved medially; genital structures not visible.

Measurements (in mm): Body length 1.67; head length 0.34, width 0.26; scape length 0.10, width 0.04; pedicel length 0.06; flagellomeres I–X length ca. 0.04–0.05; flagellomere XI length 0.06; mesosoma length 0.63; forewing length ca. 1.10; hind wing length 0.98; mesosoma length 0.70.

4. Discussion

Following the key to the subfamilies of Bethylidae of Azevedo et al. (2018), the new fossil keys out in Pristocerinae because of the following characters: macropterous species, forewing with Rs+M vein absent, metanotum developed medially and overlapping mesoscutellum posteriorly. This last character is usually considered as a synapomorphy of the Pristocerinae.

Following the keys to pristocerine genera of Azevedo et al. (2018), the new fossil keys near *Parascleroderma* because of the following characters: body smaller than 15 mm, hypopygium not deeply divided into two lobes, outer surface of hypopygium flat, dorsal pronotal area short, forewing with intersection of Rs&M far from pterostigma, profemur narrow, metafemur without spines, hypopygium slightly incurved. Among the diagnostic characters provided by Azevedo et al. (2018) for the genus *Parascleroderma*, the fossil clearly possesses the body flattened and elongated,

the propleuron elongated, the pronotum elongated and anteriorly narrowed, the intersection of Rs&M and Sc+R far from pterostigma and the metafemur enlarged. Because of the preservation in amber, characters on the clypeus, genitalia, and internal features of hypopygium cannot be observed. *Parascleroderma* is closely related to *Afgoiogfa* Argaman, 1988, *Dissomphalus* Ashmead, 1893, *Foenobethylus* Kieffer, 1913, *Protisobrachium* Benoit, 1957, and *Pseudisobrachium* Kieffer, 1904 (Alencar et al., 2018). These genera differ from our fossil as follows (Azevedo et al., 2018): pterostigma comma-like, maxillary palpus with 3 palpomeres and a long apical seta or 6 palpomeres with only short setae, notauli large and poorly impressed (*Afgoiogfa*); body robust, metasomal tergum II with process (*Dissomphalus*); posterior hypopygial margin abruptly v-incurved, metatrochanter and metafemur usually with spines (*Foenobethylus*); eyes with long erect setae, R1 present and long after the pterostigma, posterior hypopygial margin strongly incurved (*Protisobrachium*); forewing with intersection of Rs+M and Sc+R close to pterostigma (*Pseudisobrachium*). Therefore, our new species is excluded from these genera and, instead, displays features that are all diagnostic of *Parascleroderma*.

Parascleroderma was erected by Kieffer (1904) based on a species from Italy: *P. nigriceps* Kieffer, 1904. In the following years, the same author subsequently described species from Hungary, Italy, France and the Seychelles (e.g., Kieffer, 1908, 1912): *P. fulviceps* Kieffer, 1906, *P. fuscipennis* (Kieffer, 1905) (synonymized from *Ceratepyris* Kieffer, 1905 by Argaman, 1988), *P. remota* (Kieffer, 1912) (transferred from *Apenesia* by Azevedo et al., 2018), *P. seychellensis* (Kieffer, 1912) (transferred from *Sclerodermus* Latreille, 1809 by Evans, 1964) and *P. sulcatifrons* (Kieffer, 1908) (synonymized from *Ceratepyris* by Argaman, 1988). The genus currently includes 32 extant species and has a cosmopolitan distribution (Fig. 3), but was hitherto undocumented from the fossil record. Among major contributors to the knowledge of *Parascleroderma*, Evans (1963, 1964, 1967, 1978) described four new species from America North of Mexico: *P. arivaca* Evans, 1978, *P. carinata* Evans, 1964, *P. insolita* (Evans, 1963) (transferred from *Apenesia* by Evans, 1964), *P. minima* Evans, 1978; and one from Peru: *P. pucallpa* Evans, 1967. Argaman (1988) summarized the known species and erected six new ones from Europe, Turkey, Israel, and Iran: *P. cismora* Argaman, 1988, *P. fiturcata* Argaman, 1988, *P. hindola* Argaman, 1988, *P. norcasta* Argaman, 1988, *P. oriana* Argaman, 1988 and *P. varlinda* Argaman, 1988. Terayama (1998, 2006), Xu et al. (2002), and Lim et al.

(2011) described new species from the Oriental region (i.e., Korea, China, Taiwan, and Japan): *P. atayal* Terayama, 1998, *P. ishama* Terayama, 2006, *P. jinmo* Terayama, 2006, *P. maae* Xu, He and Terayama, 2002, *P. okajimai* Terayama, 1998, *P. renaiensis* Terayama, 1998 and *P. tetradentica* Lim and Lee in Lim et al., 2011. The vast majority of the known *Parascleroderma* species have been collected in the northern hemisphere but six species occur in the southern hemisphere: one from Peru and two from the Seychelles, mentioned above, *P. neounicolor* Alencar and Azevedo in Azevedo et al., 2018 (Marquises Islands), *P. nigra* Brues, 1910 (South Africa) and *P. azevedonis* (New Zealand). Alencar et al. (2018) also indicate potential new species in Thailand, Australia and South Africa.

The Zhangpu amber originated from a tropical rainforest with abundant Dipterocarpaceae (Wang et al., 2021) in the greenhouse context of the Mid-Miocene Climatic Optimum (Kasbohm and Schoene, 2018). Thus, the presence of *Parascleroderma* in Thailand nowadays is interesting because southeast Asian dipterocarp forests are the closest extant relatives of the mid-Miocene Zhangpu amber forest in terms of floral assemblage and climate (Wang et al., 2021, fig. 2). This further supports our attribution of the new species to *Parascleroderma*, whose extant species are present in tropical biomes. Therefore, it is not surprising to discover an extinct species of *Parascleroderma* northward in the Zhangpu biome (24.2°N), simultaneously with the megathermal climate of the MMCO that allowed the tropical forest to expand toward higher latitudes.

Acknowledgments

We warmly thank the guest editor Prof. Bo Wang for its invitation to contribute this special issue of Palaeoworld and for its precious comments on the manuscript. We are grateful to the reviewers, Prof. Celso O. Azevedo, Prof. Massimo Olmi and an anonymous reviewer for valuable comments and remarks on the earlier manuscript. Research on the Zhangpu amber was supported by the Strategic Priority Research Program of the Chinese Academy of Sciences (XDB26000000) and National Natural Science Foundation of China (42125201, 41688103). This work is part of Manuel Brazidec's Ph.D. project CHRYSIS: "The role of greenhouses on the diversification and evolution of chrysidoid wasps".

References

- Alencar, I.D.C.C., Waichert, C., Azevedo, C.O., 2018. Opening Pandora's box of Pristocerinae: molecular and morphological phylogenies of *Apenesia* (Hymenoptera, Bethylidae) reveal several hidden genera. *Systematic Entomology* 43, 481–509.
- Argaman, Q., 1988. A new subfamily of Bethylidae allied to Pristocerinae (Hymenoptera). *Societa Entomologica Italiana* 120, 139–152.
- Ashmead, W.H., 1893. A monograph of the North American Proctotrypidae. *Bulletin of the United States National Museum* 45, 1–472.
- Azevedo, C.O., Alencar, I.D.C.C., Ramos, M.S., Barbosa, D.N., Colombo, W.D., Vargas, J.M., Lim, J., 2018. Global guide of the flat wasps (Hymenoptera, Bethylidae). *Zootaxa* 4489, 1–294.
- Benoit, P.L.G., 1957. Hymenoptera – Bethylidae. Exploration du Parc National Albert. Mission G. F. De Witte, 1933–1935. Fascicule 88. Institut des Parcs Nationaux de Congo Belge, Tervuren, 57 pp.
- Bouju, V., Perrichot, V., 2020. A review of amber and copal occurrences in Africa and their paleontological significance. *BSGF – Earth Sciences Bulletin* 191, Article Number 17, doi: 10.1051/bsgf/2020018.
- Branstetter, M.G., Danforth, B.N., Pitts, J.P., Faircloth, B.C., Ward, P.S., Buffington, M.L., Gates, M.W., Kula, R.R., Brady, S.G., 2017. Phylogenomic insights into the evolution of stinging wasps and the origins of ants and bees. *Current Biology* 27, 1019–1025.
- Brothers, D.J., 2011. A new Late Cretaceous family of Hymenoptera, and phylogeny of the Plumariidae and Chrysoidea (Aculeata). *Zookeys* 130, 515–542.
- Brothers, D.J., Carpenter, J.M., 1993. Phylogeny of Aculeata: Chrysoidea and Vespoidea (Hymenoptera). *Journal of Hymenoptera Research* 2, 227–304.
- Brothers, D.J., Melo, G.A.R., 2021. Re-evaluation of the Cretaceous family Plumalexiidae and its relationships (Hymenoptera: Chrysoidea). *Palaeoentomology* 4, 584–591.
- Brues, C.T., 1910. Some notes on the geological history of the parasitic Hymenoptera. *Journal of New York Entomological Society* 18, 1–22.
- Carpenter, J.M., 1999. What do we know about chrysidoid (Hymenoptera) relationships? *Zoologica Scripta* 28, 215–231.

- Colombo, W.D., Azevedo, C.O., 2021. Synopsis of the fossil Scleroderminae (Hymenoptera, Bethylidae) with description of a new genus and four new species from Baltic amber. *Historical Biology* 33, 630–638.
- Colombo, W.D., Perkovsky, E.E., Azevedo, C.O., 2020. Phylogenetic overview of flat wasps (Hymenoptera, Bethylidae) reveals Elektroepyrinae, a new fossil subfamily. *Palaeoentomology* 3, 269–283.
- Colombo, W.D., Gobbi, F.T., Perkovsky, E.E., Azevedo, C.O., 2021a. Synopsis of the fossil Pristocerinae (Hymenoptera, Bethylidae), with description of two new genera and six species from Burmese, Taimyr, Baltic and Rovno ambers. *Historical Biology* 33, 1736–1752.
- Colombo, W.D., Perkovsky, E.E., Waichert, C., Azevedo, C.O., 2021b. Synopsis of the fossil flat wasps Epyrinae (Hymenoptera, Bethylidae), with description of three new genera and ten new species. *Journal of Systematic Palaeontology* 19, 29–89.
- Engel, M.S., 2019. A holopsenelline wasp in mid-Cretaceous amber from Myanmar (Hymenoptera: Bethylidae). *Palaeoentomology* 2, 199–204.
- Engel, M.S., Ortega-Blanco, J., Azevedo, C.O., 2016. A new bethylid wasp in Lebanese Early Cretaceous amber (Hymenoptera: Chrysidoidea), with comments on other Mesozoic taxa. *American Museum Novitates* 3855, 1–14.
- Evans, H.E., 1963. A revision of the *Apenesia* in the Americas (Hymenoptera, Bethylidae). *Bulletin of the Museum of Comparative Zoology* 129, 241–290.
- Evans, H.E., 1964. A synopsis of the American Bethylidae (Hymenoptera, Aculeata). *Bulletin of the Museum of Comparative Zoology* 132, 1–222.
- Evans, H.E., 1967. New generic records of Bethylidae from South America (Hymenoptera). *Proceedings of the Entomological Society of Washington* 69, 269–272.
- Evans, H.E., 1978. The Bethylidae of America North of Mexico. *Memoirs of the American Entomologist Institute* 27, 1–332.
- Falières, E., Nel, A., 2019a. A new bethylid wasps from the Lowermost Eocene Oise amber (France) (Insecta: Hymenoptera). *Palaeoentomology* 2, 211–214.
- Falières, E., Nel, A., 2019b. The first representative of the extinct flat wasp subfamily Protopristocerinae (Insecta, Hymenoptera, Bethylidae). *Palaeoentomology* 2, 322–236.

- Falières, E., Nel, A., 2019c. The first sclerodermine and pristocerine flat wasps in Lowermost Eocene amber of France (Hymenoptera: Bethyridae). *Comptes Rendus Palevol* 18, 509–515.
- Falières, E., Nel, A., 2020. First fossil of the extant flat wasp genus *Dissomphalus* Ashmead, 1893 (Hymenoptera: Bethyridae: Pristocerinae). *Zoosystema* 42, 207–213.
- Finnamore, A.T., Brothers, D.J., 1993. Superfamily Chrysoidea. In: Goulet, H., Huber, J.T. (Eds.), *Hymenoptera of the World: An Identification Guide to Families*. Agriculture Canada Publications, Ottawa, pp. 130–159.
- Grimaldi, D.A., Engel, M.S., 2005. *Evolution of the Insects*. Cambridge University Press, New York, 725 pp.
- Haliday, A.H., 1839. *Hymenoptera Britannica, Part 1. Oxyura*. Addendum to Vol. 1. *Hymenopterorum synopsis ad methodum fallenii ut plurimum accommodata*. Hippolytas Bailliere, London, 28 pp.
- Jouault, C., Brazidec, M., 2021. A new protopristocerine wasp (Hymenoptera: Bethyridae) from the mid-Cretaceous Burmese amber. *Annales de Paléontologie* 107, 102522.
- Jouault, C., Ngô-Muller, V., Pouillon, J., Nel, A., 2020. New Burmese amber fossils clarify the evolution of bethyrid wasps (Hymenoptera: Chrysoidea). *Zoological Journal of the Linnean Society* 191, 1044–1058.
- Jouault, C., Perrichot, V., Nel, A., 2021. New flat wasps from mid-Cretaceous Burmese amber deposits highlight the bethyrid antiquity and paleobiogeography (Hymenoptera: Chrysoidea). *Cretaceous Research* 123, 104772.
- Kasbohm, J., Schoene, B., 2018. Rapid eruption of the Columbia River flood basalt and correlation with the mid-Miocene climate optimum. *Science Advances* 4, eaat8223, doi: 10.1126/sciadv.aat8223.
- Kieffer, J.J., 1904. Description de nouveaux Dryininae et Bethyridae du Musée Civique de Gênes. *Annali del Museo Civico di Storia Naturale di Genova, Serie 3*, 41, 351–412.
- Kieffer, J.J., 1905. Description de nouveaux Hyménoptères exotiques. *Bulletin de la Société d'Histoire Naturelle de Metz* 24, 85–114.
- Kieffer, J.J., 1906. Description de nouveaux Hyménoptères. *Annales de la Société Scientifique de Bruxelles* 20, 113–178.

- Kieffer, J.J., 1908. Bethylidae. In: Wytsman, P. (Ed.), *Genera Insectorum* 76. Desmet-Verteneuil, Bruxelles, pp. 1–50.
- Kieffer, J.J., 1912. Hymenoptera, Proctotrupoidea. *Transactions of the Linnean Society of London, Series 2, Zoology* 15, 45–80.
- Kieffer, J.J., 1913. Nouveaux microhyménoptères de l’Afrique équatoriale. *Bollettino di Laboratorio di Zoologia Generale e Agraria della R. Scuola Superiore d’Agricoltura in Portici* 7, 105–112.
- Lanes, G.O., Kawada, R., Azevedo, C.O., Brothers, D.J., 2020. Revisited morphology applied for Systematics of flat wasps (Hymenoptera, Bethylidae). *Zootaxa* 4752, 1–127.
- Latreille, P.A., 1802. *Histoire naturelle, générale et particulière des Crustacés et des insectes*. Vol. 3. Dufort, Paris, 348 pp.
- Latreille, P.A., 1809. *Genera Crustaceorum et Insectorum secundum ordinem naturalem in familias desposita, iconibus exemplisque plurimis explicata*. Parislis et Argentorati, Armand Konig, Paris, 399 pp.
- Lepeco, A., Melo, G.A.R., 2022. The wasp genus †*Holopsenella* in mid-Cretaceous Burmese amber (Hymenoptera: †Holopsenellidae stat. nov.). *Cretaceous Research* 131, 105089.
- Lim, J., Terayama, M., Koh, S., Lee, J., Lee, S., 2011. A taxonomic review of the subfamily Pristocerinae (Hymenoptera: Chrysidoidea: Bethylidae) from Korea with descriptions of two new species. *Journal of Natural History* 45, 435–460.
- Martynova, K.V., Perkovsky, E.E., Olmi, M., Vasilenko, D.V., 2019. New records of Upper Eocene chrysidoid wasps from basins of Styr and Stokhod rivers (Rovno amber). *Paleontological Journal* 53, 998–1023.
- Mocsáry, S., 1881. A magyar fauna másnejű darázsai. (Heterogynidae fauna Hungaricae). *Mathematikai s Természettudományi Közlemények* 17, 1–96.
- Peters, R.S., Krogmann, L., Mayer, C., Donath, A., Gunkel, S., Meusemann, K., Kozlov, A., Podsiadlowski, L., Petersen, S., Lanfear, R., Diez, P.A., Heraty, J., Kjer, K.M., Klopstein, S., Meier, R., Polidori, C., Schmitt, T., Liu, S., Zhou, X., Wappler, T., Rust, J., Misof, B., Niehuis, O., 2017. Evolutionary history of the Hymenoptera. *Current Biology* 27, 1013–1018.
- Rubink, W.L., Evans, H.E., 1979. Notes on the nesting behavior of the bethylid wasps, *Epyris eriogoni* Kieffer, in southern Texas. *Psyche* 86, 313–320.

- Terayama, M., 1998. Discovery of the genus *Parascleroderma* Kieffer from the Oriental region, with descriptions of four new species (Hymenoptera: Bethyridae). *Entomological Science* 1, 129–132.
- Terayama, M., 2006. *Insects of Japan, Bethyridae (Hymenoptera)*. Vol. 1. Touka Shobo Co, Fukuoka, 317 pp.
- Tribull, C.M., Pankowski, M.V., Colombo, W.D., 2021. A new genus and species of *Pristocerinae* (Hymenoptera, Bethyridae) from Upper Eocene Baltic amber with a review of conspecific association from insect fossils. *Journal of Hymenoptera Research* 85, 119–133.
- Wang, B., Shi, G., Xu, C., Spicer, R.A., Perrichot, V., Schmidt, A.R., Feldberg, K., Heinrichs, J., Chény, C., Pang, H., Liu, X., Gao, T., Wang, Z., Ślipiński, A., Solórzano-Kraemer, M.M., Heads, S.W., Thomas, M.J., Sadowski, E., Szwed, J., Azar, D., Nel, A., Liu, Y., Chen, J., Zhang, Q., Zhang, Q., Luo, C., Yu, T., Zheng, D., Zhang, H., Engel, M.S., 2021. The mid-Miocene Zhangpu biota reveals an outstandingly rich rainforest biome in East Asia. *Science Advances* 7, eabg0625, doi: 10.1126/sciadv.abg0625.
- Xu, Z.F., He, J., Terayama, M., 2002. A new species of the genus *Parascleroderma* Kieffer, 1904 from China (Hymenoptera: Bethyridae). *Acta Zootaxonomica Sinica* 27, 794–797.
- Zheng, D., Shi, G., Hemming, S.R., Zhang, H., Wang, W., Wang, B., Chang, S.C., 2019. Age constraint on a Neogene tropical rainforest in China and its relation to the Middle Miocene Climatic Optimum. *Palaeogeography, Palaeoclimatology, Palaeoecology* 518, 82–88.

Figure captions

Fig. 1. *Parascleroderma palaeosinica* n. sp., holotype, NIGP178180; (A) habitus in lateroventral view; (B) habitus in dorsal view; (C) head in dorsal view; (D) line drawing of head and mesosoma in dorsal view; scale bar = 0.5 mm.

Fig. 2. Line drawing of *Parascleroderma palaeosinica* n. sp. in lateroventral view; scale bar = 0.5 mm.

Fig. 3. Distribution map of extant and extinct species of *Parascleroderma* showing type localities.

