



HAL
open science

A charging model for the Rosetta spacecraft

Fredrik Leffe Johansson, Anders Eriksson, Nicolas Gilet, Pierre Henri, Gaëtan Wattieaux, Matt Taylor, Christian Imhof, Fabrice Cipriani

► **To cite this version:**

Fredrik Leffe Johansson, Anders Eriksson, Nicolas Gilet, Pierre Henri, Gaëtan Wattieaux, et al.. A charging model for the Rosetta spacecraft. vEGU21, 2021, Online, France. 10.5194/egusphere-egu21-9508 . insu-03559304

HAL Id: insu-03559304

<https://insu.hal.science/insu-03559304v1>

Submitted on 7 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

EGU21-9508

<https://doi.org/10.5194/egusphere-egu21-9508>

EGU General Assembly 2021

© Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.



A charging model for the Rosetta spacecraft

Fredrik Leffe Johansson¹, Anders Eriksson¹, Nicolas Gilet², Pierre Henri^{2,3}, Gaëtan Wattieaux⁴, Matt Taylor⁵, Christian Imhof⁶, and Fabrice Cipriani⁵

¹Swedish Institute of Space Physics, Uppsala, Uppsala, Sweden (frejon@irfu.se)

²LPC2E, CNRS, Orléans, France

³Laboratoire Lagrange, OCA, CNRS, Nice, France

⁴University Paul Sabatier Toulouse III, Toulouse, France

⁵ESA/ESTEC, Noordwijk, The Netherlands

⁶Airbus Defence and Space GmbH, Friedrichshafen, Germany

Context. The electrostatic potential of a spacecraft, V_S , is important for the capabilities of in situ plasma measurements. Rosetta has been found to be negatively charged during most of the comet mission and even more so in denser plasmas.

Aims. Our goal is to investigate how the negative V_S correlates with electron density and temperature and to understand the physics of the observed correlation.

Methods. We applied full mission comparative statistics of V_S , electron temperature, and electron density to establish V_S dependence on cold and warm plasma density and electron temperature. We also used Spacecraft-Plasma Interaction System (SPIS) simulations and an analytical vacuum model to investigate if positively biased elements covering a fraction of the solar array surface can explain the observed correlations.

Results. Here, the V_S was found to depend more on electron density, particularly with regard to the cold part of the electrons, and less on electron temperature than was expected for the high flux of thermal (cometary) ionospheric electrons. This behaviour was reproduced by an analytical model which is consistent with numerical simulations.

Conclusions. Rosetta is negatively driven mainly by positively biased elements on the borders of the front side of the solar panels as these can efficiently collect cold plasma electrons. Biased elements distributed elsewhere on the front side of the panels are less efficient at collecting electrons apart from locally produced electrons (photoelectrons). To avoid significant charging, future spacecraft may minimise the area of exposed bias conductors or use a positive ground power system.

