



The Radio and Plasma Waves (RPW) Instrument on Solar Orbiter : Capabilities, Performance and First results



M. Maksimovic, J. Soucek, S. D. Bale, X. Bonnin, T. Chust,
Y. Khotyaintsev, M. Kretzschmar, D. Plettemeier, M. Steller,
Š. Štverák and the RPW Team






EGU 2020



Laboratoire d'Études Spatiales et d'Instrumentation en Astrophysique





Measuring electric & magnetic fields from DC to 16 MHz



- Electric & Magnetic Turbulence up to the e- scales
- Plasma waves up to Fp (k-vector, polarisation)
- Shocks & discontinuities

- Continuously operating (~5 kBps)
- On-Board shock & in-situ Type III detections (waveforms up to 3 kHz)
- Continuous 16 Hz Efield & S/C potential waveforms
- Burst modes (10 mins per day) with waveforms up to 1 kHz



- Quasi-Thermal Noise (Ne & Te) in dense & cold plasmas
- Electrostatic Langmuir waves
- Dusts



- Solar Radio emissions
- Interplanetary shocks (Type II)
- Strong Synergy with STIX, EUI, METIS, Solo HI




$$+X_{SCM} = \begin{bmatrix} 0.50 \\ 0.74 \\ -0.44 \end{bmatrix}$$



$$+Y_{SCM} = \begin{bmatrix} 0.60 \\ -0.67 \\ -0.44 \end{bmatrix}$$

$$+Z_{SCM} = \begin{bmatrix} -0.62 \\ -0.04 \\ -0.78 \end{bmatrix}$$







Instrument Boom deployment




MY antenna deployment: TDS MAMP versus platform data



Data shifted by 70 seconds



Detail of antenna deployment (shifted by 70 sec)




February 13, 2020



RPW-BIAS: IP shock





- Probe-to-spacecraft potential – proxy for plasma density



Courtesy Y. Khotyaintsev

LFR snapshots : Courtesy T. Chust

L1




For all SWFs one has AC channels:


E1 = V12_AC
E2 = V23_AC

Courtesy J. Soucek

Interplanetary Dust





Langmuir Waves





Ant 1

Ant 2


Ant 3




Possibly first
whistlers
observations
Courtesy M.
Kretzschmar



Solar Orbiter






WIND @ L1




Very first QTN fit attempt (M. Martinović, K. Issautier)

- $T_e = 10^5 K$
- Two Maxwellians populations (core and halo)
- possibly effect of the « hole » between the monopoles)






We still see weak Type IIIs « in between spurious lines »



NDA, courtesy L. Lamy



Courtesy
A. Vecchio

Conclusions

- **The RPW instrument is operating extremely well ... but sitting on a very noisy spacecraft**
- **Many S/C (and possibly instrument) EMC perturbations, especially for the radio receiver**
- **First Langmuir waves and dust measurements**
- **Probably first shock & whistlers waves**
- **The THR can still see Quasi-Thermal Noise from time to time and Type IIIs for very few frequencies not completely corrupted. Cleaning of data is necessary**
- **Data available for the CoIs for the time being**
- **Hope to be ready for the first public delivery of cruise phase data**