



HAL
open science

2.5 years of TROPOMI S5P total ozone column data: geophysical global ground-based validation and inter-comparison with other satellite missions

Katerina Garane, Maria-Elissavet Koukouli, Tijn Verhoelst, Christophe Lerot,
Klaus-Peter Heue, Dimitrios Balis, Alberto Redondas, Andrea Pazmino,
Ariane Bazureau, Fabian Romahn, et al.

► To cite this version:

Katerina Garane, Maria-Elissavet Koukouli, Tijn Verhoelst, Christophe Lerot, Klaus-Peter Heue, et al.. 2.5 years of TROPOMI S5P total ozone column data: geophysical global ground-based validation and inter-comparison with other satellite missions. EGU General Assembly 2020, 2020, Online, France. pp. EGU2020-8109, 10.5194/egusphere-egu2020-8109 . insu-03562306

HAL Id: insu-03562306

<https://insu.hal.science/insu-03562306>

Submitted on 8 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

EGU2020-8109

<https://doi.org/10.5194/egusphere-egu2020-8109>

EGU General Assembly 2020

© Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.



2.5 years of TROPOMI S5P total ozone column data: geophysical global ground-based validation and inter-comparison with other satellite missions

Katerina Garane¹, **Maria-Elissavet Koukoulis**¹, Tijn Verhoelst², Christophe Lerot², Klaus-Peter Heue³, Dimitrios Balis¹, Alberto Redondas⁴, Andrea Pazmino⁵, Ariane Bazureau⁵, Fabian Romahn³, Walter Zimmer³, Jian Xu³, Jean-Christopher Lambert², Diego Loyola³, Michel Van Roozendael², Florence Goutail⁵, and Jean-Pierre Pommereau⁵

¹Laboratory of Atmospheric Physics, Aristotle University of Thessaloniki, Thessaloniki, Greece (agarane@auth.gr)

²Royal Belgian Institute for Space Aeronomy (BIRA-IASB), Uccle, Belgium

³German Aerospace Centre, Remote Sensing Institute (DLR-IMF), Oberpfaffenhofen, Germany

⁴Izaña Atmospheric Research Center (IARC), State Meteorological Agency (AEMET), Tenerife, Canary Islands, Spain

⁵LATMOS, CNRS, University Versailles St Quentin, Guyancourt, France

The Sentinel-5 Precursor (S5P) mission, launched in October 2017, carries the TROPOspheric Monitoring Instrument (TROPOMI), which provides a daily global coverage at a spatial resolution as high as 5.5 km x 3.5 km and will extend the European atmospheric composition record initiated with GOME/ERS-2 in 1995. Due to the ongoing need to understand and monitor the recovery of the ozone layer, as well as the evolution of tropospheric pollution, ozone remains one of the leading species of interest during this mission.

In this work, two and a half years of TROPOMI near real time (NRTI) and offline (OFFL) total ozone column (TOC) products are presented and compared to daily and individual, globally distributed, ground-based quality assured Brewer and Dobson TOC measurements. The daily ground-based ozone measurements used here are deposited in the World Ozone and Ultraviolet Radiation Data Centre (WOUDC). The individual Brewer measurements are made available by the European Brewer Network (Eubrewnet). Furthermore, twilight zenith-sky measurements obtained with ZSL-DOAS (Zenith Scattered Light Differential Optical Absorption Spectroscopy) instruments, which form part of the SAOZ network (Système d'Analyse par Observation Zénitale), are used for the validation.

The quality of the TROPOMI TOC data is evaluated in terms of the influence of various geophysical quantities such as location, solar zenith angle, viewing angle, season, effective temperature, surface albedo and clouds. The overall statistical analysis of the global comparison shows that the mean bias and the mean standard deviation of the percentage difference between TROPOMI and ground-based TOC is within 0 –1.5% and 2.5 %–4.5 %, respectively. Moreover, based on the full available dataset, a first attempt is made for a drift investigation.

Additionally, the TROPOMI OFFL and NRTI products are evaluated against already known spaceborne sensors, namely, the Ozone Mapping Profiler Suite, on board the Suomi National Polar-orbiting Partnership (OMPS/Suomi-NPP), NASA, and the Global Ozone Monitoring Experiment 2 (GOME-2), on board the Metop-A (GOME-2/Metop-A) and Metop-B (GOME-2/Metop-B) satellites. This analysis shows a very good agreement for both TROPOMI products with well-established instruments, with the absolute differences in mean bias and mean standard deviation being below +0.7% and 1%, respectively.