



Trace gas profiles and anthropogenic plumes from metropolitan areas in West Africa during DACCIWA - Airborne measurements on board the DLR Falcon 20

Greta Stratmann, Hans Schlager, Daniel Sauer, Vanessa Brocchi, Valéry Catoire, Robert Baumann

► To cite this version:

Greta Stratmann, Hans Schlager, Daniel Sauer, Vanessa Brocchi, Valéry Catoire, et al.. Trace gas profiles and anthropogenic plumes from metropolitan areas in West Africa during DACCIWA - Airborne measurements on board the DLR Falcon 20. 20th EGU General Assembly, Apr 2018, Vienne, Austria. pp.19210. insu-03566317

HAL Id: insu-03566317

<https://insu.hal.science/insu-03566317>

Submitted on 11 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Trace gas profiles and anthropogenic plumes from metropolitan areas in West Africa during DACCIWA - Airborne measurements on board the DLR Falcon 20

Greta Stratmann (1), Hans Schlager (1), Daniel Sauer (1), Vanessa Brocchi (2), Valéry Catoire (2), and Robert Baumann (1)

(1) German Aerospace Center (DLR), Institute of Atmospheric Physics (IPA) , Germany (greta.stratmann@dlr.de), (2) 2 Laboratoire de Physique et Chimie de l'Environnement et de l'Espace (LPC2E), CNRS – Université Orléans, 45071 Orléans cedex 2, France

The DACCIWA (Dynamics-Aerosol-Chemistry-Cloud Interactions over West Africa) airborne field campaign was conducted in Southern West Africa in June/July 2016. The atmosphere in South West Africa is exposed to numerous influences like the Monsoon, biomass burning and oil rigs. Its chemical composition is modified by urbanisation, a growing population, complex meteorological influences and air pollution. During DACCIWA, three European research aircraft (DLR - Falcon 20, SAFIRE - ATR 42 and BAS - Twin Otter) were deployed from Lomé/Togo and conducted research flights across Ivory Coast, Ghana, Togo and Benin. On board the DLR Falcon O₃, SO₂, CO, NO₂ and aerosol fine mode particle number concentration and size distribution were measured during a total of 12 scientific flights.

Distinct trace gas layers were found during vertical soundings. An O₃ layer at ~ 3000 m altitude was observed during 6 flights with concentrations up to 85 ppb compared to background concentrations of about 35 ppb due to southern hemispheric biomass burning products which were transported to the DACCIWA study area. Furthermore, enhanced trace gas and particle number concentrations were detected in city pollution plumes of Lomé/Togo, Accra/Ghana and Kumasi/Ghana. O₃ enhancements of 5 – 10 ppb were measured downstream of the cities relative to the upstream observations. Measured SO₂ pollution plumes with concentrations up to 0.9 ppb near Lomé are combined with HYPSLIT dispersion simulations and are compared to emission databases.