



**HAL**  
open science

## Effect of Blue Jets on Stratospheric Ozone - a model study

Chen Xu, Nathalie Huret, Kevin M. A. Ihaddadene, Sebastien Celestin

► **To cite this version:**

Chen Xu, Nathalie Huret, Kevin M. A. Ihaddadene, Sebastien Celestin. Effect of Blue Jets on Stratospheric Ozone - a model study. 20th EGU General Assembly, Apr 2018, Vienne, Austria. pp.14678. insu-03566323

**HAL Id: insu-03566323**

**<https://insu.hal.science/insu-03566323>**

Submitted on 11 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution| 4.0 International License



## **Effect of Blue Jets on Stratospheric Ozone - a model study**

Chen Xu (1), Nathalie Huret (1), Kevin M.A. Ihaddadene (2), and Sebastien Celestin (1)

(1) LPC2E, University of Orleans, CNRS, Orleans, France, (2) Department of Physics, Space Science Center, University of New Hampshire, Durham, USA

In the framework of the TARANIS space mission dedicated to the study of transient luminous events, a detailed ion-neutral chemistry model has been developed to simulate the impact of blue jet streamers on stratospheric chemistry. It is based on the MIPLASMO model (Microphysical and Photochemical Lagrangian Model of Ozone) widely used over the last 20 years to interpret balloon and satellite measurements associated with stratospheric ozone. In this study, instead of using a constant reduced electric field value during a streamer pulse time, we used a time-dependent reduced electric field coming from an explicit streamer model. The simulations are performed on day-time/night-time and their duration is two days. Among 117 species in the model, we put the focus on nitrogen, oxygen, chlorine and bromine species, and ozone perturbations. The model results indicate that the impact on both ions and neutral species volume mixing ratios considering a quasi-real streamer (i.e. using a time-varying electric field) is different from previous model studies considering streamer as a simple pulse. These differences will be analyzed in detail as a function of altitude. Validation exercise and preliminary results will be presented.