

Hyperpectral infrared imaging of volcanic plume at Mt Etna during IMAGETNA campaign

Charlotte Segonne, Nathalie Huret, Sébastien Payan, Valéry Catoire, Giuseppe Salerno, Tjarda J. Roberts, Alessandro La Spina, Tommaso Caltabiano

▶ To cite this version:

Charlotte Segonne, Nathalie Huret, Sébastien Payan, Valéry Catoire, Giuseppe Salerno, et al.. Hyperpectral infrared imaging of volcanic plume at Mt Etna during IMAGETNA campaign. 20th EGU General Assembly, Apr 2018, Vienne, Austria. pp.12927. insu-03566325

HAL Id: insu-03566325 https://insu.hal.science/insu-03566325

Submitted on 11 Feb2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Hyperpectral infrared imaging of volcanic plume at Mt Etna during IMAGETNA campaign

Charlotte Segonne (1), Nathalie Huret (1), Sébastien Payan (2), Valéry Catoire (1), Giuseppe Salerno (3), Tjarda Roberts (1), Alessandro La Spina (3), and Tommaso Caltabiano (3)

(1) Orléans University and LPC2E/CNRS and, 3A avenue de la recherche scientifique,F4500 Orléans (charlotte.segonne@cnrs-orleans.fr), (2) LATMOS, CNRS, Sorbonne Université, France, (3) Istituto Nazionale di Geofisica e Vulcanologia, Osservatorio Etneo, Catania, Italy

Knowledge of the composition and the spatial evolution of volcanic plumes provides insights to processes occurring in the Earth's interior. On the other hand, quantification of gaseous emission fluxes is also a fundamental task in the framework of climate change in order to refine the contribution of natural emissions. UV cameras allow us to image volcanic plumes and evaluate SO_2 fluxes, although can be subject to uncertainties in the retrieval. Another technique of imaging is now available in the infra-red. Such infrared hyperspectral imager (pixel-by-pixel spectra) might represent a major step forward in volcanology due to its potential to allow SO_2 flux measurements during the night and gives access to additional relevant species but has to be tested and validated as a first step. In June 2015 a campaign of measurements - IMAGETNA – was performed at Mt Etna (Pizzi Deneri Volcano Observatory) with the intent to explore the application of these techniques for volcanic gas measurements all together. Over five days the volcanic plume was remotely observed simultaneously by employing three different hyperspectral imagers (commercial and under development), FTIR instrument, UV LWIR cameras, and radiometer. Results gathered from IR hyperspectral imageur will be presented and compared to SO_2 UV-Vis camera measurements.