



HAL
open science

Are Supermassive Hot Jupiters Providing More Favourable Conditions for Generation of Radio Emission via the Cyclotron Maser Instability? - A Case Study based on Tau Bootis b

Christof Weber, Nikolai V. Erkaev, Petra Odert, Jean-Mathias Griessmeier, Helmut Lammer, Helmut O. Rucker

► To cite this version:

Christof Weber, Nikolai V. Erkaev, Petra Odert, Jean-Mathias Griessmeier, Helmut Lammer, et al.. Are Supermassive Hot Jupiters Providing More Favourable Conditions for Generation of Radio Emission via the Cyclotron Maser Instability? - A Case Study based on Tau Bootis b. 20th EGU General Assembly, 2018, Vienne, Austria. pp.8411. insu-03566333

HAL Id: insu-03566333

<https://insu.hal.science/insu-03566333>

Submitted on 11 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution| 4.0 International License



Are Supermassive Hot Jupiters Providing More Favourable Conditions for Generation of Radio Emission via the Cyclotron Maser Instability? - A Case Study based on Tau Bootis b

Christof Weber (1,2), Nikolai V. Erkaev (3,4), Petra Odert (2), Jean-Mathias Grießmeier (5,6), Helmut Lammer (2), and Helmut O. Rucker (7)

(1) Institute of Atmospheric Physics, Czech Academy of Sciences, Prague, Czech Republic (cw@ufa.cas.cz), (2) Space Research Institute, Graz, Graz, Austria (christof.weber@oeaw.ac.at), (3) Institute for Computational Modelling SB RAS, 660036, Krasnoyarsk, Russian Federation, (4) Siberian Federal University, Krasnoyarsk, Russian Federation, (5) LPC2E - Université d'Orléans/CNRS, France, (6) Station de Radioastronomie de Nançay, Observatoire de Paris, PSL Research University, CNRS/INSU, USR 704 - Univ. Orléans, OSUC, route de Souesmes, 18330 Nançay, France, (7) Comm. Astron. ÖAW, Graz, Austria

We check if supermassive Hot Jupiters can maintain source regions for radio emission, and whether the emission can propagate to an observer outside the system. Planets identical to the Hot Jupiter Tau Bootis b are investigated, but located at different orbital distances, starting at the actual orbit of 0.046 AU up to 0.2 AU. We find that due to the strong gravity of the supermassive planet Tau Bootis b additional coolings would become very effective. Including these cooling effects, the exobase is very close to the planet, which makes Tau Bootis b a good candidate for radio observations. The case of hydrodynamic escape without cooling effects is considered as well for comparison and we check at which orbital distance the upper atmosphere is in the hydrodynamic rather than in the hydrostatic regime, i.e. where conditions for the Cyclotron Maser Instability (CMI) are favourable. In the hydrodynamic regime the ionosphere is extended up to the magnetopause and can constitute an obstacle for possibly generated radio waves, or the generation via the CMI might even be prevented completely. We find that the transition to more favourable conditions for the CMI occurs at 0.056 AU, where the exobase distance is already close to the magnetosphere standoff distance. Another interesting finding is that, even though the exobase is more extended for smaller orbits, the stronger stellar wind of the young star Tau Bootis causes a higher outflow velocity for the planetary mass loss due to the higher XUV flux. This effect lowers the electron density peak for the actual orbit of Tau Bootis b compared to the larger orbits and could thus make generation and escape of radio waves possible. In a further step of this investigation the star Tau Bootis is replaced by a Sun-like star. In this case the highest electron density peak is found for the closest orbit.