



A fully kinetic perspective of electron acceleration around a weakly outgassing comet: Ohm's law

Andrey Divin, Pierre Henri, Anders Eriksson, Vyacheslav Olshevsky, Stefano Markidis, Mihály Horányi

► To cite this version:

Andrey Divin, Pierre Henri, Anders Eriksson, Vyacheslav Olshevsky, Stefano Markidis, et al.. A fully kinetic perspective of electron acceleration around a weakly outgassing comet: Ohm's law. 20th EGU General Assembly, 2018, Vienne, Austria. pp.6571. insu-03566334

HAL Id: insu-03566334

<https://insu.hal.science/insu-03566334>

Submitted on 11 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



A fully kinetic perspective of electron acceleration around a weakly outgassing comet: Ohm's law

Jan Deca (1,2,3), Andrey Divin (4,5), Pierre Henri (6), Anders Eriksson (5), Vyacheslav Olshevsky (7), Stefano Markidis (8), Mihály Horányi (1,2)

(1) University of Colorado Boulder, Laboratory for Atmospheric and Space Physics, Boulder, United States (jandeca@gmail.com), (2) Institute for Modeling Plasma, Atmospheres and Cosmic Dust, NASA/SSERVI, USA, (3) Laboratoire Atmosphères, Milieux, Observations Spatiales, Université de Versailles à Saint Quentin, Guyancourt, France, (4) St. Petersburg State University, St. Petersburg, Russia, (5) Swedish Institute of Space Physics, Uppsala, Sweden, (6) LPC2E, CNRS, Orléans, France, (7) Centre for mathematical Plasma Astrophysics, Department of Mathematics, KU Leuven, Leuven, Belgium, (8) Computational Science and Technology, KTH Royal Institute of Technology, Stockholm, Sweden

When a comet is sufficiently close to the Sun, the sublimation of ice leads to outgassing and the formation of a coma of gas and dust. Ionisation of the outgassing material then results in mass-loading of the solar wind and magnetic field draping around the cometary nucleus. Here we present three-dimensional fully kinetic simulations of the solar wind interaction with comet 67P/Churyumov-Gerasimenko at a low-activity regime, before collisions have any impact on the plasma dynamics. The interaction scales are well below the relevant ion gyroradii. Non-equilibrium electron distributions develop. To first order, the dynamical interaction is representative of a four-fluid coupled system [Deca et al., PRL 2017, Divin et al., submitted]. Our approach is self-consistent and allows to distill Ohm's law directly from the electron dynamics in the simulation, rather than imposing it beforehand.

In the vicinity of the cometary nucleus, the balance changes between the different terms in the equation. Deciphering the relative importance of each term allows to identify the driving physics in the various regions of the cometary plasma environment. For example, we find that close to the outgassing nucleus the electron pressure gradient dominates; that at sub-ion scales, the total electric field is a superposition of the solar wind convective electric field, where electrons are frozen-in, and the ambipolar electric field; that the latter accelerates electrons parallel to the magnetic field and is the source of/provides feedback to the electron pressure gradient that balances the Ohm's law perpendicular to the magnetic field, and that the role of electron inertia is negligible to balance the electric field.

In conclusion, Ohm's law shows us what happens to all ion and electron species, and why.