

## Links between the plasmapause and the radiation belts boundaries as observed by Cluster instruments

Fabien Darrouzet, Viviane Pierrard, Sylvie Benck, Pierrette Décréau, Natalia

Ganushkina, Johan de Keyser

## ▶ To cite this version:

Fabien Darrouzet, Viviane Pierrard, Sylvie Benck, Pierrette Décréau, Natalia Ganushkina, et al.. Links between the plasmapause and the radiation belts boundaries as observed by Cluster instruments. 20th EGU General Assembly, Apr 2018, Vienne, Austria. pp.6269. insu-03566335

## HAL Id: insu-03566335 https://insu.hal.science/insu-03566335

Submitted on 11 Feb2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



## Links between the plasmapause and the radiation belts boundaries as observed by Cluster instruments

Fabien Darrouzet (1), Viviane Pierrard (1), Sylvie Benck (2), Pierrette Décréau (3), Natalia Ganushkina (4), and Johan De Keyser (1)

(1) Royal Belgian Institute for Space Aeronomy (IASB-BIRA), Brussels, Belgium (Fabien.Darrouzet@aeronomie.be), (2) Center for Space Radiations (CSR), Louvain-la-Neuve, Belgium, (3) Laboratoire de Physique et Chimie de l'Environnement et de l'Espace (LPC2E), Orléans, France, (4) Finnish Meteorological Institute (FMI), Helsinki, Finland

The Cluster mission offers the exceptional opportunity to study the relations between the position of the plasmapause and the position of the radiation belt boundaries with identical sensors on multiple spacecraft. We compare the positions of the radiation belt edges deduced from CIS observations (electrons with energy larger than 2 MeV) with the positions of the plasmapause derived from WHISPER data (electron plasma frequency). In addition, we compare those results with the boundary positions determined from RAPID observations (electrons with energy between 244.1 and 406.5 keV).

The period of 1 April 2007 to 31 March 2009 has been chosen for the analysis because at that time Cluster's perigee was located at lower radial distances than during the earlier part of the mission (as close as 2 Re, deep inside the plasmasphere and the radiation belts). This time period corresponds to a long solar activity minimum.

Differences are observed between the radiation belt boundary positions obtained from the two different instruments: The radiation belt positions are related to the energy bands. The plasmapause position is more variable than the radiation belt boundary positions, especially during small geomagnetic activity enhancements. A correspondence is observed between the plasmapause position determined by WHISPER and the outer edge of the outer radiation belt of energetic electrons (> 2 MeV) observed by CIS. There may be an apparent contradiction with previous studies that indicated a correlation between the inner edge of the outer radiation belt and the plasmapause. Radiation belt losses through plasmaspheric waves, however, occur on a longer time scale and are longitudinally averaged because of energetic particle drift motion, so the radiation belt boundaries should not necessarily reflect the instantaneous plasmapause position at one particular local time. Moreover, during higher geomagnetic activity time periods, the plasmapause is located closer to the inner boundary of the outer radiation belt.