



# Controlled meteorological (CMET) balloon profiling of the Arctic atmospheric boundary layer

Tjarda J Roberts, Lars Hole, Paul Voss

## ► To cite this version:

Tjarda J Roberts, Lars Hole, Paul Voss. Controlled meteorological (CMET) balloon profiling of the Arctic atmospheric boundary layer. 19th EGU General Assembly, Apr 2017, Vienne, Austria. pp.13431. insu-03568130

**HAL Id: insu-03568130**

**<https://insu.hal.science/insu-03568130>**

Submitted on 13 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

## **Controlled meteorological (CMET) balloon profiling of the Arctic atmospheric boundary layer**

Tjarda Roberts (1), Lars Hole (2), and Paul Voss (3)

(1) CNRS/INSU, LPC2E/Université d'Orléans, Orléans, France (TjardaRoberts@gmail.com), (2) Norwegian Meteorological Institute, Bergen, Norway, (3) Smith College, Picker Engineering Program, Northampton, MA, USA

We demonstrate profiling of the atmospheric boundary layer over Arctic ice-free and sea-ice covered regions by free-floating controllable CMET balloons. The CMET observations (temperature, humidity, wind-speed, pressure) provide in-situ meteorological datasets in very remote regions for comparison to atmospheric models.

Controlled Meteorological (CMET) balloons are small airborne platforms that use reversible lift-gas compression to regulate altitude. These balloons have approximately the same payload mass as standard weather balloons but can float for many days, change altitude on command, and transmit meteorological and system data in near-real time via satellite.

Five Controlled Meteorological (CMET) balloons were launched from Ny-Ålesund in Svalbard (Spitsbergen) over 5–12 May 2011 and measured vertical atmospheric profiles (temperature, humidity, wind) over coastal and remote areas to both the east and west. One notable CMET flight achieved a suite of 18 continuous soundings that probed the Arctic atmospheric boundary layer (ABL) over a period of more than 10 h. Profiles from two CMET flights are compared to model output from ECMWF Era-Interim reanalysis (ERA-I) and to a high-resolution (15 km) Arctic System Reanalysis (ASR) product. To the east of Svalbard over sea-ice, the CMET observed a stable ABL profile with a temperature inversion that was reproduced by ASR but not captured by ERA-I. In a coastal ice-free region to the west of Svalbard, the CMET observed a stable ABL with strong wind-shear. The CMET profiles document increases in ABL temperature and humidity that are broadly reproduced by both ASR and ERA-I. The ASR finds a more stably stratified ABL than observed but captured the wind shear in contrast to ERA-I. Detailed analysis of the coastal CMET-automated soundings identifies small-scale temperature and humidity variations with a low-level flow and provides an estimate of local wind fields.

We show that CMET balloons are a valuable approach for profiling the free atmosphere and atmospheric boundary layer in remote regions such as the Arctic, where few other in-situ observations are available to trace processes and for model evaluation.

### **References:**

Roberts, T. J., Dütsch, M., Hole, L. R., and Voss, P. B.: Controlled meteorological (CMET) free balloon profiling of the Arctic atmospheric boundary layer around Spitsbergen compared to ERA-Interim and Arctic System Reanalyses. *Atmos. Chem. Phys.*, 16, 12383-12396, doi:10.5194/acp-16-12383-2016, 2016.

Hole L. R., Bello A. P., Roberts T. J., Voss P. B., Vihma T.: Measurements by controlled meteorological balloons in coastal areas of Antarctica. *Antarctic Science*, 1-8, doi:10.1017/S0954102016000213, 2016.

Voss P. B., Hole L. R., Helbling E. F., Roberts T. J.: Continuous in-situ soundings in the arctic boundary layer: a new atmospheric measurement technique using controlled meteorological balloons. *Journal of Intelligent Robot Systems*, 70, 609-617, doi 10.1007/s10846-012-9758-6, 2013.