



# Modeling Thunderstorm Ground Enhancements and Gamma-ray Glows

Eric Cramer, Bagrat Mailyan, Sebastien Celestin, Joseph Dwyer

## ► To cite this version:

Eric Cramer, Bagrat Mailyan, Sebastien Celestin, Joseph Dwyer. Modeling Thunderstorm Ground Enhancements and Gamma-ray Glows. 19th EGU General Assembly, Apr 2017, Vienne, Austria. pp.11532. insu-03568141

**HAL Id: insu-03568141**

**<https://insu.hal.science/insu-03568141>**

Submitted on 13 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

## **Modeling Thunderstorm Ground Enhancements and Gamma-ray Glows**

Eric Cramer (1), Bagrat Mailyan (1,2), Sebastien Celestin (3), and Joseph Dwyer (4)

(1) Center for Space Plasma & Aeronomic Research, The University of Alabama in Huntsville, Huntsville, United States (Eric.Cramer@uah.edu), (2) Cosmic Ray Division, Yerevan Physics Institute, Armenia (Bagrat.Mailyan@uah.edu), (3) LPC2E, University of Orleans, CNRS, Orleans, France (sebastien.celestin@cnrs-orleans.fr), (4) Department of Physics, University of New Hampshire, Durham, United States (Joseph.Dwyer@unh.edu)

Energetic radiation from thunderstorms is currently being measured by ground-based particle detectors worldwide, e.g. Torri et al., JGR, 107, 2002; Khaerdinov et al., Atmospheric Research, 76, 2005; Chilingarian et al., Phys. Rev. D, 82, 2010; and Tsuchiya et al., JGR, 116, 2011. Additionally, long bursts of radiation lasting more than one second (gamma-ray glows), have been observed from aircraft and balloons, e.g. McCarthy and Parks, GRL, 12, 1985; Eack et al., GRL, 27, 2000; Smith et al., JGR, 116, 2011; and Kelley et al., Nature Communications, 6, 2015. The Relativistic Runaway Electron Avalanche (RREA) model explains that the observed particle fluxes are due to electron acceleration and consequent gamma-ray emission in the large-scale atmospheric electric fields [Dwyer, GRL, 30, 2003].

Since the primary RREA electrons rapidly attenuate in the atmosphere, most of the information on the nature of the electric field inside the thundercloud can come from detection of the emitted gamma rays. In this work, we use the Relativistic Runaway Electron Avalanche Model (REAM) developed by Dwyer, GRL, 30, 2003, to model various layouts of the electric field configuration and study the predicted photon spectra. The comparison of the electron avalanche length calculated with REAM and with that of the model developed by Celestin and Pasko, JGR, 116, 2011, is also discussed.