



## Rosetta spacecraft potential and electron temperature in the coma of comet 67P

Elias Odelstad, Gabriella Stenberg-Wieser, Martin Wieser, Anders Eriksson, Hans Nilsson, Pierre Henri

### ► To cite this version:

Elias Odelstad, Gabriella Stenberg-Wieser, Martin Wieser, Anders Eriksson, Hans Nilsson, et al.. Rosetta spacecraft potential and electron temperature in the coma of comet 67P. 19th EGU General Assembly, Apr 2017, Vienne, Austria. pp.9258. insu-03568143

**HAL Id: insu-03568143**

**<https://insu.hal.science/insu-03568143>**

Submitted on 13 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

## **Rosetta spacecraft potential and electron temperature in the coma of comet 67P**

Elias Odelstad (1,2), Gabriella Stenberg-Wieser (3), Martin Wieser (3), Anders Eriksson (1), Hans Nilsson (3), and Pierre Henri (4)

(1) Swedish Institute of Space Physics, Uppsala, Sweden (elias.odelstad@irfu.se), (2) Department of Physics and Astronomy, Uppsala University, Uppsala, Sweden, (3) Swedish Institute of Space Physics, Kiruna, Sweden, (4) LPC2E, Orléans, France

We present and compare measurements of the spacecraft potential ( $V_{S/C}$ ) of ESA's Rosetta spacecraft during its  $\sim 2$  year stay in the inner coma of comet 67P/Churyumov-Gerasimenko, by the Rosetta Langmuir probe (RPC-LAP) and Ion Composition Analyzer (RPC-ICA) instruments.  $V_{S/C}$  was mostly negative, driven so by the high ( $\sim 5$ - $10$  eV) electron temperature of the coma photoelectrons. LAP only picks up a portion of the full  $V_{S/C}$  since the two probes, mounted on booms of length 2.2 and 1.6 m, respectively, are generally inside the potential field of the negatively charged spacecraft. Comparison of the minimum energy of positive ions collected by ICA shows that this portion varies between 0.7 and 1  $V_{S/C}$ , with generally good correspondence between the two instruments except when local ion production is weak and accelerated ions dominate the flux. We map  $V_{S/C}$  and its evolution during the mission and use it to constrain the electron density  $n_e$  and temperature  $T_e$  in the inner coma. Comparison of  $V_{S/C}$  measurements to electron density measurements by the Mutual Impedance Probe (RPC-MIP) allows more accurate estimates of the electron temperature and a cross-calibration of the  $n_e$  and  $T_e$  measurements by the different instruments.