



## Halogen Chemistry in Volcanic Plumes (Invited)

Tjarda J Roberts

### ► To cite this version:

Tjarda J Roberts. Halogen Chemistry in Volcanic Plumes (Invited). 19th EGU General Assembly, Apr 2017, Vienne, Austria. pp.7578. insu-03568148

**HAL Id: insu-03568148**

**<https://insu.hal.science/insu-03568148>**

Submitted on 13 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

## Halogen Chemistry in Volcanic Plumes (Invited)

Tjarda Roberts

CNRS-INSU, LPC2E-Université d'Orléans, Orléans, France (Tjarda.Roberts@cnrs-orleans.fr)

Volcanoes release vast amounts of gases and particles in the atmosphere. Volcanic halogens (HF, HCl, HBr, HI) are co-emitted alongside SO<sub>2</sub>, and observations show rapid formation of BrO and OCIO in the plume as it disperses into the troposphere. The development of 1D and Box models (e.g. *PlumeChem*) that simulate volcanic plume halogen chemistry aims to characterise how volcanic reactive halogens form and quantify their atmospheric impacts.

Following recent advances, these models can broadly reproduce the observed downwind BrO/SO<sub>2</sub> ratios using “bromine-explosion” chemistry schemes, provided they use a “high-temperature initialisation” to inject radicals (OH, Cl, Br and possibly NO<sub>x</sub>) which “kick-start” the low-temperature chemistry cycles that convert HBr into reactive bromine (initially as Br<sub>2</sub>). The modelled rise in BrO/SO<sub>2</sub> and subsequent plateau/decline as the plume disperses downwind reflects cycling between reactive bromine, particularly Br-BrO, and BrO-HOBr-BrONO<sub>2</sub>. BrCl is produced when aerosol becomes HBr-depleted. Recent model simulations suggest this mechanism for reactive chlorine formation can broadly account for OCIO/SO<sub>2</sub> reported at Mt Etna.

Predicted impacts of volcanic reactive halogen chemistry include the formation of HNO<sub>3</sub> from NO<sub>x</sub> and depletion of ozone. This concurs with HNO<sub>3</sub> widely reported in volcanic plumes (although the source of NO<sub>x</sub> remains under question), as well as observations of ozone depletion reported in plumes from several volcanoes (Mt Redoubt, Mt Etna, Eyjafjallajökull). The plume chemistry can transform mercury into more easily deposited and potentially toxic forms, for which observations are limited. Recent incorporation of volcanic halogen chemistry in a 3D regional model of degassing from Ambrym (Vanuatu) also predicts how halogen chemistry causes depletion of OH to lengthen the SO<sub>2</sub> lifetime, and highlights the potential for halogen transport from the troposphere to the stratosphere.

However, the model parameter-space is vast and only partially constrained by available observations. Reactions on aerosol are a key driver of the chemistry and are affected by uncertainties in both the HOBr reactive uptake coefficient and the aerosol surface area. Recent work has explored the reactive uptake of HOBr on sulfate-rich aerosol, whilst field-measurements at Mt Etna have aimed to quantify the size-resolved primary aerosol emission, towards improving model representations of this highly non-linear volcanic plume halogen chemistry.