



**HAL**  
open science

# Effects of the geomagnetic field on the beaming geometry of TGFs

Sebastien Celestin

► **To cite this version:**

Sebastien Celestin. Effects of the geomagnetic field on the beaming geometry of TGFs. EGU General Assembly 2016, Apr 2016, Vienne, Austria. insu-03573508

**HAL Id: insu-03573508**

**<https://insu.hal.science/insu-03573508>**

Submitted on 15 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



## **Effects of the geomagnetic field on the beaming geometry of TGFs**

Sebastien Celestin

LPC2E, University of Orleans, CNRS, Orleans, France (sebastien.celestin@cnrs-orleans.fr)

Terrestrial gamma-ray flashes (TGFs) are bursts of high-energy photons originating from the Earth's atmosphere in association with thunderstorm activity [e.g., Briggs et al., JGR, 118, 3805, 2013]. Although TGFs are believed to be produced inside thunderclouds (below 15 km altitude), the underlying physical mechanisms are still debated. Large-scale relativistic runaway electron avalanches (RREAs) along with relativistic feedback caused by positrons and photons have been proposed to occur in thunderclouds and to produce TGFs [e.g., Dwyer et al., Space Sci. Rev., 173, 133, 2012]. It has also been found that the production of thermal runaway electrons by stepping lightning leaders and their further acceleration could explain the TGF spectra and fluences for intracloud (IC) lightning electric potentials above  $\sim 100$  MV [Xu et al., GRL, 39, L08801, 2012; Celestin et al., JGR, 120, 2015]. In both scenarios, runaway electron avalanches take place and the related bremsstrahlung produces the TGF.

The impact of the geomagnetic field on RREAs has been seldom studied (with the notable exceptions of Lehtinen et al. [JGR, 104, 24699, 1999], Babich et al. [Geom. Aeron., 44, 243, 2004] and Cramer et al. [AGU Fall Meeting, abstract AE33A-0472, San Francisco, USA, 2015]), particularly in view of recent knowledge acquired about TGF sources properties. In this work, we study the effects of the geomagnetic field on the runaway electron beam geometry in large-scale RREAs and in the vicinity of lightning leaders and the corresponding impact on TGF observations using analytical and numerical means.