

On the use of hyperpectral infrared imagers for studying volcano plumes: IMAGETNA campaign

Sébastien Payan, Nathalie Huret, Valery Catoire, Stéphane Langlois, Giuseppe Salerno, Tjarda J Roberts, Delphy Rodriguez, Armande Pola Fossi, Alessandro La Spina, Tomasso Caltabiano, et al.

▶ To cite this version:

Sébastien Payan, Nathalie Huret, Valery Catoire, Stéphane Langlois, Giuseppe Salerno, et al.. On the use of hyperpectral infrared imagers for studying volcano plumes: IMAGETNA campaign. EGU General Assembly 2016, Apr 2016, Vienne, Austria. insu-03573533

HAL Id: insu-03573533 https://insu.hal.science/insu-03573533

Submitted on 14 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Geophysical Research Abstracts Vol. 18, EGU2016-4877, 2016 EGU General Assembly 2016 © Author(s) 2016. CC Attribution 3.0 License.



On the use of hyperpectral infrared imagers for studying volcano plumes: IMAGETNA campaign

sebastien payan (1), Nathalie Huret (2), Valery Catoire (2), Stéphane Langlois (3), Giuseppe Salerno (4), Tjarda Roberts (2), Delphy Rodriguez (1), Armande Pola Fossi (3), Alessandro La Spina (4), Tomasso Caltabiano (4), Yann Ferrec (3), Stépnane Chevrier (2), and Mike Burton (5)

(1) LATMOS, UPMC Case 102, Paris cedex 05, France (sebastien.payan@upmc.fr), (2) Orléans University and LPC2E/CNRS and, 3A avenue de la recherche scientifique,F4500 Orléans, (3) ONERA French Aerospace Lab, Palaiseau, France, (4) Istituto Nazionale di Geofisica e Vulcanologia, Osservatorio Etneo, Catania, Italy, (5) University of Manchester

Knowledge of the composition and the spatial evolution of volcanic plumes provides insights to processes occurring in the Earth's interior. On the other hand, quantification of gaseous emission fluxes is also a fundamental task in the framework of climate change in order to refine the contribution of natural emissions. UV cameras allow us to image volcanic plumes and evaluate SO₂ fluxes, although can be subject to uncertainties in the retrieval. Another technique of imaging is now available in the infra-red. Such infrared hyperspectral imager (pixel-by-pixel spectra) might represent a major step forward in volcanology due to its potential to allow SO₂ flux measurements during the night and gives access to additional relevant species but has to be tested and validated as a first step. In June 2015 a campaign of measurements - IMAGETNA - was performed at Mt Etna (Pizzi Deneri Volcano Observatory) with the intent to explore the application of these techniques for volcanic gas measurements all together. Over five days the volcanic plume was remotely observed simultaneously by employing three different hyperspectral imagers (commercial and under development), FTIR instrument, UV LWIR cameras, and radiometer. Results gathered from the different instruments will be compared and by performing sensitivity tests on the retrieval codes the reliability of applying these techniques to volcanic gas observations will be evaluated. The campaign, the characteristics of the different instruments involved as well as the instrumental deployment strategy will be presented. Direct comparisons of spectral radiance in the infrared obtained by the different infrared instruments will be shown for several selected fields of view. First results for SO₂ from UV and in the infrared imagers will be shown, as well as first investigations on other species detected in the infrared.