



**HAL**  
open science

## Oxygen isotope mass balance of atmospheric nitrate at Dome C, East Antarctica, during the OPALE campaign

Joel Savarino, William Vicars, Matthieu Legrand, Susanne Preunkert, Bruno Jourdain, Markus Michael Frey, Alexandre Kukui, Nicolas Caillon, Jaime Gil Roca

### ► To cite this version:

Joel Savarino, William Vicars, Matthieu Legrand, Susanne Preunkert, Bruno Jourdain, et al.. Oxygen isotope mass balance of atmospheric nitrate at Dome C, East Antarctica, during the OPALE campaign. Atmospheric Chemistry and Physics Discussions, 2015, 15, pp.24041-24083. 10.5194/acpd-15-24041-2015 . insu-03577055

**HAL Id: insu-03577055**

**<https://insu.hal.science/insu-03577055v1>**

Submitted on 16 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

# Oxygen isotope mass balance of atmospheric nitrate at Dome C, East Antarctica, during the OPALE campaign

J. Savarino<sup>1,2</sup>, W. C. Vicars<sup>1,2,a</sup>, M. Legrand<sup>1,2</sup>, S. Preunkert<sup>1,2</sup>, B. Jourdain<sup>1,2</sup>, M. M. Frey<sup>3</sup>, A. Kukui<sup>4,5</sup>, N. Caillon<sup>1,2</sup>, and J. Gil Roca<sup>4,5</sup>

<sup>1</sup>Université Grenoble Alpes, Laboratoire de Glaciologie et Géophysique de l'Environnement (LGGE), 38000 Grenoble, France

<sup>2</sup>CNRS, Laboratoire de Glaciologie et Géophysique de l'Environnement (LGGE), 38000 Grenoble, France

<sup>3</sup>British Antarctic Survey, Natural Environment Research Council, Cambridge, UK

<sup>4</sup>Laboratoire Atmosphère, Milieux et Observations Spatiales (LATMOS), UMR8190, CNRS-Université de Versailles Saint Quentin, Université Pierre et Marie Curie, Paris, France

<sup>5</sup>Laboratoire de Physique et Chimie de l'Environnement et de l'Espace (LPC2E), UMR6115 CNRS-Université d'Orléans, 45071 Orléans CEDEX 2, France

<sup>a</sup>now at: Technical Services Program, Air Pollution Control Division, Colorado Department of Public Health and Environment, Denver, CO, USA

Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Received: 26 June 2015 – Accepted: 11 August 2015 – Published: 7 September 2015

Correspondence to: J. Savarino (jsavarino@ujf-grenoble.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

ACPD

15, 24041–24083, 2015

**Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign**

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Variations in the stable oxygen isotope composition of atmospheric nitrate act as novel tools for studying oxidative processes taking place in the troposphere. They provide both qualitative and quantitative constraints on the pathways determining the fate of atmospheric nitrogen oxides ( $\text{NO} + \text{NO}_2 = \text{NO}_x$ ). The unique and distinctive  $^{17}\text{O}$ -excess ( $\Delta^{17}\text{O} = \delta^{17}\text{O} - 0.52 \times \delta^{18}\text{O}$ ) of ozone, which is transferred to  $\text{NO}_x$  via oxidation, is a particularly useful isotopic fingerprint in studies of  $\text{NO}_x$  transformations. Constraining the propagation of  $^{17}\text{O}$ -excess within the  $\text{NO}_x$  cycle is critical in polar areas where there exists the possibility of extending atmospheric investigations to the glacial/interglacial time scale using deep ice core records of nitrate. Here we present measurements of the comprehensive isotopic composition of atmospheric nitrate collected at Dome C (East Antarctic plateau) during the austral summer of 2011/12. Nitrate isotope analysis has been here combined for the first time with key precursors involved in nitrate production ( $\text{NO}_x$ ,  $\text{O}_3$ ,  $\text{OH}$ ,  $\text{HO}_2$ ,  $\text{RO}_2$ , etc.) and direct observations of the transferrable  $\Delta^{17}\text{O}$  of surface ozone, which was measured at Dome C throughout 2012 using our recently developed analytical approach. Assuming that nitrate is mainly produced in Antarctica in summer through the  $\text{OH} + \text{NO}_2$  pathway and using concurrent measurements of  $\text{OH}$  and  $\text{NO}_2$ , we calculated a  $\Delta^{17}\text{O}$  signature for nitrate in the order of  $(21\text{--}22 \pm 3)\text{‰}$ . These values are lower than the measured values that ranged between 27 and 31 ‰. This discrepancy between expected and observed  $\Delta^{17}\text{O}(\text{NO}_3^-)$  values suggests the existence of an unknown process that contributes significantly to the atmospheric nitrate budget over this east Antarctic region.

## 1 Introduction

The search for ice core proxies related to past change of oxidative properties of the atmosphere is motivated by the need to model simulations of ozone and  $\text{OH}$  changes over preindustrial–industrial and glacial–interglacial timescales. Early ice

## Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 core reconstructions of oxidants based on  $\text{H}_2\text{O}_2$  (Sigg and Neftel, 1991) and HCHO (Staffelbach et al., 1991) measurements were hampered by the occurrence of post-depositional alteration of  $\text{H}_2\text{O}_2$  and HCHO concentrations in the upper snowpack prior to preservation in the ice (Hutterli et al., 2003). More recently, the  $^{17}\text{O}$ -excess ( $\Delta^{17}\text{O} = \delta^{17}\text{O} - 0.52 \times \delta^{18}\text{O}$ ) of nitrate, a unique isotopic signature inherited from ozone via bimolecular chemical reactions in the atmosphere, has shown promise as a conserved proxy for past oxidant concentrations (McCabe et al., 2005; Alexander et al., 2004). The  $\Delta^{17}\text{O}$  signal of nitrate reflects the relative importance of  $\text{NO}_x$  transformation mechanisms and recent studies suggest that the measurement of this isotopic signal in 10 ice cores and ancient sediments may provide relevant information regarding the role of ozone in the overall oxidative capacity of the paleo-atmosphere (McCabe et al., 2007; Michalski et al., 2004, 2003; Savarino et al., 2007; Thiemens, 2006). Parallel studies of the nitrogen isotope ratios ( $\delta^{15}\text{N}$ ) of nitrate in polar ice and snow suggest that this isotopic tracer may serve as a proxy for past variations in natural sources of atmospheric  $\text{NO}_x$  (Hastings et al., 2009, 2005; Jarvis et al., 2008). However, the chemical and physical factors governing the oxygen and nitrogen isotopic composition of atmospheric nitrate and its nitrogen oxide precursors are not fully understood (Alexander et al., 2009; Morin et al., 2009). This is particularly true in the polar troposphere, where the UV-photolysis of trace species present in the snowpack or marine aerosols initiates 20 complex boundary layer oxidation processes involving reactive halogen species (Bloss et al., 2010; Grannas et al., 2007; Wang et al., 2007) and results in distinctive  $\Delta^{17}\text{O}$  and  $\delta^{15}\text{N}$  signatures in atmospheric nitrate (Morin et al., 2012, 2007, 2008).

25 Here we present measurements of  $\delta^{15}\text{N}$  and  $\delta^{17}\text{O}$  and  $\delta^{18}\text{O}$  isotopic composition of atmospheric nitrate collected at Dome C between October 2011 and January 2012. These measurements were conducted within the framework of the OPALE project (Oxidant Production over Antarctic Land and its Export, Preunkert et al., 2012), which has provided an opportunity to combine nitrate isotopic observations with a wealth of meteorological and chemical observations, including measurements of species involved in nitrate production ( $\text{NO}_x$ ,  $\text{O}_3$ , OH,  $\text{HO}_2$ ,  $\text{RO}_2$ , etc.). The primary objective of this study

## Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



was to reconcile observations of  $\Delta^{17}\text{O}$  for atmospheric nitrate at Dome C with quantitative predictions based on nitrate isotope mass balance and atmospheric chemistry parameters, a unique opportunity offered by the OPALE campaign.

## 2 Methods

### 2.1 Site description and scientific context

Dome C is situated 3233 m a.s.l. on the East Antarctic Plateau (75°06' S, 123°23' E), approximately 1100 km from the coastal research station Dumont d'Urville and 560 km from the Vostok station. Deep ice cores were extracted at Dome C in the framework of the European Project for Ice Coring in Antarctica (EPICA) covering approximately 800 000 yr (EPICA-community-members, 2004) and Vostok covering the last 420 000 years (Petit et al., 1999). In parallel, studies aiming to understand the meteorological, chemical, and physical factors governing the variability of trace constituents preserved in the ice were initiated. This was done for aerosol in the framework of the French environmental observation service CESOA (Etude du cycle atmosphérique du Soufre en relation avec le climat aux moyennes et hautes latitudes Sud, <http://cesoa.ore.fr>) dedicated to the study of the sulfur cycle (Jourdain et al., 2008; Preunkert et al., 2008).

Although the Antarctic plateau is extraordinarily dry, cold, and far removed from sources of anthropogenic emissions, first atmospheric measurements of oxidants conducted in 1998–1999 during the ISCAT (Investigation of Sulfur Chemistry in the Antarctic Troposphere) field campaign revealed a high level of photochemical activity. For example, the average summertime OH concentration ( $2 \times 10^6 \text{ cm}^{-3}$ ) over the South Pole was found to be similar to that of the tropical marine boundary layer (MBL) (Mauldin et al., 2001). Unexpectedly high levels of nitric oxide (NO) were also detected, with concentrations one to two orders of magnitude higher than that typically observed in other remote regions (Davis et al., 2001). Model simulations revealed that the large OH

## Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

**Oxygen isotope mass  
balance of  
atmospheric nitrate  
at Dome C during the  
OPALE campaign****J. Savarino et al.**

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

concentrations observed at South Pole were a result of the elevated NO level, which catalyzes a rapid cycling of HO<sub>2</sub> to OH (Chen et al., 2004, 2001). The high concentrations of NO<sub>x</sub> were also inferred to drive in situ photochemical production of ozone during the ISCAT campaign (Crawford et al., 2001). Surface ozone and NO<sub>x</sub> measurements at Dome C suggest a similar level of enhanced oxidant production during November–January (Frey et al., 2013, 2014; Legrand et al., 2009).

The high levels of photochemical activity observed at South Pole and Dome C are now understood in terms of NO<sub>x</sub> release from the snowpack (Honrath et al., 2000, 1999; Jones et al., 2001, 2000; Zhou et al., 2001). This process is initiated by the photolysis of nitrate, which can lead to large fluxes of NO<sub>2</sub>, NO, and HONO from permanently sunlit snow (Anastasio and Chu, 2009; Grannas et al., 2007; Jacobi and Hilker, 2007; Legrand et al., 2014; Frey et al., 2013). Observed and modeled NO<sub>x</sub> production rates are largely capable of explaining the high levels of photochemical activity observed on the Antarctic plateau during spring (France et al., 2011; Liao and Tan, 2008; Wang et al., 2007) although detailed and speciation of nitrogen oxides chemistry remain largely unknown in this rich NO<sub>x</sub>/poor VOCs environment (Kukui et al., 2014; Frey et al., 2014; Legrand et al., 2014; Davis et al., 2008).

## 2.2 High-volume sampling

Bulk atmospheric samples were collected at Dome C on glass fiber filters using a high-volume air sampler (HVAS), which was installed on a platform 1 m above the ground. The HVAS was run by applying an average STP flow rate of 0.7 m<sup>3</sup> min<sup>-1</sup> that ensures the collection of a sufficient amount of nitrate for isotopic analysis. The atmospheric NO<sub>3</sub><sup>-</sup> collected on glass fiber filters represents the sum of atmospheric particulate NO<sub>3</sub><sup>-</sup> (pNO<sub>3</sub><sup>-</sup>) and gaseous nitric acid (HNO<sub>3</sub>) (Frey et al., 2009; Savarino, unpublished data). 11 HVAS samples were obtained during the OPALE campaign (from October 2011 to January 2012). After each collection period, filters were removed from the HVAS and placed in clean 50 mL centrifuge tubes, which were sealed in plastic bags and stored at -20 °C. Upon arrival at our laboratory in Grenoble, atmospheric filter samples were

extracted in 40 mL of ultra-pure water via centrifugation using Millipore Centricon™ filter units. Nitrate concentration was then determined for each filter extract solution using a colorimetric technique (Frey et al., 2009).

### 2.3 Ozone collection

The nitrite-coated filter technique for ozone isotope analysis has been described in detail in (Vicars et al., 2012; Vicars and Savarino, 2014). The principle of ozone collection underlying this technique is the filter-based chemical trapping of ozone via aqueous phase reaction with nitrite (Adon et al., 2010; Geyh et al., 1997; Koutrakis et al., 1993; Krzyzanowski, 2004):



By coupling this routine ozone measurement technique with recent analytical developments in the  $\Delta^{17}\text{O}$  analysis of nanomole quantities of nitrate (Kaiser et al., 2007), the  $^{17}\text{O}$ -excess transferred by one of the two  $\text{O}_3$  terminal atoms through bimolecular chemical reactions, denoted  $\Delta^{17}\text{O}(\text{O}_3)_{\text{term}}$ , as well as ozone's bulk  $\Delta^{17}\text{O}$  value, denoted  $\Delta^{17}\text{O}(\text{O}_3)_{\text{bulk}}$  ( $= 2/3 \times \Delta^{17}\text{O}(\text{O}_3)_{\text{term}}$  since all  $^{17}\text{O}$ -excess is located at the two  $\text{O}_3$  terminal atoms (Bhattacharya et al., 2008; Janssen and Tuzson, 2006), can be inferred from the oxygen isotopic composition of the nitrate produced via R1 within the coated filter matrix.

Ambient ozone collections were performed using an active air sampler consisting of 1/4" Teflon™ (PFA) tubing connecting three main sampler components: (i) a standard low-volume vacuum pump (Welch™, Model 2522C-02) equipped with a volume counter and needle valve (or flow meter) for flow rate regulation, (ii) a closed PFA filter holder assembly (Savillex™) containing a pre-coated 47 mm glass fiber sampling substrate (Whatman™, GF/A type), and (iii) an open-faced PFA filter holder assembly containing a 47 mm PTFE membrane filter (Zylon™, 5 μm pore size) for the removal of particulate species upstream of the coated filter. Glass fiber sampling substrates were

Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign**

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

coated prior to sample collection with 1 mL of a nitrite-based ozone sampling solution (0.1 M NaNO<sub>2</sub>, 0.05 M K<sub>2</sub>CO<sub>3</sub>, 0.1 M glycerol) (Koutrakis et al., 1993), allowed to dry at 75 °C for approximately 10 min, and then stored frozen in the dark in individual plastic PetriSlide™ containers (Millipore™). Samples were collected by loading pre-coated filters into the sampling filter holder, which was then connected to the prefilter (upstream) and needle valve/pump (downstream) and covered in aluminum foil to limit light exposure, a step that is necessary to limit the blank production rate during sample collection (Vicars et al., 2012). Air was then pumped through the sampling system at a target flow rate of 3.0 L min<sup>-1</sup>.

Sampling was conducted during the OPALE campaign (December 2011–January 2012) from a climate-controlled shelter, and a total of 28 samples were collected. However, due to difficulties in the application of our analytical technique to the unique environmental conditions encountered in Antarctica, the results obtained from these samples were inconclusive (i.e. unrealistic variability) due to the lack of light protection of the filter holder (Vicars et al., 2012, 2013). Sampling and isotopic analysis of ozone was therefore repeated in 2012, and a complete annual record of Δ<sup>17</sup>O(O<sub>3</sub>) was obtained (*n* = 60). Procedural filter blanks were also collected at regular intervals and were subjected to all of the same preparation, storage, handling, and analytical procedures as field samples. After sample collection, filter samples and procedural blanks were returned to their containers, which were covered in aluminum foil and stored at –20 °C before processing and analysis.

Filter samples were extracted in 18 mL of deionized water (18.2 MΩ, hereafter referred to as “MQ water”). In order to remove the excess (i.e., unreacted) nitrite reagent from the sample extracts, the solutions were treated with 1 mL of a 1 M sulfamic acid solution and then neutralized with a corresponding addition of high-purity sodium hydroxide (Granger and Sigman, 2009; Vicars et al., 2012). Extract solutions were then filtered via centrifugation using Millipore Centricon™ assemblies. The nitrate extracted from the coated filter samples was then subjected to isotopic analysis, as described in the following section.

## 2.4 Isotopic analysis

The comprehensive isotopic composition of nitrate ( $^{15}\text{N}/^{14}\text{N}$ ,  $^{17}\text{O}/^{16}\text{O}$ ,  $^{18}\text{O}/^{16}\text{O}$ ) was measured on a Finnigan<sup>TM</sup> MAT253 isotope ratio mass spectrometer (IRMS), equipped with a GasBench II<sup>TM</sup> and coupled to an in-house built nitrate interface (Morin et al., 2009). Nitrate in both the nitrite-coated filter and aerosol sample extracts was prepared for isotopic analysis by conversion to  $\text{N}_2\text{O}$  via the bacterial denitrifier method (Casciotti et al., 2002; Kaiser et al., 2007; Michalski et al., 2002; Sigman et al., 2001). The detailed analytical procedure has been described elsewhere (see Morin et al., 2009) and is here briefly presented.

Denitrifying bacteria (*Pseudomonas aureofaciens*) were cultured in nitrate-amended soy broth and incubated for 5 days in stoppered glass bottles. Bacterial cultures, after concentration by centrifugation and re-suspension, were dispensed as 2 mL aliquots into 20 mL glass vials, which were then crimped and purged with helium for 3 h. Approximately 100 nmol of sample nitrate was then injected into the purged vials and conversion of the sample nitrate to nitrous oxide ( $\text{N}_2\text{O}$ ) via bacterial denitrification was allowed to proceed overnight. The  $\text{N}_2\text{O}$  sample was then cryo-focused in a liquid nitrogen trap and introduced into a gold furnace where it was thermally decomposed at 900 °C into  $\text{O}_2$  and  $\text{N}_2$ . Following separation via gas chromatography, the  $\text{O}_2$  and  $\text{N}_2$  sample gases were directed into the ionization chamber of the IRMS. All analytical steps were simultaneously performed on nitrate isotopic standards and their equimolar mixtures (International Atomic Energy Agency USGS 32, USGS 34, and USGS 35), which were prepared in an identical background matrix as the samples. Individual analyses were normalized through comparison with these three nitrate reference materials (Coplen, 2011; Werner and Brand, 2001). All isotopic enrichment values for nitrate are reported relative to VSMOW and air  $\text{N}_2$  for oxygen and nitrogen, respectively. The overall accuracy of the method is estimated as the standard deviation of the residuals from the linear regression between the measured reference materials and

### Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

their expected values. For the results reported here, the average uncertainty obtained for  $\delta^{18}\text{O}$ ,  $\Delta^{17}\text{O}$ , and  $\delta^{15}\text{N}$  were 1.6, 0.5, and 1.0‰, respectively.

## 2.5 Complementary measurements

Concurrent chemical measurements conducted at Dome C during the campaign include HONO (Legrand et al., 2014),  $\text{HO}_x$  (= OH +  $\text{HO}_2$  +  $\text{RO}_2$ ) radicals (Kukui et al., 2014),  $\text{O}_3$  (Legrand et al., in preparation) and NO and  $\text{NO}_2$  (Frey et al., 2014). Photolysis rate coefficients and meteorological parameters were also documented.

## 3 Results and discussion

### 3.1 Isotope ratios of ozone and atmospheric nitrate

Atmospheric nitrate concentrations observed at Dome C during the campaign are presented in Fig. 1, the corresponding nitrate  $\Delta^{17}\text{O}$  and  $\delta^{15}\text{N}$  values in Fig. 2. Atmospheric nitrate concentrations ranged between 20 and 90  $\text{ng m}^{-3}$ , with the maximum values occurring in mid December 2011, concurrent with the period of intensive of atmospheric sampling of the OPALE field campaign. These values are in good agreement with those observed during the 2007–2008 and 2009–2010 field studies conducted at Dome C by Frey et al. (2009) and Erbland et al. (2013), respectively.

$\Delta^{17}\text{O}$  values for atmospheric nitrate ranged between 27.3 and 32.4‰,  $\delta^{15}\text{N}$  between  $-42.8$  and 1.7‰. The observed strongly depleted  $\delta^{15}\text{N}(\text{NO}_3^-)$  values are in good agreement with those previously reported and having unambiguously attributed to the transformation of local snowpack  $\text{NO}_x$  emissions via photochemistry in the boundary layer, which led to peaks in atmospheric nitrate concentration during the period from October to December (Erbland et al., 2013). As seen in Fig. 2, variations in  $\Delta^{17}\text{O}$  and  $\delta^{15}\text{N}$  were negatively correlated ( $r$  value of  $-0.86$ ) and again show similar amplitude and phase than those reported in previous studies (Erbland et al., 2013; Frey et al., 2009).

24050

Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

⏪

⏩

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

A time series showing the year-round record of  $\Delta^{17}\text{O}(\text{O}_3)_{\text{bulk}}$  at Dome C in 2012 is presented in Fig. 3.  $\Delta^{17}\text{O}(\text{O}_3)_{\text{bulk}}$  averaged  $24.9 \pm 1.9\text{‰}$  over 2012, corresponding to  $\Delta^{17}\text{O}(\text{O}_3)_{\text{term}}$  values of  $37.4 \pm 1.9\text{‰}$ . As shown in Fig. 4, these  $\Delta^{17}\text{O}(\text{O}_3)_{\text{bulk}}$  values are consistent with those observed in Grenoble (France), as well as with measurements conducted along a latitudinal transect from  $50^\circ\text{S}$  to  $50^\circ\text{N}$  in the Atlantic Ocean (Vicars and Savarino, 2014). Although the  $\Delta^{17}\text{O}(\text{O}_3)_{\text{bulk}}$  seasonal cycle reveals some interesting features, like the winter maximum, probably in response of the permanent winter darkness and stratospheric air mass intrusions, a complete description is beyond the scope of the present paper. What should be kept in mind here is the pretty stable  $\Delta^{17}\text{O}(\text{O}_3)_{\text{bulk}}$  value close to  $26\text{‰}$  that can be considered as representative for the OPALE campaign hold in November–January.

### 3.2 Nitrate isotope mass balance

The availability of a large database of trace chemical species measurements at Dome C during a portion of the OPALE field campaign (December 2011) offers a unique opportunity to compare observed  $\Delta^{17}\text{O}(\text{NO}_3^-)$  values in the atmosphere to estimated ones calculated from concurrent observations. As discussed at length in recent studies (for example, by Morin et al., 2011 and Vicars et al., 2013), the  $^{17}\text{O}$ -excess transfer functions associated with the various nitrate production pathways (i.e.,  $\Delta^{17}\text{O}(\text{NO}_3^-)_i$  values) can be estimated as a function of the  $\Delta^{17}\text{O}$  of nitrate precursor gases (i.e.,  $\text{NO}_x$ ,  $\text{O}_3$ ,  $\text{OH}$ , etc.) using mass balance calculations that trace the origin of oxygen atoms transferred during the chemical transformation of  $\text{NO}_x$  in the atmosphere. All atmospheric nitrate production channels involve either  $\text{NO}_2$  or a  $\text{NO}_x$  reservoir species derived from  $\text{NO}_2$  (e.g.,  $\text{N}_2\text{O}_5$ ). The first step in determining the  $\Delta^{17}\text{O}$  signature of each pathway is therefore a quantitative assessment of the steady state  $\Delta^{17}\text{O}$  value of  $\text{NO}_2$ , which is typically calculated as a function of the  $\Delta^{17}\text{O}$  value of  $\text{O}_3$  and the reaction dynamics involved in the conversion of  $\text{NO}$  to  $\text{NO}_2$ . As Dome C in summer is

permanently under sunlight, photochemical inter-conversion of  $\text{NO}_x$  is permanent:



5 At photochemical steady state (i.e. Reactions R2–R4 being faster than  $\text{NO}_2$  net sink reactions), an assumption that can be reasonably applied throughout the day at Dome C during summer, we have (Morin et al., 2011):

$$\Delta^{17}\text{O}(\text{NO}_2) = \alpha \times (1.18 \times \Delta^{17}(\text{O}_3)_{\text{bulk}} + 6.6) \quad (1)$$

10 where the term in bracket represents the laboratory deduced anomaly transfer function of the  $\text{NO} + \text{O}_3$  reaction (Savarino et al., 2008),  $\Delta^{17}\text{O}(\text{O}_3)_{\text{bulk}}$  the  $^{17}\text{O}$ -excess of the bulk  $\text{O}_3$  and  $\alpha$  represents the fraction of the atmospheric  $\text{NO}_2$  reservoir that has been produced through oxidation by  $\text{O}_3$  rather than  $\text{HO}_2/\text{RO}_2$  at photochemical equilibrium (Alexander et al., 2009; Michalski et al., 2003; Morin et al., 2011; Röckmann et al., 2001):

$$\alpha = \frac{k_{\text{NO}+\text{O}_3}[\text{NO}][\text{O}_3]}{k_{\text{NO}+\text{O}_3}[\text{NO}][\text{O}_3] + k_{\text{NO}+\text{HO}_2}[\text{NO}][\text{HO}_2]^*} \quad (2)$$

15 with  $[\text{HO}_2]^* = [\text{HO}_2] + [\text{RO}_2]$ .

It is important to note here that Eqs. (1) and (2) although established under the  $\text{NO}_x$  steady-state approximation, are independent of  $\text{NO}_2$  concentration for which a bias in measurement cannot be ruled out. Indeed, as discussed by Frey et al. (2013, 2014), bias in  $\text{NO}_2$  measurements is suspected partly because it remains difficult to explain the observed ratio of  $\text{NO}_2/\text{NO}$  which is systematically higher (up to a factor of 7) than predicted by calculations made by assuming photochemical steady state considering the  $\text{NO}_2$  photolysis and reaction of  $\text{NO}$  with  $\text{O}_3$ ,  $\text{HO}_2/\text{RO}_2$  and  $\text{BrO}$ . Equation (2) also assumes that  $[\text{HO}_2]^*$  is predominantly formed by the reaction  $\text{H} + \text{O}_2$

24052

Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALÉ campaign

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and  $R + O_2$  during the OPALE campaign (Kukui et al., 2014), resulting in the formation of  $[HO_2]^*$  devoid of any significant  $^{17}O$ -excess (Morin et al., 2011). Using OPALE measurements of  $NO$ ,  $O_3$ ,  $OH$  and  $HO_2/RO_2$  (Frey et al., 2014; Kukui et al., 2014), along with temperature dependent reaction kinetics data obtained from (Atkinson et al., 2004), we have calculated the diurnally mass averaged trend in  $\alpha$  for the month of December 2011 at Dome C. Measurements of  $\Delta^{17}O(O_3)_{bulk}$  at Dome C during the OPALE campaign averaged  $25 \pm 2\%$ , corresponding to  $\Delta^{17}O(O_3)_{term}$  values of  $37 \pm 2\%$  (Fig. 4). Samples collected in December indicate  $\Delta^{17}O(O_3)_{bulk}$  values close to  $26\%$  ( $\Delta^{17}O(O_3)_{term} = 3/2 \times \Delta^{17}O(O_3)_{bulk} = 39\text{--}40\%$ , Fig. 3), and we have therefore adopted a  $\Delta^{17}O(O_3)_{term}$  value of  $40\%$  in the subsequent mass balance calculations, in good agreement with the predicted value from a 1-D atmospheric model (Zahn et al., 2006). The diurnally mass average trend in  $\Delta^{17}O(NO_2)$  calculated using a  $\Delta^{17}O(O_3)_{bulk}$  value of  $26\%$  and Eq. (2) is shown in Fig. 5. No trend is observed during the OPALE campaign and on average the predicted value is  $\Delta^{17}O(NO_2) = 31 \pm 2\%$  throughout December, corresponding to average  $\alpha$  value of 0.83. In other words, at steady state, the concentrations of  $O_3$  and  $HO_2^*$  measured during OPALE predicts that around 83% of  $NO_2$  is formed via Reaction (R3) (see also Table 1). In the absence of the  $\alpha$  dilution effect introduced by the  $HO_2^*$  reaction,  $\Delta^{17}O(NO_2)$  would equal  $37\%$ , a value  $8\%$  lower than an estimation obtained from modeling only  $NO_x$ - $O_3$  chemistry at standard temperature and pressure (Michalski et al., 2014). This difference is essentially explained by the use of different  $\Delta^{17}O(O_3)_{bulk}$  ( $32\%$  Michalski's simulation,  $26\%$  for our observations), which possibly correspond to different conditions of the two studies.

By accounting for the origin of the oxygen atom transferred during the conversion of  $NO_2$  to nitrate, the  $\Delta^{17}O$  signature of the nitrate produced through different reaction mechanisms can be calculated. For summer conditions at Dome C, it is reasonable to assume that the dominant atmospheric nitrate formation pathway is the gas-phase association of  $NO_2$  and the  $OH$  radical (Alexander et al., 2009):



leading to the following  $^{17}\text{O}$ -excess mass balance (Michalski et al., 2003 ; Morin et al., 2011):

$$\Delta^{17}\text{O}(\text{NO}_3^-) = \frac{2}{3}\Delta^{17}\text{O}(\text{NO}_2) + \frac{1}{3}\Delta^{17}\text{O}(\text{OH}) \quad (3)$$

In order to predict the  $\Delta^{17}\text{O}$  value of the nitrate produced through Reaction (R5) by mass balance, the isotopic composition of tropospheric OH must be known. The OH radical participates in a rapid isotopic exchange with atmospheric water vapor, which represents a very large oxygen reservoir relative to OH, with a  $\Delta^{17}\text{O}$  that is negligible compared to ozone or nitrate (Luz and Barkan, 2010). This exchange tends to erase the  $^{17}\text{O}$ -excess of OH under humidity and temperature conditions typical of the mid-latitudes (Dubey et al., 1997); therefore, the  $\Delta^{17}\text{O}$  of OH is normally assumed to be zero in modeling studies applied to these regions. As discussed by Morin et al. (2007), this assumption of  $\Delta^{17}\text{O}(\text{OH}) = 0$  is not valid under the low humidity conditions encountered in the polar atmosphere. The degree of isotopic equilibration between OH and  $\text{H}_2\text{O}$  can be determined as a function of the relative rates of the isotope exchange reaction and the main OH sink reactions:

$$\beta = \frac{L}{L + k_{\text{H}_2\text{O}+\text{OH}}[\text{H}_2\text{O}]} \quad (4)$$

where  $L$  represents the total chemical loss rate of OH.  $\beta$  is the factor relating the initial  $\Delta^{17}\text{O}$  transferred to OH upon its formation, denoted  $\Delta^{17}\text{O}(\text{OH})_{\text{prod.}}$ , to its steady state  $\Delta^{17}\text{O}$  value (Morin et al., 2007):

$$\Delta^{17}\text{O}(\text{OH}) = \beta \times \Delta^{17}\text{O}(\text{OH})_{\text{prod.}} \quad (5)$$

In plain words, Eqs. (4) and (5) predict that when the isotopic exchange reaction dominates over OH chemical losses (i.e.,  $\beta \ll 1$ ), the steady state  $\Delta^{17}\text{O}$  value of OH will be equal to that of water (i.e.,  $\Delta^{17}\text{O} \approx 0\text{‰}$ ). Conversely, when water vapor concentrations are low and the rate of chemical loss is large relative to the rate of the isotopic

## Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

exchange, then  $\Delta^{17}\text{O}(\text{OH}) = \Delta^{17}\text{O}(\text{OH})_{\text{prod.}}$ . Kukui et al. (2014), using a Master Chemical Mechanism box model, constrained by the OPALE meteorological conditions and concurrent chemical observations, give the rate of the OH chemical sources and sinks.  $\text{NO}_2$  as measured by Frey et al. (2014) represents at most only ca. 10 % (equivalent of ca. 1 ‰) of the total sink of OH, which is predominantly dominated by reactions with  $\text{CO}$ ,  $\text{CH}_4$ , aldehydes and to a lesser extent by reactions with  $\text{O}_3$ ,  $\text{H}_2$ , and  $\text{NO}$ . Thus, the possible overestimation of  $\text{NO}_2$  concentration has only a minor effect on  $\beta$  calculation and is well embedded within the total uncertainty of such calculation. To assess the value of  $\Delta^{17}\text{O}(\text{OH})$ , we have computed  $\beta$  for the conditions found during the OPALE campaign using the same 0-D box model that is used to evaluate the budget of OH and  $\text{RO}_2$  during the OPALE campaign (see Kukui et al., 2014 and Table 1) and used the exchange kinetic rates given in Dubey et al. (1997). The absolute water vapor concentration is deduced from relative humidity and temperature measurements using Bolton, (1980) (i.e.  $P_{\text{water}} = 6.112 \times e^{\frac{(17.67 \times (T-273))}{T-29.5}}$ , with  $P_{\text{water}}$  in hPa and  $T$  in K). The results of this calculation (Fig. 6), indicate that  $\beta$  varies between  $0.70 \pm 0.10$  ( $1 \sigma$ ) and  $0.30 \pm 0.10$  from midnight to noon for conditions prevailing during the OPALE campaign, suggesting that on daily average basis approximately 43 % of the  $\Delta^{17}\text{O}$  value originally present in OH is preserved from exchange with  $\text{H}_2\text{O}$ , consistent with estimates for an Arctic site described by Morin et al. (2007).

The value of  $\Delta^{17}\text{O}(\text{OH})_{\text{prod.}}$  is more difficult to assess because of the interplay of  $\text{HO}_x$  family and the different sources involved in OH formation. In the rich  $\text{NO}_x$  atmosphere at Dome C in summer, the  $\text{O}(^1\text{D}) + \text{H}_2\text{O}$  reaction forming OH is a minor reaction pathway. When multiple pathways are involved in the production of OH,  $\Delta^{17}\text{O}(\text{OH})_{\text{prod.}}$  can be estimated by a simple isotope mass balance equation where  $\Delta^{17}\text{O}(\text{OH})_{\text{prod.}} = \sum_i P_i \times \Delta^{17}\text{O}_i$  with  $P_i$  the relative production rate of the  $i$ th reaction pathway with respect to the total production rate and  $\Delta^{17}\text{O}_i$  its associated  $^{17}\text{O}$ -excess (Morin et al., 2011). Observations at Dome C during the OPALE campaign indicate that the photolysis of HONO and the  $\text{HO}_2 + \text{NO}$  reaction may represent the most sig-



## Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

nificant sources of OH at Dome C during the period of seasonal snowpack emissions (Kukui et al., 2014). However, the measurement of HONO (around  $\text{pmol mol}^{-1}$ ) during OPALE, probably biased by  $\text{HO}_2\text{NO}_2$  interference (Legrand et al., 2014) is incompatible with the  $\text{HO}_x$  ( $= \text{OH} + \text{HO}_2/\text{RO}_2$ ) radical budget. Best agreement is achieved when HONO at Dome C is assumed to originate from snow emissions with the emission strength evaluated by Legrand et al. (2014). Using a 1-D model, Kukui et al. (2014) show that the concentrations of HONO corresponding to about 20–30 % of measured HONO are consistent with those calculated from the budget analysis of OH radicals with the concentrations of  $\text{NO}_2$  either calculated from NO measurements assuming PSS or observed by Frey et al. (2014). Therefore, the production of OH by HONO photolysis is adjusted consequently and the 0-D box model (Kukui et al., 2014) is used to calculate all other production rates of OH. Note that even lowering HONO to 20–30 % of the measured values, this species remains the major primary source of radicals at Dome C. Applying the isotope  $^{17}\text{O}$ -excess transfer (Morin et al., 2011) and the  $\text{OH}_{\text{prod.}}$  isotope mass balance,  $\Delta^{17}\text{O}(\text{OH})_{\text{prod.}}$  on average equals  $5 \pm 2\%$  ( $1 \sigma$ ). Because the major process leading to the emission of HONO from the snowpack is the photolysis of nitrate, which possesses a  $\Delta^{17}\text{O}$  value of approximately 32‰, both in the snow “skin layer” (Erbland et al., 2013) and in the top 10 cm of snow (Frey et al., 2009), we have assumed that  $\Delta^{17}\text{O}(\text{HONO})_{\text{atm}} = \Delta^{17}\text{O}(\text{NO}_3^-)_{\text{snow}}$  as both oxygen atoms of HONO can be tracked back to the nitrate. An example of the isotope mass balance calculation is given in Table 1. Figure 7 shows the diurnally-integrated average of the  $\Delta^{17}\text{O}(\text{OH})$ .  $\Delta^{17}\text{O}(\text{OH})$  varies in a narrow range, between 1 and 3‰. An estimation of the  $\Delta^{17}\text{O}$  signature for the  $\text{NO}_2 + \text{OH}$  channel,  $\Delta^{17}\text{O}(\text{NO}_3^-)_{\text{R2}}$ , that accounts for the  $^{17}\text{O}$ -excess carried by the OH radical results in values ranging between 20 and 23‰. Averaging over the same time period as the nitrate atmospheric sampling, diurnally-integrated average  $\Delta^{17}\text{O}(\text{NO}_3^-)$  values of  $21\text{--}22\% \pm 3\%$  can be estimated for December (Table 2). These values are 6–8‰ lower than the observed atmospheric values for  $\Delta^{17}\text{O}(\text{NO}_3^-)$  (27–30‰ during OPALE, Fig. 2 and Table 2). The source of discrepancy between ob-

served and modeled  $\Delta^{17}\text{O}(\text{NO}_3^-)$  is presently unknown. A critical evaluation may offer nevertheless some clues.

## 4 Discussion

### 4.1 Alternative sources of $\text{NO}_2$

5 A possible explanation for the underestimation of  $\Delta^{17}\text{O}(\text{NO}_3^-)$  involves the potential  
role of reactive halogen chemistry in the troposphere over the Antarctic plateau (Bloss  
et al., 2010; Morin et al., 2008). Reactive halogen oxides ( $X\text{O} = \text{BrO}, \text{ClO}, \text{IO}, \text{etc.}$ )  
are produced through the reaction of halogen radicals ( $X$ ) with ozone, a pathway that  
10 plays an important role in the catalytic process responsible for ozone depletion events  
(ODEs) observed in the Arctic boundary layer since the 1980s (Fan and Jacob, 1992;  
Simpson et al., 2007):



In terms of the chemical budget of  $\text{NO}_x$ , the impact of  $X\text{O}$  can occur via two chem-  
ical mechanisms (see Sect. 4.2 for the second mechanism involving the formation of  
15 halogen nitrate,  $X\text{ONO}_2$ ). First,  $X\text{O}$  can oxidize  $\text{NO}$  to  $\text{NO}_2$ , a pathway that competes  
with the  $\text{NO} + \text{O}_3$  and  $\text{NO} + \text{HO}_2/\text{RO}_2$  reactions in terms of  $\text{NO}$  oxidation:



For conditions typical of the Antarctic boundary layer,  $1 \text{ pmol mol}^{-1}$  of  $X\text{O}$  has roughly  
the same oxidizing power as  $4 \text{ nmol mol}^{-1}$  of ozone in terms of  $\text{NO}$  oxidation (Atkinson  
20 et al., 2007). Therefore, when halogen oxides are present at relevant levels, the R7  
reaction can result in concentrations of  $\text{NO}_2$  that are higher than that predicted from the  
balance between  $\text{NO}_2$  destruction via photolysis and production through the reaction

of NO with O<sub>3</sub> or HO<sub>2</sub>/RO<sub>2</sub> (i.e., the extended Leighton mechanism):

$$\frac{[\text{NO}_2]}{[\text{NO}]} = \frac{k_{\text{NO}+\text{O}_3}[\text{O}_3] + k_{\text{NO}+\text{HO}_2}[\text{HO}_2] + k_{\text{NO}+\text{RO}_2}[\text{RO}_2] + k_{\text{NO}+\text{XO}}[\text{XO}]}{j_{\text{NO}_2}} \quad (6)$$

The interaction of XO in the NO<sub>x</sub> cycle at Dome C would have important implications for the Δ<sup>17</sup>O of atmospheric nitrate. The production of halogen oxide radicals proceeds through a direct transfer of a terminal oxygen atom from ozone to the XO product (Zhang et al., 1997). Therefore, it is expected that the Δ<sup>17</sup>O of XO is equal to Δ<sup>17</sup>O(O<sub>3</sub>)<sub>term.</sub>, which means that the reaction of NO with XO is roughly equivalent to the NO + O<sub>3</sub> reaction in terms of Δ<sup>17</sup>O transfer to NO<sub>2</sub> (note that the NO + XO transfer is greater than NO + O<sub>3</sub> as in the later case, part of the central O<sub>3</sub> atom participates in the reaction). The participation of XO species in the oxidation of NO thus leads to a greater Δ<sup>17</sup>O transfer to NO<sub>2</sub> by effectively increasing the value of α. However, on the Antarctic plateau, BrO did not exceed 2–3 pmol mol<sup>-1</sup> at most during OPALE campaign (Frey et al., 2014). Including BrO chemistry would only increase α by 2% (due to the specific form of α ≡ 1/(1+x)) of which is too low to significantly influence Δ<sup>17</sup>O(NO<sub>2</sub>) and ultimately Δ<sup>17</sup>O(NO<sub>3</sub><sup>-</sup>). In the absence of measurements of other halogens we cannot completely rule out a role of the halogen chemistry there. However, even with α = 1, its maximum but unrealistic value due to the high concentration of HO<sub>2</sub>, Δ<sup>17</sup>O(NO<sub>3</sub><sup>-</sup>) would reach the range of 23–25‰, in better agreement with the observations but still significantly lower. Similarly, in the event of a non isotopic steady state of NO<sub>2</sub> (Michalski et al., 2014), it is very unlikely that Δ<sup>17</sup>O(NO<sub>2</sub>) could reach values greater than its primary snow nitrate source (i.e. Δ<sup>17</sup>O(NO<sub>2</sub>) > Δ<sup>17</sup>O(NO<sub>3</sub><sup>-</sup>)<sub>snow</sub> = 30–35‰ in summer at Dome C, (Erland et al., 2013; Frey et al., 2009), still leaving the predicted Δ<sup>17</sup>O(NO<sub>3</sub><sup>-</sup>) underestimated with respect to atmospheric observations.

Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 4.2 Alternative oxidation pathways of NO<sub>2</sub>

Considering Reaction (R5) as the main source of HNO<sub>3</sub>, an alternative approach is to consider that OH bears a higher Δ<sup>17</sup>O than the estimate calculated previously. Assuming a β of one, which seems again unrealistic, will increase Δ<sup>17</sup>O(NO<sub>3</sub><sup>-</sup>) by 1‰ at most (Table 2), still insufficient to explained atmospheric observations as NO + HO<sub>2</sub> remain a major source of OH, independently of the assumed β.

Alternatively, if measured HONO concentrations are considered instead of those assumed to constrain by the HO<sub>x</sub> budget (i.e. 4 times lower than measured), average Δ<sup>17</sup>O(NO<sub>3</sub><sup>-</sup>) values of 23–24‰ are calculated (Table 2), again systematically lower than the observed range of 27–30‰. However, given the significant uncertainty surrounding the isotopic composition of HONO and its relative contribution to total OH production at Dome C, it is not possible to make a firm conclusion in this regard.

Therefore, neither the common sources of NO<sub>2</sub> nor the daytime formation of HNO<sub>3</sub> seems to be able to explain the high Δ<sup>17</sup>O(NO<sub>3</sub><sup>-</sup>) values of atmospheric nitrate observed at Dome C in summer. When this observation is taken together with the high NO<sub>2</sub>/NO ratio observed by Frey et al. during two summer seasons at Dome C (Frey et al., 2013, 2014), clearly our current understanding of the NO<sub>x</sub> chemistry on the Antarctic plateau seems to be incomplete.

There are several other processes that possibly account for the disagreement between the measurements and mass balance calculations. Indeed, in addition to its impact on NO<sub>x</sub> cycling through the R7 pathway, an increasing body of evidence points towards reactive halogen chemistry as a major NO<sub>x</sub> sink and source of nitrate via the production and subsequent hydrolysis of XNO<sub>3</sub> species (Sander et al., 1999; Savarino et al., 2013; Vogt et al., 1996):



## Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

A critical analysis of the CHABLIS data led Bauguitte et al. (2012) to conclude that the Reactions (R8) and (R9) pathway exerted predominant control over the chemical loss rate of  $\text{NO}_x$  during the campaign, despite the significant uncertainties involved in the parameterization of the uptake processes (Finlayson-Pitts, 2009). This implies that  $X\text{NO}_3$  uptake may also represent a significant source of nitrate at Dome C should halogen oxide radicals ( $X\text{O}$ ) be present at the required concentration. Experimental (Gane et al., 2001) and theoretical (McNamara and Hillier, 2001) studies suggest that the oxygen atom initially associated with  $X\text{O}$  combines with the N atom of  $\text{NO}_2$  to form nitrate, thus transferring the isotopic signature of both  $X\text{O}$  and  $\text{NO}_2$ . The specific  $\Delta^{17}\text{O}$  value induced by  $X\text{NO}_3$  hydrolysis can thus be expressed as follows (Morin et al., 2007):

$$\Delta^{17}\text{O}(\text{NO}_3^-)_{\text{R6}} = \frac{2}{3}\Delta^{17}\text{O}(\text{NO}_2) + \frac{1}{3}\Delta^{17}\text{O}(\text{O}_3)_{\text{term}} \quad (7)$$

efficiently bypassing the  $\text{OH}^{17}\text{O}$ -excess budget. Through consideration of the increased  $\Delta^{17}\text{O}$  transfer associated with Reaction (R8), the observations of  $\Delta^{17}\text{O}(\text{NO}_3^-)$  during December can be reconciled with the values calculated by mass-balance if approximately 10–20 % of total nitrate production is assumed to occur via  $X\text{NO}_3$  hydrolysis. However, no sufficient halogen concentration has been observed on the Antarctic plateau to sustain such chemical pathway but we note that chlorine chemistry has never been probed on the Antarctic plateau.

There is increasing body of evidence that heterogeneous hydrolysis of  $\text{NO}_2$  can be a possible source of  $\text{HONO}$  and  $\text{HNO}_3$  in acidic conditions (Finlayson-Pitts, 2009), with the potential to explain the difference between the calculated and measured atmospheric  $\Delta^{17}\text{O}(\text{NO}_3^-)$  values. This mechanism would represent a source of nitrate with a  $\Delta^{17}\text{O}$  value roughly equivalent to the nitrate originally present in the surface snow (i.e., 30–35‰), a signature significantly higher than that induced by Reaction (R5). If this production mechanism is active at the air–snow interface at Dome C and results in the slow emission of nitrate to the atmosphere via physical release after its formation,

## Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALÉ campaign

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

it would act to increase the  $\Delta^{17}\text{O}$  value of nitrate in the boundary layer relative to the local  $\Delta^{17}\text{O}(\text{NO}_3^-)_{\text{R5}}$  oxidation signature. However, considering the propensity of nitric acid to stick on snow (Crowley et al., 2010), the snowpack to act as a sink rather than a source of nitric acid (Dibb, 2004; Erbland et al., 2013) and the fast  $\text{NO}_x$  recycling that should take place within the snowpack, it is very unlikely that  $\Delta^{17}\text{O}(\text{NO}_3^-)$  could be explained by a direct nitric acid emissions from snow, which has been ultimately shown to be limited (Slusher et al., 2010; Erbland et al., 2013; Berhanu et al., 2014).

A critical analysis of  $\Delta^{17}\text{O}(\text{NO}_3^-)$  shows in fact that such high values correspond mainly to the nighttime chemistry of  $\text{NO}_x$  (Michalski et al., 2003; Morin et al., 2008). Nighttime chemistry involves species like  $\text{N}_2\text{O}_5$  and  $\text{NO}_3$  in the process of forming  $\text{HNO}_3$  and again efficiently by-pass the OH pathway. It is conceivable that below the photic zone, within the snowpack,  $\text{N}_2\text{O}_5$  and  $\text{NO}_3$  could be produced when  $\text{O}_3$  and  $\text{NO}_2$  are transported at depth but there is no reason to think that such dark  $\text{NO}_x$  chemistry could in a way or in another survive the photic zone transition and thus influences the overlying atmosphere.

Stratospheric nitrate deposited to the surface snow during winter, which has been observed to possess  $\Delta^{17}\text{O}(\text{NO}_3^-)$  values in the range of 35–41‰ (Erbland et al., 2013) and possibly more, may act to buffer the  $\Delta^{17}\text{O}$  of the atmospheric nitrate reservoir via evaporation late into the spring and summer. However, this seems again unlikely given the rapidity of  $\text{NO}_x$  cycling and oxidative loss at Dome C during this time (Frey et al., 2013; Legrand et al., 2009).

While it is presently difficult to determine the precise nature of the process(es) leading to the relatively large  $^{17}\text{O}$ -excess values observed for atmospheric nitrate at Dome C, the correlation observed between the  $\delta^{15}\text{N}$  and  $\Delta^{17}\text{O}$  values of atmospheric nitrate (see Sect. 3.1) provides at least one direct line of evidence that the high  $\Delta^{17}\text{O}(\text{NO}_3^-)$  values observed during spring and early summer could be associated with snowpack emissions of  $\text{NO}_x$ . Considering only samples collected at Dome C between October and December, both those reported here and those collected in 2009 and described by Erbland et al. (2013), a strong anticorrelation ( $r = -0.90$ ) is observed between the

$\delta^{15}\text{N}$  and  $\Delta^{17}\text{O}$  values of atmospheric nitrate (Fig. 8). In other words, the atmospheric nitrate sampled in early spring, which is heavily depleted in  $^{15}\text{N}$  due its formation from snowpack  $\text{NO}_x$  emissions, possessed consistently higher  $\Delta^{17}\text{O}$  values than the nitrate sampled directly after this period of maximum snow photochemistry. This finding suggests that the mechanism producing enhanced  $\Delta^{17}\text{O}(\text{NO}_3^-)$  values observed during this time is tightly coupled in time and space with the intensity of  $\text{NO}_x$  emissions from the snowpack, an observation very similar to that of Morin et al. (2012), who detected a similar relationship between  $\delta^{15}\text{N}$  and  $\Delta^{17}\text{O}$  for atmospheric nitrate in the springtime boundary layer over Barrow, Alaska ( $71^\circ\text{N}$ ). The authors of this study attributed the observed correlation to the coupling of snowpack  $\text{NO}_x$  emissions and reactive halogen chemistry, suggesting that these two processes were interrelated and mutually strengthening. In the case of the OPALE 2011–2012 data, the correlation between  $\delta^{15}\text{N}$  and  $\Delta^{17}\text{O}$  could arise from any of the potential pathways previously discussed. For example, as proposed by Morin et al. (2012), the R8 and R9 pathways may be enhanced under periods of intense snowpack emissions. Alternatively, a correlation could result from an increased contribution to total OH production from the photolysis of HONO, which is co-emitted with  $\text{NO}_x$  via nitrate photochemistry (Grannas et al., 2007) and may induce a larger  $^{17}\text{O}$ -excess in OH as compared to the conventional  $\text{O}(^1\text{D}) + \text{H}_2\text{O}$  pathway. Furthermore, the hydrolysis of  $\text{NO}_2$  in snow, should it contribute significantly to nitrate production at Dome C, is likely amplified during periods when concentrations  $\text{NO}_2$  are high in the snowpack interstitial air due to nitrate/nitrite photochemistry. Therefore, while the processes responsible for driving the formation of atmospheric nitrate at Dome C during summer cannot be unambiguously identified, the isotopic results presented here clearly indicate that snowpack emissions result in enhanced  $\Delta^{17}\text{O}$  transfer to nitrate. Our understanding of  $\text{NO}_x$  chemistry above the snow surface at Dome C is therefore incomplete.

## Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 5 Conclusions

Constraining the propagation of ozone's  $^{17}\text{O}$ -excess signature within the  $\text{NO}_x$  cycle is critical in polar areas where the opportunity is offered to extend atmospheric investigations based on  $\Delta^{17}\text{O}$  measurements to the glacial/interglacial time scale using deep ice core records of nitrate. However, the factors governing the present-day isotopic composition of atmospheric nitrate over the Antarctic plateau remain poorly understood, primarily due to the complex nature of the boundary layer photochemistry initiated during spring by  $\text{NO}_x$  emissions from the snowpack.

An isotopic mass balance performed for atmospheric nitrate during December 2011, informed by in situ oxidant concentration measurements conducted within the framework of the OPALE field study, suggests the existence of an unexpected process bypassing the commonly accepted daytime chemistry of  $\text{NO}_2$  (i.e.  $\text{NO}_2 + \text{OH}$ ) that contributes significantly to the atmospheric nitrate budget over Dome C. The strong negative correlation observed between the  $\delta^{15}\text{N}$  and  $\Delta^{17}\text{O}$  values of nitrate between October and December suggests that this unknown process is enhanced during periods of intense emissions from the snowpack. Potential explanations for this observation include: (i) an increased  $\Delta^{17}\text{O}$  transfer from OH due to its formation from the photolysis of HONO released from the snowpack, (ii) heterogeneous hydrolysis of  $\text{NO}_2$  due to the high concentrations of  $\text{NO}_2$  in the snowpack interstitial air, and (iii) the co-emission of reactive halogen species that act as an intermediate in the transfer of  $\Delta^{17}\text{O}$  from ozone to nitrate. Further research is needed to solve the many inconsistencies (e.g. high  $\text{NO}_2/\text{NO}$  ratio, high concentration of  $\text{NO}_2$ , unresolved HONO atmospheric concentration, interference such as  $\text{HO}_2\text{NO}_2$ , isotope mass balance) observed during the OPALE experiments.

*Acknowledgements.* The research leading to these results has received funding from the European Community's Seventh Framework Programme (FP7/2007–2013) under the grant agreement number 237 890. We would like to thank INSU for its financial support for lab experiments through its LEFE program. The Agence nationale de la recherche (ANR) is gratefully acknowledged for its financial support through the OPALE project (contract NT09-451281). The Insti-

24063

ACPD

15, 24041–24083, 2015

### Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





tute Polaire Paul-Emile Victor (IPEV) supported the research and polar logistics through the program SUNITEDC No. 1011. This work has been partially supported by a grant from Labex OSUG@2020 (Investissements d'avenir – ANR10 LABX56. We would also like to thank all the field team members present during the OPALE campaign. Meteorological data were obtained from “IPEV/PNRA: Routine Meteorological Observation at Station Concordia”.

## References

- Adon, M., Galy-Lacaux, C., Yoboué, V., Delon, C., Lacaux, J. P., Castera, P., Gardrat, E., Pienaar, J., Al Ourabi, H., Laouali, D., Diop, B., Sigha-Nkamdjou, L., Akpo, A., Tathy, J. P., Lavenu, F., and Mougín, E.: Long term measurements of sulfur dioxide, nitrogen dioxide, ammonia, nitric acid and ozone in Africa using passive samplers, *Atmos. Chem. Phys.*, 10, 7467–7487, doi:10.5194/acp-10-7467-2010, 2010.
- Alexander, B., Savarino, J., Kreutz, K., and Thiemens, M. H.: Impact of preindustrial biomass-burning emissions on the oxidative pathways of tropospheric sulfur and nitrogen, *J. Geophys. Res.*, 109, D08303, doi:10.1029/2003JD004218, 2004.
- Alexander, B., Hastings, M. G., Allman, D. J., Dachs, J., Thornton, J. A., and Kunasek, S. A.: Quantifying atmospheric nitrate formation pathways based on a global model of the oxygen isotopic composition ( $\Delta^{17}\text{O}$ ) of atmospheric nitrate, *Atmos. Chem. Phys.*, 9, 5043–5056, doi:10.5194/acp-9-5043-2009, 2009.
- Anastasio, C. and Chu, L.: Photochemistry of nitrous acid (HONO) and nitrous acid-ium ion ( $\text{H}_2\text{ONO}^+$ ) in aqueous solution and ice, *Environ. Sci. Technol.*, 43, 1108–1114, doi:10.1021/es802579a, 2009.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume I – gas phase reactions of  $\text{O}_x$ ,  $\text{HO}_x$ ,  $\text{NO}_x$  and  $\text{SO}_x$  species, *Atmos. Chem. Phys.*, 4, 1461–1738, doi:10.5194/acp-4-1461-2004, 2004.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume III – gas phase reactions of inorganic halogens, *Atmos. Chem. Phys.*, 7, 981–1191, doi:10.5194/acp-7-981-2007, 2007.

## Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign**

J. Savarino et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Bauguitte, S. J.-B., Bloss, W. J., Evans, M. J., Salmon, R. A., Anderson, P. S., Jones, A. E., Lee, J. D., Saiz-Lopez, A., Roscoe, H. K., Wolff, E. W., and Plane, J. M. C.: Summertime  $\text{NO}_x$  measurements during the CHABLIS campaign: can source and sink estimates unravel observed diurnal cycles?, *Atmos. Chem. Phys.*, 12, 989–1002, doi:10.5194/acp-12-989-2012, 2012.
- Berhanu, T. A., Meusinger, C., Erbland, J., Jost, R., Bhattacharya, S. K., Johnson, M. S., and Savarino, J.: Laboratory study of nitrate photolysis in Antarctic snow, II. Isotopic effects and wavelength dependence, *J. Chem. Phys.*, 140, 244305, doi:10.1063/1.4882899, 2014.
- Bhattacharya, S. K., Pandey, A., and Savarino, J.: Determination of intramolecular isotope distribution of ozone by oxidation reaction with silver metal, *J. Geophys. Res.*, 113, D03303, doi:10.1029/2006jd008309, 2008.
- Bloss, W. J., Camredon, M., Lee, J. D., Heard, D. E., Plane, J. M. C., Saiz-Lopez, A., Bauguitte, S. J.-B., Salmon, R. A., and Jones, A. E.: Coupling of  $\text{HO}_x$ ,  $\text{NO}_x$  and halogen chemistry in the antarctic boundary layer, *Atmos. Chem. Phys.*, 10, 10187–10209, doi:10.5194/acp-10-10187-2010, 2010.
- Bolton, D.: The computation of equivalent potential temperature, *Mon. Weather Rev.*, 108, 1046–1053, 1980.
- Casciotti, K. L., Sigman, D. M., Galanter Hastings, M., Böhlke, J. K., and Hilkert, A.: Measurement of the oxygen isotopic composition of nitrate in seawater and freshwater using the denitrifier method, *Anal. Chem.*, 74, 4905–4912, 2002.
- Chen, G., Davis, D., Crawford, J., Nowak, J. B., Eisele, F., Mauldin, R. L., Tanner, D., Buhr, M., Shetter, R., Lefer, B., Arimoto, R., Hogan, A., and Blake, D.: An investigation of South Pole  $\text{HO}_x$  chemistry: comparison of model results with ISCAT observations, *Geophys. Res. Lett.*, 28, 3633–3636, doi:10.1029/2001GL013158, 2001.
- Chen, G., Davis, D., Crawford, J., Hutterli, L. M., Huey, L. G., Slusher, D., Mauldin, L., Eisele, F., Tanner, D., Dibb, J., Buhr, M., McConnell, J., Lefer, B., Shetter, R., Blake, D., Song, C. H., Lombardi, K., and Arnoldy, J.: A reassessment of  $\text{HO}_x$  South Pole chemistry based on observations recorded during ISCAT 2000, *Atmos. Environ.*, 38, 5451–5461, doi:10.1016/j.atmosenv.2003.07.018, 2004.
- Coplen, T. B.: Guidelines and recommended terms for expression of stable-isotope-ratio and gas-ratio measurement results, *Rapid Commun. Mass Sp.*, 25, 2538–2560, doi:10.1002/rcm.5129, 2011.

**Oxygen isotope mass  
balance of  
atmospheric nitrate  
at Dome C during the  
OPALE campaign**

J. Savarino et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Crawford, J. H., Davis, D. D., Chen, G., Buhr, M., Oltmans, S., Weller, R., Mauldin, L., Eisele, F., Shetter, R., Lefer, B., Arimoto, R., and Hogan, A.: Evidence for photochemical production of ozone at the South Pole surface, *Geophys. Res. Lett.*, 28, 3641–3644, doi:10.1029/2001GL013055, 2001.
- 5 Crowley, J. N., Ammann, M., Cox, R. A., Hynes, R. G., Jenkin, M. E., Mellouki, A., Rossi, M. J., Troe, J., and Wallington, T. J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume V – heterogeneous reactions on solid substrates, *Atmos. Chem. Phys.*, 10, 9059–9223, doi:10.5194/acp-10-9059-2010, 2010.
- Davis, D., Nowak, J. B., Chen, G., Buhr, M., Arimoto, R., Hogan, A., Eisele, F., Mauldin, L.,  
10 Tanner, D., Shetter, R., Lefer, B., and McMurry, P.: Unexpected high levels of NO observed at South Pole, *Geophys. Res. Lett.*, 28, 3625–3628, doi:10.1029/2000GL012584, 2001.
- Davis, D., Seelig, J., Huey, G., Crawford, J., Chen, G., Wang, Y., Buhr, M., Helmig, D., Neff, W., and Blake, D.: A reassessment of Antarctic plateau reactive nitrogen based on ANTICI 2003 airborne and ground based measurements, *Atmos. Environ.*, 42, 2831–2848,  
15 doi:10.1016/j.atmosenv.2007.07.039, 2008.
- Dibb, J.: Soluble reactive nitrogen oxides at South Pole during ISCAT 2000, *Atmos. Environ.*, 38, 5399–5409, doi:10.1016/j.atmosenv.2003.01.001, 2004.
- Dubey, M. K., Mohrschladt, R., Donahue, N. M., and Anderson, J. G.: Isotope specific kinetics of hydroxyl radical (OH) with water (H<sub>2</sub>O): testing models of reactivity and atmospheric  
20 fractionation, *J. Phys. Chem.-US*, 101, 1494–1500, 1997.
- EPICA-community-members: Eight glacial cycles from an Antarctic ice core, *Nature*, 429, 623–628, doi:10.1038/nature02599, 2004.
- Erbland, J., Vicars, W. C., Savarino, J., Morin, S., Frey, M. M., Frosini, D., Vince, E., and Martins, J. M. F.: Air–snow transfer of nitrate on the East Antarctic Plateau – Part 1: Isotopic  
25 evidence for a photolytically driven dynamic equilibrium in summer, *Atmos. Chem. Phys.*, 13, 6403–6419, doi:10.5194/acp-13-6403-2013, 2013.
- Fan, S.-M. and Jacob, D. J.: Surface ozone depletion in Arctic spring sustained by bromine reactions on aerosols, *Nature*, 359, 522–524, 1992.
- Finlayson-Pitts, B. J.: Reactions at surfaces in the atmosphere: integration of experiments and theory as necessary (but not necessarily sufficient) for predicting the physical chemistry of  
30 aerosols, *Phys. Chem. Chem. Phys.*, 11, 7760–7779, doi:10.1039/b906540g, 2009.
- France, J. L., King, M. D., Frey, M. M., Erbland, J., Picard, G., Preunkert, S., MacArthur, A., and Savarino, J.: Snow optical properties at Dome C (Concordia), Antarctica; implications for

**Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign**

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

snow emissions and snow chemistry of reactive nitrogen, *Atmos. Chem. Phys.*, 11, 9787–9801, doi:10.5194/acp-11-9787-2011, 2011.

Frey, M. M., Savarino, J., Morin, S., Erbland, J., and Martins, J. M. F.: Photolysis imprint in the nitrate stable isotope signal in snow and atmosphere of East Antarctica and implications for reactive nitrogen cycling, *Atmos. Chem. Phys.*, 9, 8681–8696, doi:10.5194/acp-9-8681-2009, 2009.

Frey, M. M., Brough, N., France, J. L., Anderson, P. S., Traulle, O., King, M. D., Jones, A. E., Wolff, E. W., and Savarino, J.: The diurnal variability of atmospheric nitrogen oxides (NO and NO<sub>2</sub>) above the Antarctic Plateau driven by atmospheric stability and snow emissions, *Atmos. Chem. Phys.*, 13, 3045–3062, doi:10.5194/acp-13-3045-2013, 2013.

Frey, M. M., Roscoe, H. K., Kukui, A., Savarino, J., France, J. L., King, M. D., Legrand, M., and Preunkert, S.: Atmospheric nitrogen oxides (NO and NO<sub>2</sub>) at Dome C, East Antarctica, during the OPALE campaign, *Atmos. Chem. Phys.*, 15, 7859–7875, doi:10.5194/acp-15-7859-2015, 2015.

Gane, M. P., Williams, N. A., and Sodeau, J. R.: A reflection-absorption infrared spectroscopy (RAIRS) investigation of the low-temperature heterogeneous hydrolysis of bromine nitrate, *J. Phys. Chem.-US*, 105, 4002–4009, 2001.

Geyh, A., Wolfson, J. M., and Koutrakis, P.: Development and evaluation of a small active ozone sampler, *Environ. Sci. Technol.*, 31, 2326–2330, 1997.

Granger, J. and Sigman, D. M.: Removal of nitrite with sulfamic acid for nitrate N and O isotope analysis with the denitrifier method, *Rapid Commun. Mass Sp.*, 23, 3753–3762, doi:10.1002/rcm.4307, 2009.

Grannas, A. M., Jones, A. E., Dibb, J., Ammann, M., Anastasio, C., Beine, H. J., Bergin, M., Bottenheim, J., Boxe, C. S., Carver, G., Chen, G., Crawford, J. H., Dominé, F., Frey, M. M., Guzmán, M. I., Heard, D. E., Helmig, D., Hoffmann, M. R., Honrath, R. E., Huey, L. G., Hutterli, M., Jacobi, H. W., Klán, P., Lefer, B., McConnell, J., Plane, J., Sander, R., Savarino, J., Shepson, P. B., Simpson, W. R., Sodeau, J. R., von Glasow, R., Weller, R., Wolff, E. W., and Zhu, T.: An overview of snow photochemistry: evidence, mechanisms and impacts, *Atmos. Chem. Phys.*, 7, 4329–4373, doi:10.5194/acp-7-4329-2007, 2007.

Hastings, M. G., Sigman, D. M., and Steig, E. J.: Glacial/interglacial changes in the isotopes of nitrate from the Greenland Ice Sheet Project 2 (GISP2) ice core, *Global Biogeochem. Cy.*, 19, GB4024, doi:10.1029/2005gb002502, 2005.

**Oxygen isotope mass  
balance of  
atmospheric nitrate  
at Dome C during the  
OPALE campaign**

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Hastings, M. G., Jarvis, J. C., and Steig, E. J.: Anthropogenic impacts on nitrogen isotopes of ice-core nitrate, *Science*, 324, 1288–1288, doi:10.1126/science.1170510, 2009.
- Honrath, R. E., Peterson, M. C., Guo, S., Dibb, J. E., Shepson, P. B., and Campbell, B.: Evidence of NO<sub>x</sub> production within or upon ice particles in the Greenland snowpack, *Geophys. Res. Lett.*, 26, 695–698, 1999.
- Honrath, R. E., Guo, S., Peterson, M. C., Dziobak, M. P., Dibb, J. E., and Arsenault, M. A.: Photochemical production of gas phase NO<sub>x</sub> from ice crystal NO<sub>3</sub>, *J. Geophys. Res.*, 105, 24183–24190, 2000.
- Hutterli, M. A., McConnell, J. R., Bales, R. C., and Stewart, R. W.: Sensitivity of hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) and formaldehyde (HCHO) preservation in snow to changing environmental conditions: implications for ice core records, *J. Geophys. Res.-Atmos.*, 108, D1, doi:10.1029/2002JD002528, 2003.
- Jacobi, H. W. and Hilker, B.: A mechanism for the photochemical transformation of nitrate in snow, *J. Photoch. Photobio. A*, 185, 371–382, doi:10.1016/j.jphotochem.2006.06.039, 2007.
- Janssen, C. and Tuzson, B.: A diode laser spectrometer for symmetry selective detection of ozone isotopomers, *Appl. Phys. B*, 82, 487–494, doi:10.1007/s00340-005-2044-6, 2006.
- Jarvis, J. C., Steig, E. J., Hastings, M. G., and Kunasek, S. A.: Influence of local photochemistry on isotopes of nitrate in Greenland snow, *Geophys. Res. Lett.*, 35, L21804, doi:10.1029/2008gl035551, 2008.
- Jones, A. E., Weller, R., Wolff, E. W., and Jacobi, H.-W.: Speciation and rate of photochemical NO and NO<sub>2</sub> production in Antarctica snow, *Geophys. Res. Lett.*, 27, 345–348, 2000.
- Jones, A. E., Weller, R., Anderson, P. S., Jacobi, H. W., Wolff, E. W., Schrems, O., and Miller, H.: Measurements of NO<sub>x</sub> emissions from the Antarctic snowpack, *Geophys. Res. Lett.*, 28, 1499–1502, 2001.
- Jourdain, B., Preunkert, S., Cerri, O., Castebrunet, H., Udisti, R., and Legrand, M.: Year-round record of size-segregated aerosol composition in central Antarctica (Concordia station): implications for the degree of fractionation of sea-salt particles, *J. Geophys. Res.*, 113, D14308, doi:10.1029/2007jd009584, 2008.
- Kaiser, J., Hastings, M. G., Houlton, B. Z., Röckmann, T., and Sigman, D. M.: Triple oxygen isotope analysis of nitrate using the denitrifier method and thermal decomposition of N<sub>2</sub>O, *Anal. Chem.*, 79, 599–607, 2007.
- Koutrakis, P., Wolfson, J. M., Bunyaviroch, A., Froehlich, S. E., Hirano, K., and Muliki, J. D.: Measurement of ambient ozone using a nitrite-coated filter, *Anal. Chem.*, 65, 209–214, 1993.

**Oxygen isotope mass  
balance of  
atmospheric nitrate  
at Dome C during the  
OPALE campaign**

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Krzyzanowski, J.: Ozone variation with height in a forest canopy – results from a passive sampling field campaign, *Atmos. Environ.*, 38, 5957–5962, doi:10.1016/j.atmosenv.2004.07.017, 2004.

5 Kukui, A., Legrand, M., Preunkert, S., Frey, M. M., Loisil, R., Gil Roca, J., Jourdain, B., King, M. D., France, J. L., and Ancellet, G.: Measurements of OH and RO<sub>2</sub> radicals at Dome C, East Antarctica, *Atmos. Chem. Phys.*, 14, 12373–12392, doi:10.5194/acp-14-12373-2014, 2014.

10 Legrand, M., Preunkert, S., Jourdain, B., Gallée, H., Goutail, F., Weller, R., and Savarino, J.: Year-round record of surface ozone at coastal (Dumont d'Urville) and inland (Concordia) sites in East Antarctica, *J. Geophys. Res.*, 114, D20306, doi:10.1029/2008jd011667, 2009.

15 Legrand, M., Preunkert, S., Frey, M., Bartels-Rausch, Th., Kukui, A., King, M. D., Savarino, J., Kerbrat, M., and Jourdain, B.: Large mixing ratios of atmospheric nitrous acid (HONO) at Concordia (East Antarctic Plateau) in summer: a strong source from surface snow?, *Atmos. Chem. Phys.*, 14, 9963–9976, doi:10.5194/acp-14-9963-2014, 2014.

20 Liao, W. and Tan, D.: 1-D Air-snowpack modeling of atmospheric nitrous acid at South Pole during ANTCI 2003, *Atmos. Chem. Phys.*, 8, 7087–7099, doi:10.5194/acp-8-7087-2008, 2008.

Luz, B. and Barkan, E.: Variations of <sup>17</sup>O/<sup>16</sup>O and <sup>18</sup>O/<sup>16</sup>O in meteoric waters, *Geochim. Cosmochim. Ac.*, 74, 6276–6286, doi:10.1016/j.gca.2010.08.016, 2010.

25 Mauldin, R. L., Eisele, F. L., Tanner, D. J., Kosciuch, E., Shetter, R., Lefer, B., Hall, S. R., Nowak, J. B., Buhr, M., Chen, G., Wang, P., and Davis, D.: Measurements of OH, H<sub>2</sub>SO<sub>4</sub>, and MSA at the South Pole during ISCAT, *Geophys. Res. Lett.*, 28, 3629–3632, 2001.

McCabe, J. R., Boxe, C. S., Colussi, A. J., Hoffman, M. R., and Thiemens, M. H.: Oxygen isotopic fractionation in the photochemistry of nitrate in water and ice, *J. Geophys. Res.*, 110, D15310, doi:10.1029/2004jd005484, 2005.

30 McCabe, J. R., Thiemens, M. H., and Savarino, J.: A record of ozone variability in South Pole Antarctic snow: role of nitrate oxygen isotopes, *J. Geophys. Res.*, 112, D12303, doi:10.1029/2006jd007822, 2007.

McNamara, J. P. and Hillier, I. H.: Mechanism of the hydrolysis of halogen nitrates in small water clusters studied by electronic structure methods, *J. Phys. Chem.-US*, 105, 7011–7024, 2001.

Michalski, G., Savarino, J., Bohlke, J. K., and Thiemens, M.: Determination of the total oxygen isotopic composition of nitrate and the calibration of a Delta O-17 nitrate reference material, *Anal. Chem.*, 74, 4989–4993, doi:10.1021/ac0256282, 2002.

## Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Michalski, G., Scott, Z., Kabling, M., and Thiemens, M. H.: First measurements and modeling of  $\Delta^{17}\text{O}$  in atmospheric nitrate, *Geophys. Res. Lett.*, 30, 1870, doi:10.1029/2003gl017015, 2003.

Michalski, G., Böhlke, J. K., and Thiemens, M.: Long term atmospheric deposition as the source of nitrate and other salts in the Atacama Desert, Chile: new evidence from mass-independent oxygen isotopic compositions, *Geochim. Cosmochim. Ac.*, 68, 4023–4038, doi:10.1016/j.gca.2004.04.009, 2004.

Michalski, G., Bhattacharya, S. K., and Girsch, G.:  $\text{NO}_x$  cycle and the tropospheric ozone isotope anomaly: an experimental investigation, *Atmos. Chem. Phys.*, 14, 4935–4953, doi:10.5194/acp-14-4935-2014, 2014.

Morin, S., Savarino, J., Bekki, S., Gong, S., and Bottenheim, J. W.: Signature of Arctic surface ozone depletion events in the isotope anomaly ( $\Delta^{17}\text{O}$ ) of atmospheric nitrate, *Atmos. Chem. Phys.*, 7, 1451–1469, doi:10.5194/acp-7-1451-2007, 2007.

Morin, S., Savarino, J., Yan, N., Frey, M. M., Bottenheim, J., Bekki, S., and Martins, J.: Tracing the origin and fate of  $\text{NO}_x$  in the Arctic atmosphere using stable isotopes, *Science*, 322, 730–732, doi:10.1126/science.1161910, 2008.

Morin, S., Savarino, J., Frey, M. M., Domine, F., Jacobi, H. W., Kaleschke, L., and Martins, J. M. F.: Comprehensive isotopic composition of atmospheric nitrate in the Atlantic Ocean boundary layer from 65° S to 79° N, *J. Geophys. Res.*, 114, D05303, doi:10.1029/2008jd010696, 2009.

Morin, S., Sander, R., and Savarino, J.: Simulation of the diurnal variations of the oxygen isotope anomaly ( $\Delta^{17}\text{O}$ ) of reactive atmospheric species, *Atmos. Chem. Phys.*, 11, 3653–3671, doi:10.5194/acp-11-3653-2011, 2011.

Morin, S., Erbland, J., Savarino, J., Domine, F., Bock, J., Friess, U., Jacobi, H. W., Sihler, H., and Martins, J. M. F.: An isotopic view on the connection between photolytic emissions of  $\text{NO}_x$  from the Arctic snowpack and its oxidation by reactive halogens, *J. Geophys. Res.-Atmos.*, 117, D00R08, doi:10.1029/2011jd016618, 2012.

Petit, J. R., Jouzel, J., Raynaud, D., Barkov, N. I., Barnola, J. M., Basile, I., Bender, M., Chappellaz, J., Davis, M., Delaygue, G., Delmotte, M., Kotlyakov, V. M., Legrand, M., Lipenkov, V. Y., Lorius, C., Pepin, L., Ritz, C., Saltzman, E., and Stievenard, M.: Climate and atmospheric history of the past 420,000 years from the Vostok ice core, *Antarctica, Nature*, 399, 429–436, 1999.

**Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign**

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Preunkert, S., Jourdain, B., Legrand, M., Udisti, R., Becagli, S., and Cerri, O.: Seasonality of sulfur species (dimethyl sulfide, sulfate, and methanesulfonate) in Antarctica: inland versus coastal regions, *J. Geophys. Res.*, 113, D15302, doi:10.1029/2008jd009937, 2008.
- Preunkert, S., Ancellet, G., Legrand, M., Kukui, A., Kerbrat, M., Sarda-Estève, R., Gros, V., and Jourdain, B.: Oxidant Production over Antarctic Land and its Export (OPALE) project: an overview of the 2010–2011 summer campaign, *J. Geophys. Res.-Atmos.*, 117, D15307, doi:10.1029/2011JD017145, 2012.
- Röckmann, T., Kaiser, J., Brenninkmeijer, C. A. M., Crowley, J. N., Borchers, R., Brand, W. A., and Crutzen, P. J.: Isotopic enrichment of nitrous oxide ( $^{15}\text{N}^{14}\text{NO}$ ), ( $^{14}\text{N}^{15}\text{NO}$ ), ( $^{14}\text{N}^{14}\text{N}^{18}\text{O}$ ) in the stratosphere and in the laboratory, *J. Geophys. Res.*, 106, 10403–10410, 2001.
- Sander, R., Rudich, Y., von Glasow, R., and Crutzen, P. J.: The role of  $\text{BrNO}_3$  in marine tropospheric chemistry: a model study, *Geophys. Res. Lett.*, 26, 2857–2860, 1999.
- Savarino, J., Kaiser, J., Morin, S., Sigman, D. M., and Thiemens, M. H.: Nitrogen and oxygen isotopic constraints on the origin of atmospheric nitrate in coastal Antarctica, *Atmos. Chem. Phys.*, 7, 1925–1945, doi:10.5194/acp-7-1925-2007, 2007.
- Savarino, J., Bhattacharya, S. K., Morin, S., Baroni, M., and Doussin, J. F.: The  $\text{NO} + \text{O}_3$  reaction: a triple oxygen isotope perspective on the reaction dynamics and atmospheric implications for the transfer of the ozone isotope anomaly, *J. Chem. Phys.*, 128, 194303, doi:10.1063/1.2917581, 2008.
- Savarino, J., Morin, S., Erbland, J., Grannec, F., Patey, M. D., Vicars, W., Alexander, B., and Achterberg, E. P.: Isotopic composition of atmospheric nitrate in a tropical marine boundary layer, *P. Natl. Acad. Sci. USA*, 110, 17668–17673, doi:10.1073/pnas.1216639110, 2013.
- Sigg, A. and Neffel, A.: Evidence for a 50-percent increase in  $\text{H}_2\text{O}_2$  over the past 200 years from a Greenland ice core, *Nature*, 351, 557–559, 1991.
- Sigman, D. M., Casciotti, K. L., Andreani, M., Barford, C., Galanter, M., and Böhlke, J. K.: A bacterial method for the nitrogen isotopic analysis of nitrate in seawater and freshwater, *Anal. Chem.*, 73, 4145–4153, 2001.
- Simpson, W. R., von Glasow, R., Riedel, K., Anderson, P., Ariya, P., Bottenheim, J., Burrows, J., Carpenter, L. J., Frieß, U., Goodsite, M. E., Heard, D., Hutterli, M., Jacobi, H.-W., Kaleschke, L., Neff, B., Plane, J., Platt, U., Richter, A., Roscoe, H., Sander, R., Shepson, P., Sodeau, J., Steffen, A., Wagner, T., and Wolff, E.: Halogens and their role in polar boundary-layer ozone depletion, *Atmos. Chem. Phys.*, 7, 4375–4418, doi:10.5194/acp-7-4375-2007, 2007.



**Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign**

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Slusher, D. L., Neff, W. D., Kim, S., Huey, L. G., Wang, Y., Zeng, T., Tanner, D. J., Blake, D. R., Beyersdorf, A., Lefer, B. L., Crawford, J. H., Eisele, F. L., Mauldin, R. L., Kosciuch, E., Buhr, M. P., Wallace, H. W., and Davis, D. D.: Atmospheric chemistry results from the ANTCI 2005 Antarctic plateau airborne study, *J. Geophys. Res.*, 115, D07304, doi:10.1029/2009jd012605, 2010.
- Staffelbach, T., Neftel, A., Stauffer, B., and Jacob, D.: A record of the atmospheric methane sink from formaldehyde in polar ice cores, *Nature*, 349, 603–605, 1991.
- Thiemens, M. H.: History and applications of mass-independent isotope effects, *Annu. Rev. Earth Pl. Sc.*, 34, 217–262, 2006.
- Vicars, W. C. and Savarino, J.: Quantitative constraints on the  $^{17}\text{O}$ -excess ( $\Delta^{17}\text{O}$ ) signature of surface ozone: ambient measurements from  $50^\circ\text{N}$  to  $50^\circ\text{S}$  using the nitrite-coated filter technique, *Geochim. Cosmochim. Ac.*, 135, 270–287, doi:10.1016/j.gca.2014.03.023, 2014.
- Vicars, W. C., Bhattacharya, S. K., Erbland, J., and Savarino, J.: Measurement of the  $^{17}\text{O}$ -excess ( $\Delta^{17}\text{O}$ ) of tropospheric ozone using a nitrite-coated filter, *Rapid Commun. Mass Sp.*, 26, 1219–1231, doi:10.1002/rcm.6218, 2012.
- Vicars, W. C., Morin, S., Savarino, J., Wagner, N. L., Erbland, J., Vince, E., Martins, J. M. F., Lerner, B. M., Quinn, P. K., Coffman, D. J., Williams, E. J., and Brown, S. S.: Spatial and diurnal variability in reactive nitrogen oxide chemistry as reflected in the isotopic composition of atmospheric nitrate: results from the CalNex 2010 field study, *J. Geophys. Res.-Atmos.*, 118, 10567–10588, doi:10.1002/jgrd.50680, 2013.
- Vogt, R., Crutzen, P. J., and Sander, R.: A mechanism for halogen release from sea-salt aerosol in the remote marine boundary layer, *Nature*, 383, 327–330, 1996.
- Wang, Y., Choi, Y., Zeng, T., Davis, D., Buhr, M., Gregoryhuey, L., and Neff, W.: Assessing the photochemical impact of snow  $\text{NO}_x$  emissions over Antarctica during ANTCI 2003, *Atmos. Environ.*, 41, 3944–3958, doi:10.1016/j.atmosenv.2007.01.056, 2007.
- Werner, R. A. and Brand, W. A.: Referencing strategies and techniques in stable isotope ratio analysis, *Rapid Commun. Mass Sp.*, 15, 501–519, doi:10.1002/rcm.258, 2001.
- Zahn, A., Franz, P., Bechtel, C., Groß, J.-U., and Röckmann, T.: Modelling the budget of middle atmospheric water vapour isotopes, *Atmos. Chem. Phys.*, 6, 2073–2090, doi:10.5194/acp-6-2073-2006, 2006.
- Zhang, J. S., Miao, T. T., and Lee, Y. T.: Crossed molecular beam study of the reaction  $\text{Br} + \text{O}_3$ , *J. Phys. Chem.-US*, 101, 6922–6930, 1997.

Zhou, X., Beine, H. J., Honrath, R. E., Fuentes, J. D., Simpson, W., Shepson, P. B., and Bottenheim, J. W.: Snowpack photochemical production of HONO: a major source of OH in the Arctic boundary layer in springtime, *Geophys. Res. Lett.*, 28, 4087–4090, doi:10.1029/2001GL013531, 2001.

ACPD

15, 24041–24083, 2015

**Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign**

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPAL campaign

J. Savarino et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 1.** Rate of production and sink of OH and mass balance calculation of  $\Delta^{17}\text{O}$ .

Conditions for 19 Dec 2011, 3:45 p.m. OH = 3.96 molecules cm <sup>-3</sup>	<sup>a</sup> Median rate in 10 <sup>5</sup> molecules cm <sup>-3</sup> s <sup>-1</sup>	$\Delta^{17}\text{O}_i^c$ in ‰
<b>Net sources of OH</b>		
P1 HONO + <i>hν</i> → OH + NO	5.1 <sup>b</sup>	32
P2 H <sub>2</sub> O <sub>2</sub> + <i>hν</i> → 2OH	1.7	2
P3 O <sub>3</sub> + <i>hν</i> + H <sub>2</sub> O → 2OH	0.6	20
P4 CH <sub>3</sub> OOH + <i>hν</i> → HO <sub>2</sub> + OH	0.3	0
<b>Recycling RO<sub>2</sub> → OH</b>		
P5 NO + HO <sub>2</sub> → NO <sub>2</sub> + OH	7.7	0
P6 HO <sub>2</sub> + O <sub>3</sub> → OH + 2O <sub>2</sub>	0.4	0
<b>Net sink of OH</b>		
L1 CO + OH → HO <sub>2</sub> + CO <sub>2</sub>	6.3	
L2 CH <sub>4</sub> + OH → CH <sub>3</sub> O <sub>2</sub> + H <sub>2</sub> O	2.6	
L2 HCHO + OH → HO <sub>2</sub> + CO	0.8	
L4 CH <sub>3</sub> CHO + OH → CH <sub>3</sub> CO <sub>3</sub>	0.9	
L5 O <sub>3</sub> + OH → HO <sub>2</sub> + O <sub>2</sub>	0.6	
L6 H <sub>2</sub> + OH + O <sub>2</sub> → HO <sub>2</sub> + H <sub>2</sub> O	0.60	
L7 CH <sub>3</sub> OOH + OH → CH <sub>3</sub> O <sub>2</sub> + H <sub>2</sub> O	0.5	
L8 H <sub>2</sub> O <sub>2</sub> + OH → HO <sub>2</sub> + H <sub>2</sub> O	0.3	
<b>Net OH losses</b>		
L9 NO <sub>2</sub> + OH → HNO <sub>3</sub>	3.9	
L10 NO + OH → HONO	0.6	
L11 OH + RO <sub>2</sub> → products	0.5	
L12 OH + RO <sub>2</sub> NO <sub>2</sub> → products	0.6	
L13 OH + HONO → NO <sub>2</sub> + H <sub>2</sub> O	0.2	
L14 OH + HNO <sub>3</sub> → H <sub>2</sub> O + NO <sub>3</sub>	0.0	
<b>Isotope exchange</b>		
E <sub>1</sub> HQ + H <sub>2</sub> O ⇌ HO + H <sub>2</sub> Q	24.3	
<b>NO<sub>2</sub> main source</b>		
N1 NO + O <sub>3</sub> → NO <sub>2</sub> + O <sub>2</sub>	27.0	37
<b><sup>17</sup>O-excess NO<sub>2</sub></b>		
$\alpha = (N1 / (N1 + P5))$	0.78	
$\Delta^{17}\text{O}(\text{NO}_2)$		29
<b><sup>17</sup>O-excess OH</b>		
$\Delta^{17}\text{O}(\text{OH})_{\text{prod}} = (\sum P_i \times \Delta^{17}\text{O}_i) / \sum P_i$		5.8
$\beta = \sum L_i / (\sum L_i + E_1)$	0.43	
$\Delta^{17}\text{O}(\text{OH})$		2.5

<sup>a</sup>: Production rates obtained from a 0-D box model (see Kukui et al., 2014 for details).

<sup>b</sup>: HONO production rate divided by a factor 4 to balance the HO<sub>2</sub> radical budget (see Kukui et al., 2014; Legrand et al., 2014 for justification).

<sup>c</sup>: HONO is assumed to be formed by the photodissociation of nitrate in snow.  $\Delta^{17}\text{O}(\text{NO}_3)_{\text{snow}}$  is therefore assigned to HONO. The rest of the <sup>17</sup>O-excess transfer (i.e. P<sub>2</sub> to P<sub>6</sub> and N<sub>1</sub>) follows the rules established in (Morin et al., 2011) and a  $\Delta^{17}\text{O}(\text{O}_3)_{\text{bulk}} = 26\%$ .

## Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign

J. Savarino et al.

**Table 2.** Comparison of measured and calculated  $\Delta^{17}\text{O}(\text{NO}_3^-)$  values.

Sampling Period	Measured	Calculated			
		$\alpha$ constrained by observations $\Delta^{17}\text{O}(\text{OH})^*$ based on $\text{HO}_x$ budget	$\alpha = 1$ $\Delta^{17}\text{O}(\text{OH})^*$ based $\text{HO}_x$ budget	$\alpha$ constrained by observations $\beta = 1$	$\alpha$ constrained by observations $\Delta^{17}\text{O}(\text{OH})$ based on observed HONO
10 Dec–16 Dec	<b>29.6</b>	21.9	25.6	22.6	27.0
16 Dec–23 Dec	<b>29.0</b>	21.0	25.6	21.7	26.3
23 Dec–30 Dec	<b>27.8</b>	21.6	25.4	22.0	25.7
30 Dec–2 Jan	<b>27.3</b>	21.5	25.3	22.4	24.9

\* HONO production rate divided by a factor 4 to balance the  $\text{HO}_x$  radical budget (see Kukui et al. (2014) and Legrand et al. (2014) for justification).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign


J. Savarino et al.



**Figure 1.** Atmospheric nitrate concentrations observed between October 2011 and January 2012. The samples collected during the intensive measurement period of the OPALE campaign (December 2011) are indicated with open circles.

## Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign


J. Savarino et al.



**Figure 2.**  $\Delta^{17}\text{O}$  (primary y axis) and  $\delta^{15}\text{N}$  (secondary y axis) of atmospheric nitrate collected between October 2011 and January 2012. The samples collected during the intensive measurement period of the OPALE campaign (December 2011) are indicated with open symbols.

## Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALÉ campaign

J. Savarino et al.




**Figure 3.**  $\Delta^{17}\text{O}(\text{O}_3)_{\text{bulk}}$  values for the 60 ambient air samplings done at Dome C throughout 2012. Vertical error bars refer to the total uncertainty estimated for the technique ( $\pm 1.7\text{‰}$ ).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign

J. Savarino et al.




**Figure 4.** Comparison of  $\Delta^{17}\text{O}(\text{O}_3)_{\text{bulk}}$  values obtained at Dome C with those previously reported by Vicars and Savarino (2014) at other sites. The box plot indicates the interquartile range (box) and the median (line), maximum, and minimum values. The mean value is denoted by a circle.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



## Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALÉ campaign

J. Savarino et al.




**Figure 5.** Quantitative assessment of the daily averaged trend in the  $\Delta^{17}\text{O}$  of  $\text{NO}_2$  at Dome C during December 2011 derived from concurrent measurements of ozone,  $\text{NO}$ , and  $\text{HO}_2/\text{RO}_2$ .

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

## Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALE campaign

J. Savarino et al.



**Figure 6.** December 2011 time-series for  $\beta$ , the fraction of the  $^{17}\text{O}$ -excess originally associated with the OH radical that is preserved against isotopic exchange with water.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Figure 7. Same as Fig. 5 but for  $\Delta^{17}\text{O}$  of OH.

Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALÉ campaign

J. Savarino et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Oxygen isotope mass balance of atmospheric nitrate at Dome C during the OPALÉ campaign

J. Savarino et al.



**Figure 8.** Relationship observed between the  $\delta^{15}\text{N}$  and  $\Delta^{17}\text{O}$  values of atmospheric nitrate present at Dome C between October and December during both the 2009 and 2011 summer campaigns. Error bars indicate the typical analytical uncertainties associated with the measurements.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)