



HAL
open science

Day-Night Cloud Asymmetry Inhibits Early Ocean Formation on Venus

M. Turbet, E. Bolmont, G. Chaverot, D. Ehrenreich, J. Leconte, Emmanuel Marcq

► **To cite this version:**

M. Turbet, E. Bolmont, G. Chaverot, D. Ehrenreich, J. Leconte, et al.. Day-Night Cloud Asymmetry Inhibits Early Ocean Formation on Venus. Venera-D: Venus Cloud Habitability System Workshop, Nov 2021, Virtual Meeting, Unknown Region. pp.LPI Contribution No. 2629, id.4051. insu-03582672

HAL Id: insu-03582672

<https://insu.hal.science/insu-03582672>

Submitted on 21 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DAY-NIGHT CLOUD ASYMMETRY INHIBITS EARLY OCEAN FORMATION ON VENUS. M. Turbet¹, E. Bolmont¹, G. Chaverot¹, D. Ehrenreich¹, J. Leconte² and E. Marcq³, ¹Observatoire astronomique de l'Université de Genève, Versoix, Switzerland (martin.turbet@unige.ch), ²Laboratoire d'astrophysique de Bordeaux, Université de Bordeaux, CNRS, B18N, Pessac, France, ³LATMOS/IPSL, UVSQ, Université Paris-Saclay, Sorbonne Université, CNRS, Guyancourt, France.

Abstract: In this contribution, we will present the results of new 3-D numerical simulations of the atmosphere of young rocky planets (supposedly rich in water vapor) carried out with the LMD-Generic Global Climate Model (GCM). We will first show how and why clouds tend to accumulate preferentially on the night side of these planets as soon as water vapor is a major component of the atmosphere. We will then show how and why this result has major consequences on the conditions of formation of early oceans on Venus, as well as on Earth and exoplanets.

Acknowledgments: This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement number 832738/ESCAPE. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement numbers 724427/FOUR ACES and 679030/WHIPLASH). This work has been carried out within the framework of the National Centre of Competence in Research PlanetS supported by the Swiss National Science Foundation. We acknowledge the financial support of the SNSF. M.T. thanks the Gruber Foundation for its support to this research.